

Condensation in Globally Coupled Populations of Chaotic Dynamical Systems

D.H. Zanette * and A. S. Mikhailov

Fritz-Haber-Institut der Max-Planck-Gesellschaft

Faradayweg 4-6, 14195 Berlin, Germany

(October 22, 2018)

Abstract

The condensation transition, leading to complete mutual synchronization in large populations of globally coupled chaotic Rössler oscillators, is investigated. Statistical properties of this transition and the cluster structure of partially condensed states are analyzed.

PACS: 05.45+b, 05.20.-y, 05.90.+m

Typeset using REVTeX

*Permanent address: Consejo Nacional de Investigaciones Científicas y Técnicas, Centro Atómico Bariloche, 8400 Bariloche, Argentina

Appearance of condensates is an important property of quantum and classical systems. When a population of interacting Bose particles (such as, e.g., superfluid liquid He⁴) is considered, a condensate is formed by a finite fraction of particles occupying the same quantum state [1]. Condensation of electron-hole pairs in metals is responsible for the phenomenon of superconductivity [2]. Condensation phenomena in classical systems are closely related to the effect of mutual synchronization of oscillations. In large populations of globally coupled limit-cycle oscillators, one or several groups of oscillators following exactly the same phase trajectory can spontaneously appear [3–7]. A similar behavior in populations of globally coupled logistic or circle maps has been found by K. Kaneko [8,9]. Mutual synchronization of several coupled chaotic oscillators has been studied (see, e.g., [10]). To analyze properties of the condensation transition, which involves progressive formation of coherent oscillator groups, larger systems must be however considered. In this Letter we investigate condensation phenomena in a population of a thousand globally coupled chaotic Rössler oscillators.

Not every form of global coupling between elements can lead to mutual synchronization. Synchronous chaotic oscillations are possible if ‘vector’ coupling (with identical coupling coefficients for all oscillator variables) is employed [11]. Asymptotic synchronization conditions in the limit of high coupling intensities have also been constructed for the case when coupling involves only one of the oscillator variables [10]. However, only the linear stability of synchronous oscillations is proven in both of these cases and therefore even for a strong coupling the presence of other dynamical attractors, corresponding to persistent asynchronous oscillations, cannot be excluded [12].

In this Letter a different form of global coupling that yields more robust synchronization is introduced. We take an arbitrary dynamical system with m variables $\mathbf{w}(t) = \{\mathbf{w}_1(t), \dots, \mathbf{w}_m(t)\}$ whose evolution is determined by equations

$$\dot{\mathbf{w}} = \mathbf{f}(\mathbf{w}), \tag{1}$$

where $\mathbf{f}(\mathbf{w})$ are some known functions. A globally coupled population consisting of N such

dynamical systems ($i = 1, 2, \dots, N$) is constructed by adding the coupling terms, so that the resulting evolution equations are

$$\dot{\mathbf{w}}_i = \mathbf{f}(\mathbf{w}_i) + \varepsilon \hat{\mathbf{A}}(\bar{\mathbf{w}} - \mathbf{w}_i) + \varepsilon' [\mathbf{f}(\bar{\mathbf{w}}) - \mathbf{f}(\mathbf{w}_i)], \quad (2)$$

where $\bar{\mathbf{w}}$ is the global average, $\bar{\mathbf{w}}(t) = N^{-1} \sum_{i=1}^N \mathbf{w}_i(t)$, the positive parameters ε and ε' specify the intensity of global coupling, and $\hat{\mathbf{A}}$ is a constant matrix. Note that the previously used forms of coupling correspond to a choice $\varepsilon' = 0$; the ‘vector’ coupling is obtained for $\hat{\mathbf{A}} = \mathbf{I}$ where \mathbf{I} is the identity matrix.

When $\varepsilon' = 1$ equations (2) reduce to

$$\dot{\mathbf{w}}_i = \varepsilon \hat{\mathbf{A}}(\bar{\mathbf{w}} - \mathbf{w}_i) + \mathbf{f}(\bar{\mathbf{w}}). \quad (3)$$

They describe linear motion of elements forming the population under action of a force which is collectively produced by all of them. The evolution of the deviations $\delta \mathbf{w}_i(t) = \mathbf{w}_i(t) - \bar{\mathbf{w}}(t)$ is then determined by an *exact* linear set of equations $\delta \dot{\mathbf{w}}_i = -\varepsilon \hat{\mathbf{A}} \delta \mathbf{w}_i$. Therefore, these deviations exponentially decrease with time, and global stability of synchronous oscillations is thus guaranteed, if all real eigenvalues of the matrix $\hat{\mathbf{A}}$ and the real parts of all its complex eigenvalues are positive.

Since globally stable synchronous oscillations are present in the system (2) at $\varepsilon' = 1$, they may also be expected in a neighborhood of this point, i.e. for an interval of coupling intensities $\varepsilon_0 < \varepsilon' < 1$. As shown below, this interval can be so wide that even weak coupling leads to the synchronization.

We consider a population of chaotic Rössler oscillators. Each elementary oscillator is described now by three variables $\mathbf{w}(t) = \{x(t), y(t), z(t)\}$, and we have $f_x = -y - z$, $f_y = x + ay$, and $f_z = b - cz + xz$, where a , b and c are fixed parameters. The collective dynamics of a globally coupled population of N identical Rössler oscillators ($i = 1, 2, \dots, N$) is governed by equations

$$\begin{aligned}
\dot{x}_i &= -y_i - z_i, \\
\dot{y}_i &= x_i + ay_i + K(\bar{y} - y_i), \\
\dot{z}_i &= b - cz_i + x_iz_i + K(\bar{x}\bar{z} - x_iz_i),
\end{aligned} \tag{4}$$

where $\bar{x}(t)$, $\bar{y}(t)$ and $\bar{z}(t)$ are global averages, such as $\bar{x}(t) = N^{-1} \sum_{i=1}^N x_i(t)$, and K is the intensity of global coupling. This form of coupling is derived from (2) by taking $\varepsilon = \varepsilon' = K$ and choosing a matrix $\hat{\mathbf{A}}$ whose elements are zero except for $A_{xy} = A_{xz} = -A_{yx} = 1$, $A_{zz} = c$, and $A_{yy} = 1 - a$. The system (4) reduces to a set of independent identical Rössler oscillators [13] for $K = 0$.

The eigenvalues of the matrix $\hat{\mathbf{A}}$ are $\lambda_1 = c$ and $\lambda_{2,3} = \frac{1}{2}(1 - a) \pm \frac{1}{2}\sqrt{(1 - a)^2 - 4}$. All three eigenvalues (or their real parts) are positive if $a < 1$. Therefore, for $K = 1$ all deviations from the global averages exponentially decrease with time and the states of all elements in the population asymptotically converge. We see that at $K = 1$ the oscillator population (4) possesses a global attractor which corresponds to coherent motion of all elements of the system. This attractor coincides with the attractor of a single Rössler oscillator and is chaotic for the parameters $a = 0.15$, $b = 0.4$ and $c = 8.5$ used in our simulations. Thus, the population (4) of globally coupled chaotic Rössler oscillators should undergo condensation as the coupling intensity K is gradually increased towards $K = 1$. To investigate the condensation phenomena, we have performed numerical simulations of a population of $N = 1000$ such oscillators under varying intensity K of global coupling. Below we first analyze statistical properties of the condensation transition and later consider how condensation is influenced by heterogeneities.

To characterize condensation, distances $d_{ij} = [(x_i - x_j)^2 + (y_i - y_j)^2 + (z_i - z_j)^2]^{1/2}$ between the states of all $N(N - 1)/2$ possible pairs of elements (i, j) have been computed. When condensation occurs, some of the elements have exactly the same coordinates and distances between the states of these elements are zero. The ratio of the number of pairs with zero distances to the total number of pairs can thus be chosen as the *order parameter* r of the condensation transition. In absence of a condensate, $r = 0$. On the other hand, $r = 1$ when

complete mutual synchronization of the whole population takes place.

To prepare the initial condition, the system was first allowed to evolve up to $t = 100$ without global coupling ($K = 0$) so that its elements get uniformly distributed over the Rössler attractor. Global coupling was then introduced, time was reset to zero, and the evolution of the coupled population was started. During a transient T , identical element pairs were appearing. The transient was over when the number of these pairs did not further increase with time. In our simulations, the transients were always shorter than $T = 1000$.

Figure 1 shows the dependence of the order parameter r on the global coupling intensity K obtained by averaging over 20 realizations at $t = 2000$. Condensation begins at $K_c \approx 0.017$ when a nonvanishing fraction of identical pairs first appears. In the interval from K_c to $K \approx 0.06$, nonmonotonous dependence of the order parameter on the coupling intensity is observed. At higher coupling intensities K , the fraction of identical pairs steadily grows until full condensation ($r = 1$) is established for $K > K_0$, with $K_0 \approx 0.093$. Under conditions of partial condensation, the number of identical pairs varies from one realization to another. Bars in Fig. 1 indicate statistical dispersion of the simulation data at a few selected values of the global coupling intensity. Note that, once a certain number of identical pairs has been formed at the end of the transient period, it remains later constant in any given simulation. Therefore, the dispersion reveals a degeneracy in the asymptotic properties of this system in the partially condensed state.

The order parameter r characterizes the overall degree of condensation, but is not sensitive to the condensate structure. To determine the detailed structure of the population, distributions over pair distances between the states of all elements at a fixed time moment ($t = 2000$) have been analyzed. Histograms of such distributions at four different coupling intensities K are shown in Fig. 2. They are constructed by counting pairs with distances lying within subsequent intervals of width $\Delta d = 0.25$. The number of pairs inside each next interval, divided by the total number of pairs $N(N - 1)/2$, yields the height of the respective bar. In absence of global coupling, elements are relatively uniformly distributed over the single-oscillator attractor and a smooth distribution over distances d is thus observed

(Fig. 2a). When global coupling is introduced but remains below the critical point K_c of the condensation transition, the distribution is modified (Fig. 2b). Now, pairs with small distances have already appeared, though identical pairs are still absent (cf. Fig. 1). Slightly above the critical point, a strongly nonuniform distribution with many narrow peaks is seen (Fig. 2c). The distributions found at larger intensities of global coupling (e.g., Fig. 2d) are formed by several distinct lines. When complete condensation has taken place ($r = 1$), the histogram has only one line located at $d = 0$ (not shown). Distributions formed by several lines are characteristic for situations when the whole population breaks down into a number of coherent clusters. All elements in a cluster follow the same dynamical trajectory and, hence, distances between pairs of elements belonging to the same cluster are zero. When elements from two different clusters are chosen, their distances are all equal. If M such clusters are present, the distribution over pair distances would consist of $M(M - 1)/2 + 1$ individual lines. Hence, four lines seen in Fig. 2d correspond to a condensate with three clusters. In our simulations, pure three-cluster regimes were typically found starting from $K \approx 0.045$. They were replaced by pure two-cluster regimes at $K \approx 0.058$.

Hence, the condensate appearing above the transition point includes only a small fraction of elements of the population, so that the order parameter is small in this region. The condensed elements are distributed over a large number of small clusters. The rest of the population has asynchronous dynamics, though it is already essentially influenced by global coupling between the elements. When the coupling intensity is increased, the number of clusters decreases. However, the typical sizes of clusters become at the same time larger and therefore the condensate contains a greater fraction of the population. Starting from a certain intensity of global coupling, all elements belong to one of a few big clusters. As the coupling intensity is further increased, relative sizes of clusters vary and some of them disappear. This process eventually leads to complete condensation which can be viewed as a single-cluster regime.

Fig. 3 shows the typical dependence of the order parameter r of the condensation transition on the total population size N , where each point is obtained by averaging over 20

independent realizations and intensity of global coupling is kept constant ($K = 0.08$). We see that a strong size dependence is characteristic for relatively small populations ($N < 500$). Starting from a population size of about $N = 1000$, the dependence displays saturation, with the remaining small variations lying within the statistical dispersion range. Hence, the results of our study of a population with 1000 oscillators might already be representative for the asymptotic behavior in the infinite-size limit $N \rightarrow \infty$.

The complete condensation ($r = 1$) has been found within a wide interval of the global coupling intensity. Starting in this region with random initial conditions for all elements, we see that the elements soon form a compact cloud that rapidly shrinks with time. The characteristic radius $R(t)$ of this cloud can be defined by

$$R^2(t) = \frac{1}{N} \sum_{i=1}^N [\delta x_i^2(t) + \delta y_i^2(t) + \delta z_i^2(t)], \quad (5)$$

where δx_i , δy_i and δz_i are the deviations from the respective global averages. The radius R decreases with time, as the trajectories of all elements asymptotically converge to the same orbit (in our simulations the convergence was followed until the trajectories of all elements became identical up to the computer precision of about 10^{-6}). An important quantitative property of this regime is the characteristic time which the system needs to reach the condensed state; the inverse of this time represents the condensation exponent (closely related to the transverse Lyapunov exponents discussed in Ref. [10]):

$$\gamma = - \lim_{t \rightarrow \infty} \frac{1}{t} \ln \frac{R(t)}{R(0)} \quad (6)$$

The simulations show that γ starts to grow from zero at $K = K_0$, has a maximum at $K \approx 0.65$ and then slowly decreases approaching the value $\gamma \approx 0.38$ at $K = 1$.

The robustness of the complete condensation suggests that it might persist in some form even in heterogeneous populations. To test this, we have carried out simulations where the elements forming the population were not identical. Heterogeneity was introduced by replacing the constant parameter c in Eqs. (4) by random numbers c_i that were uniformly distributed inside the segment $[c - \sigma_c, c + \sigma_c]$. Alternatively, the same procedure was applied

to the parameter a . Starting from random initial conditions, a compact cloud of elements was again formed in the heterogeneous case. However, the cloud did not shrink until it transformed into a single point. Instead, after an initial decrease, its radius $R(t)$ fluctuated around a certain mean value. Fig. 4 shows the mean radius $\langle R \rangle$ as function of dispersions σ_c or σ_a for $K = 0.2$. We see that in a wide interval of heterogeneities the mean radius of the cloud is approximately linear proportional to the dispersion. Moreover, even for relatively large dispersions this radius is significantly smaller than the characteristic attractor diameter D , representing the maximal possible pair distance at $K = 0$ (i.e. $D \approx 25$, cf. Fig. 2a). Therefore, even though the states of all elements are no longer identical (i.e. the complete condensation is destroyed), their variations remain small. The elements form a little cloud whose center moves along a definite trajectory.

This behavior can be understood by analyzing the effect of weak heterogeneities at the maximal strength of global coupling $K = 1$. If the parameter c is replaced by $c_i = c + \Delta c_i$ with random variations Δc_i , the population dynamics is described by

$$\begin{aligned}\dot{x}_i &= -y_i - z_i, \\ \dot{y}_i &= x_i + (a - 1)y_i + \bar{y}, \\ \dot{z}_i &= b - c_i z_i + \bar{x}\bar{z}.\end{aligned}\tag{7}$$

Hence, motion of each element is governed by linear equations of motion in the presence of global driving forces \bar{y} and $\bar{x}\bar{z}$, which are the same for all elements. Therefore, deviations δz_i from the global average obey the equation

$$\dot{\delta z}_i = -c\delta z_i - \Delta c_i \bar{z} - \Delta c_i \delta z_i + \overline{\Delta c_i \delta z_i}.\tag{8}$$

If heterogeneities are weak, i.e. $\Delta c_i = \theta \zeta_i$ where θ is a small parameter, the last two terms in this equation have order θ^2 and can be neglected. Integrating the resulting linear equation, we obtain

$$\delta z_i(t) = \theta \zeta_i \int_0^\infty \exp(-c\tau) \bar{z}(t - \tau) d\tau.\tag{9}$$

Considering the behavior of other deviations δx_i and δy_i , one can similarly show that they are also proportional to $\Delta c_i = \theta \zeta_i$. This means that the radius $\langle R \rangle$ of the population cloud

depends as $\langle R \rangle \sim \theta$ on the heterogeneity strength. Note also that in this case all deviations are scaling-identical, i.e., for example, $\delta z_i(t) = \theta \zeta_i Q(t)$ where the function $Q(t)$ is the same for any element i . The same results are obtained when weak random variations of the parameter a are considered.

Our simulations have revealed that the linear dependence of the mean radius $\langle R \rangle$ on the dispersions σ_c and σ_a , that has been analytically shown above only at $K = 1$, persists in a wide range of coupling intensities. Moreover, this dependence is found even for relatively large dispersions, where the parameter θ would not be small. This behavior is apparently a consequence of a strong compression provided in the entire interval of complete condensation by global coupling.

The condensation transition has been studied above for a particular kind of chaotic oscillators and for a certain form of global coupling. We have however checked that similar results are also obtained when a different ('vector') form of global coupling is used. This suggests that observed statistical properties may be typical for various large populations of globally coupled chaotic oscillators undergoing a condensation transition.

Financial support from Fundación Antorchas, Argentina, and from Alexander von Humboldt-Stiftung, Germany, is gratefully acknowledged. D.H.Z. wishes to thank the Fritz Haber Institute for hospitality during his stay in Berlin.

REFERENCES

- [1] N.N. Bogoliubov, *Lectures on Quantum Statistics*, vol. I (Gordon and Breach, New York, 1967).
- [2] H. Umezawa, H. Matsumoto and M. Tachiki, *Thermo-Field Dynamics and Condensed States* (North-Holland, Amsterdam, 1982).
- [3] Y. Kuramoto, *Chemical Oscillations, Waves and Turbulence* (Springer, Berlin, 1984).
- [4] N. Nakagawa and Y. Kuramoto, *Physica D* **75**, 74 (1994).
- [5] H. Sakaguchi and Y. Kuramoto, *Prog. Theor. Phys* **76**, 576 (1986).
- [6] V. Hakim and W.-J. Rappel, *Phys. Rev. A* **46**, R7347 (1992).
- [7] D. Golomb, D. Hansel, B. Shraiman and H. Sompolinsky, *Phys. Rev. A* **45**, 3516 (1992).
- [8] K. Kaneko, *Physica D* **23**, 436 (1986); *ibid.* **37**, 60 (1989).
- [9] K. Kaneko, *Physica D* **54**, 5 (1991).
- [10] J. F. Heagy, T. L. Carrol, and L. M. Pecora, *Phys. Rev. E* **50**, 1874 (1994).
- [11] H. Fujisaka and T. Yamada, *Progr. Theor. Phys.* **69**, 32 (1983).
- [12] In the complex Ginzburg-Landau equation describing a population of locally coupled limit-cycle oscillators, both linearly stable synchronous uniform oscillations and complex spatiotemporal patterns, such as spiral waves or turbulence, can often be found depending on the initial conditions at the same system parameters; see: B. I. Shraiman, A. Pumir, W. von Saarloos, P. C. Hohenberg, H. Chaté and M. Holen, *Physica D* **57**, 241 (1995).
- [13] O. E. Röessler, *Phys. Lett. A* **57**, 97 (1976).

FIGURES

FIG. 1. Order parameter r of the condensation transition as function of the coupling constant K . Averaging over 20 independent realizations is performed; bars indicate statistical dispersion of data at several selected points.

FIG. 2. Normalized histograms of the distribution over pair distances d between elements in a population of $N = 1000$ globally coupled Rössler oscillators at different stages of the condensation transition: (a) $K = 0$; (b) $K = 0.01$; (c) $K = 0.02$; (d) $K = 0.05$.

FIG. 3. Dependence of the order parameter r on the population size N at a fixed intensity of global coupling $K = 0.08$. Averaging over 20 independent realizations; bars indicate statistical dispersion of data.

FIG. 4. Radius $\langle R \rangle$ of the cloud formed by a heterogeneous population as function of dispersions σ_a (empty circles) and σ_c (filled circles) at $K = 0.2$.




Fig. 1, Zanette & Mikhailov (PRL)




Fig. 2, Zanette & Mikhailov (PRL)




Fig. 3, Zanette & Mikhailov (PRL)




Fig. 4, Zanette & Mikhailov (PRL)