arXiv:alg-geom/9311007v1 19 Nov 1993

ON THE PICARD NUMBER OF FANO 3-FOLDS WITH TERMINAL SINGULARITIES

VIACHESLAV V. NIKULIN

ABSTRACT. We prove: Let X be a Fano 3-fold with terminal Q-factorial singularities and X does not have a small extremal ray and a face of Kodaira dimension 1 or 2 for Mori polyhedron $\overline{NE}(X)$. Then Picard number $\rho(X) \leq 7$.

INTRODUCTION.

Here we continue investigations started in [N6], [N7].

Algebraic varieties we consider are defined over the field \mathbb{C} of complex numbers. In this paper, we get a final result on estimating of the Picard number $\rho = \dim N_1(X)$ of a Fano 3-fold X with terminal Q-factorial singularities if X does not have small extremal rays and its Mori polyhedron does not have faces with Kodaira dimension 1 or 2. One can consider this class as a generalization of the class of Fano 3-folds with Picard number 1. There are many non-singular Fano 3-folds satisfying this condition and with Picard number 2 (see [Mo-Mu] and also [Ma]). We also think that studying of the Picard number of this class may be important for studying of Fano 3-folds with Picard number 1 too (see Corollary 2 below).

Let X be a Fano 3-fold with \mathbb{Q} -factorial terminal singularities. Let R be an extremal ray of the Mori polyhedron $\overline{NE}(X)$ of X. We say that R has the type (I) (respectively (II)) if curves of R fill an irreducible divisor D(R) of X and the contraction of the ray R contracts the divisor D(R) in a point (respectively on a curve). An extremal ray R is called *small* if curves of this ray fill a curve on X.

A pair $\{R_1, R_2\}$ of extremal rays has the type \mathfrak{B}_2 if extremal rays R_1, R_2 are different, both have the type (II), and have the same divisor $D(R_1) = D(R_2)$.

We recall that a face γ of Mori polyhedron $\overline{NE}(X)$ defines a contraction f_{γ} : $X \to X'$ (see [Ka1] and [Sh]) such that f(C) is a point for an irreducible curve C iff C belongs to γ . The dim X' is called the Kodaira dimension of the γ . A set \mathcal{E} of extremal rays is called extremal if it is contained in a face of Mori polyhedron.

Basic Theorem. Let X be a Fano 3-fold with terminal \mathbb{Q} -factorial singularities. Assume that X does not have a small extremal ray, and Mori polyhedron $\overline{NE}(X)$ does not have a face of Kodaira dimension 1 or 2.

Then we have the following statements for the X:

(1) The X does not have a pair of extremal rays of the type \mathfrak{B}_2 and Mori polyhedron $\overline{NE}(X)$ is simplicial;

(2) The X does not have more than one extremal ray of the type (I).

(3) If \mathcal{E} is an extremal set of k extremal rays of X, then the \mathcal{E} has one of the types: $\mathfrak{A}_1 \amalg (k-1)\mathfrak{C}_1$, $\mathfrak{D}_2 \amalg (k-2)\mathfrak{C}_1$, $\mathfrak{C}_2 \amalg (k-2)\mathfrak{C}_1$, $k\mathfrak{C}_1$ (we use notation of Theorem 2.3.3).

(4) We have the inequality for the Picard number of the X:

$$\rho(X) = \dim N_1(X) \le 7.$$

Proof. See Theorem 2.5.8.

It follows from (4):

Corollary 1. Let X be a Fano 3-fold with terminal \mathbb{Q} -factorial singularities and $\rho(X) > 7$.

Then X has either a small extremal ray or a face of Kodaira dimension 1 or 2 for Mori polyhedron.

We mention that non-singular Fano 3-folds do not have a small extremal ray (by Mori [Mo1]), and the maximal Picard number for them is equal to 10 according to their classification by Mori and Mukai [Mo-Mu]. Thus, all these statements already work for non-singular Fano 3-folds.

From the statement (2) of the Theorem, we also get the following application of Basic Theorem to geometry of Fano 3-folds.

Let us consider a Fano 3-fold X and blow-ups X_p in different non-singular points $\{x_1, ..., x_p\}$ of X. We say that this is a Fano blow-up if X_p is Fano. We have the following very simple

Proposition. Let X be a Fano 3-fold with terminal \mathbb{Q} -factorial singularities and without small extremal rays. Let X_p be a Fano blow up of X.

Then for any small extremal ray S on X_p , the S has a non-empty intersection with one of exceptional divisors $E_1, ..., E_p$ of this blow up and does not belong to any of them. The divisors $E_1, ..., E_p$ define p extremal rays of the type (I) on X_p .

Proof. See Proposition 2.5.14.

It is known that a contraction of a face of Kodaira dimension 1 or 2 of NE(X) of a Fano 3-fold X has a general fiber which is a rational surface or curve respectively, because this contraction has relatively negative canonical class. See [Ka1], [Sh]. It is known that a small extremal ray is rational [Mo2].

Then, using Basic Theorem and Proposition, we can divide Fano 3-folds of Basic Theorem on the following 3 classes:

Corollary 2. Let X be a Fano 3-fold with terminal Q-factorial singularities and without small extremal rays, and without faces of Kodaira dimension 1 or 2 for Mori polyhedron. Let ϵ be the number of extremal rays of the type (I) on X (by Basic Theorem, the $\epsilon \leq 1$).

Then there exists $p, 1 \le p \le 2 - \epsilon$, such that X belongs to one of classes (A), (B) or (C) below:

(A) There exists a Fano blow-up X_p of X with a face of Kodaira dimension 1 or 2. Thus, birationally, X is a fibration on rational surfaces over a curve or rational curves over a surface.

(B) There exist Fano blow-ups X_p of X for general p points on X such that for all these blow-ups the X_p has a small extremal ray S. Then images of curves of S on X give a system of rational curves on X which cover a Zariski open subset of X.

(C) There do not exist Equal blow one Y of Y for concerd remaints

ON THE PICARD NUMBER OF FANO 3-FOLDS WITH TERMINAL SINGULARITIES 3

We remark that for Fano 3-folds with Picard number 1 the $\epsilon = 0$. Thus, $1 \le p \le 2$.

Using statements (2), (3) and (4) of Basic Theorem, one can formulate similar results for Fano blow ups in curves.

To prove Basic Theorem, we classify appropriate so called extremal sets and E-sets of extremal rays of the type (I) or (II). We use so called diagram method to deduce from this classification the statement (4) of the Basic Theorem.

A set \mathcal{E} of extremal rays is called *extremal* if it is contained in a face of Mori polyhedron. The \mathcal{E} has *Kodaira dimension* 3 if a contraction of this face gives a morphism on a 3-fold. For Fano 3-folds with Q-factorial terminal singularities, we give a description of extremal sets \mathcal{E} of Kodaira dimension 3 which contain extremal rays of the types (I) or (II) only.

A set \mathcal{L} of extremal rays is called *E-set* if \mathcal{L} is not extremal, but any proper subset of \mathcal{L} is extremal. Thus, the \mathcal{L} is minimal non-extremal. For Fano 3-folds with Q-factorial terminal singularities, we give a description of *E*-sets \mathcal{L} such that \mathcal{L} contains extremal rays of the types (I) or (II) only, and any proper subset of \mathcal{L} is extremal of Kodaira dimension 3.

I am grateful to Profs. Sh. Ishii, M. Reid and J. Wiśniewski for useful discussions. I am grateful to referee for useful comments.

I am grateful to Professor Igor R. Shafarevich for his constant interest to and support of these my studies.

This paper was prepared in Steklov Mathematical Institute, Moscow; Max-Planck Institut für Matematik, Bonn, 1990; Kyoto University, 1992–1993; Mathematical Sciences Research Institute, Berkeley, 1993. I thank these Institutes for hospitality.

Preliminary variant of this paper was published as a preprint [N8] (see also [N9] about connected results).

VIACHESLAV V. NIKULIN

CHAPTER 1. DIAGRAM METHOD.

Here we give a simplest variant of the diagram method for multi-dimensional algebraic varieties. We shall use this method in the next chapter. This part also contains some corrections and generalizations to the corresponding part of our papers [N6] and [N7].

Let X be a projective algebraic variety with Q-factorial singularities over an algebraically closed field. Let dim $X \ge 2$. Let $N_1(X)$ be the R-linear space generated by all algebraic curves on X by the numerical equivalence, and let $N^1(X)$ be the R-linear space generated by all Cartier (or Weil) divisors on X by the numerical equivalence. Linear spaces $N_1(X)$ and $N^1(X)$ are dual to one another by the intersection pairing. Let NE(X) be a convex cone in $N_1(X)$ generated by all effective curves on X. Let $\overline{NE}(X)$ be the closure of the cone NE(X) in $N_1(X)$. It is called *Mori cone* (or polyhedron) of X. A non-zero element $x \in N^1(X)$ is called nef if $x \cdot \overline{NE}(X) \ge 0$. Let NEF(X) be the set of all nef elements of X and the zero. It is the convex cone in $N^1(X)$ dual to Mori cone $\overline{NE}(X)$. A ray $R \subset \overline{NE}(X)$ with origin 0 is called *extremal* if from $C_1 \in \overline{NE}(X)$, $C_2 \in \overline{NE}(X)$ and $C_1 + C_2 \in R$ it follows that $C_1 \in R$ and $C_2 \in R$.

We consider a condition (i) for a set \mathcal{R} of extremal rays on X.

(i) If $R \in \mathcal{R}$, then all curves $C \in R$ fill out an irreducible divisor D(R) on X.

In this case, we can correspond to \mathcal{R} (and subsets of \mathcal{R}) an oriented graph $G(\mathcal{R})$ in the following way: Two different rays R_1 and R_2 are joined by an arrow R_1R_2 with the beginning in R_1 and the end in R_2 if $R_1 \cdot D(R_2) > 0$. Here and in what follows, for an extremal ray R and a divisor D we write $R \cdot D > 0$ if $r \cdot D > 0$ for $r \in R$ and $r \neq 0$. (The same for the symbols $\leq \geq$ and <.)

A set \mathcal{E} of extremal rays is called *extremal* if it is contained in a face of $\overline{NE}(X)$. Equivalenty, there exists a nef element $H \in N^1(X)$ such that $\mathcal{E} \cdot H = 0$. Evidently, a subset of an extremal set is extremal too.

We consider the following condition (ii) for extremal sets \mathcal{E} of extremal rays.

(ii) An extremal set $\mathcal{E} = \{R_1, ..., R_n\}$ satisfies the condition (i), and for any real numbers $m_1 \ge 0, ..., m_n \ge 0$ which are not all equal to 0, there exists a ray $R_j \in \mathcal{E}$ such that $R_j \cdot (m_1 D(R_1) + m_2 D(R_2) + ... + m_n D(R_n)) < 0$. In particular, the effective divisor $m_1 D(R_1) + m_2 D(R_2) + ... + m_n D(R_n)$ is not nef.

A set \mathcal{L} of extremal rays is called *E-set* (extremal in a different sense) if the \mathcal{L} is not extremal but every proper subset of \mathcal{L} is extremal. Thus, \mathcal{L} is a minimal non-extremal set of extremal rays. Evidently, an *E*-set \mathcal{L} contains at least two elements.

We consider the following condition (iii) for *E*-sets \mathcal{L} .

(iii) Any proper subset of an E-set $\mathcal{L} = \{Q_1, ..., Q_m\}$ satisfies the condition (ii), and there exists a non-zero effective nef divisor $D(\mathcal{L}) = a_1 D(Q_1) + a_2 D(Q_2) + ... + a_m D(Q_m)$.

The following statement is very important.

Lemma 1.1. An *E*-set \mathcal{L} satisfying the condition (iii) is connected in the following sense: For any decomposition $\mathcal{L} = \mathcal{L}_1 \coprod \mathcal{L}_2$, where \mathcal{L}_1 and \mathcal{L}_2 are non-empty, there exists an arrow Q_1Q_2 such that $Q_1 \in \mathcal{L}_1$ and $Q_2 \in \mathcal{L}_2$.

If \mathcal{L} and \mathcal{M} are two different E-sets satisfying the condition (iii), then there exists an arrow LM where $L \in \mathcal{L}$ and $M \in \mathcal{M}$.

Dreach Let $\mathcal{L} = \{ O = O \}$ Dre (:::) there exists a perfective on $D(\mathcal{L}) = \sigma D(O) \}$

 $a_2D(Q_2) + \ldots + a_mD(Q_m)$. If one of the coefficients a_1, \ldots, a_m is equal to zero, we get a contradiction with the conditions (ii) and (iii). It follows that all the coefficients a_1, \ldots, a_m are positive. Let $\mathcal{L} = \mathcal{L}_1 \coprod \mathcal{L}_2$ where $\mathcal{L}_1 = \{Q_1, \ldots, Q_k\}$ and $\mathcal{L}_2 = \{Q_{k+1}, \ldots, Q_m\}$. The divisors $D_1 = a_1D(Q_1) + \ldots + a_kD(Q_k)$ and $D_2 =$ $a_{k+1}D(Q_{k+1}) + \ldots + a_mD(Q_m)$ are non-zero. By (ii), there exists a ray $Q_i, 1 \leq i \leq k$, such that $Q_i \cdot D_1 < 0$. On the other hand, $Q_i \cdot D(\mathcal{L}) = Q_1 \cdot (D_1 + D_2) \geq 0$. It follows, that there exists $j, k+1 \leq j \leq m$, such that $Q_i \cdot D(Q_j) > 0$. It means that Q_iQ_j is an arrow.

Let us prove the second statement. By the condition (iii), for every ray $R \in \mathcal{L}$, we have the inequality $R \cdot D(\mathcal{M}) \geq 0$. If $R \cdot D(\mathcal{M}) = 0$ for any $R \in \mathcal{L}$, then the set \mathcal{L} is extremal, and we get the contradiction. It follows that there exists a ray $R \in \mathcal{L}$ such that $R \cdot D(\mathcal{M}) > 0$. It follows the statement.

Let $NEF(X) = \overline{NE}(X)^* \subset N^1(X)$ be the cone of nef elements of X and $\mathcal{M}(X) = NEF(X)/\mathbb{R}^+$ its projectivization. We use usual relations of orthogonality between subsets of $\mathcal{M}(X)$ and $\overline{NE}(X)$. So, for $U \subset \mathcal{M}(X)$ and $V \subset \overline{NE}(X)$ we write $U \perp V$ if $x \cdot y = 0$ for any $\mathbb{R}^+ x \in U$ and any $y \in V$. Thus, for $U \subset \mathcal{M}(X)$, $V \subset \overline{NE}(X)$ we denote

$$U^{\perp} = \{ y \in \overline{NE}(X) \mid U \perp y \}, \quad V^{\perp} = \{ x \in \mathcal{M}(X) \mid x \perp V \}.$$

A subset $\gamma \subset \mathcal{M}(X)$ is called a *face* of $\mathcal{M}(X)$ if there exists a non-zero element $r \in \overline{NE}(X)$ such that $\gamma = r^{\perp}$.

A convex set is called a *closed polyhedron* if it is a convex hull of a finite set of points. A convex closed polyhedron is called *simplicial* if all its faces are simplexes. A convex closed polyhedron is called *simple* (equivalently, it has simplicial angles) if it is dual to a similicial one. In other words, any its face of codimension k is contained exactly in k faces of γ of the highest dimension.

We need some relative notions to notions above.

We say that $\mathcal{M}(X)$ is a closed polyhedron in its face $\gamma \subset \mathcal{M}(X)$ if γ is a closed polyhedron and $\mathcal{M}(X)$ is a closed polyhedron in a neighbourhood T of γ . Thus, there should exist a closed polyhedron \mathcal{M}' such that $\mathcal{M}' \cap T = \mathcal{M}(X) \cap T$.

We will use the following notation. Let $\mathcal{R}(X)$ be the set of all extremal rays of X. For a face $\gamma \subset \mathcal{M}(X)$,

$$\mathcal{R}(\gamma) = \{ R \in \mathcal{R}(X) \mid \exists \mathbb{R}^+ H \in \gamma : R \cdot H = 0 \}$$

and

$$\mathcal{R}(\gamma^{\perp}) = \{ R \in \mathcal{R}(X) \mid \gamma \perp R \}.$$

Let us assume that $\mathcal{M}(X)$ is a closed polyhedron in its face γ . Then sets $\mathcal{R}(\gamma_1)$ and $\mathcal{R}(\gamma_1^{\perp})$ are finite for any face $\gamma_1 \subset \gamma$. Evidently, the face γ is simple if

(1)
$$\#\mathcal{R}(\gamma_1^{\perp}) - \#\mathcal{R}(\gamma^{\perp}) = \operatorname{codim}_{\gamma}\gamma_1$$

for any face γ_1 of γ . Then we say that the polyhedron $\mathcal{M}(X)$ is simple in its face γ . Evidently, this condition is equivalent to the condition:

(2)
$$\operatorname{dim} [\mathcal{D}] = \operatorname{dim} [\mathcal{D}(\alpha, \pm)] \quad \#\mathcal{D}(\alpha, \pm)$$

for any extremal set \mathcal{E} such that $R(\gamma^{\perp}) \subset \mathcal{E}$. Here $[\cdot]$ denotes a linear hull. (In [N6], we required a more strong condition for a polyhedron $\mathcal{M}(X)$ to be simple in its face $\gamma: \ \sharp \mathcal{R}(\gamma_1^{\perp}) = \dim \mathcal{M}(X) - \dim \gamma_1$ for any face γ_1 of γ .)

Let A, B are two vertices of an oriented graph G. The distance $\rho(A, B)$ in G is a length (the number of links) of a shortest oriented path of the graph G with the beginning in A and the end in B. The distance is $+\infty$ if this path does not exist. The diameter diam G of an oriented graph G is the maximum distance between ordered pairs of its vertices. By the Lemma 1.1, the diameter of an E-set is a finite number if this set satisfies the condition (iii).

The Theorem 1.2 below is an analog for algebraic varieties of arbitrary dimension of the Lemma 3.4 from [N2] and the Lemma 1.4.1 from [N5], which were devoted to surfaces.

Theorem 1.2. Let X be a projective algebraic variety with \mathbb{Q} -factorial singularities and dim $X \geq 2$. Let us suppose that $\mathcal{M}(X)$ is closed and simple in its face γ . Assume that the set $\mathcal{R}(\gamma)$ satisfies the condition (i) above. Assume that there are some constants d, C_1, C_2 such that the conditions (a) and (b) below hold:

(a) For any E-set $\mathcal{L} \subset \mathcal{R}(\gamma)$ such that \mathcal{L} contains at least two elements which don't belong to $\mathcal{R}(\gamma^{\perp})$ and for any proper subset $\mathcal{L}' \subset \mathcal{L}$ the set $R(\gamma^{\perp}) \cup \mathcal{L}'$ is extremal, the condition (iii) is valid and

diam
$$G(\mathcal{L}) \leq d$$
.

(b) For any extremal subset \mathcal{E} such that $\mathcal{R}(\gamma^{\perp}) \subset \mathcal{E} \subset \mathcal{R}(\gamma)$, we have: the \mathcal{E} satisfies the condition (ii) and for the distance in the oriented graph $G(\mathcal{E})$

$$\sharp\{(R_1, R_2) \in (\mathcal{E} - \mathcal{R}(\gamma^{\perp})) \times (\mathcal{E} - \mathcal{R}(\gamma^{\perp})) \mid 1 \le \rho(R_1, R_2) \le d\} \le C_1 \sharp(\mathcal{E} - \mathcal{R}(\gamma^{\perp}));$$

and

$$\sharp\{(R_1, R_2) \in (\mathcal{E} - \mathcal{R}(\gamma^{\perp})) \times (\mathcal{E} - \mathcal{R}(\gamma^{\perp})) \mid d+1 \le \rho(R_1, R_2) \le 2d+1\} \\
\le C_2 \sharp (\mathcal{E} - \mathcal{R}(\gamma^{\perp})).$$

Then dim $\gamma < (16/3)C_1 + 4C_2 + 6$.

Proof. We use the following Lemma 1.3 which was proved in [N1]. The Lemma was used in [N1] to get a bound (≤ 9) on the dimension of a hyperbolic (Lobachevsky) space admitting an action of an arithmetic reflection group with a field of definition of the degree > N. Here N is some absolute constant.

Lemma 1.3. Let \mathcal{M} be a convex closed simple polyhedron of a dimension n, and $A_n^{i,k}$ the average number of i-dimensional faces of k-dimensional faces of \mathcal{M} .

Then for n > 2k - 1

$$A_n^{i,k} < \frac{\binom{n-i}{n-k} \cdot \left(\binom{[n/2]}{i} + \binom{n-[n/2]}{i}\right)}{\binom{[n/2]}{k} + \binom{n-[n/2]}{k}}$$

In particular, if n > 3

$$A_n^{0,2} < \begin{cases} \frac{4(n-1)}{n-2} & \text{if } n \text{ is even,} \\ 4n & \text{if } n \text{ is outd} \end{cases}$$

ON THE PICARD NUMBER OF FANO 3-FOLDS WITH TERMINAL SINGULARITIES 7

Proof. See [N1]. We mention that the right side of the inequality of the Lemma 1.3 decreases and tends to the number $2^{k-i} \binom{k}{i}$ of *i*-dimensional faces of *k*-dimensional cube if *n* increases.

From the estimate of $A_n^{0,2}$ of the Lemma, it follows the following analog of Vinberg's Lemma from [V]. Vinberg's Lemma was used by him to obtain an estimate (dim < 30) for the dimension of a hyperbolic space admitting an action of a discrete reflection group with a bounded fundamental polyhedron.

By definition, an *angle* of a polyhedron T is an angle of a 2-dimensional face of T. Thus, the angle is defined by a vertex A of T, a plane containing A and a 2-dimensional face γ_2 of T, and two rays with the beginning at A which contain two corresponding sides of the γ_2 . To define an *oriented angle* of T, one should in addition put in order two rays of the angle.

Lemma 1.4. Let \mathcal{M} be a convex simple polyhedron of a dimension n. Let C and D are some numbers. Suppose that oriented angles (2-dimensional, plane) of \mathcal{M} are supplied with weights and the following conditions (1) and (2) hold:

(1) The sum of weights of all oriented angles at any vertex of M is not greater than Cn + D.

(2) The sum of weights of all oriented angles of any 2-dimensional face of \mathcal{M} is at least 5 - k where k is the number of vertices of the 2-dimensional face.

Then

$$n < 8C + 5 + \begin{cases} 1 + 8D/n & \text{if } n \text{ is even} \\ (8C + 8D)/(n-1) & \text{if } n \text{ is odd} \end{cases}$$

In particular, for $C \ge 0$ and D = 0, we have

$$n < 8C + 6.$$

Proof. We correspond to a non-oriented plane angle of \mathcal{M} a weight which is equal to the sum of weights of two corresponding oriented angles. Evidently, the conditions of the Lemma hold for the weights of non-oriented angles too if we forget about the word "oriented". Then we obtain Vinberg's lemma from [V] which we formulate a little bit more precisely here. Since the proof is simple, we give the proof here.

Let Σ be the sum of weights of all (non-oriented) angles of the polyhedron \mathcal{M} . Let α_0 be the number of vertices of \mathcal{M} and α_2 the number of 2-dimensional faces of \mathcal{M} . Since \mathcal{M} is simple,

$$\alpha_0 \frac{n(n-1)}{2} = \alpha_2 A_n^{0,2}.$$

From this equality and conditions of the Lemma, we get inequalities

$$(Cn+D)\alpha_0 \ge \Sigma \ge \sum \alpha_{2,k}(5-k) = 5\alpha_2 - \alpha_2 A_n^{0,2} =$$
$$= \alpha_2(5-A_n^{0,2}) = \alpha_0(n(n-1)/2)(5/A_n^{0,2}-1).$$

Here $\alpha_{2,k}$ is the number of 2-dimensional faces with k vertices of \mathcal{M} . Thus, from this inequality and Lemma 1.3, we get

$$Cn + D \ge (n(n-1)/2)(5/A_n^{0,2} - 1) > \begin{cases} n(n-6)/8 & \text{if } n \text{ is even,} \\ (n(n-1)/2)(5/A_n^{0,2} - 1) > (n(n-1)/2)(2n(n-1)/2) \end{cases}$$

From this calculations, Lemma 1.4 follows.

The proof of Theorem 1.2. (Compare with [V].) Let \angle be an oriented angle of γ . Let $\mathcal{R}(\angle) \subset \mathcal{R}(\gamma)$ be the set of all extremal rays of $\mathcal{M}(X)$ which are orthogonal to the vertex of \angle . Since $\mathcal{M}(X)$ is simple in γ , the set $\mathcal{R}(\angle)$ is a disjoint union

$$\mathcal{R}(\angle) = \mathcal{R}(\angle^{\perp}) \cup \{R_1(\angle)\} \cup \{R_2(\angle)\}$$

where $\mathcal{R}(\angle^{\perp})$ contains all rays orthogonal to the plane of the angle \angle , the rays $R_1(\angle)$ and $R_2(\angle)$ are orthogonal to the first and second side of the oriented angle \angle respectively. Evidently, the set $\mathcal{R}(\angle)$ and the ordered pair of rays $(R_1(\angle), R_2(\angle))$ define the oriented angle \angle uniquely. We define the weight $\sigma(\angle)$ by the formula:

$$\sigma(\angle) = \begin{cases} 2/3, & \text{if } 1 \le \rho(R_1(\angle), R_2(\angle)) \le d, \\ 1/2, & \text{if } d+1 \le \rho(R_1(\angle), R_2(\angle)) \le 2d+1, \\ 0, & \text{if } 2d+2 \le \rho(R_1(\angle), R_2(\angle)). \end{cases}$$

Here we take the distance in the graph $G(\mathcal{R}(\angle))$. Let us prove conditions of the Lemma 1.4 with the constants $C = (2/3)C_1 + C_2/2$ and D = 0.

The condition (1) follows from the condition (b) of the Theorem. We remark that rays $R_1(\angle), R_2(\angle)$ do not belong to the set $R(\gamma^{\perp})$.

Let us prove the condition (2).

Let γ_3 be a 2-dimensional triangle face (triangle) of γ . The set $\mathcal{R}(\gamma_3)$ of all extremal rays orthogonal to points of γ_3 is the union of the set $\mathcal{R}(\gamma_3^{\perp})$ of external rays, which are orthogonal to the plane of the triangle γ_3 , and rays R_1, R_2, R_3 , which are orthogonal to sides of the triangle γ_3 . Union of the set $\mathcal{R}(\gamma_3^{\perp})$ with any two rays from R_1, R_2, R_3 is extremal, since it is orthogonal to a vertex of γ_3 . On the other hand, the set $\mathcal{R}(\gamma_3) = \mathcal{R}(\gamma_3^{\perp}) \cup \{R_1, R_2, R_3\}$ is not extremal, since it is not orthogonal to a point of $\mathcal{M}(X)$. Indeed, the set of all points of $\mathcal{M}(X)$, which are orthogonal to the set $\mathcal{R}(\gamma_3^{\perp}) \cup \{R_2, R_3\}, \mathcal{R}(\gamma_3^{\perp}) \cup \{R_1, R_3\}, \text{ or } \mathcal{R}(\gamma_3^{\perp}) \cup \{R_1, R_2\}$ is the vertex A_1, A_2 or A_3 respectively of the triangle γ_3 , and the intersection of these sets of vertices is empty. Thus, there exists an E-set $\mathcal{L} \subset \mathcal{R}(\gamma_3)$, which contains the set of rays $\{R_1, R_2, R_3\}$. By the condition (a), the graph $G(\mathcal{L})$ contains a shortest oriented path s of the length $\leq d$ which connects the rays R_1, R_3 . If this path does not contain the ray R_2 , then the oriented angle of γ_3 defined by the set $\mathcal{R}(\gamma_3^{\perp}) \cup \{R_1, R_3\}$ and the pair (R_1, R_3) has the weight 2/3. If this path contains the ray R_2 , then the oriented angle of γ_3 defined by the set $\mathcal{R}(\gamma_3^{\perp}) \cup \{R_1, R_2\}$ and the pair (R_1, R_2) has the weight 2/3. Thus, we proved that the side A_2A_3 of the triangle γ_3 defines an oriented angle of the triangle with the weight 2/3 and the first side A_2A_3 of the oriented angle. The triangle has three sides. It follows the condition (2) of the Lemma 1.4 for the triangle.

Let γ_4 be a 2-dimensional quadrangle face (quadrangle) of γ . In this case,

$$\mathcal{R}(\gamma_4) = \mathcal{R}(\gamma_4^{\perp}) \cup \{R_1, R_2, R_3, R_4\}$$

where $\mathcal{R}(\gamma_4^{\perp})$ is the set of all extremal rays which are orthogonal to the plane of the quadrangle and the rays R_1, R_2, R_3, R_4 are orthogonal to the consecutive sides of the quadrangle. As above, one can see that the sets $\mathcal{R}(\gamma_4^{\perp}) \cup \{R_1, R_3\}$, $\mathcal{R}(\mathbf{x}_4^{\perp}) \cup \{R_2, R_3\}$, $\mathcal{R}(\mathbf{x}_4^{\perp}) \cup \{R_3\}$,

ON THE PICARD NUMBER OF FANO 3-FOLDS WITH TERMINAL SINGULARITIES 9

 $\mathcal{R}(\gamma_4^{\perp}) \cup \{R_3, R_4\}$, and $\mathcal{R}(\gamma_4^{\perp}) \cup \{R_4, R_1\}$ are extremal. It follows that there are *E*-sets \mathcal{L}, \mathcal{N} such that $\{R_1, R_3\} \subset \mathcal{L} \subset \mathcal{R}(\gamma_4^{\perp}) \cup \{R_1, R_3\}$ and $\{R_2, R_4\} \subset \mathcal{N} \subset$ $\mathcal{R}(\gamma_4^{\perp}) \cup \{R_2, R_4\}$. By the Lemma 1.1, there exist rays $R \in \mathcal{L}$ and $Q \in \mathcal{N}$ such that RQ is an arrow. By the condition (a) of the Theorem, one of the rays R_1, R_3 is joined by an oriented path s_1 of the length $\leq d$ with the ray R and this path does not contain another ray from R_1, R_3 (here R is the terminal of the path s_1). We can suppose that this ray is R_1 (otherwise, one should replace the ray R_1 by the ray R_3). As above, we can suppose that the ray Q is connected by the oriented path s_2 of the length $\leq d$ with the ray R_2 and this path does not contain the ray R_4 . The path $s_1 R Q s_2$ is an oriented path of the length $\leq 2d+1$ in the oriented graph $G(\mathcal{R}(\gamma_4^{\perp})) \cup \{R_1, R_2\})$. It follows that the oriented angle of the quadrangle γ_4 , such that consecutive sides of this angle are orthogonal to the rays R_1 and R_2 respectively, has the weight $\geq 1/2$. Thus, we proved that for a pair of opposite sides of γ_4 there exists an oriented angle with weight $\geq 1/2$ such that the first side of this oriented angle is one of this opposite sides of the quadrangle. A quadrangle has two pairs of opposite sides. It follows that the sum of weights of oriented angles of γ_4 is ≥ 1 . It proves the condition (2) of the Lemma 1.4 and the Theorem.

Below, we apply the Theorem 1.2 to 3-folds.

CHAPTER 2. THREEFOLDS

1. Contractible extremal rays.

We consider normal projective 3-folds X with \mathbb{Q} -factorial singularities.

Let R be an extremal ray of Mori polyhedron NE(X) of X. A morphism f: $X \to Y$ on a normal projective variety Y is called the *contraction* of the ray R if for an irreducible curve C of X the image f(C) is a point iff $C \in R$. The contraction f is defined by a linear system H on X (H gives the nef element of $N^1(X)$, which we denote by H also). It follows that an irreducible curve C is contracted iff $C \cdot H = 0$. We assume that the contraction f has properties: $f_* \mathcal{O}_X = \mathcal{O}_Y$ and the sequence

$$(1-1) 0 \to \mathbb{R}R \to N_1(X) \to N_1(Y) \to 0$$

is exact where the arrow $N_1(X) \to N_1(Y)$ is f_* . An extremal ray R is called *contractible* if there exists its contraction f with these properties.

The number $\kappa(R) = \dim Y$ is called *Kodaira dimension* of the contractible extremal ray R.

A face γ of NE(X) is called *contractible* if there exists a morphism $f: X \to Y$ on a normal projective variety Y such that $f_*\gamma = 0$, $f_*\mathcal{O}_X = \mathcal{O}_Y$ and f contracts curves lying in γ only. The $\kappa(\gamma) = \dim Y$ is called *Kodaira dimension of* γ .

Let *H* be a general nef element orthogonal to a face γ of Mori polyhedron. Numerical Kodaira dimension of γ is defined by the formula

$$\kappa_{num}(\gamma) = \begin{cases} 3, & \text{if } H^3 > 0; \\ 2, & \text{if } H^3 = 0 \text{ and } H^2 \neq 0; \\ 1, & \text{if } H^2 \equiv 0. \end{cases}$$

It is obvious that for a contractible face γ we have $\kappa_{num}(\gamma) \geq \kappa(\gamma)$. In particular, $\kappa_{num}(\gamma) = \kappa(\gamma)$ for a contractible face γ of Kodaira dimension $\kappa(\gamma) = 3$.

2. Pairs of extremal rays of Kodaira dimension three lying in contractible faces of $\overline{NE}(X)$ of Kodaira dimension three.

Further X is a projective normal threefold with \mathbb{Q} -factorial singularities.

Lemma 2.2.1. Let R be a contractible extremal ray of Kodaira dimension 3 and $f: X \to Y$ its contraction.

Then there are three possibilities:

(I) All curves $C \in R$ fill an irreducible Weil divisor D(R), the contraction f contracts D(R) in a point and $R \cdot D(R) < 0$.

(II) All curves $C \in R$ fill an irreducible Weil divisor D(R), the contraction f contracts D(R) on an irreducible curve and $R \cdot D(R) < 0$.

(III) (small extremal ray) All curves $C \in R$ give a finite set of irreducible curves and the contraction f contracts these curves in points.

Proof. Assume that some curves of R fill an irreducible divisor D. Then $R \cdot D < 0$ (this inequality follows from the Proposition 2.2.6 below). Suppose that $C \in R$ and D does not contain C. It follows that $R \cdot D \ge 0$. We get a contradiction. It follows the Lemma.

According to the Lemma 2.2.1, we say that an extremal ray R has the type (I), (II) or (III) (small) if it is contractible of Kodaira dimension 3 and the statements (I) (II) or (III) respectively hold

Lemma 2.2.2. Let R_1 and R_2 are two different extremal rays of the type (I). Then divisors $D(R_1)$ and $D(R_2)$ do not intersect one another.

Proof. Otherwise, $D(R_1)$ and $D(R_2)$ have a common curve and the rays R_1 and R_2 are not different.

For an irreducible Weil divisor D on X let

 $\overline{NE}(X, D) = (\text{image } \overline{NE}(D)) \subset \overline{NE}(X).$

Lemma 2.2.3. Let R be an extremal ray of the type (II), and f its contraction. Then $\overline{NE}(X, D(R)) = R + \mathbb{R}^+S$, where $\mathbb{R}^+f_*S = \mathbb{R}^+(f(D))$.

Proof. This follows at once from the exact sequence (1.1).

Lemma 2.2.4. Let R_1 and R_2 are two different extremal rays of the type (II) such that the divisors $D(R_1) = D(R_2)$.

Then for $D = D(R_1) = D(R_2)$ we have:

$$\overline{NE}(X,D) = R_1 + R_2.$$

In particular, do not exist three different extremal rays of the type (II) such that their divisors are coincided.

Proof. This follows from the Lemma 2.2.3.

Lemma 2.2.5. Let R be an extremal ray of the type (II) and f its contraction.

Then there does not exist more than one extremal ray Q of the type (I) such that $D(R) \cap D(Q)$ is not empty. If Q is this ray, then $D(R) \cap D(Q)$ is a curve and any irreducible component of this curve is not contained in fibers of f.

Proof. The last statement is obvious. Let us proof the first one. Suppose that Q_1 and Q_2 are two different extremal rays of the type (I) such that $D(Q_1) \cap D(R)$ and $D(Q_2) \cap D(R)$ are not empty. Then the plane angle $\overline{NE}(X, D(R))$ (see the Lemma 2.2.3) contains three different extremal rays: Q_1, Q_2 and R. It is impossible.

The following key Proposition is very important.

Proposition 2.2.6. Let X be a projective 3-fold with \mathbb{Q} -factorial singularities, $D_1, ..., D_m$ irreducible divisors on X and $f: X \to Y$ a surjective morphism such that dim $X = \dim Y$ and dim $f(D_i) < \dim D_i$. Let $y \in f(D_1) \cap ... \cap f(D_m)$.

Then there are $a_1 > 0, ..., a_m > 0$ and an open $U, y \in U \subset f(D_1) \cup ... \cup f(D_m)$, such that

$$C \cdot (a_1 D_1 + \dots + a_m D_m) < 0$$

if a curve $C \subset D_1 \cup ... \cup D_m$ belongs to a non-trivial algebraic family of curves on $D_1 \cup ... \cup D_m$ and $f(C) = point \in U$.

Proof. It is the same as for the well-known case of surfaces (but, for surfaces, it is not necessary to suppose that C belongs to a nontrivial algebraic family). Let Hbe an irreducible ample divisor on X and $H' = f_*H$. Since dim $f(D_i) < \dim D_i$, it follows that $f(D_1) \cup \ldots \cup f(D_m) \subset H'$. Let ϕ be a non-zero rational function on Ywhich is regular in a neighbourhood U of y on Y and is equal to zero on the divisor H'. In the open set $f^{-1}(U)$ the divisor

$$(f^*\phi) = \sum^m a_i D_i + \sum^n b_j Z_j$$

where all $a_i > 0$ and all $b_j > 0$. Here every divisor Z_j is different from any divisor D_i . We have

$$0 = C \cdot \sum_{i=1}^{m} a_i D_i + C \cdot \sum_{j=1}^{n} b_j Z_j.$$

Here $C \cdot (\sum_{j=1}^{n} b_j Z_j) > 0$ since C belongs to a nontrivial algebraic family of curves on a surface $D_1 \cup \ldots \cup D_m$ and one of the divisors Z_j is the hyperplane section H.

Lemma 2.2.7. Let R_1, R_2 are two extremal rays of the type (II), divisors $D(R_1)$, $D(R_2)$ are different and $D(R_1) \cap D(R_2) \neq \emptyset$. Assume that R_1, R_2 belong to a contractible face of $\overline{NE}(X)$ of Kodaira dimension 3. Let $0 \neq F_1 \in R_1$ and $0 \neq F_2 \in R_2$.

Then

$$(F_1 \cdot D(R_2))(F_2 \cdot D(R_1)) < (F_1 \cdot D(R_1))(F_2 \cdot D(R_2)).$$

Proof. Let f be the contraction of a face of Kodaira dimension 3, which contains both rays R_1, R_2 . By the proposition 2.2.6, there are $a_1 > 0, a_2 > 0$ such that

$$a_1(F_1 \cdot D(R_1)) + a_2(F_1 \cdot D(R_2)) < 0$$
 and $a_1(F_2 \cdot D(R_1)) + a_2(F_2 \cdot D(R_2)) < 0$.

Or

$$-a_1(F_1 \cdot D(R_1)) > a_2(F_1 \cdot D(R_2))$$
 and $-a_2(F_2 \cdot D(R_2)) > a_1(F_2 \cdot D(R_1))$

where $F_1 \cdot D(R_1) < 0$, $F_2 \cdot D(R_2) < 0$ and $F_1 \cdot D(R_2) > 0$, $F_2 \cdot D(R_1) > 0$. Multiplying inequalities above, we obtain the Lemma.

3. A classification of extremal sets of extremal rays which contain extremal rays of the type (I) and simle extremal rays of the type (II).

As above, we assume that X is a projective normal 3-fold with \mathbb{Q} -factorial singularities.

Definition 2.3.1. An extremal ray R of the type (II) is called *simple* if

$$R \cdot (D(R) + D) \ge 0$$

for any irreducible divisor D such that $R \cdot D > 0$.

The following statement gives a simple sufficient condition for an extremal ray to be simple.

Proposition 2.3.2. Let R be an extremal ray of the type (II) and $f: X \to Y$ the contraction of R. Suppose that the curve f(D(R)) is not contained in the set of singularities of Y.

Then

(1) the ray R is simple;

(2) if X has only isolated singularities, then a general element C of the ray R (a general fiber of the morphism $f \mid D(R)$) is isomorphic to \mathbb{P}^1 and the divisor D(R) is non-singular along C. If additionally $R \cdot K_X < 0$, then $C \cdot D(R) = C \cdot K_X = -1$.

(3) In particular, both statements (1) and (2) are true if X has terminal singularities and $R \cdot K_X < 0$.

Proof. Let D be an irreducible divisor on X such that $R \cdot D > 0$. Since $R \cdot D(R) < 0$, the divisor D is different from D(R) and the intersection $D \cap D(R)$ is a surrouted by D(R) = 0.

which does not belong to R. Then $D' = f_*(D)$ is an irreducible divisor on Y and $\Gamma = f(D(R))$ is a curve on D'. Let $y \in \Gamma$ be a non-singular point of Y. Then the divisor D' is defined by some local equation ϕ in a neighbourhood U of y. Evidently, in the open set $f^{-1}(U)$ the divisor

$$(f^*\phi) = D + m(D(R))$$

where the integer $m \ge 1$. Let a curve $C \in R$ and $f(C) = y \in U \cap f(D(R))$. Then $0 = C \cdot (D + m(D(R))) = C \cdot (D + D(R)) + C \cdot (m - 1)(D(R))$. Since $m \ge 1$ and $C \cdot D(R) < 0$, it follows that $C \cdot (D + D(R)) \ge 0$.

Let us prove (2). Let us consider a linear system |H| of hyperplane sections on Y and the corresponding linear systems on the resolutions of singularities of Y and X. Let us apply Bertini's theorem (see, for example, [H, ch. III, Corollary 10.9 and the Exercise 11.3]) to this linear systems. Singularities of X and Y are isolated. Then by Bertini theorem, for a general element H of |H| we obtain that (a) H and $f^{-1}(H)$ are irreducible and non-singular; (b) H intersects Γ transversely in non-singular points of Γ . Let us consider the corresponding birational morphism $f' = f | H' : H' \to H$ of the non-singular irreducible surfaces. It is a composition of blowing ups in non-singular points. Thus, fibers of f' over $H \cap \Gamma$ are trees of non-singular rational curves. The exceptional curve of the first of these blowing ups is identified with the fiber of the projectivization of the normal bundle $\mathbb{P}(\mathcal{N}_{\Gamma/Y})$. Thus, we obtain a rational map over the curve Γ

$$\phi: \mathbb{P}(\mathcal{N}_{\Gamma/Y}) \to D(R)$$

of the irreducible surfaces. Evidently, it is the injection in the general point of $\mathbb{P}(\mathcal{N}_{\Gamma/Y})$. It follows that ϕ is a birational isomorphism of the surfaces. Since ϕ is a birational map over the curve Γ , it follows that the general fibers of this maps are birationally isomorphic. It follows that a general fiber of f' is $C \simeq \mathbb{P}^1$. Since C is non-singular and is an intersection of the non-singular surface H' with the surface D(R), and since X has only isolated singularities, it follows that D(R) is non-singular along the general curve C.

The X and D(R) are non-singular along $C \simeq \mathbb{P}^1$ and the curve C is non-singular. Then the canonical class $K_C = (K_X + D(R)) | C$ where both divisors K_X and D(R) are Cartier divisors on X along C. It follows that $-2 = \deg K_C = K_X \cdot C + D(R) \cdot C$, where the both numbers $K_X \cdot C$ and $D(R) \cdot C$ are negative integers. Then $D(R) \cdot C = K_X \cdot C = -1$.

If X has terminal sungularities and $R \cdot K_X < 0$, then Y has terminal singularities too (see, for example, [Ka1]). Moreover, 3-dimensional terminal singularities are isolated. From (1), (2), the last statement of the Proposition follows.

In connection with Proposition 2.3.2, also see [Mo2, 1.3 and 2.3.2] and [I, Lemma 1].

Let R_1, R_2 are two extremal rays of the type (I) or (II). They are joined if $D(R_1) \cap D(R_2) \neq \emptyset$. It defines *connected components* of a set of extremal rays of the type (I) or (II).

We recall (see Chapter I) that a set \mathcal{E} of extremal rays is called *extremal* if it is contained in a face of $\overline{NE}(X)$. We say that \mathcal{E} is *extremal of Kodaira dimension 3* if it is contained in a face of numerical Kodaira dimension 3 of $\overline{NE}(X)$.

We prove the following elegification regult

Theorem 2.3.3. Let $\mathcal{E} = \{R_1, R_2, ..., R_n\}$ be an extremal set of extremal rays of the type (I) or (II). Suppose that every extremal ray of \mathcal{E} of the type (II) is simple. Assume that \mathcal{E} is contained in a contractible face with Kodaira dimension 3 of $\overline{NE}(X)$. (Thus, \mathcal{E} is extremal of Kodaira dimension 3.)

Then every connected component of \mathcal{E} has a type $\mathfrak{A}_1, \mathfrak{B}_2, \mathfrak{C}_m$ or \mathfrak{D}_2 below (see figure 1).

 (\mathfrak{A}_1) One extremal ray of the type (I).

 (\mathfrak{B}_2) Two different extremal rays S_1, S_2 of the type (II) such that their divisors $D(S_1) = D(S_2)$ are coincided.

 $(\mathfrak{C}_m) \ m \geq 1$ extremal rays $S_1, S_2, ..., S_m$ of the type (II) such that their divisors $D(S_2), D(S_3), ..., D(S_m)$ do not intersect one another, and $S_1 \cdot D(S_i) = 0$ and $S_i \cdot D(S_1) > 0$ for i = 2, ..., m.

 (\mathfrak{D}_2) Two extremal rays S_1, S_2 , where S_1 is of the type (II) and S_2 of the type (I), $S_1 \cdot D(S_2) > 0$ and $S_2 \cdot D(S_1) > 0$. Either $S_1 \cdot (b_1D(S_1) + b_2D(S_2)) < 0$ or $S_2 \cdot (b_1D(S_1) + b_2D(S_2)) < 0$ for any b_1, b_2 such that $b_1 \ge 0, b_2 \ge 0$ and one of b_1, b_2 is not zero.

The following inverse statement is true: If $\mathcal{E} = \{R_1, R_2, ..., R_n\}$ is a connected set of extremal rays of the type (I) or (II) and \mathcal{E} has a type $\mathfrak{A}_1, \mathfrak{B}_2, \mathfrak{C}_m$ or \mathfrak{D}_2 above, then \mathcal{E} generates a simplicial face $R_1 + ... + R_n$ of the dimension n and numerical Kodaira dimension 3 of $\overline{NE}(X)$. In particular, extremal rays of the set \mathcal{E} are linearly independent.

Proof. Let us prove the first statement. We can suppose that \mathcal{E} is connected. We have to prove that then \mathcal{E} has the type $\mathfrak{A}_1, \mathfrak{B}_2, \mathfrak{C}_m$ or \mathfrak{D}_2 . If n = 1, this is obvious.

Let n = 2. From Lemma 2.2.2, it follows that one of the rays R_1, R_2 has the type (II). Let R_1 has the type (II) and R_2 the type (I). Since $D(R_1) \cap D(R_2) \neq \emptyset$, then evidently $R_2 \cdot D(R_1) > 0$. If $R_1 \cdot D(R_2) = 0$, then the curve $D(R_1) \cap D(R_2)$ belongs to the ray R_1 . It follows that the rays R_1 and R_2 contain the same curve. We get a contradiction. Thus, $R_1 \cdot D(R_2) > 0$. The rays R_1, R_2 belong to a contractible face of Kodaira dimension 3 of Mori polyhedron. Let f be a contraction of this face. By the Lemma 2.2.3, f contracts the divisors $D(R_1), D(R_2)$ in a same point. By Proposition 2.2.6, there exist positive a_1, a_2 such that $R_1 \cdot (a_1 D(R_1) + a_2 D(R_2)) < 0$ and $R_2 \cdot (a_1 D(R_1) + a_2 D(R_2)) < 0$. Now suppose that for some $b_1 > 0$ and $b_2 > 0$ the inequalities $R_1 \cdot (b_1 D(R_1) + b_2 D(R_2)) \ge 0$ and $R_2 \cdot (b_1 D(R_1) + b_2 D(R_2)) \ge 0$ hold. There exists $\lambda > 0$ such that $\lambda b_1 \le a_1, \lambda b_2 \le a_2$ and one of these inequalities is an equality. For example, let $\lambda b_1 = a_1$. Then

$$R_1 \cdot (a_1 D(R_1) + a_2 D(R_2)) = R_1 \cdot \lambda (b_1 D(R_1) + b_2 D(R_2)) + R_1 \cdot (a_2 - \lambda b_2) D(R_2) \ge 0.$$

We get a contradiction. It proves that in this case \mathcal{E} has the type \mathfrak{D}_2 . Figure 1.

Now assume that both rays R_1, R_2 have the type (II). Since the rays R_1, R_2 are simple, from Lemma 2.2.7, it follows that either $R_1 \cdot D(R_2) = 0$ or the $R_2 \cdot D(R_1) = 0$. If both these equalities hold, the rays R_1, R_2 have a common curve. We get a contradiction. Thus, in this case, \mathcal{E} has the type \mathfrak{C}_2 .

Let n = 3. Every proper subset of \mathcal{E} has connected components of types $\mathfrak{A}_1, \mathfrak{B}_2, \mathfrak{C}_m$ or \mathfrak{D}_2 . Using Lemmas 2.2.2 —2.2.5, one can see very easily that either \mathcal{E} has the type \mathfrak{C}_3 or we have the following case:

The rays R_1, R_2, R_3 have the type (II), every two element subset of \mathcal{E} has the type \mathfrak{C} and we can find a numeration such that $R = D(R) \ge 0$, $R = D(R) \ge 0$

 $0, R_3 \cdot D(R_1) > 0$. Let f be a contraction of the face γ . By Lemma 2.2.3, f contracts the divisors $D(R_1), D(R_2), D(R_3)$ in a one point. By Proposition 2.2.6, there are positive a_1, a_2, a_3 such that

$$R_i \cdot (a_1 D(R_1) + a_2 D(R_2) + a_3 D(R_3)) < 0$$

for i = 1, 2, 3. On the other hand, from simplicity of the rays R_1, R_2, R_3 , it follows that

$$R_i \cdot (D(R_1) + D(R_2) + D(R_3)) \ge 0.$$

Let $a_1 = \min\{a_1, a_2, a_3\}$. From the last inequality,

$$R_1 \cdot (a_1 D(R_1) + a_2 D(R_2) + a_3 D(R_3)) =$$

 $= R_1 \cdot a_1(D(R_1) + D(R_2) + D(R_3)) + R_1((a_2 - a_1)D(R_2) + (a_3 - a_1)D(R_3)) \ge 0.$

We get a contradiction with the inequality above.

Let n > 3. We have proven that every two or three element subset of \mathcal{E} has connected components of types $\mathfrak{A}_1, \mathfrak{B}_2, \mathfrak{C}_m$ or \mathfrak{D}_2 . It follows very easily that then \mathcal{E} has the type \mathfrak{C}_n (we suppose that \mathcal{E} is connected).

Let us prove the inverse statement.

For the type \mathfrak{A}_1 this is obvious.

Let \mathcal{E} has the type \mathfrak{B}_2 . Since the rays S_1, S_2 are extremal of Kodaira dimension 3, there are *nef* elements H_1, H_2 such that $H_1 \cdot S_1 = H_2 \cdot S_2 = 0, H_1^3 > 0, H_2^3 > 0$. Let $0 \neq C_1 \in S_1$ and $0 \neq C_2 \in S_2$. Let D be a divisor of the rays S_1 and S_2 . Let us consider a map

(3.1)
$$(H_1, H_2) \to H =$$
$$= (-D \cdot C_2)(H_2 \cdot C_1)H_1 + (-D \cdot C_1)(H_1 \cdot C_2)H_2 + (H_2 \cdot C_1)(H_1 \cdot C_2)D.$$

For a fixed H_1 , we get a linear map $H_2 \to H$ of the set of nef elements H_2 orthogonal to S_2 into the set of nef elements H orthogonal to S_1 and S_2 . This map has a one dimensional kernel, generated by $(-D \cdot C_2)H_1 + (H_1 \cdot C_2)D$. It follows that $S_1 + S_2$ is a 2-dimensional face of $\overline{NE}(X)$.

For a general *nef* element $H = a_1H_1 + a_2H_2 + bD$ orthogonal to this face, where $a_1, a_2, b > 0$, we have $H^3 = (a_1H_1 + a_2H_2 + bD)^3 \ge (a_1H_1 + a_2H_2 + bD)^2 \cdot (a_1H_1 + a_2H_2 + bD) \cdot (a_1H_1 + a_2H_2 + bD) \cdot (a_1H_1 + a_2H_2 + bD) = (a_1H_1 + a_2H_2 + bD) \cdot (a_1H_1 + a_2H_2 + bD) \cdot (a_1H_1 + a_2H_2 + bD) = (a_1H_1 + a_2H_2 + bD) \cdot (a_1H_1 + a_2H_2 + bD) \cdot (a_1H_1 + a_2H_2 + bD) = (a_1H_1 + a_2H_2 + bD) \cdot (a_1H_1 + a_2H_2 + bD) \cdot (a_1H_1 + a_2H_2 + bD) = (a_1H_1 + a_2H_2 + bD) \cdot (a_1H_1 + a_2H_2 + bD) = (a_1H_1 + a_2H_2 + bD) \cdot (a_1H_1 + a_2H_2 + bD) = (a_1H_1 + a_2H_2 + bD) \cdot (a_1H_1 + a_2H_2 + bD) = (a_1H_1 + a_2H_2 + bD) \cdot (a_1H_1 + a_2H_2 + bD) = (a_1H_1 + a_2H_2 + bD) = (a_1H_1 + a_2H_2 + bD) \cdot (a_1H_1 + a_2H_2 + bD) = (a_1H_1 + a_2H_2 + bD$ $(a_1H_1+a_2H_2)^2 \cdot (a_1H_1+a_2H_2+bD) \ge (a_1H_1+a_2H_2)^3 > 0$, since $a_1H_1+a_2H_2+bD$ and $a_1H_1+a_2H_2$ are *nef*. It follows that the face S_1+S_2 has numerical Kodaira dimension 3.

Let \mathcal{E} has the type \mathfrak{C}_m . Let H be a *nef* element orthogonal to the ray S_1 . Let $0 \neq C_i \in S_i$. Let us consider a map

(3.2)
$$H \to H' = H + \sum_{i=2}^{m} \left(-(H \cdot C_i) / (C_i \cdot D(S_i)) \right) D(S_i).$$

It is a linear map of the set of nef elements H orthogonal to S_1 into the set of nef elements H' orthogonal to the rays $S_1, S_2, ..., S_m$. The kernel of the map has the dimension m-1. It follows that the rays $S_1, S_2, ..., S_m$ belong to a face of $\overline{NE}(X)$ of a dimension $\leq m$. On the other hand, multiplying rays $S_1, ..., S_m$ on the divisors $D(S_1), ..., D(S_m)$, one can see very easily that the rays $S_1, ..., S_m$ are linearly independent. Thus, they generate a m-dimensional face of $\overline{NE}(X)$. Let us show that this face is $S_1 + S_2 + ... + S_m$. To prove this, we show that every m-1 subset of \mathcal{E} is contained in a face of $\overline{NE}(X)$ of a dimension $\leq m-1$.

If this subset contains the ray S_1 , this subset has the type \mathfrak{C}_{m-1} . By induction, we can suppose that this subset belongs to a face of $\overline{NE}(X)$ of the dimension m-1. Let us consider the subset $\{S_2, S_3, ..., S_m\}$. Let H be an ample element of X. For the element H, the map (3.2) gives an element H' which is orthogonal to the rays $S_2, ..., S_m$, but is not orthogonal to the ray S_1 . It follows that the set $\{S_2, ..., S_m\}$ belongs to a face of the Mori polyhedron of the dimension < m. Like above, one can see that for a general H orthogonal to S_1 the element H' has $(H')^3 \ge H^3 > 0$.

Let \mathcal{E} has the type \mathfrak{D}_2 . Let H be a *nef* element orthogonal to the ray S_2 . Let $0 \neq C_i \in S_i$. Let us consider a map

(3.3)
$$H \to H' = H + \frac{(H \cdot C_1)((-D(S_2) \cdot C_2)D(S_1) + (D(S_1) \cdot C_2)D(S_2))}{(D(S_2) \cdot C_2)(D(S_1) \cdot C_1) - (D(S_1) \cdot C_2)(D(S_2) \cdot C_1)}$$

Evidently, $C_2 \cdot ((-D(S_2) \cdot C_2) D(S_1) + (D(S_1) \cdot C_2) D(S_2)) = 0$. From this equality and the inequality from the definition of the system \mathfrak{D}_2 , it follows that $C_1 \cdot ((-D(S_2) \cdot C_2) D(S_1) + (D(S_1) \cdot C_2) D(S_2)) < 0$. Thus, the denominator from the formula (3.3) is positive. Then (3.3) is a linear map of the set of *nef* elements H orthogonal to the ray S_2 into the set of *nef* elements H' orthogonal to the rays S_1, S_2 . Evidently, the map has a one dimensional kernel. Thus, the rays S_1 and S_2 generate a two dimensional face $S_1 + S_2$ of Mori polyhedron. As above, for a general element Horthogonal to S_2 we have $(H')^3 \geq H^3 > 0$.

Corollary 2.3.4. Let $\mathcal{E} = \{R_1, R_2, ..., R_n\}$ be an extremal set of extremal rays of the type (I) or (II) and every extremal ray of \mathcal{E} of the type (II) is simple. Assume that \mathcal{E} is contained in a contractible face with Kodaira dimension 3 of the $\overline{NE}(X)$. Let $m_1 \geq 0, m_2 \geq 0, ..., m_n \geq 0$ and at least one of $m_1, ..., m_n$ is positive.

Then there exists $i, 1 \leq i \leq n$, such that

$$R_i \cdot (m_1 D(R_1) + \dots + m_n D(R_n)) < 0.$$

Thus, the condition (ii) from Chapter I is valid.

Proof. It is sufficient to prove this statement for the connected \mathcal{E} . For every type $\mathfrak{A}_1, \mathfrak{B}_2, \mathfrak{C}_m$ and \mathfrak{D}_2 of the Theorem 2.3.3, one can prove it very easily.

Unfortunately, in general, the inverse statement of the Theorem 2.3.3 holds only for connected extremal sets \mathcal{E} . We will give two cases when it is true for a non-

Definition 2.3.5. A threefold X is called *strongly projective* (respectively *very strongly projective*) if the following statement holds: a set $\{Q_1, ..., Q_n\}$ of extremal rays of the type (II) is extremal of Kodaira dimension 3 (respectively generates the simplicial face $Q_1 + ... + Q_n$ of $\overline{NE}(X)$ of the dimension n and Kodaira dimension 3) if its divisors $D(Q_1), ..., D(Q_n)$ do not intersect one another.

Theorem 2.3.6. Let $\mathcal{E} = \{R_1, R_2, ..., R_n\}$ be a set of extremal rays of the type (I) or (II) such that every connected component of \mathcal{E} has the type $\mathfrak{A}_1, \mathfrak{B}_2, \mathfrak{C}_m$ or \mathfrak{D}_2 . Then:

(1) \mathcal{E} is extremal of numerical Kodaira dimension 3 if and only if the same is true for any subset of \mathcal{E} containing only extremal rays of the type (II) whose divisors do not intersect one another. In particular, it holds if X is strongly projective.

(2) \mathcal{E} generates a simplicial face $R_1 + ... + R_n$ with numerical Kodaira dimension 3 of the Mori polyhedron if and only if the same is true for any subset of \mathcal{E} containing only extremal rays of the type (II) whose divisors do not intersect one another. In particular, it is true if X is very strongly projective.

Proof. Let us prove (1). Only the inverse statement is non-trivial. We prove it using an induction by n. For n = 1, the statement is obvious.

Assume that some connected component of \mathcal{E} has the type \mathfrak{A}_1 . Suppose that this component contains the ray R_1 . By induction, there exists a nef element H such that $H^3 > 0$ and $H \cdot R_i = 0$ if i > 1. Then there exists $k \ge 0$, such that $H' = H + kD(R_1)$ is nef and $H' \cdot \mathcal{E} = 0$. As above, one can prove that $(H')^3 \ge H^3 > 0$.

Assume that some connected component of \mathcal{E} has the type \mathfrak{B}_2 . Suppose that this component contains the rays R_1, R_2 and $D(R_1) = D(R_2) = D$. Then, by induction, there are nef elements H_1 and H_2 such that $H_1^3 > 0, H_2^3 > 0$ and $H_1 \cdot \{R_1, R_3, ..., R_n\} = 0, H_2 \cdot \{R_2, R_3, ..., R_n\} = 0$. As for the proof of the inverse statement of the Theorem 2.3.3 in the case \mathfrak{B}_2 , there are $k_1 \ge 0, k_2 \ge 0, k_3 \ge 0$ such that the element $H = k_1H_1 + k_2H_2 + k_3D$ is nef, $H \cdot \mathcal{E} = 0$ and $H^3 > 0$.

Assume that some connected component of \mathcal{E} has the type $\mathfrak{C}_m, m > 1$. We use notation of the Theorem 2.3.3 for this connected component. Let this is $\{S_1, S_2, ..., S_m\}$. By the induction, there exists a nef element H such that H is orthogonal to $\mathcal{E} - \{S_2, ..., S_m\}$ and $H^3 > 0$. As for the proof of the inverse statement of the Theorem 2.3.3 in the case \mathfrak{C}_m , there are $k_2 \geq 0, ..., k_m \geq 0$ such that $H' = H + k_2 D(S_2) + ... + k_m D(S_m)$ is nef, $H' \cdot \mathcal{E} = 0$ and $(H')^3 \geq H^3 > 0$.

Assume that some connected component of \mathcal{E} has the type \mathfrak{D}_2 . We use notation of the Theorem 2.3.3 for this connected component. Let this is $\{S_1, S_2\}$. By the induction, there exists nef element H such that $H^3 > 0$ and H is orthogonal to $\mathcal{E} - \{S_1\}$. As for the theorem 2.3.3, there are $k_1 \geq 0, k_2 \geq 0$ such that H' = $H + k_1 D(S_1) + k_2 D(S_2)$ is nef, $H' \cdot \mathcal{E} = 0$ and $(H')^3 \geq H^3 > 0$.

If every connected component of \mathcal{E} has the type \mathfrak{C}_1 , then the statement holds by the condition of the Theorem.

Let us prove (2). Only the inverse statement is non-trivial. We prove it using an induction by n. For n = 1 the statement is true. It is sufficient to prove that \mathcal{E} is contained in a face of a dimension $\leq n$ of Mori polyhedron because, by the induction, any its n-1 element subset generates a simplicial face of the dimension n-1 of Mori polyhedron.

A gauge that some connected component of \mathcal{L} has the type \mathfrak{A} . Suppose that the

ray R_1 belongs to this component and $0 \neq C_1 \in R_1$. Let us consider the map

$$H \to H' = H + ((H \cdot C_1)/(-D(R_1) \cdot C_1))D(R_1)$$

of the set of nef elements H orthogonal to the set $\{R_2, ..., R_n\}$ into the set of nef elements H' orthogonal to the \mathcal{E} . It is the linear map with one dimensional kernel. Since, by the induction, the set $\{R_2, ..., R_n\}$ is contained in a face of Mori polyhedron of the dimension n - 1, it follows that \mathcal{E} is contained in a face of the dimension n.

If \mathcal{E} has a connected component of the type $\mathfrak{B}_2, \mathfrak{C}_m, m > 1$, or \mathfrak{D}_2 , a proof is the same if one uses the maps (3.1), (3.2) and (3.3) above.

If all connected components of \mathcal{E} have the type \mathfrak{C}_1 , the statement holds by the condition.

Remark 2.3.7. Like the statement (1) of the Theorem 2.3.6, one can prove that a set \mathcal{E} of extremal rays with connected components of the type $\mathfrak{A}_1, \mathfrak{B}_2, \mathfrak{C}_m$ or \mathfrak{D}_2 is extremal if and only if the same is true for any subset of \mathcal{E} containing only extremal rays of the type (II) whose divisors do not intersect one another.

The next statement is simple but important. To simplify notation, we say that for a fixed $a_1, ..., a_n$ we have a *linear dipendence condition*

$$a_1R_1 + \dots + a_nR_n = 0$$

between extremal rays $R_1, ..., R_n$ if there exist non-zero $C_i \in R_i$ such that

$$a_1C_1 + \ldots + a_nC_n = 0.$$

Proposition 2.3.8. Assume that a set $\mathcal{E} = \{R_1, R_2, ..., R_m\}$ of extremal rays has connected components of the type $\mathfrak{A}_1, \mathfrak{B}_2, \mathfrak{C}_m$ or \mathfrak{D}_2 and there exists a linear dependence condition $a_1R_1 + a_2R_2 + ... + a_mR_m = 0$ with all $a_i \neq 0$.

Then all connected components of \mathcal{E} have the type \mathfrak{B}_2 . Let these components are $\mathfrak{B}^1, ..., \mathfrak{B}^t$. Then $t \geq 2$, and we can choose a numeration such that $\mathfrak{B}^i = \{R_{i1}, R_{i2}\}$ and the linear dependence has a form

$$a_{11}R_{11} + a_{21}R_{21} + \dots + a_{t1}R_{t1} = a_{12}R_{12} + a_{22}R_{22} + \dots + a_{t2}R_{t2}.$$

where all $a_{ij} > 0$.

Proof. Let us multiply the equality $a_1R_1 + a_2R_2 + ... + a_mR_m = 0$ on divisors $D(R_1), ..., D(R_m)$. Then we get that $a_k = 0$ if the ray R_k belongs to a connected component of the type $\mathfrak{A}_1, \mathfrak{C}_m$ or \mathfrak{D}_2 . Thus, all connected components of \mathcal{E} have the type \mathfrak{B}_2 . Let these components are

$$\mathfrak{B}^1 = \{R_{11}, R_{12}\}, \mathfrak{B}^2 = \{R_{21}, R_{22}\}, \dots, \mathfrak{B}^t = \{R_{t1}, R_{t2}\}.$$

Obviously, $t \geq 2$, and we can rewrite the linear dependence as

$$a_{11}R_{11} + a_{12}R_{12} + a_{21}R_{21} + a_{22}R_{22} + \dots + a_{t1}R_{t1} + a_{t2}R_{t2} = 0,$$

where all $a_{ij} \neq 0$. Multiplying this equation on all divisors $D(R_{ij})$ and using inequalities $R_{ij} \cdot D(R_{ij}) < 0$, we get the last statement of Proposition.

4. A classification of *E*-sets of extremal rays of the type (I) or (II).

As above, we suppose that X is a projective normal 3-fold with \mathbb{Q} -factorial singularities.

We recall that a set \mathcal{L} of extremal rays is called *E-set* if it is not extremal but any proper subset of \mathcal{L} is extremal (it is contained in a face of $\overline{NE}(X)$). Thus, an *E*-set is a minimal non-extremal set of extremal rays.

Theorem 2.4.1. Let \mathcal{L} be a E-set of extremal rays of the type (I) or (II). Suppose that every ray of the type (II) of \mathcal{L} is simple and every proper subset of \mathcal{L} is contained in a contractible face of Kodaira dimension 3 of Mori polyhedron.

Then we have one of the following cases:

(a) \mathcal{L} is connected and $\mathcal{L} = \{R_1, R_2, R_3\}$, where any R_i has the type (II) and every of 2-element subsets $\{R_1, R_2\}, \{R_2, R_3\}, \{R_3, R_1\}$ of \mathcal{L} has the type \mathfrak{C}_2 . Here $R_1 \cdot D(R_2) > 0, R_2 \cdot D(R_3) > 0, R_3 \cdot D(R_1) > 0$ but $R_2 \cdot D(R_1) = R_3 \cdot D(R_2) =$ $R_1 \cdot D(R_3) = 0$. The divisor $D(\mathcal{L}) = D(R_1) + D(R_2) + D(R_3)$ is nef.

(b) \mathcal{L} is connected and $\mathcal{L} = \{R_1, R_2\}$, where at least one of the rays R_1, R_2 has the type (II). There are positive m_1, m_2 such that $R \cdot (m_1 D(R_1) + m_2 D(R_2)) \geq 0$ for any extremal ray R of the type (I) or simple extremal ray of the type (II) on X. If the divisor $m_1 D(R_1) + m_2 D(R_2)$ is not nef, both the extremal rays R_1, R_2 have the type (II).

(c) \mathcal{L} is connected and $\mathcal{L} = \{R_1, R_2\}$ where both R_1 and R_2 have the type (II) and there exists the simple extremal ray S_1 of the type (II) such that the rays R_1, S_1 define the extremal set of the type \mathfrak{B}_2 (it means that $S_1 \neq R_1$ but the divisors $D(S_1) = D(R_1)$) and the rays S_1, R_2 define the extremal set of the type \mathfrak{C}_2 , where $S_1 \cdot D(R_2) = 0$ but $R_2 \cdot D(S_1) > 0$. Here there do not exist positive m_1, m_2 such that the divisor $m_1D(R_1)+m_2D(R_2)$ is nef, since evidently $S_1 \cdot (m_1D(R_1)+m_2D(R_2)) < 0$. See figure 2 below.

(d) $\mathcal{L} = \{R_1, ..., R_k\}$ where $k \geq 2$, all rays $R_1, ..., R_k$ have the type (II) and the divisors $D(R_1), ..., D(R_k)$ do not intersect one another. Any proper subset of \mathcal{L} is contained in a contractible face of Kodaira dimension 3 of Mori polyhedron but \mathcal{L} is not contained in a face of Mori polyhedron.

Proof. Let $\mathcal{L} = \{R_1, ..., R_n\}$ be a *E*-set of extremal rays satisfying the conditions

The case 1. Let \mathcal{L} is not connected. Then every connected component of \mathcal{L} is extremal and, by the theorem 2.3.3, it has the type $\mathfrak{A}_1, \mathfrak{B}_2, \mathfrak{C}_m$ or \mathfrak{D}_2 . If some of these components does not have the type \mathfrak{C}_1 , then, by the statement (1) of the Theorem 2.3.6, \mathcal{L} is extremal and we get the contradiction. Thus, we get the case (d) of the Theorem.

The case 2. Let $\mathcal{L} = \{R_1, ..., R_n\}$ is connected.

Let $n \geq 4$. By the Theorem 2.3.3, any proper subset of \mathcal{L} has connected components of the type $\mathfrak{A}_1, \mathfrak{B}_2, \mathfrak{C}_m$ or \mathfrak{D}_2 . Like for the proof of the Theorem 2.3.3, it follows that \mathcal{L} has the type \mathfrak{C}_n . By the Theorem 2.3.3, then \mathcal{L} is extremal. We get the contradiction.

Let n = 3. Then, like for the proof of the Theorem 2.3.3, we get that \mathcal{L} has the type (a).

Let n = 2 and $\mathcal{L} = \{R_1, R_2\}$. If both rays R_1, R_2 have the type (I), then, by the Lemma 2.2.2, \mathcal{L} is not connected and we get the contradiction.

Let R_1 has the type (I) and R_2 has the type (II). Since the set \mathcal{L} is not extremal, by the Theorem 2.3.3, there are positive m_1, m_2 such that $R_1 \cdot (m_1 D(R_1) + m_2 D(R_2)) \geq 0$ and $R_2 \cdot (m_1 D(R_1) + m_2 D(R_2)) \geq 0$. By the Lemma 2.2.3, it follows that $C \cdot (m_1 D(R_1) + m_2 D(R_2)) \geq 0$ if the curve C is contained in the $D(R_1) \cup D(R_2)$. If C is not contained in $D(R_1) \cup D(R_2)$, then obviously $C \cdot (m_1 D(R_1) + m_2 D(R_2)) \geq 0$. It follows, that the divisor $m_1 D(R_1) + m_2 D(R_2)$ is nef. Thus, we get the case (b).

Let both rays R_1, R_2 have the type (II). If $D(R_1) = D(R_2)$, then we get an extremal set $\{R_1, R_2\}$ by the Theorem 2.3.3. Thus, the divisors $D(R_1)$ and $D(R_2)$ are different. By the Lemma 2.2.1, the curve $D(R_1) \cap D(R_2)$ does not have an irreducible component which belongs to both rays R_1 and R_2 . Since rays R_1, R_2 are simple, it follows that $R_1 \cdot (D(R_1) + D(R_2)) \ge 0$ and $R_2 \cdot (D(R_1) + D(R_2)) \ge 0$. Let R be an extremal ray of the type (I) or simple extremal ray of the type (II). If the divisor D(R) does not coincide with the divisor $D(R_1)$ or $D(R_2)$, then obviously $R \cdot (D(R_1) + D(R_2)) \ge 0$. Thus, if there does not exist an extremal ray R which has the same divisor as the ray R_1 or R_2 , we get the case (b).

Assume that $D(R) = D(R_1)$. Then, by the Lemma 2.2.5, the ray R has the type (II) too. If $R \cdot D(R_2) = 0$, we get the case (c) of the Theorem where $S_1 = R$. If $R \cdot D(R_2) > 0$, then $R \cdot (D(R_1) + D(R_2)) \ge 0$ since the ray R is simple. Then we get the case (b) of the Theorem.

5. An application of the diagram method to Fano 3-folds with terminal singularities.

We restrict ourselves by considering Fano 3-folds with \mathbb{Q} -factorial terminal singularities, but it is possible to formulate and prove corresponding results for a negative part of Mori cone of 3-dimensional variety with \mathbb{Q} -factorial terminal singularities like in [N7].

We recall that an algebraic 3-fold X over \mathbb{C} with \mathbb{Q} -factorial singularities is called Fano if the anticanonical class $-K_X$ is ample. By results of Kawamata [Ka1] and Shokurov [Sh], any face of $\overline{NE}(X)$ is contractible and $\overline{NE}(X)$ is generated by a finite set of extremal rays if X is a Fano 3-fold with terminal \mathbb{Q} -factorial singularities.

5.1. Preliminary results.

Lemma 2.5.1. Let X be a Fano 3-fold with Q-factorial terminal singularities. Let $\mathcal{E} = \{R_1, ..., R_n\}$ be a set of n extremal rays on of the type (II) and with disjoint divisors $D(R_1), ..., D(R_n)$ on X. (Thus, \mathcal{E} has the type $n\mathfrak{C}_1$).

If we suppose that the set \mathcal{E} is not extremal, then there exists a small extremal ray S and i, $1 \leq i \leq n$, such that $S \cdot (-K_X + D(R_i)) < 0$ and $S \cdot D(R_j) = 0$ if $j \neq i$.

It follows that any curve of the ray S belongs to the divisor $D(R_i)$.

Proof. By Proposition 2.3.2, the divisor $H = -K_X + D(R_1) + ... D(R_n)$ is orthogonal to \mathcal{E} . Besides, H is *nef* and $H^3 > 0$ if there does not exist a small extremal ray S with the property above. Then, \mathcal{E} is extremal of Kodaira dimension 3.

Definition 2.5.2. A set $\{R, S\}$ of extremal rays has the type \mathfrak{E}_2 if the ray R has type (II), the extremal ray S is small and $S \cdot D(R) < 0$. (See Figure 3.)

Thus, by Lemma 2.5.1, the set $R_1, ..., R_n, S$ of extremal rays contains a subset of the type \mathfrak{E}_2 .

From Proposition 2.3.2, any extremal ray of X of the type (II) is simple, and from Sections 3 and 4 we get a classification of extremal sets and E-sets of extremal rays of the type (I) and (II) on X.

Figure 3.

We have the following general Theorem.

Theorem 2.5.3. Let X be a Fano 3-fold with \mathbb{Q} -factorial terminal singularities. Let α be a face of $\overline{NE}(X)$. Then we have the following possibilities:

(1) There exists a small extremal ray S such that $\alpha + S$ is contained in a face of $\overline{NE}(X)$ of Kodaira dimension 3.

(2) There are extremal rays R_1, R_2 of the type (II) and a small extremal ray S such that $\alpha + R_1$ and $\alpha + R_2$ are contained in faces of $\overline{NE}(X)$ of Kodaira dimension 3, the ray R_2 does not belong to α , and one of the sets $\{R_1, S\}$ or $\{R_2, S\}$ has the type \mathfrak{E}_2 .

(3) The face α is contained in a face of $\overline{NE}(X)$ of Kodaira dimension 1 or 2.

(4) There exists an E-set $\mathcal{L} = \{R_1, R_2\}$ such that $R_1 \not\subset \alpha, R_2 \not\subset \alpha$, but $\alpha + R_1$ and $\alpha + R_2$ are contained in faces of $\overline{NE}(X)$ of Kodaira dimension 3. The \mathcal{L} satisfies the condition (c) of Theorem 2.4.1: Thus, both extremal rays R_1, R_2 have the type (II) $R_1 \cdot D(R_2) > 0$ and $R_2 \cdot D(R_1) > 0$ and there exists an extremal ray R'_1 of the type (II) exclusion that $D(R_2) = D(R'_1) = 0$ (5) There are extremal rays $R_1, ..., R_n$ of the type (II) such that any of them does not belong to α , $\alpha + R_1 + ... + R_n$ is contained in a face of $\overline{NE}(X)$ of Kodaira dimension 3 and

$$\dim \alpha + R_1 + \dots + R_n < \dim \alpha + n.$$

(6) dim $N_1(X)$ – dim $\alpha \leq 12$.

Proof. Let us consider the face $\gamma = \alpha^{\perp}$ of $\mathcal{M}(X)$ and apply Theorem 1.2 to this face γ . We have dim $\gamma = \dim N_1(X) - 1 - \dim \alpha$.

Assume that α does not satisfy the conditions (1), (3) and (5). Then $\mathcal{R}(\gamma)$ contains extremal rays of the type (I) or (II) only and $\mathcal{M}(X)$ is closed and simple in the face γ . By Proposition 2.3.2 and Theorem 2.3.3, any extremal subset \mathcal{E} of $\mathcal{R}(\gamma)$ has connected components of the types $\mathfrak{A}_1, \mathfrak{B}_2, \mathfrak{C}_n$ or \mathfrak{D}_2 . By Corollary 2.3.4, the condition (ii) is valid for extremal subsets of $R(\gamma)$. Let $\mathcal{L} \subset \mathcal{R}(\gamma)$ be a *E*-set. Assume that at least two elements $R_1, R_2 \in \mathcal{L}$ don't belong to $\mathcal{R}(\gamma^{\perp})$ and for any proper subset $\mathcal{L}' \subset \mathcal{L}$ we have that $\mathcal{L}' \cup \mathcal{R}(\gamma^{\perp})$ is extremal. Let us apply Theorem 2.4.1 to \mathcal{L} .

Assume that \mathcal{L} has the type (d). By Lemma 2.5.1, one of extremal rays R_1 of \mathcal{L} together with some small extremal ray S define a set of the type \mathfrak{E}_2 . Since $\{R_1\} \subset \mathcal{L}$ is a proper subset of \mathcal{L} , the $\mathcal{R}(\gamma^{\perp}) \cup \{R_1\}$ is extremal. Or $\alpha + R_1$ is contained in a face of $\overline{NE}(X)$. Since \mathcal{L} has at least 2 elements which do not belong to $\mathcal{R}(\gamma^{\perp})$, there exists another extremal ray R_2 of \mathcal{L} which does not belong to $\mathcal{R}(\gamma^{\perp})$. Like above, $\alpha + R_2$ is contained in a face of $\overline{NE}(X)$ of Kodaira dimension 3. By definition of the case (d), both extremal rays R_1, R_2 have the type (II). Thus, we get the case (2) of Theorem.

Assume that \mathcal{L} has the type (c). Then we get the case (4) of Theorem.

Assume that $\mathcal{L} = \{R_1, R_2\}$ has the type (b). Suppose that the divisor $m_1 D(R_1) + m_2(D(R_2))$ is not *nef* (see the case (b) of Theorem 2.4.1). Then there exists a small extremal ray S such that $S \cdot (m_1 D(R_1) + m_2 D(R_2)) < 0$. It follows that one of the sets $\{R_1, S\}$ or $\{R_2, S\}$ has the type \mathfrak{E}_2 . Thus, we get the case (2).

Assume that $\mathcal{L} = \{R_1, R_2, R_3\}$ has the type (a). Then the divisor $D(R_1) + D(R_2) + D(R_3)$ is *nef*.

Thus, if we additionally exclude the cases (2) and (4), then all conditions of the Theorem 1.2 are satisfied. By Theorems 2.4.1 and 2.3.3, we can take d = 2, $C_1 = 1$ and $C_2 = 0$. (See Figure 4 for graphs $G(\mathcal{E})$ corresponding to extremal sets \mathcal{E} of the types $\mathfrak{A}_1, \mathfrak{B}_2, \mathfrak{C}_m$ and \mathfrak{D}_2 .) Thus, by Theorem 1.2, dim $\gamma < 34/3$. It follows that dim $N_1(X) - \dim \alpha \leq 12$.

Figure 4

5.2. General properties of configurations of extremal rays of the type \mathfrak{B}_2 .

Let $\{R_{11}, R_{12}\}$ be a set of extremal rays of the type \mathfrak{B}_2 . By Theorem 2.3.3, they define a 2-dimensional face $R_{11} + R_{12}$ of $\overline{NE}(X)$. Let $\{R_{21}, R_{22}\}$ be another set of extremal rays of the type \mathfrak{B}_2 . Since two different 2-dimensional faces of $\overline{NE}(X)$ may have only a common extremal ray, the divisors $D(R_{11}) = D(R_{12})$ and $D(R_{21}) = D(R_{22})$ don't have a common point. There exists the maximal set $\{R_{11}, R_{12}\}, \{R_{21}, R_{22}\}, ..., \{R_{n1}, R_{n2}\}$ of pairs of extremal rays of the type \mathfrak{B}_2 .

Lemma 2.5.4. Any t pairs $\{R_{11}, R_{12}\}, \{R_{21}, R_{22}\}, ..., \{R_{t1}, R_{t2}\}$ of extremal rays of the type \mathfrak{B}_2 generate a face

$$\sum_{i=1}^{t} \sum_{j=1}^{2} R_{ij} \subset \overline{NE}(X) \subset N_1(X)$$

of the Kodaira dimension 3 of NE(X).

Proof. This face is orthogonal to the nef divisor $H = -K_X + D(R_{11}) + ... + D(R_{t1})$ with $H^3 \ge (-K_X)^3 > 0$.

Lemma 2.5.5. In notation above, there exists a changing of numeration of pairs of extremal rays R_{i1}, R_{i2} such that $R_{11} + \cdots + R_{t1}$ is a simplicial face of $\overline{NE}(X)$.

Proof. For t = 1, it is obvious. Let us suppose that $\theta = R_{11} + \cdots + R_{(t-1)1}$ is a simplicial face of the face

$$\alpha_{t-1} = \sum_{i=1}^{t-1} \sum_{j=1}^{2} R_{ij}.$$

The face

$$\alpha_t = \sum_{i=1}^t \sum_{j=1}^2 R_{ij}$$

has α_{t-1} as its face and does not coincide with the face α_{t-1} . It follows that there exists a face β of α_t of the dimension t such that $\beta \not\subset \alpha_{t-1}$ but $\theta \subset \beta$ is a face of β . It follows that all extremal rays of β are the extremal rays $R_{11}, ..., R_{(t-1)1}$ and some of extremal rays R_{t1}, R_{t2} . Assume that both extremal rays R_{t1}, R_{t2} belong to β . Then the extremal rays $R_{11}, ..., R_{(t-1)1}, R_{t1}, R_{t2}$ are linearly dependent, since dim $\beta = t$. By Proposition 2.3.8, it is impossible. Thus, only one of extremal rays R_{t1}, R_{t2} belongs to the face β . Suppose that this is R_{t1} . Then $\beta = R_{11} + \cdots + R_{(t-1)1} + R_{t1}$ will be the face we were looking for.

We divide the maximal set $\{R_{11}, R_{12}\}, \{R_{21}, R_{22}\}, ..., \{R_{n1}, R_{n2}\}$ of pairs of extremal rays of the type \mathfrak{B}_2 on two parts:

$$\{R_{11}, R_{12}\}, \{R_{21}, R_{22}\}, .., \{R_{m1}, R_{m2}\}$$

and

$$\{R_{(m+1)1},R_{(m+1)2}\},\{R_{(m+2)1},R_{(m+2)2}\},..,\{R_{(m+k)1},R_{(m+k)2}\}$$

where n = m + k. By definition, here the extremal rays R_{i1} , R_{i2} belong to the first part if and only if they are linearly independent from other extremal rays from the set $\{R_{11}, R_{12}\}, \{R_{21}, R_{22}\}, ..., \{R_{n1}, R_{n2}\}$. Thus, extremal rays R_{j1}, R_{j2} belong to the second part if they are linearly dependent from other extremal rays from the set $\{R_{n1}, R_{n2}\}, \{R_{n2}, R_{n2}\}, ..., \{R_{n2}, R_{n2}\}$. **Lemma 2.5.6.** Let S be an extremal ray of the type (II) such that $\{R_{i1}, S\}$ define a configuration (c) of the Theorem 2.4.1. Thus: $R_{i1} \cdot D(S) > 0$, $S \cdot D(R_{i1}) > 0$ and $R_{i2} \cdot D(S) = 0$. Then the extremal rays R_{i1}, R_{i2} are linearly independent from all other extremal rays in $\{R_{11}, R_{12}\}, \{R_{21}, R_{22}\}, ..., \{R_{n1}, R_{n2}\}$. Thus, $1 \le i \le m$. There does not exist a configuration of this type with the ray R_{i2} . Thus, there does not exist an extremal ray S' of the type (II) such that $R_{i2} \cdot D(S') > 0$, $S' \cdot D(R_{i2}) > 0$ and $R_{i1} \cdot D(S') = 0$.

Proof. The $R_{i1} + R_{i2}$ and $R_{i2} + S$ are 2-dimensional faces of NE(X) with intersection by the extremal ray R_{i2} . It follows that any curve of D(S) belongs to the face $R_{i2} + S$ (by Lemma 2.2.3). It follows that the divisor D(S) has no common point with the divisor $D(R_{j1})$ for any other pair R_{j1}, R_{j2} for $j \neq i$. Multiplying on D(S) a linear relation of extremal rays R_{i1}, R_{i2} with other extremal rays $\{R_{11}, R_{12}\}, \{R_{21}, R_{22}\}, ..., \{R_{n1}, R_{n2}\}$ and using Proposition 2.3.8, we get that this linear relation does not exist.

Let us suppose that there exists an extremal ray S' (see formulation of the Lemma). Then $R_{i1} + S'$ is another 2-dimensional face of $\overline{NE}(X)$. Evidently, divisors D(S) and D(S') have a non-empty intersection. Thus, faces $R_{i2} + S$ and $R_{i1} + S'$ have a common ray. But it is possible only if S = S'. Thus, we get a contradiction, because $R_{i1} \cdot D(S) > 0$ but $R_{i1} \cdot D(S') = 0$.

Using this Lemma 2.5.6, we can subdivide the first set

$$\{R_{11}, R_{12}\}, \{R_{21}, R_{22}\}, ..., \{R_{m1}, R_{m2}\}.$$

on sets

$$\{R_{11}, R_{12}\}, \{R_{21}, R_{22}\}, .., \{R_{m_11}, R_{m_12}\}$$

and

$$\{R_{(m_1+1)1}, R_{(m_1+1)2}\}, ..., \{R_{(m_1+m_2)1}, R_{(m_1+m_2)2}\}$$

where $m_1 + m_2 = m$. Here R_{i1}, R_{i2} belong to the first part if and only if there exists an extremal ray S such that R_{i1}, S satisfies the condition of Lemma 2.5.6. By Lemma 2.5.6, the numeration between extremal rays R_{i1} and R_{i2} is then canonical.

Let us consider the second set

$$\{R_{(m+1)1}, R_{(m+1)2}\}, \{R_{(m+2)1}, R_{(m+2)2}\}, ..., \{R_{(m+k)1}, R_{(m+k)2}\}.$$

We introduce an invariant

$$\delta = \dim \sum_{i=m+1}^{m+k} \sum_{j=1}^{2} R_{ij} - k$$

of X. Evidently,

either
$$k = \delta = 0$$
 or $k \ge 2$ and $1 \le \delta < k$.

Thus,

dim
$$\sum_{k=1}^{m+k} \sum_{j=1}^{2} R_{ij} = k + \delta.$$

Let

$$\rho_0(X) = \dim N_1(X) - \dim \sum_{i=1}^{n=m+k} \sum_{j=1}^2 R_{ij}.$$

Then

$$\rho(X) = \dim N_1(X) = \rho_0(X) + 2m + k + \delta.$$

The invariants: $\rho_0(X), n, m, k, \delta, m_1, m_2$ are important invariants of a Fano 3-fold X.

The following Lemma will be very useful:

Lemma 2.5.7. Let \mathcal{E} be an extremal set of extremal rays. Let

$$\{R_{11}, R_{12}\} \cup \ldots \cup \{R_{t1}, R_{t2}\}$$

be a set of different pairs of extremal rays of the type \mathfrak{B}_2 . Assume that $R \cdot D(R_{i1}) = 0$ for any $R \in \mathcal{E}$ and any $i, 1 \leq i \leq t$.

Then there are extremal rays $Q_1, ..., Q_r$ such that the following conditions are valid:

(a) $r \leq t$;

(b) For any $i, 1 \leq i \leq r$, there exists $j, 1 \leq j \leq t$, such that $Q_i \cdot D(R_{j1}) > 0$ (in particular, Q_i is different from extremal rays of pairs of extremal rays $\{R_{u1}, R_{u2}\}$ of the type \mathfrak{B}_2);

(c) For any $j, 1 \leq j \leq t$, there exists an extremal ray $Q_i, 1 \leq i \leq r$, such that

$$Q_i \cdot D(R_{j1}) > 0;$$

(d) The set $\mathcal{E} \cup \{Q_1, ..., Q_r\}$ of extremal rays is extremal, and extremal rays $\{Q_1, ..., Q_r\}$ are linearly independent.

Proof. Let us consider a linear subspace $V \,\subset N_1(X)$ generated by all extremal rays \mathcal{E} . Let us consider a factorisation map $\pi : N_1(X) \to N_1(X)/V$. The cone $\pi(\overline{NE}(X))$ is generated by images of extremal rays T which together with all extremal rays from \mathcal{E} belong to faces of $\overline{NE}(X)$. There exists a curve C on X such that $C \cdot D(R_{11}) > 0$. This curve C (as any element $x \in \overline{NE}(X)$) is a linear combination of extremal rays T with non-negative coefficients and extremal rays from \mathcal{E} with real coefficients. We have $R \cdot D(R_{11}) = 0$ for any extremal ray $R \in \mathcal{E}$. Thus, there exists an extremal ray T above such that $T \cdot D(R_{11}) > 0$. It follows that Tis different from extremal rays of pairs of the type \mathfrak{B}_2 . We take $Q_1 = T$. By our construction, the set $\mathcal{E} \cup \{Q_1\}$ is extremal since it is contained in a face of $\overline{NE}(X)$. If $Q_1 \cdot D(R_{j1}) > 0$ for any j such that $1 \leq j \leq t$, then r = 1, and the set $\{Q_1\}$ gives the set we were looking for. Otherwise, there exists a minimal j such that $2 \leq j \leq t$ and $Q_1 \cdot D(R_{j1}) = 0$. Then we replace \mathcal{E} by $\mathcal{E} \cup \{Q_1\}$ and the set

$$\{R_{11}, R_{12}\} \cup \ldots \cup \{R_{t1}, R_{t2}\}$$

by

$$\{R_{i1}, R_{i2} \mid 1 \le j \le t, \ Q_1 \cdot D(R_{i1}) = 0\}$$

and repeat this procedure.

5.3. Basic Theorems.

We ment to prove the following basic statement

Basic Theorem 2.5.8. Let X be a Fano 3-fold with terminal \mathbb{Q} -factorial singularities. Assume that X does not have a small extremal ray, and Mori polyhedron $\overline{NE}(X)$ does not have a face of Kodaira dimension 1 or 2.

Then we have the following statements for the X:

(1) The X does not have a pair of extremal rays of the type \mathfrak{B}_2 (thus, in notation above, the invariant n = 0) and Mori polyhedron $\overline{NE}(X)$ is simplicial.

(2) The X does not have more than one extremal ray of the type (I).

(3) If \mathcal{E} is an extremal set of k extremal rays of X, then the \mathcal{E} has one of the types: $\mathfrak{A}_1 \amalg (k-1)\mathfrak{C}_1$, $\mathfrak{D}_2 \amalg (k-2)\mathfrak{C}_1$, $\mathfrak{C}_2 \amalg (k-2)\mathfrak{C}_1$, $k\mathfrak{C}_1$ (we use notation of Theorem 2.3.3).

(4) We have the inequality for the Picard number of X:

$$\rho(X) = \dim N_1(X) \le 7.$$

Proof. We use notations introduced in the Section 5.2. We devide the proof on several steps.

Let us consider extremal rays

$$\mathcal{E}_0 = \{R_{11}, R_{12}\} \cup \{R_{21}, R_{22}\} \cup \dots \cup \{R_{n1}, R_{n2}\}.$$

Let

$$\mathcal{E}_0^{ind} = \{R_{11}, R_{12}\} \cup \{R_{21}, R_{22}\} \cup \dots \cup \{R_{m1}, R_{m2}\},\$$

and

$$\mathcal{E}_0^{dep} = \{R_{(m+1)1}, R_{(m+1)2}\} \cup \{R_{(m+2)1}, R_{(m+2)2}\} \cup \dots \cup \{R_{n1}, R_{n2}\}.$$

By Lemma 2.5.4, the set \mathcal{E}_0 is extremal. Let \mathcal{E} be a maximal extremal set of extremal rays which contains \mathcal{E}_0 . Let $\mathcal{E}_1 = \mathcal{E} - \mathcal{E}_0$. By Proposition 2.3.8, $\sharp \mathcal{E}_1 = \rho(X) - 1 - \dim[\mathcal{E}_0]$. By Theorem 2.3.3, for $S \in \mathcal{E}_1$, the divisor D(S) has no a common point with divisors $D(R_{i1})$, $1 \leq i \leq n$.

Lemma 2.5.9. Let X satisfies the conditions of Theorem 2.5.8. Let Q be an extremal ray such that Q is different from extremal rays R_{ij} , $1 \le i \le n$, $1 \le j \le 2$, and the set $\mathcal{E}_1 \cup \{Q\}$ is extremal. Then the Q has the type (II) and there exists exactly one i such that $1 \le i \le n$ and $Q \cdot D(R_{i1}) > 0$ and $D(Q) \cap D(R_{j1}) = \emptyset$ if $j \ne i$.

Proof. Assume that Q has the type (I). Then the divisor D(Q) has no a common point with the divisors $D(R_{i1})$, $1 \leq i \leq n$. By Theorems 2.3.3, 2.3.6 and Lemma 2.5.1, the set $\{Q\} \cup \mathcal{E}_1 \cup \mathcal{E}_0$ is extremal. We then get a contradiction with the condition that $\mathcal{E}_1 \cup \mathcal{E}_0$ is a maximal extremal set. Thus, the extremal ray Q has the type (II).

If D(Q) has no a common point with divisors $D(R_{i1})$, $1 \leq i \leq n$, we get a contradiction by the same way. Thus, there exists *i* such that $1 \leq i \leq n$ and $D(Q) \cap D(R_{i1}) \neq \emptyset$. Let us consider a projectivisation $P\overline{NE}(X)$. By Lemma 2.2.2, $P\overline{NE}(X, D(Q))$ is an interval with two ends. Its first end is the vertex PQ and its second end is a point of the edge $P(R_{i1} + R_{i2})$ of the convex polyhedron $P\overline{NE}(X)$. Thus, the *i* is defined by the extremal ray Q. Evidently, $Q \cdot D(R_{i1}) > 0$.

Lemma 2.5.10. With conditions of Lemma 2.5.9 above, assume that $m + 1 \le i \le n$. Then there exists exactly one extremal ray $Q = Q_i$ with conditions of Lemma 2.5.9: thus, the set $\mathcal{E}_1 \cup \{Q_i\}$ is extremal and $Q_i \cdot D(R_{i1}) > 0$, and $D(Q_i) \cap D(R_{j1}) = \emptyset$ if $j \ne i$.

Proof. The

$$\beta = \sum_{S \in \mathcal{E}_1} S + \sum_{R \in \mathcal{E}_0} R$$

is a face of $\overline{N}E(X)$ of highest dimension $\rho(X) - 1$, and

$$\beta_i = \sum_{S \in \mathcal{E}_1} S + \sum_{R \in \mathcal{E}_0 - \{R_{i1}, R_{i2}\}} R$$

is a face $\beta_i \subset \beta \subset \overline{NE}(X)$ of the dimension $\rho(X) - 2$ and of the codimension one in β . (Here we use that $m + 1 \leq i \leq m + k$). It follows that there exists exactly one face β'_i of $\overline{NE}(X)$ such that β'_i contains β_i , dim $\beta'_i = \rho(X) - 1$, and $\beta'_i \neq \beta$. By Theorems 2.3.3 and 2.3.6, and Lemma 2.5.9, $\beta'_i = \beta_i + Q_i$ where Q_i is an extremal ray such that the set $\mathcal{E}_1 \cup \{Q_i\} \cup (\mathcal{E}_0 - \{R_{i1}, R_{i2}\})$ is extremal, and the ray Q_i has properties of Lemma 2.5.10. It follows that the Q_i is unique and does exist.

Lemma 2.5.11. In notation above, the set $\mathcal{E}_1 \cup \mathcal{E}_0^{ind} \cup \{Q_{m+1}, ..., Q_n\}$ is extremal.

Proof. We apply Lemma 2.5.7 to $\mathcal{E} = \mathcal{E}_1 \cup \mathcal{E}_0^{ind}$ and \mathcal{E}_0^{dep} . By Lemma 2.5.7, there are extremal rays $Q'_{m+1}, \dots, Q'_{m+r}$ such that the set $\mathcal{E}_1 \cup \mathcal{E}_0^{ind} \cup \{Q'_{m+1}, \dots, Q'_{m+r}\}$ is extremal and for any $i, m+1 \leq i \leq m+r$, there exists $j, m+1 \leq j \leq n$, such that $Q'_i \cdot D(R_{j1}) > 0$. Moreover, for any $j, m+1 \leq j \leq n$, there exists an extremal ray $Q_i, m+1 \leq i \leq m+r$, such that

$$Q_i' \cdot D(R_{j1}) > 0.$$

By Lemmas 2.5.9 and 2.5.10, r = k and $\mathcal{E}_1 \cup \mathcal{E}_0^{ind} \cup \{Q'_{m+1}, \dots, Q'_{m+r}\} = \mathcal{E}_1 \cup \mathcal{E}_0^{ind} \cup \{Q_{m+1}, \dots, Q_n\}.$

Lemma 2.5.12. The set \mathcal{E}_0^{dep} is empty.

Proof. By Lemmas 2.5.9, 2.5.10 and 2.5.11, the set of extremal rays $U = \mathcal{E}_1 \cup \mathcal{E}_0^{ind} \cup \{Q_{m+1}, ..., Q_n\}$ is a maximal extremal set which contains $\mathcal{E}_1 \cup \mathcal{E}_0^{ind}$ and does not contain extremal rays from \mathcal{E}_0^{dep} . Assume that $k = n - m \neq 0$. Then $k \geq 2$ and dim $U = \rho_0(X) - 1 + 2m + k$. But the dimension of a face of $\overline{NE}(X)$ of highest dimension is equal to $\rho(X) - 1 = \rho_0(X) - 1 + 2m + k + \delta$ where $\delta \geq 1$. Thus, the extremal set U is not maximal, and there exists another extremal ray S such that $U \cup \{S\}$ is extremal. By definition of U, the $S \in \mathcal{E}_0^{dep}$. Let $S = R_{i1}$ where $m + 1 \leq i \leq n$. Since $Q_i \cdot D(R_{i1}) > 0$, by Theorem 2.3.3, the extremal set $\{Q_i, R_{i1}\}$ has the type \mathfrak{C}_2 . Thus, $R_{i1} \cdot D(Q_i) = 0$. By definition of the set \mathcal{E}_0^{dep} , there exists a linear dependence $\sum_{l=m+1}^{l=n} a_{l1}R_{l1} + a_{l2}R_{l2} = 0$ where $a_{i1} \neq 0$ and $a_{i2} \neq 0$. Multiplying the equality above on $D(Q_i)$, we get $a_{i2} = 0$. Thus, we get a contradiction. (Compair with Lemma 2.5.6.)

Lemma 2.5.13. The set \mathcal{E}_0^{ind} is empty.

Proof. Since $\mathcal{E}_0^{dep} = \emptyset$, the set $U = \mathcal{E}_1 \cup \mathcal{E}_0^{ind} = \mathcal{E}_1 \cup \{R_{11}, R_{12}\} \cup ... \cup \{R_{m1}, R_{m2}\}$ is a maximal extremal set. It follows that U generates a simplicial face of $\overline{NE}(X)$ of codimension 1. Thus, $U_1 = \mathcal{E}_1 \cup \mathcal{E}_0^{ind} - \{R_{m2}\} = \mathcal{E}_1 \cup \{R_{11}, R_{12}\} \cup ... \cup \{R_{(m-1)1}, R_{(m-1)2}\} \cup \{R_{m1}\}$ generates a simplicial face of $\overline{NE}(X)$ of codimension 2. It follows that there exists an extremal ray Q_{m2} such that $U'_1 = \mathcal{E}_1 \cup \{R_{11}, R_{12}\} \cup ... \cup \{R_{(m-1)1}, R_{(m-1)2}\} \cup \{R_{m1}\} \cup \{Q_{m2}\}$ generates a simplicial face of $\overline{NE}(X)$ of codimension 1, and Q_{m2} is different from R_{m2} . By Lemma 2.5.9, $Q_{m2} \cdot D(R_{m1}) > 0$. Thus, by Theorem 2.3.3, $\{Q_{m2}, R_{m1}\}$ is an extremal set of the type \mathfrak{C}_2 where $R_{m1} \cdot D(Q_{m2}) = 0$.

Similarly, we can find an extremal ray Q_{m1} such that the set $\{Q_{m1}, R_{m2}\}$ is extremal of the type \mathfrak{C}_2 where $R_{m2} \cdot D(Q_{m1}) = 0$. Then we get a contradiction with Lemma 2.5.6. Thus, m = 0, and the set $\mathcal{E}_0^{ind} = \emptyset$.

Thus, we proved that X does not have a pair of extremal rays of the type \mathfrak{B}_2 . By Theorem 2.3.3 and Proposition 2.3.8, the Mori polyhedron $\overline{NE}(X)$ is then simplicial. Thus, we have proven the statement (1).

Now let us prove (2): X does not have more than one extremal ray of the type (I). We need some definitions.

Let P be a set of divisorial extremal rays. We say that P is divisorially connected if there is no decomposition $P = P_1 \cup P_2$ such that both P_1 and P_2 are non-empty and for any $R \in P_1$ and any $Q \in P_2$ divisors D(R) and D(Q) do not have a common point. It defines divisorially connected components of a set of extremal rays (we had used this definition before, see definition before Theorem 2.3.3). For example, for our case, a set of extremal rays is extremal iff its divisorially connected components have the type $\mathfrak{A}_1, \mathfrak{B}_2, \mathfrak{C}_m$ or \mathfrak{D}_2 . Also, we can say what does it mean that two sets P_1 and P_2 of extremal rays are divisorially joint: this means that there exist extremal rays $Q_1 \in P_1$ and $Q_2 \in P_2$ such that divisors $D(Q_1)$ and $D(Q_2)$ have a common point (in particular, this divisors or even extremal rays Q_1, Q_2 may coincide).

We say that a set P of divisorial extremal rays is single arrows connected if for any two diffent extremal rays $R_1, R_2 \in P$ there exists an oriented path in the graph G(P) with the beginning in R_1 and terminal in R_2 . This defines single arrows connected components of P. Like above, we can say what does it mean that two sets P_1 and P_2 of divisorial extremal rays are single arrows joint: either they have a common extremal ray, or there exist extremal rays $Q_1, Q'_1 \in P_1$ and $Q_2, Q'_2 \in P_2$ such that there are an oriented path joining Q_1 and Q_2 and an oriented path joining Q'_2 and Q'_1 in the set $P_1 \cup P_2$. For example, we can reformulate Lemma 1.1: for our situation, an E-set \mathcal{L} is single arrows connected, two E-sets \mathcal{L} and \mathcal{M} are single arrows joint (using Lemma 2.5.1, Theorem 2.4.1 and the statement (1) we have proven).

By Lemma 2.2.2, divisors of different extremal rays of the type (I) do not have a common point. By Theorem 2.3.6, any set of extremal rays of the type (I) generates a simplicial face of $\overline{NE}(X)$ of Kodaira dimension 3. It follows that the set of extremal rays of the type (I) is finite. Let

$$\{R_1, ..., R_s\}$$

be the hole set of divisorial extremal rays of the type (I) on X. We should prove that $a \leq 1$

ON THE PICARD NUMBER OF FANO 3-FOLDS WITH TERMINAL SINGULARITIES29

Let \mathcal{E} be a maximal extremal set of extremal rays on X containing $\{R_1, ..., R_s\}$. Let $T_1, ..., T_s$ are divisorially connected components of \mathcal{E} containing $R_1, ..., R_s$ respectively. By Theorem 2.3.3, these connected components are different. The connected component T_i either contains one ray R_i (has the type \mathfrak{A}_1) or contains two rays: the ray R_i and another ray of the type (II). Then this connected component has the type \mathfrak{D}_2 . Let

$$\mathcal{E}_0 = \mathcal{E} - (T_1 \cup \dots \cup T_s).$$

Thus, extremal sets $\mathcal{E}_0, T_1, ..., T_s$ are divisorially disjoint.

By [Ka1] and [Sh], any face of $\overline{NE}(X)$ is contractible, and by our condition it has Kodaira dimension 3. By Proposition 2.2.6, for any $1 \le i \le s$, there exists an effective divisor $D(T_i)$ which is a linear combination of divisors of rays from T_i with positive coefficients, such that $R \cdot D(T_i) < 0$ for any $R \in T_i$.

Using the divisors $D(T_i)$, similarly to Lemma 2.5.7, we can find extremal rays

$$\{Q_1, \dots, Q_r\}$$

with properties:

(a) $r \leq s$;

(b) For any $i, 1 \leq i \leq r$, there exists $j, 1 \leq j \leq t$, such that $Q_i \cdot D(T_j) > 0$ (in particular, Q_i is different from extremal rays of \mathcal{E} and do not have the type (I));

(c) For any $j, 1 \leq j \leq s$, there exists an extremal ray $Q_i, 1 \leq i \leq r$, such that

$$Q_i \cdot D(T_j) > 0;$$

(d) The set $\mathcal{E}_0 \cup \{Q_1, ..., Q_r\}$ of extremal rays is extremal, and extremal rays $\{Q_1, ..., Q_r\}$ are linearly independent.

By our condition, all extremal rays on X are divisorial. Thus, by (b), the extremal rays $Q_1, ..., Q_r$ have the type (II).

Let us take a ray Q_i and let $Q_i \cdot D(T_j) > 0$. By Theorem 2.3.3, the T_j generates a simplicial face γ_j of $\overline{NE}(X)$. Since T_j has the type \mathfrak{A}_1 or \mathfrak{D}_2 , one can see easily using Lemma 2.2.3, that any curve of divisors of rays from T_j belongs to this face. It follows that $\overline{NE}(X, D(Q_i))$ is a 2-dimensional angle bounded by the ray Q_i and a ray from the face γ_j since the divisor $D(Q_i)$ evidently has a common curve with one of divisors D(R) of $R \in T_j$. Since any two sets from T_1, \ldots, T_s do not have a common extremal ray, the faces $\gamma_1, \ldots, \gamma_s$ do not have a common ray (not necessarily extremal). It follows that the angle $\overline{NE}(X, D(Q_i))$ does not have a common ray with the face γ_k for $k \neq j$. Thus, the divisor $D(Q_i)$ do not have a common point with divisors of rays T_k . It follows that r = s and we can choose numeration $Q_1, \ldots Q_s$ such that $Q_i \cdot D(T_i) > 0$ but $D(Q_i)$ do not have a common point with divisors of extremal rays T_j if $i \neq j$ (the Q_i is divisorially disjoint with T_j if $i \neq j$).

Let us fix $i, 1 \leq i \leq n$. The set $\mathcal{E} \cup \{Q_i\}$ is not extremal since \mathcal{E} is maximal extremal. Since Q_i is divisorially disjoint with divisorially connected components T_j of \mathcal{E} if $j \neq i$, it follows that the set $\mathcal{E}_0 \cup T_i \cup \{Q_i\}$ is not extremal also. Then, the last set contains an E-set \mathcal{L}_i . Since $\mathcal{E}_0 \cup T_i$ is extremal, the set \mathcal{L}_i contains the ray Q_i . The \mathcal{L}_i is single arrows connected (we have mentioned this above). Let $\mathcal{E}_{0i} = \mathcal{L}_i \cap \mathcal{E}_0$. We claim that the subset

$$U \in \mathcal{C} \cup \{\mathcal{O}\}$$

of \mathcal{L}_i is single arrows connected too. Really, the set $\mathcal{L}'_i = \mathcal{L}_i - (\mathcal{E}_{0i} \cup \{Q_i\})$ is not empty since the set $\mathcal{E}_{0i} \cup \{Q_i\}$ is extremal as a subset of the extremal set $\mathcal{E}_0 \cup \{Q_1, ..., Q_s\}$. Since \mathcal{L}_i is single arrows connected, there exists a shortest oriented path in \mathcal{L}_i connecting any $S \in \mathcal{E}_{0i}$ with \mathcal{L}'_i . But \mathcal{E}_{0i} and \mathcal{L}'_i are divisorially (and then single arrows) disjoint. Thus, the beginning of this path is a path in $\mathcal{E}_{0i} \cup \{Q_i\}$ joining S with the ray Q_i . The same considerations prove that there exists an oriented path in $\mathcal{E}_{0i} \cup \{Q_i\}$ joining Q_i with S.

By Lemma 1.1, for $i \neq j$, the sets \mathcal{L}_i and \mathcal{L}_j are single arrows joint. Using the fact that \mathcal{L}'_i and \mathcal{L}'_j are divisorially disjoint, one similarly can prove that the sets $U_i = \mathcal{E}_{0i} \cup \{Q_i\}$ and $U_j = \mathcal{E}_{0j} \cup \{Q_j\}$ are single arrows joint.

Let U be the union of all sets $U_i = \mathcal{E}_{0i} \cup \{Q_i\}, i = 1, ..., s$. Then U is single arrows connected extremal set of extremal rays of the type (II). By Theorem 2.3.3, it is possible only if U is either empty or has the type \mathfrak{C}_1 . It follows that $s \leq 1$. This proves the statement (2).

Let us prove (3). We use the following

Statement. The contraction of a ray R of the type (II) on X gives a Fano 3-fold X' with terminal Q-factorial singularities and without small extremal rays and without faces of Kodaira dimension 1 or 2 for $\overline{NE}(X')$. Extremal sets \mathcal{E}' on X' are in one to one correspondence with extremal sets \mathcal{E} on X which contain the ray R.

Proof. Let $\sigma : X \to X'$ be a contraction of R. The X' has terminal Q-factorial singularities by [Ka1] and [Sh]. We have, $K_X = \sigma^*(K_{X'}) + dD(R)$. Multiplying this equality on R and using Proposition 2.3.2, we get that d = 1. By the statement (1), it follows that $\sigma^*(-K_{X'}) = -K_X + D(R)$ is nef and only contracts the extremal ray R. Then $-K_{X'}$ is ample on X' and X' is a Fano 3-fold with terminal Q-factorial singularities. Faces of $\overline{NE}(X')$ are in one to one correspondence with faces of $\overline{NE}(X)$ which contain the R. Contractions of faces of $\overline{NE}(X')$ are dominated by contractions of the corresponding faces of $\overline{NE}(X)$. It follows the last statement.

Let $\mathcal{E} = \{R_1, ..., R_k\}$ be an extremal set on X. By Theorem 2.3.3, it has connected components of the type $\mathfrak{A}_1, \mathfrak{B}_2, \mathfrak{C}_m$ or \mathfrak{D}_2 . Moreover, by (1) and (2), it does not have a connected component of the type \mathfrak{B}_2 and does not have more than one connected component of the type \mathfrak{A}_1 . By Statement above, the same should be true for the extremal set \mathcal{E}' one gets by the contraction of any extremal ray R_i of the type (II) of \mathcal{E} . It follows the statement (3).

Now we prove (4): $\rho(X) \leq 7$.

First, we show how to prove $\rho(X) \leq 8$ applying Theorem 1.2 to the face $\gamma = \mathcal{M}(X)$ of dim $\mathcal{M}(X) = m = \rho(X) - 1$. By the statement (1) of Theorem 2.5.8 and Theorems 2.3.3 and 2.4.1, the $\mathcal{M}(X)$ is simple and all conditions of Theorem 1.2 are valid for some constants d, C_1, C_2 . By Theorem 2.4.1, we can take d = 2. By the proof of Theorem 1.2, we should find the constants C_1 and C_2 for maximal extremal sets \mathcal{E} only (only this sets we really use). Thus, $\sharp \mathcal{E} = m$. By the statement (3), then the constants $C_1 \leq 2/m$ and $C_2 = 0$. Thus, we get m < (16/3)2/m + 6. Then, $m = \rho(X) - 1 \leq 7$, and $\rho(X) \leq 8$.

To prove the better inequality $\rho(X) \leq 7$, we should analyze the proof of Theorem 1.2 for our case more carefully. We will show that the conditions of Lemma 1.4 hold for the $\mathcal{M}(X)$ with the constants C = 0 and D = 2/3. By Lemma 1.4, we then get

ON THE PICARD NUMBER OF FANO 3-FOLDS WITH TERMINAL SINGULARITIES31

Like for the proof of Theorem 1.2, we introduce a weight of an oriented angle, but using a new formula: $\sigma(\angle) = 2/3$ if $\rho(R_1(\angle), R_2(\angle)) = 1$, and $\sigma(\angle) = 0$ otherwise.

By the statement (3) of Theorem 2.5.8, the condition (1) of Lemma 1.4 holds with constants C = 0 and D = 2/3.

Let us prove the condition (2) of Lemma 1.4. For k = 3 (triangle) it is true since an *E*-set which has at least 3 elements has the type (a) of Theorem 2.4.1 (see the proof of Theorem 1.2). Thus, the triangle has at least three oriented angles with the weight 2/3. For k = 4 (quadrangle), we proved (when we were proving Theorem 1.2) that one can find at least two oriented angles of the quadrangle such that for any of them $\rho(R_1(\angle), R_2(\angle))$ is finite. By the statement (3) of Theorem 2.5.8, then $\rho(R_1(\angle), R_2(\angle)) = 1$. Thus, the quadrangle has at least two oriented angles of the weight 2/3. This finishes the proof of Theorem 2.5.8.

Now, we give an application of the statement (2) of Theorem 2.5.8 to geometry of Fano 3-folds.

Let us consider a Fano 3-fold X and blow-ups X_p in different non-singular points $\{x_1, ..., x_p\}$ of X. We say that this is a Fano blow-up if X_p is Fano. We have the following very simple

Proposition 2.5.14. Let X be a Fano 3-fold with terminal \mathbb{Q} -factorial singularities and without small extremal rays. Let X_p be a Fano blow up of X.

Then for any small extremal ray S on X_p , the S has a non-empty intersection with one of exceptional divisors $E_1, ..., E_p$ of this blow up and does not belong to any of them. Moreover, the exceptional divisors $E_1, ..., E_p$ define p extremal rays $Q_1, ..., Q_p$ of the type (I) on X_p such that $E_i = D(Q_i)$.

Proof. The last statement is clear. Let S be a small extremal ray on X_p which does not intersect divisors $E_1, ..., E_p$. Let H be a general nef element orthogonal to S. Let $l_1, ..., l_n$ are lines which generate extremal rays $Q_1, ..., Q_p$. Then the divisor $H' = H + (l_1 \cdot H)/(-l_1 \cdot E_1)E_1 + ... + (l_p \cdot H)/(-l_p \cdot E_p)E_p$ is a nef divisor on X_p orthogonal to all extremal rays $Q_1, ..., Q_p, S$, and $(H')^3 > H^3 > 0$. This proves that the extremal rays $Q_1, ..., Q_p, S$ generate a face of $\overline{NE}(X_p)$ of Kodaira dimension 3. Then, by contraction of extremal rays $Q_1, ..., Q_p$, the image of S gives a small extremal ray on X.

This gives a contradiction.

It is known that a contraction of a face of Kodaira dimension 1 or 2 of $\overline{NE}(Y)$ of a Fano 3-fold Y has a general fiber which is a rational surface or curve respectively, because this contraction has relatively negative canonical class. See [Ka1], [Sh]. It is known that a small extremal ray is rational [Mo2].

Then, using the statement (2) of Theorem 2.5.8 and Proposition 2.5.14, we can divide Fano 3-folds of Theorem 2.5.8 on the following 3 classes:

Corollary 2.5.15. Let X be a Fano 3-fold with terminal Q-factorial singularities and without small extremal rays, and without faces of Kodaira dimension 1 or 2 for Mori polyhedron. Let ϵ be the number of extremal rays of the type (I) on X (by Theorem 2.5.8, the $\epsilon < 1$).

Then there exists $p, 1 \le p \le 2 - \epsilon$, such that X belongs to one of classes (A), (B) or (C) below:

(A) There exists a Fano blow-up X_p of X with a face of Kodaira dimension 1 or 2. Thus, birationally, X is a fibration on rational surfaces over a curve or rational surfaces.

VIACHESLAV V. NIKULIN

(B) There exist Fano blow-ups X_p of X for general p points on X such that for all these blow-ups the X_p has a small extremal ray S. Then images of curves of S on X give a system of rational curves on X which cover a Zariski open subset of X.

(C) There do not exist Fano blow-ups X_p of X for general p points.

We remark that for Fano 3-folds with Picard number 1 the $\epsilon = 0$. Thus, $1 \le p \le 2$.

We mention that the statements (3) and (4) of Theorem 2.5.8 give similar information for blow ups of X at curves. Of course, it is more difficult to formulate these statements.

ON THE PICARD NUMBER OF FANO 3-FOLDS WITH TERMINAL SINGULARITIES33

References

- [A] V. A. Alekseev, Fractional indices of log del Pezzo surfaces, Izv. Akad. Nauk SSSR Ser. Mat. 52 (1988), no. 6, 1288 – 1304; English transl. in Math. USSR Izv. 33 (1989).
- [A-N1] V. A. Alekseev and V. V. Nikulin, The classification of Del Pezzo surfaces with logterminal singularities of the index ≤ 2, involutions of K3 surfaces and reflection groups in Lobachevsky spaces, Russian, Doklady po matematike i prilogeniyam, MIAN 2 (1988), no. 2, 51–151.
- [A-N2] _____, The classification of Del Pezzo surfaces with log-terminal singularities of the index ≤ 2 and involutions of K3 surfaces, Dokl. AN SSSR **306** (1989), no. 3, 525 528; English transl. in Soviet Math. Dokl. **39** (1989).
- [G–H] P. Griffiths and J. Harris, *Principles of algebraic geometry*, A. Wiley Interscience, New York, 1978.
- [Ha] R. Hartshorne, Algebraic geometry, Springer, 1977.
- Sh. Ishii, Quasi-Gorenstein Fano 3-folds with isolated non-rational loci, Compositio Math. 77 (1991), 335-341.
- [Ka1] Yu. Kawamata, The cone of curves of algebraic varieties, Ann. of Math. 119 (1984), no. 2, 603–633.
- [Ka2] Yu. Kawamata, Boundness of Q-Fano threefolds, Preprint (1989).
- [Kh] A. G. Khovanskii, Hyperplane sections of polyhedra, toric varieties and discrete groups in Lobachevsky space, Functional Anal. i Prilozhen 20 (1986), no. 1, 50–61; English transl. in Functional Anal. Appl. 20 (1986), no. 1.
- [Ma] K. Matsuki, Weyl groups and birational transformations among minimal models, Preprint (1993).
- [Mo1] S. Mori, *Threefols whose canonical bundles are not numerically effective*, Ann. of Math. **116** (1982), no. 2, 133–176.
- [Mo2] _____, Flip theorem and the existence of minimal models for 3-folds, Journal of Amer. Math. Society 1 (1988), no. 1, 117–253.
- [Mo–Mu] S. Mori and S. Mukai, On Fano 3-folds with $B_2 \ge 2$, Algebraic varieties and Analytic value value (Tokyo, 1981), Studies in Pure and Math., vol. 1, Kinikuniya, Tokyo, 1983, pp. 101–129.
- [N1] V. V. Nikulin, On the classification of arithmetic groups generated by reflections in Lobachevsky spaces, Izv. Akad. Nauk SSSR Ser. Mat. 45 (1981), no. 1, 113 – 142; English transl. in Math. USSR Izv. 18 (1982).
- [N2] _____, Del Pezzo surfaces with log-terminal singularities, Mat. Sbor. 180 (1989), no. 2, 226 - 243; English transl. in Math. USSR Sb. 66 (1990).
- [N3] _____, Del Pezzo surfaces with log-terminal singularities. II, Izv. Akad. Nauk SSSR Ser. Mat. **52** (1988), no. 5, 1032 – 1050; English transl. in 'Math. USSR Izv. **33** (1989).
- [N4] _____, Del Pezzo surfaces with log-terminal singularities. III, Izv. Akad. Nauk SSSR Ser. Mat. **53** (1989), no. 6, 1316 – 1334; English transl. in Math. USSR Izv. **35** (1990).
- [N5] _____, Del Pezzo surfaces with log-terminal singularities and nef anticanonical class and reflection groups in Lobachevsky spaces, Preprint Max-Planck-Institut für Mathematik Bonn **89-28** (1988).
- [N6] _____, Algebraic 3-folds and diagram method, Math. USSR Izv. **37** (1991), no. 1, 157-189.
- [N7] _____, Algebraic 3-folds and diagram method. II, Preprint Max-Planck-Institut für Mathematik Bonn **90-104** (1990).
- [N8] _____, On the Picard number of Fano 3-folds with terminal singularities, Preprint Math. Sci. Res. Institute Berkeley **057-93** (1993).
- [N9] _____, A lecture on Kähler cone and Picard number of Calabi-Yau 3-folds, Preprint Math. Sci. Res. Institute Berkeley **061-93** (1993).
- M. N. Prokhorov, The absence of discrete reflection groups with non-compact fundamental polyhedron of finite volume in Lobachevsky spaces of large dimension, Izv. Akad. Nauk SSSR Ser. Mat. 50 (1986), no. 2, 413 – 424; English transl. in Math. USSR Izv. 28 (1987).
- [Sh] V. V. Shokurov, *The nonvanishing theorem*, Izv. Akad. Nauk SSSR Ser. Mat. **49** (1985),

VIACHESLAV V. NIKULIN

[V] E. B. Vinberg, The absence of crystallographic reflection groups in Lobachevsky spaces of large dimension, Trudy Moscow. Mat. Obshch. 47 (1984), 68 – 102; English transl. in Trans. Moscow Math. Soc. 47 (1985).

STEKLOV MATHEMATICAL INSTITUTE, UL. VAVILOVA 42, MOSCOW 117966, GSP-1, RUSSIA. E-mail address: slava@nikulin.mian.su

34