

IMAGES OF ℓ -ADIC REPRESENTATIONS AND AUTOMORPHISMS OF ABELIAN VARIETIES

A. SILVERBERG AND YU. G. ZARHIN

1. INTRODUCTION

Suppose that F is either a global field or a finitely generated extension of \mathbf{Q} , A is an abelian variety over F , ℓ is a prime number, and $\ell \neq \text{char}(F)$. Let $\mathfrak{G}_\ell(F, A)$ denote the algebraic envelope of the image of the absolute Galois group of F under the ℓ -adic representation associated to A , and let $\mathfrak{G}_\ell(F, A)^0$ denote its identity connected component. In §3 we prove that the intersection of $\mathfrak{G}_\ell(F, A)^0(\mathbf{Q}_\ell)$ with the torsion subgroup of the center of $\text{End}(A) \otimes \mathbf{Q}$ is independent of ℓ . In the case where F is a finitely generated extension of \mathbf{Q} , this would follow from the Mumford-Tate Conjecture. Our results do not assume the Mumford-Tate Conjecture, and apply even in the positive characteristic case, where there is no analogue of the Mumford-Tate Conjecture. The result in the characteristic zero case can therefore be viewed as providing evidence in the direction of the Mumford-Tate conjecture.

Let $F_{\Phi, \ell}(A)$ be the smallest extension F' of F such that $\mathfrak{G}_\ell(F', A)$ is connected. We call this extension the ℓ -connectedness extension, or connectedness extension.

The algebraic group $\mathfrak{G}_\ell(F, A)$ and the field $F_{\Phi, \ell}(A)$ were introduced by Serre ([13], [14], [16]), who proved that if F is a global field or a finitely generated extension of \mathbf{Q} , then $F_{\Phi, \ell}(A)$ is independent of ℓ (see [13], [14], Proposition 1.1 of [6], and Proposition 6.14 of [5]). In such cases, we will denote the field $F_{\Phi, \ell}(A)$ by $F_\Phi(A)$. Our results rely heavily on the ℓ -independence results of Serre and their generalizations [5].

Let $F(\text{End}(A))$ denote the smallest extension of F over which all the endomorphisms of A are defined. Then (see Proposition 2.10 of [17]),

$$F(\text{End}(A)) \subseteq F_{\Phi, \ell}(A).$$

By enlarging the ground field, we may assume that $F = F(\text{End}(A)) = F_{\Phi, \ell}(A)$. We then consider twists of the ℓ -adic representations associated to A . The results of this paper follow from the ℓ -independence of the connectedness extensions associated to these twists.

See [18] for a study of the connectedness extensions attached to twists of abelian varieties. See [17] for conditions for the connectedness of $\mathfrak{G}_\ell(F, A)$.

2. DEFINITIONS, NOTATION, AND LEMMAS

Let \mathbf{Z} , \mathbf{Q} , and \mathbf{C} denote respectively the integers, rational numbers, and complex numbers. If G is an algebraic group, let G^0 denote the identity connected component. If F is a field, let F^s denote a separable closure of F and let \bar{F} denote an algebraic closure of F . If A is an abelian variety over a field F , write $\text{End}_F(A)$ for the endomorphisms of A which are defined over F , write $\text{Aut}_F(A)$ for the automorphisms of A defined over F , let $\text{End}(A) = \text{End}_{F^s}(A)$, let $\text{End}^0(A) = \text{End}(A) \otimes_{\mathbf{Z}} \mathbf{Q}$,

and let $\text{End}_F^0(A) = \text{End}_F(A) \otimes_{\mathbf{Z}} \mathbf{Q}$. If ℓ is a prime number and $\ell \neq \text{char}(F)$, let $T_\ell(A) = \varprojlim A_{\ell^r}$ (the Tate module), let $V_\ell(A) = T_\ell(A) \otimes_{\mathbf{Z}} \mathbf{Q}_\ell$, and let $\rho_{A,\ell}$ denote the ℓ -adic representation

$$\rho_{A,\ell} : \text{Gal}(F^s/F) \rightarrow \text{Aut}(T_\ell(A)) \subseteq \text{Aut}(V_\ell(A)).$$

Let $\mathfrak{G}_\ell(F, A)$ denote the algebraic envelope of the image of $\rho_{A,\ell}$, i.e., the Zariski closure in $\text{Aut}(V_\ell(A))$ of the image of $\text{Gal}(F^s/L)$ under $\rho_{A,\ell}$. Let $F_{\Phi,\ell}(A)$ be the smallest extension F' of F in F^s such that $\mathfrak{G}_\ell(F', A)$ is connected.

Lemma 2.1 (Lemma 2.7 of [17]). *If A is an abelian variety over a field F , L is a finite extension of F in F^s , and ℓ is a prime number, then*

$$\mathfrak{G}_\ell(L, A) \subseteq \mathfrak{G}_\ell(F, A) \text{ and } \mathfrak{G}_\ell(L, A)^0 = \mathfrak{G}_\ell(F, A)^0.$$

In particular, if $\mathfrak{G}_\ell(F, A)$ is connected, then $\mathfrak{G}_\ell(F, A) = \mathfrak{G}_\ell(L, A)$.

Lemma 2.2. *Suppose A and B are abelian varieties over a field F , L is a finite extension of F in F^s , ℓ is a prime number, $\ell \neq \text{char}(F)$, $\mathfrak{G}_\ell(F, A)$ is connected, and A and B are isomorphic over L . Then:*

- (i) $\mathfrak{G}_\ell(F, B)^0 = \mathfrak{G}_\ell(F, A)$, and
- (ii) $\mathfrak{G}_\ell(L, B)$ is connected, i.e., $F_{\Phi,\ell}(B) \subseteq L$.

Proof. Since A and B are isomorphic over L , and $\mathfrak{G}_\ell(F, A)$ is connected, we have

$$\begin{aligned} \mathfrak{G}_\ell(L, B) &= \mathfrak{G}_\ell(L, A) = \mathfrak{G}_\ell(F, A) = \mathfrak{G}_\ell(F, A)^0 \\ &= \mathfrak{G}_\ell(L, A)^0 = \mathfrak{G}_\ell(L, B)^0 = \mathfrak{G}_\ell(F, B)^0, \end{aligned}$$

using Lemma 2.1. The result follows. \square

Lemma 2.3. *Suppose A is an abelian variety over a global field F and*

$$c : \text{Gal}(F^s/F) \rightarrow \text{End}_F^0(A)^\times$$

is a continuous homomorphism of finite order. For each prime $\ell \neq \text{char}(F)$ let

$$\rho_{\ell,c} : \text{Gal}(F^s/F) \rightarrow \text{Aut}(V_\ell(A)), \quad \sigma \mapsto c(\sigma)\rho_{A,\ell}(\sigma)$$

be the twist of $\rho_{A,\ell}$. Then $\{\rho_{\ell,c}\}$ constitutes a strictly compatible system of integral ℓ -adic representations of $\text{Gal}(F^s/F)$. More precisely, suppose M is a finite Galois extension of F such that c factors through $\text{Gal}(M/F)$. Let S_ℓ be the set of finite places v of F such that either A has bad reduction at v , v is ramified in M/F , or the residue characteristic of v is ℓ . Let v be a finite place of F , w a place of F^s lying over v , and κ_v and κ_w the residue fields at v and w , respectively. Let $\tau \in \text{Gal}(F^s/F)$ be an element that acts as the Frobenius automorphism of κ_w/κ_v . Suppose that $v \notin S_\ell$, and let

$$\varphi_w = \rho_{\ell,c}(\tau) \in \text{Aut}(V_\ell(A)).$$

Then $\rho_{\ell,c}$ is unramified at v , the characteristic polynomial $P_v(t)$ of φ_w lies in $\mathbf{Z}[t]$ and does not depend on the choice of w and ℓ , and the roots of $P_v(t)$ all have complex absolute value $\sqrt{\#\kappa_v}$.

Proof. We use that $\{\rho_{A,\ell}\}$ is a strictly compatible system of integral ℓ -adic representations of $\text{Gal}(F^s/F)$ (see [19] and I.2 of [15]). Let A_v be the abelian variety over κ_v which is the reduction of A at v . The choice of w allows us to identify the Tate modules $V_\ell(A)$ and $V_\ell(A_v)$, and this identification is compatible with the natural embedding $\text{End}^0(A) \hookrightarrow \text{End}^0(A_v)$. Let

$$Fr_w = \rho_{A,\ell}(\tau) \in \text{Aut}(V_\ell(A)).$$

Then

$$\varphi_w = c(\tau)Fr_w \in \text{Aut}(V_\ell(A)) \subseteq \text{Aut}(V_\ell(A_v)),$$

and the identification of $\text{Aut}(V_\ell(A))$ with $\text{Aut}(V_\ell(A_v))$ identifies Fr_w with the Frobenius endomorphism of A_v inside $\text{Aut}(V_\ell(A_v))$. It follows from Weil's results on endomorphisms of abelian varieties that $P_v(t)$ has rational coefficients which do not depend on the choice of w and ℓ . If $m = [M : F]$ then $(c(\tau)Fr_w)^m = (Fr_w)^m \in \text{End}(A_v)$ and therefore all roots of $P_v(t)$ are algebraic integers. Therefore, $P_v(t) \in \mathbf{Z}[t]$. Further, Weil's results imply that the eigenvalues of φ_w have absolute value $\sqrt{\#\kappa_v}$. \square

Theorem 2.4. *If F is either a finitely generated extension of \mathbf{Q} or a function field in one variable over a finite field, then every finite abelian group occurs as a Galois group over F .*

Proof. See Theorem 3.12c of [11], and IV.2.1 and IV.1.2 of [7]. \square

Next we define the Mumford-Tate group of a complex abelian variety A (see §2 of [10] or §6 of [22]). If A is a complex abelian variety, let $V = H_1(A(\mathbf{C}), \mathbf{Q})$ and consider the Hodge decomposition $V \otimes \mathbf{C} = H_1(A(\mathbf{C}), \mathbf{C}) = H^{-1,0} \oplus H^{0,-1}$. Define a homomorphism $\mu : \mathbf{G}_m \rightarrow GL(V)$ as follows. For $z \in \mathbf{C}$, let $\mu(z)$ be the automorphism of $V \otimes \mathbf{C}$ which is multiplication by z on $H^{-1,0}$ and is the identity on $H^{0,-1}$.

Definition 2.5. The *Mumford-Tate group* MT_A of A is the smallest algebraic subgroup of $GL(V)$, defined over \mathbf{Q} , which after extension of scalars to \mathbf{C} contains the image of μ .

If A is an abelian variety over a subfield F of \mathbf{C} , we fix an embedding of \bar{F} in \mathbf{C} . This gives an identification of $V_\ell(A)$ with $H_1(A, \mathbf{Q}) \otimes \mathbf{Q}_\ell$, and allows us to view $MT_A \times \mathbf{Q}_\ell$ as a linear \mathbf{Q}_ℓ -algebraic subgroup of $GL(V_\ell(A))$. Let

$$MT_{A,\ell} = MT_A \times_{\mathbf{Q}} \mathbf{Q}_\ell.$$

Then $MT_A(\mathbf{Q}_\ell) = MT_{A,\ell}(\mathbf{Q}_\ell)$. The Mumford-Tate conjecture for abelian varieties (see [12]) may be reformulated as the equality of \mathbf{Q}_ℓ -algebraic groups, $\mathfrak{S}_\ell(F, A)^0 = MT_{A,\ell}$.

Conjecture 2.6 (Mumford-Tate Conjecture). *If A is an abelian variety over a finitely generated extension F of \mathbf{Q} , then $\mathfrak{S}_\ell(F, A)^0 = MT_{A,\ell}$.*

The inclusion $\mathfrak{S}_\ell(F, A)^0 \subseteq MT_{A,\ell}$ was proved by Piatetski-Shapiro [9], Deligne [2], and Borovoi [1].

It is well-known that MT_A contains the homotheties \mathbf{G}_m and that the centralizer of MT_A in $\text{End}(V)$ is $\text{End}^0(A)$. Therefore, the center of $MT_A(\mathbf{Q})$ contains -1 and is contained in the center of $\text{End}^0(A)$.

3. ℓ -INDEPENDENCE

Suppose that F is either a finitely generated extension of \mathbf{Q} or a global field. Suppose $F = F_{\Phi}(A)$, so that $\mathfrak{G}_{\ell}(F, A) = \mathfrak{G}_{\ell}(F, A)^0 = \mathfrak{G}_{\ell}(L, A)$ for all finite extensions L of F . It follows from [3], [4], [20], [21], and VI.5 and XII.2 of [8] that $\mathfrak{G}_{\ell}(F, A)$ is a reductive \mathbf{Q}_{ℓ} -algebraic group, whose centralizer in $\text{End}(V_{\ell}(A))$ is $\text{End}(A) \otimes \mathbf{Q}_{\ell}$. This implies that the center of $\mathfrak{G}_{\ell}(F, A)(\mathbf{Q}_{\ell})$ is contained in $(Z \otimes \mathbf{Q}_{\ell})^{\times}$, where Z is the center of $\text{End}^0(A)$. Let μ_A denote the group of elements of finite order in the center of $\text{End}^0(A)$. In the case where $F \subset \mathbf{C}$, the Mumford-Tate Conjecture (Conjecture 2.6) would imply that $\mu_A \cap \mathfrak{G}_{\ell}(F, A)(\mathbf{Q}_{\ell})$ is the torsion subgroup of the center of $MT_A(\mathbf{Q})$, and therefore is independent of ℓ . In the following two results we prove that $\mu_A \cap \mathfrak{G}_{\ell}(F, A)(\mathbf{Q}_{\ell})$ is independent of ℓ (without assuming the Mumford-Tate Conjecture).

It follows from Weil's results on abelian varieties [19] (as was pointed out by Deligne; see 2.3 of [12]) that $\mathfrak{G}_{\ell}(F, A)$ contains the homotheties \mathbf{G}_m . In particular,

$$-1 \in \mathfrak{G}_{\ell}(F, A)(\mathbf{Q}_{\ell}).$$

Theorem 3.1. *Suppose A is an abelian variety over a finitely generated extension F of \mathbf{Q} , and $F = F_{\Phi}(A)$. Let μ_A denote the group of elements of finite order in the center of $\text{End}^0(A)$. Then $\mu_A \cap \mathfrak{G}_{\ell}(F, A)(\mathbf{Q}_{\ell})$ is independent of the prime ℓ .*

Proof. Over \mathbf{C} , we can view A as \mathbf{C}^d/L with L a lattice in \mathbf{C}^d . Then $L' = \sum_{\gamma \in \mu_A} \gamma(L)$ is a μ_A -invariant lattice in \mathbf{C}^d that contains L as a subgroup of finite index. The complex abelian variety \mathbf{C}^d/L' has a model A' defined over a finite extension F' of F such that A and A' are F' -isogenous and μ_A coincides with the set of elements of finite order in the center of $\text{End}(A')$. Since $\mathfrak{G}_{\ell}(F', A') = \mathfrak{G}_{\ell}(F', A) = \mathfrak{G}_{\ell}(F, A)$, we may assume without loss of generality that μ_A coincides with the set of elements of finite order in the center of $\text{End}(A)$. By Theorem 2.4, we can choose an abelian extension M of F such that $\text{Gal}(M/F)$ is isomorphic to μ_A . Let

$$\chi : \text{Gal}(M/F) \rightarrow \mu_A$$

be an isomorphism, let $c : \text{Gal}(F^s/F) \rightarrow \mu_A$ be the composition of χ with the projection $\text{Gal}(F^s/F) \rightarrow \text{Gal}(M/F)$, and let B denote the twist of A by the cocycle induced by c . By Lemma 2.2i, $\mathfrak{G}_{\ell}(F, B)^0 = \mathfrak{G}_{\ell}(F, A)$. The character c induces an isomorphism

$$\text{Gal}(M/F_{\Phi}(B)) \cong \mu_A \cap \mathfrak{G}_{\ell}(F, B)^0(\mathbf{Q}_{\ell}) = \mu_A \cap \mathfrak{G}_{\ell}(F, A)(\mathbf{Q}_{\ell}).$$

Since $\text{Gal}(M/F_{\Phi}(B))$ is independent of ℓ , we are done. \square

Theorem 3.2. *Suppose F is a function field in one variable over a finite field, A is an abelian variety over F , and ℓ is a prime number not equal to $\text{char}(F)$. Suppose $F = F_{\Phi}(A)$, and let μ_A denote the group of elements of finite order in the center of $\text{End}^0(A)$. Then $\mu_A \cap \mathfrak{G}_{\ell}(F, A)(\mathbf{Q}_{\ell})$ is independent of ℓ .*

Proof. By Theorem 2.4 we can choose an abelian extension M of F such that $\text{Gal}(M/F)$ is isomorphic to μ_A . Let

$$\chi : \text{Gal}(M/F) \rightarrow \mu_A$$

be an isomorphism, let $c : \text{Gal}(F^s/F) \rightarrow \mu_A$ be the composition of χ with the projection $\text{Gal}(F^s/F) \rightarrow \text{Gal}(M/F)$, and define $\rho_{\ell, c} : \text{Gal}(F^s/F) \rightarrow \text{Aut}(V_{\ell}(A))$

by $\rho_{\ell,c}(\sigma) = c(\sigma)\rho_{A,\ell}(\sigma)$. For $F \subseteq F' \subseteq F^s$, let $\mathfrak{G}_{\ell,c}(F')$ denote the Zariski closure of $\rho_{\ell,c}(\text{Gal}(F^s/F'))$. Let $F_{\Phi,c}$ denote the smallest extension F' of F in F^s such that $\mathfrak{G}_{\ell,c}(F')$ is connected. By Lemma 2.3, $\{\rho_{\ell,c}\}$ is a strictly compatible system of integral ℓ -adic representations. Therefore by Proposition 6.14 of [5], $F_{\Phi,c}$ is independent of ℓ . By definition,

$$\text{Gal}(M/F_{\Phi,c}) \cong \mu_A \cap \mathfrak{G}_{\ell,c}(F)^0(\mathbf{Q}_\ell) \quad \text{and} \quad \mathfrak{G}_{\ell,c}(M) = \mathfrak{G}_\ell(M, A).$$

Lemma 2.1 is valid with $\mathfrak{G}_{\ell,c}(F')$ in place of $\mathfrak{G}_\ell(F', A)$; the proof remains unchanged. Therefore,

$$\mathfrak{G}_{\ell,c}(F)^0 = \mathfrak{G}_{\ell,c}(M)^0 = \mathfrak{G}_\ell(M, A)^0 = \mathfrak{G}_\ell(F, A).$$

Since $\text{Gal}(M/F_{\Phi,c})$ is independent of ℓ , we are done. \square

REFERENCES

- [1] M. Borovoi, *The action of the Galois group on the rational cohomology classes of type (p, p) of abelian varieties* (Russian), Mat. Sbornik (N. S.) **94** (136) (1974) 649–652 = Math. USSR Sbornik **23** (1974) 613–616.
- [2] P. Deligne (notes by J. Milne), *Hodge cycles on abelian varieties*, in Hodge cycles, motives, and Shimura varieties (P. Deligne, J. Milne, A. Ogus, K.-y. Shih), Lecture Notes in Mathematics **900**, Springer-Verlag, Berlin-Heidelberg-New York, 1982, pp. 9–100.
- [3] G. Faltings, *Endlichkeitssätze für abelsche Varietäten über Zahlkörpern*, Invent. Math. **73** (1983) 349–366.
- [4] G. Faltings, *Complements to Mordell*, Chapter VI of Rational Points (G. Faltings, G. Wüstholz, et al.), Friedr. Vieweg & Sohn, Braunschweig/Wiesbaden, 1984.
- [5] M. Larsen, R. Pink, *On ℓ -independence of algebraic monodromy groups in compatible systems of representations*, Invent. math. **107** (1992) 603–636.
- [6] M. Larsen, R. Pink, *Abelian varieties, ℓ -adic representations, and ℓ -independence*, Math. Ann. **302** (1995) 561–579.
- [7] B. H. Matzat, *Konstruktive Galoistheorie*, Lecture Notes in Math. **1284**, Springer-Verlag, Berlin-Heidelberg, 1987.
- [8] L. Moret-Bailly, *Pinceaux de Variétés Abéliennes*, Astérisque **129** (1985).
- [9] I. I. Piatetski-Shapiro, *Interrelations between the Tate and Hodge conjectures for abelian varieties* (Russian), Mat. Sbornik **85** (1971) 610–620 = Math. USSR Sbornik **14** (1971) 615–625.
- [10] K. Ribet, *Hodge classes on certain types of abelian varieties*, Amer. J. Math. **105** (1983) 523–538.
- [11] D. J. Saltman, *Generic Galois Extensions and Problems in Field Theory*, Adv. in Math. **43** (1982) 250–283.
- [12] J.-P. Serre, *Représentations ℓ -adiques*, in Algebraic Number Theory (Proceedings of the International Taniguchi Symposium, Kyoto, 1976) (S. Iyanaga, ed.), Japan Society for the Promotion of Science, Tokyo, 1977, pp. 177–193 = # 112 of Œuvres, Vol. III, Springer-Verlag, Berlin-Heidelberg-New York-Tokyo, 1986, pp. 384–400.
- [13] J.-P. Serre, Letters to K. Ribet, Jan. 1, 1981 and Jan. 29, 1981.
- [14] J.-P. Serre, Résumé des cours de 1984–1985, Résumé des cours de 1985–1986, Collège de France.
- [15] J.-P. Serre, *Abelian ℓ -adic representations and elliptic curves*, Second edition, Addison-Wesley, Redwood City, CA, 1989.
- [16] J.-P. Serre, *Propriétés conjecturales des groupes de Galois motiviques et des représentations ℓ -adiques*, in Motives (U. Jannsen, S. Kleiman, J.-P. Serre, eds.), Proc. Symp. Pure Math. **55** (1994), Part 1, pp. 377–400.
- [17] A. Silverberg, Yu. G. Zarhin, *Connectedness results for ℓ -adic representations associated to abelian varieties*, Comp. math. **97** (1995) 273–284.
- [18] A. Silverberg, Yu. G. Zarhin, *Connectedness extensions for abelian varieties*, preprint.
- [19] A. Weil, *Variétés abéliennes et courbes algébriques*, Hermann, Paris, 1948.

- [20] Yu. G. Zarhin, *Endomorphisms of abelian varieties over fields of finite characteristic* (Russian), *Izv. Akad. Nauk SSSR Ser. Mat.* **39** (1975) 272–277 = *Math. USSR - Izv.* **9** (1975) 255–260.
- [21] Yu. G. Zarhin, *Abelian varieties in characteristic p* (Russian), *Mat. Zametki* **19** (1976) 393–400 = *Math. Notes* **19** (1976) 240–244.
- [22] Yu. G. Zarhin, *Weights of simple Lie algebras in the cohomology of algebraic varieties* (Russian), *Izv. Akad. Nauk SSSR Ser. Mat.* **48** (1984) 264–304 = *Math. USSR - Izv.* **24** (1985) 245–282.

DEPARTMENT OF MATHEMATICS, OHIO STATE UNIVERSITY, 231 W. 18 AVENUE, COLUMBUS, OHIO 43210–1174, USA

E-mail address: `silver@math.ohio-state.edu`

DEPARTMENT OF MATHEMATICS, PENNSYLVANIA STATE UNIVERSITY, UNIVERSITY PARK, PA 16802, USA,

INSTITUTE FOR MATHEMATICAL PROBLEMS IN BIOLOGY, RUSSIAN ACADEMY OF SCIENCES, PUSHCHINO, MOSCOW REGION, 142292, RUSSIA

E-mail address: `zarhin@math.psu.edu`