

Thermal emission from bare quark matter surfaces of hot strange stars

Vladimir V. Usov

Department of Condensed Matter Physics, Weizmann Institute, Rehovot 76100, Israel

ABSTRACT

We consider the thermal emission of photons and e^+e^- pairs from the bare quark surface of a hot strange star. The radiation of high-energy ($\gtrsim 20$ MeV) equilibrium photons prevails at the surface temperature $T_s \gtrsim 5 \times 10^{10}$ K, while below this temperature, $8 \times 10^8 \lesssim T_s < 5 \times 10^{10}$ K, emission of e^+e^- pairs created by the Coulomb barrier at the quark surface dominates. The thermal luminosity of a hot strange star in both photons and e^+e^- pairs is estimated.

Subject headings: radiation mechanisms: thermal - stars: neutron

1. Introduction

Witten (1984) pointed out that strange quark matter (SQM) composed of roughly equal numbers of up, down, and strange quarks may be the absolute ground state of the strong interaction, i.e., absolutely stable with respect to ^{56}Fe . Detailed studies showed that, with the uncertainties inherent in a nuclear-physics calculation, the existence of stable SQM is reasonable (e.g., Farhi & Jaffe 1984; Chmaj & Slomiński 1989; Terazawa 1990; Kettner et al. 1995; Weber 1999). If Witten's idea is true, at least some of the compact objects known to astronomers as pulsars, powerful, accreting X-ray sources, X-ray and γ -ray bursters, etc. might be strange stars, not neutron stars as usually assumed (Alcock, Farhi, & Olinto 1986; Glendenning 1990, 1996; Caldwell & Friedman 1991; Madsen & Olesen 1991; Cheng & Dai 1996; 1998; Xu, Qiao, & Zhang 1999). SQM with the density of $\sim 5 \times 10^{14}$ g cm $^{-3}$ can exist, by hypothesis, up to the surface of strange stars. This differs qualitatively from the case of a neutron star surface and opens observational possibilities to distinguish strange stars from neutron stars, if indeed the formers exist. In this Letter, we consider thermal emission of photons and e^+e^- pairs from the bare SQM surfaces of hot strange stars.

2. Bare quark surfaces of strange stars and thermal emission

At the bare SQM surface of a strange star the density changes abruptly from $\sim 5 \times 10^{14}$ g cm $^{-3}$ to zero. The thickness of the SQM surface is about 1 fm = 10^{-13} cm, which is a

typical strong interaction length scale. The SQM surface is a very poor radiator of thermal photons at energies less than about 20 MeV (Alcock et al. 1986). This is because the plasma frequency that is related to the particle density of quarks is very high (see below).

2.1. Thermal Equilibrium Radiation

Hot SQM is filled with electromagnetic waves in thermodynamic equilibrium with quarks. The dispersion relation of these waves may be written in the following simple form (e.g., Adams, Ruderman, & Woo 1963; Alcock et al. 1986)

$$\omega^2 = \omega_p^2 + k^2 c^2, \quad (1)$$

where ω is the frequency of electromagnetic waves, k is their wavenumber,

$$\omega_p = \left(\frac{8\pi e^2 c^2 n_b}{3 \mu} \right)^{1/2} \quad (2)$$

is the plasma frequency, n_b is the baryon number density of SQM, and $\mu \simeq \hbar c (\pi^2 n_b)^{1/3}$ is the chemical potential of quarks. Equation (1) is the familiar dispersion relation for a plasma, and its conventional interpretation may be applied to SQM. Propagating modes exist only for $\omega > \omega_p$. Taking this into account, equation (2) yields the lower limit on the energy of electromagnetic quanta that are in thermodynamic equilibrium with quarks, $\varepsilon_\gamma = \hbar\omega > \hbar\omega_p \simeq 18.5(n_b/n_0)^{1/3}$ MeV, where $n_0 = 0.17 \text{ fm}^{-3}$ is normal nuclear matter density. At the SQM surface where the pressure is zero, we expect $n_b \simeq (1.5 - 2)n_0$ and $\hbar\omega_p \simeq 20 - 25$ MeV (e.g., Farhi & Jaffe 1984; Alcock et al. 1986; Chmaj, Haensel, & Słomiński 1991). Therefore, the spectrum of thermal equilibrium photons radiated from the bare SQM surfaces of strange stars is expected to be very hard.

The energy flux emitted from the unit surface of SQM in thermal equilibrium photons is (Bekefi 1966; Chmaj et al. 1991)

$$F_{\text{eq}} = \frac{\hbar}{c^2} \int_{\omega_p}^{\infty} d\omega \frac{\omega (\omega^2 - \omega_p^2) g(\omega)}{\exp(\hbar\omega/k_B T_s) - 1}, \quad (3)$$

where

$$g(\omega) = \frac{1}{2\pi^2} \int_0^{\pi/2} d\vartheta \sin \vartheta \cos \vartheta D(\omega, \vartheta), \quad (4)$$

k_B is the Boltzmann constant, $D(\omega, \vartheta)$ is the coefficient of radiation transmission from SQM to vacuum, $D = 1 - (R_\perp + R_\parallel)/2$, and

$$R_\perp = \frac{\sin^2(\vartheta - \vartheta_0)}{\sin^2(\vartheta + \vartheta_0)}, \quad R_\parallel = \frac{\tan^2(\vartheta - \vartheta_0)}{\tan^2(\vartheta + \vartheta_0)} \quad (5)$$

$$\vartheta_0 = \arcsin[\sin \vartheta \sqrt{1 - (\omega_p/\omega)^2}] \quad (6)$$

(see, e.g., Landau, Lifshitz, & Pitaevskii 1984). Figure 1 shows the ratio of the equilibrium photon emissivity of the bare surface of hot SQM to the black body surface emissivity, $\xi_{\text{eq}} = F_{\text{eq}}/F_{\text{BB}}$, where $F_{\text{BB}} = \sigma T_s^4$, and σ is the Stefan-Boltzmann constant. From Figure 1 we can see that at $T_s \ll \hbar\omega_p/k_B \sim 10^{11}$ K the equilibrium photon radiation from the bare surface of a strange star is very small, compared to the black body one.

2.2. Emission of e^+e^- pairs

It was pointed out (Usov 1998) that the bare surface of a hot strange star may be a powerful source of e^+e^- pairs which are created in an extremely strong electric field at the quark surface and flow away from the star. The electric field is generated because there are electrons with the density $n_e \simeq (10^{-3} - 10^{-4})n_b$ in SQM to neutralize the electric charge of the quarks (e.g., Alcock et al. 1986). The electrons, being bound to SQM by the electromagnetic interaction alone, are able to move freely across the SQM surface, but clearly cannot move to infinity because of the bulk electrostatic attraction to the quarks. The electron distribution extends up to $\sim 10^3$ fm above the quark surface, and a strong electric field is generated in the surface layer to prevent the electrons from escaping to infinity, counterbalancing the degeneracy and thermal pressure. The typical magnitude of the electric field at the SQM surface is $\sim 5 \times 10^{17}$ V cm $^{-1}$ (Alcock et al. 1986). This field is a few ten times higher than the critical field $E_{\text{cr}} = m_e^2 c^3 / e \hbar \simeq 1.3 \times 10^{16}$ V cm $^{-1}$ at which vacuum is unstable to creation of e^+e^- pairs (Schwinger 1951; Reinhardt 1994). In such a strong electric field, $E \gg E_{\text{cr}}$, in vacuum, the pair creation rate is extremely high, $W_\pm \simeq 1.7 \times 10^{50} (E/E_{\text{cr}})^2$ cm $^{-3}$ s $^{-1}$. At $E \simeq 5 \times 10^{17}$ V cm $^{-1}$, we have $W_\pm \simeq 2.5 \times 10^{53}$ cm $^{-3}$ s $^{-1}$. The high-electric-field region is, however, not a vacuum. The electrons present fill up states into which electrons that would be created have to go. This reduces the pair-creation rate from the vacuum value. It can, in fact, be shown that at zero temperature the process of pair creation is suppressed altogether because there is no free levels for electrons to be created.

At finite temperatures, $T_s > 0$, in thermodynamical equilibrium electronic states are only partly filled, and pair creation by the Coulomb barrier becomes possible. Since the rate of pair production when electrons are created into the empty states is extremely high, the empty states below the pair creation threshold, $\varepsilon \leq \varepsilon_F - 2m_e c^2$, are occupied by created electrons almost instantly, where $\varepsilon_F = \hbar c (\pi^2 n_e)^{1/3}$ is the Fermi energy of electrons, and m_e is the electron mass (Usov 1998). Then, the rate of pair creation by the Coulomb barrier is determined by the process of thermalization of electrons, which favors the empty-state production below the pair creation threshold. The thermal energy of SQM is the source of energy for the process of pair creation.

The flux of e^+e^- pairs from the unit surface of SQM is (Usov 1998)

$$\dot{n}_{\pm} \simeq \Delta r_E \Delta n_e t_{\text{th}}^{-1}, \quad (7)$$

where Δr_E is the thickness of the electron surface layer with a strong electric field,

$$\Delta n_e \simeq \frac{3k_B T_s}{\varepsilon_F} \exp\left(-\frac{2m_e c^2}{k_B T_s}\right) n_e. \quad (8)$$

is the density of electronic empty states with energies below the pair creation threshold at thermodynamical equilibrium, and t_{th} is the characteristic time of thermalization of electrons.

In the surface electron layer, the spectrum of electrons is thermalized due to electron-electron collisions, and the thermalization time is about ν_{ee}^{-1} , where (e.g., Potekhin et al. 1999)

$$\nu_{ee} \simeq \frac{3}{2\pi} \left(\frac{\alpha}{\pi}\right)^{1/2} \frac{(k_B T_s)^2}{\hbar \varepsilon_F} J(\zeta) \quad (9)$$

is the frequency of electron-electron collisions for degenerate electrons with $\varepsilon_F \gg m_e c^2$, $\alpha = e^2/\hbar c = 1/137$ is the fine structure constant, (c.f. Usov 1998)

$$J(\zeta) = \frac{1}{3} \frac{\zeta^3 \ln(1 + 2\zeta^{-1})}{(1 + 0.074\zeta)^3} + \frac{\pi^5}{6} \frac{\zeta^4}{(13.9 + \zeta)^4}, \quad (10)$$

$$\zeta = 2\sqrt{\frac{\alpha}{\pi}} \frac{\varepsilon_F}{k_B T_s} \simeq 0.1 \frac{\varepsilon_F}{k_B T_s}. \quad (11)$$

For typical parameters, $\Delta r_E \simeq 500$ fm and $\varepsilon_F \simeq 18$ MeV, equations (7) - (9) yield

$$\dot{n}_{\pm} \simeq 10^{39} \left(\frac{T_s}{10^9 \text{ K}} \right)^3 \exp \left[-11.9 \left(\frac{T_s}{10^9 \text{ K}} \right)^{-1} \right] J(\zeta) \text{ s}^{-1} \quad (12)$$

within a factor of 2 or so.

The energy flux from the unit surface of SQM in e^+e^- pairs created by the Coulomb barrier is $F_{\pm} \simeq \varepsilon_{\pm} \dot{n}_{\pm}$, where $\varepsilon_{\pm} \simeq m_e c^2 + k_B T_s$ is the mean energy of created particles and n_{\pm} is given by equation (12). Figure 1 shows the ratio of the SQM surface emissivity in e^+e^- pairs to the black body surface emissivity, $\xi_{\pm} = F_{\pm}/F_{\text{BB}}$, versus the surface temperature T_s . Creation of e^+e^- pairs by the Coulomb barrier is the main mechanism of thermal emission from the surface of SQM at $8 \times 10^8 \lesssim T_s \lesssim 5 \times 10^{10}$ K, while the equilibrium radiation dominates at extremely high temperatures, $T_s > 5 \times 10^{10}$ K.

The SQM surface emissivity in photons at energies $\varepsilon_{\gamma} < \hbar\omega_p$ is strongly suppressed (see above). However, low energy photons may leave SQM if they are produced by a non-equilibrium process in the surface layer with the thickness of $\sim c/\omega_p \simeq 10^{-12}$ cm. The emissivity of SQM in non-equilibrium quark-quark bremsstrahlung radiation at low energies, $\varepsilon_{\gamma} < \hbar\omega_p$, has been estimated by Chmaj et al. (1991), $\xi_{\text{neq}} = F_{\text{neq}}/F_{\text{BB}} \simeq 10^{-4}$. Chmaj et al. (1991) made a few assumptions that led to over-estimate of ξ_{neq} , and $\xi_{\text{neq}} \simeq 10^{-4}$ is, in fact, an upper limit. Hence, our estimate of the SQM surface emissivity, $\xi = \xi_{\text{eq}} + \xi_{\pm}$, is valid for $T_s \gtrsim 8 \times 10^8$ K when ξ_{\pm} is more than $\sim 10^{-4}$ (see Fig. 1). For $T_s < 8 \times 10^8$ K the value of ξ is uncertain, $\xi_{\pm} \lesssim \xi \lesssim 10^{-4}$, especially at low energies of photons.

2.3. Thermal radiation from a hot strange star

At $T_s \gtrsim 8 \times 10^8$ K the total luminosity of a bare strange star in both photons and e^+e^- pairs is

$$L = L_{\text{eq}} + L_{\pm} = 4\pi R^2 (F_{\text{eq}} + F_{\pm}), \quad (13)$$

where $R \simeq 10^6$ cm is the radius of the strange star. Figure 2 shows the value of L as a function of the surface temperature T_s . At $T_s < 8 \times 10^8$ K the total luminosity is somewhere between $\sim 4\pi R^2 F_{\pm}$ and $\sim 10^{-3} R^2 F_{\text{BB}}$.

At $T_s \gtrsim 8 \times 10^8$ K the luminosity in e^+e^- pairs created by the Coulomb barrier at the SQM surface is very high, $L_{\pm} \gtrsim 10^{40}$ ergs s⁻¹ (see Fig. 2), that is at least four orders of magnitude higher than

$$L_{\pm}^{\max} \simeq 4\pi m_e c^3 R / \sigma_T \simeq 10^{36} \text{ ergs s}^{-1}, \quad (14)$$

where σ_T is the Thomson cross-section. In this case, the time-scale $t_{\text{ann}} \sim (n_{\pm} \sigma_T c)^{-1}$ for annihilation of e^+e^- pairs is much shorter than the time-scale $t_{\text{esc}} \sim R/c$ for their escape, $t_{\text{ann}}/t_{\text{esc}} \simeq L_{\pm}^{\max}/L_{\pm} < 10^{-4} \ll 1$, and e^+e^- pairs outflowing from the stellar surface mostly annihilate in the vicinity of the strange star, $r \sim R$ (Beloborodov 1999 and references therein). If the total luminosity is not very high, $L \lesssim 10^{42} \text{ ergs s}^{-1}$, the luminosity in e^+e^- pairs at the distance $r \gg R$ cannot be essentially more than L_{\pm}^{\max} . At $L \gg 10^{42} \text{ ergs s}^{-1}$ the outflowing e^+e^- wind is relativistic, and the luminosity in pairs at $r \gg R$ is $\sim L_{\pm}^{\max} (L/10^{42} \text{ ergs s}^{-1})^{3/4} < 10^{-6} L$ (e.g., Usov 1994; Lyutikov & Usov 2000). Hence, far from a bare strange star with the surface temperature $T_s \gtrsim 8 \times 10^8 \text{ K}$ the photon luminosity dominates irrespective of T_s and practically coincides with the total luminosity given by equation (13).

3. Discussion

In this paper, we have considered the thermal emission from the bare SQM surface of a hot strange star. The SQM surface emissivity in both photons and e^+e^- pairs is estimated for $T_s \gtrsim 8 \times 10^8 \text{ K}$. We found that at $T_s \gtrsim 1.5 \times 10^9 \text{ K}$ the SQM surface emissivity is $\gtrsim 10\%$ of the black body surface emissivity. Below this temperature, $T_s \lesssim 1.5 \times 10^9 \text{ K}$, the SQM surface emissivity decreases rapidly with decrease of T_s , and at $T_s < 8 \times 10^8 \text{ K}$ it is at least four orders of magnitude smaller than the emissivity of the black body surface.

Recently, it was argued that SQM is a superconductor if its temperature is not too high (for a review, see Rajagopal & Wilczek 2000). In this case, the non-equilibrium quark-quark bremsstrahlung radiation discussed in (Chmai et al. 1991) is completely suppressed, and equation (13) may be used at $T_s < 8 \times 10^8 \text{ K}$ as well.

”Normal” matter (ions and electrons) may be present at the SQM surface of a strange star. The ions in the inner layer are supported against the gravitational attraction to the underlying strange star by the very strong electric field of the Coulomb barrier. There is an upper limit to the amount of normal matter at the quark surface, $\Delta M \lesssim 10^{-5} M_{\odot}$ (Glendenning & Weber 1992). Such a massive envelope of normal matter with $\Delta M \sim 10^{-5} M_{\odot}$ completely obscures the quark surface. However, it was pointed out (e.g., Haensel, Paczyński, & Amsterdamski 1991) that a strange star at the moment of its formation is very hot. The temperature in the stellar interior may be as high as a few $\times 10^{11} \text{ K}$. The rate of mass ejection from such a hot compact star is very high (e.g.,

Woosley & Baron 1992; Levinson & Eichler 1993; Woosley 1993), and most probably, the normal-matter envelope is blown away by radiation pressure. High temperatures also lead to considerable reduction of the Coulomb barrier, increasing the tunneling of nuclei through the barrier toward the SQM surface (Kettner et al. 1995). Therefore, it is natural to expect the SQM surface of a very young strange star to be nearly (or completely) bare. It was argued that the normal-matter atmosphere of such a star remains optically thin until the temperature of the SQM surface is higher than $\sim 3 \times 10^7$ K (Usov 1997).

Since SQM at the surface of a bare strange star is bound via strong interaction rather than gravity, such a star can radiate at the luminosity greatly exceeding the Eddington limit at the stellar mass of $\sim M_{\odot}$ (Alcock et al. 1986; Chmaj et al. 1991; Usov 1998 and this paper). Therefore, bare strange stars are reasonable candidates for soft γ -ray repeaters (SGRs) that are the sources of short bursts of high-frequency (X-ray and γ -ray) emission with Super-Eddington luminosities, up to $\sim 10^{43} - 10^{45}$ ergs s^{-1} . The bursting activity of SGRs may be explained by fast heating of the stellar surface up to the temperature of $\sim (1 - 2) \times 10^9$ K (see Fig. 2) and its subsequent thermal emission. The heating mechanism may be either impacts of comets onto bare strange stars (Zhang, Xu, & Qiao 2000) or fast decay of superstrong ($\sim 10^{14} - 10^{15}$ G) magnetic fields (Usov 1984; Thompson & Duncan 1995; Heyl & Kulkarni 1998).

This work was supported by the Israel Science Foundation of the Israel Academy of Sciences and Humanities.

REFERENCES

- Alcock, C., Farhi, E., & Olinto, A. 1986, *ApJ*, 310, 261
- Adams, J.B., Ruderman, M.A., Woo, C.-H. 1963, *Phys. Rev.*, 129, 1383
- Bekefi, G. 1966, *Radiation Processes in Plasma* (New York: Wiley)
- Beloborodov, A.M. 1999, *MNRAS*, 305, 181
- Caldwell, R.R., & Friedman, J.L. 1991, *Phys. Lett. B*, 264, 143
- Cheng, K.S., & Dai, Z.G. 1996, *Phys. Rev. Lett.*, 77, 1210
- Cheng, K.S., & Dai, Z.G. 1998, *Phys. Rev. Lett.*, 80, 18
- Chmaj, T., Haensel, P., & Slomiński, W. 1991, *Nucl. Phys. B*, 24, 40

- Chmaj, T., & Slomiński, W. 1989, *Phys. Rev. D*, 40, 165
- Farhi, E., & Jaffe, R.L. 1984, *Phys. Rev. D*, 30, 2379
- Glendenning, N.K. 1990, *Mod. Phys. Lett. A*, 5, 2197
- Glendenning, N.K. 1996, *Compact Stars: Nuclear Physics, Particle Physics, and General Relativity* (Verlag New York: Springer)
- Glendenning, N.K., & Weber, F. 1992, *ApJ*, 400, 647
- Haensel, P., Paczyński, B., & Amsterdamski, P. 1991, *ApJ*, 375, 209
- Heyl, J.S. & Kulkarni, S.R. 1998, *ApJ*, 506, L61
- Kettner, Ch., Weber, F., Weigel, M.K., Glendenning, N.K. 1995, *Phys. Rev. D*, 51, 1440
- Landau, L.D., Lifshitz, E.M., & Pitaevskii, L.P. 1984, *Electrodynamics of continuous media* (Oxford: Pergamon Press)
- Levinson, A., & Eichler, D. 1993, *ApJ*, 418, 386
- Lyutikov, M. & Usov, V.V. 2000, *ApJ*, 543, L129
- Madsen, J., & Olesen, M.L. 1991, *Phys. Rev. D*, 43, 1069
- Potekhin, A.Y., Baiko, D.A., Haensel, P., & Yakovlev, D.G. 1999, *A&A*, 346, 345
- Rajagopal, K. & Wilczek, F. 2000, preprint (hep-ph/0011333)
- Reinhardt, G. 1994, *Quantum Electrodynamics* (Berlin: Springer)
- Schwinger, J. 1951, *Phys. Rev.*, 82, 664
- Terazawa, H. 1990, *J. Phys. Soc. Japan*, 59, 1199
- Thompson, C. & Duncan, R.C. 1995, *MNRAS*, 275, 255
- Usov, V.V. 1984, *Ap&SS*, 107, 191
- Usov, V.V. 1994, *MNRAS*, 267, 1035
- Usov, V.V. 1997, *ApJ*, 481, L107
- Usov, V.V. 1998, *Phys. Rev. Lett.*, 80, 230
- Weber, F. 1999, *J. Phys. G: Nucl. Part. Phys.*, 25, 195

Witten, E. 1984, *Phys. Rev. D*, 30, 272

Woosley, S.E. 1993, *A&AS*, 97, 205

Woosley, S.E., & Baron, E. 1992, *ApJ*, 391, 228

Xu, R.X., Qiao, G.J., & Zhang, B. 1999, *ApJ*, 522, L109

Zhang, B., Xu, R.X., & Qiao, G.J. 2000, *ApJ*, 545, L127

FIGURE CAPTIONS

Fig. 1. Total emissivity of bare SQM surface (solid line), which is the sum of emissivities in equilibrium photons (dashed line) and e^+e^- pairs (dotted line), divided by the black body emissivity, $\xi = \xi_{\text{eq}} + \xi_{\pm}$. Our calculations are valid for the surface temperature $T_s \gtrsim 8 \times 10^8$ K when ξ is more than the upper limit on ξ_{neq} , $\xi_{\text{neq}} \lesssim 10^{-4}$, which is shown by dot-dashed line.

Fig. 2. The total luminosity of a bare strange star (solid line) as a function of the surface temperature T_s , $L = L_{\text{eq}} + L_{\pm}$, where L_{eq} and L_{\pm} are the luminosities in thermal equilibrium photons (dashed line) and e^+e^- pairs (dotted line) respectively. The upper limit on the luminosity in non-equilibrium photons, $L_{\text{neq}} \lesssim 10^{-4}4\pi R^2 F_{\text{BB}}$, is shown by dot-dashed line.



