

Infrared Space Observatory Spectra of R Coronae Borealis Stars. I. Emission Features in the Interval 3 - 25 microns¹

David L. Lambert

Department of Astronomy; University of Texas; Austin, TX 78712-1083
dll@astro.as.utexas.edu

N. Kameswara Rao

Indian Institute of Astrophysics; Bangalore, 560034 India
nkrao@iiap.ernet.in

Gajendra Pandey and Inese I. Ivans

Department of Astronomy; University of Texas; Austin, TX 78712-1083
pandey@astro.as.utexas.edu, iivans@astro.as.utexas.edu

ABSTRACT

Infrared Space Observatory 3 - 25 μm spectra of the R Coronae Borealis stars V854 Cen, R CrB, and RY Sgr are presented and discussed. Sharp emission features coincident in wavelengths with the well known Unidentified Emission Features are present in the spectrum of V854 Cen but not of R CrB or RY Sgr. Since V854 Cen is not particularly H-poor and has a 1000 times more H than the other stars, the emission features are probably from a carrier containing hydrogen. There is a correspondence between the features and emission from laboratory samples of hydrogenated amorphous carbon. A search for C₆₀ in emission or absorption proved negative. Amorphous carbon particles account for the broad emission features seen between 6 - 14 μm in the spectrum of each star.

Subject headings: stars:circumstellar matter – stars: variables: other (R Coronae Borealis)

1. Introduction

The rare class of peculiar stars known as the R Coronae Borealis stars (here, R CrBs) possess two very distinctive characteristics: a propensity to fade at unpredictable times by up to about 8 magnitudes, and an F-G supergiant-like atmosphere that is very H-deficient, He-rich, with considerable amounts of carbon. The fading of the visible light is often rapid with deep minima achieved in a few days to a few weeks but the return to maximum light is generally slower occurring over a period of

several months. O’Keefe’s (1939) suggestion that a R CrB fades when a cloud of soot forms to obscure the photosphere has stood the test of time but the detailed identification of the dust particles and their mode and site of formation remain open questions.

Infrared photometry and low-resolution spectroscopy of R CrBs at maximum light and in decline have revealed several key features of the dusty circumstellar shells. Discovery of IR-excesses provided confirmation of the hypothesis that dust grains were a major constituent of the circumstellar shells (Stein et al. 1969; Lee & Feast 1969). Subsequent studies showed the dust to be at an equivalent blackbody temperature of 500 - 1000 K and present around almost all R CrB

¹Based on observations with *ISO*, an ESA project with instruments funded by ESA Member States (especially the PI countries: France, Germany, the Netherlands, and the United Kingdom) and with the participation of ISAS and NASA.

stars (Feast & Glass 1973; Feast et al. 1977; Glass 1978), but not distributed in a spherically homogeneous shell (Forrest, Gillett, & Stein 1972). Later observations including measurements of optical polarization suggested the dust may be ejected in a preferred plane (Stanford et al. 1988; Rao & Raveendran 1993; Clayton et al. 1997). Feast et al. (1977) showed that for RY Sgr, a 38 day pulsating R CrB star, the intensity of the $3.5 \mu\text{m}$ emission from the warm dust varied with the same period as the visual light showing that the dust was heated by starlight. This variation was shown to continue with no significant decline in mean intensity when the star experienced a deep visual decline. The mean intensity did, however, drop at times when the star remained at maximum light. This extended series of IR observations show that the circumstellar shell is composed of discrete clouds; should a cloud form along the line of sight to the star, a decline is witnessed but the heating of the collection of clouds comprising the circumstellar shell is little affected because only a modest fraction of the stellar surface is blocked from directly heating the shell. *IRAS* observations revealed, in addition to the warm dust detected from the ground, an extended ('fossil') shell of cold ($T \sim 30 \text{ K}$) dust (Gillett et al. 1986; Rao & Nandy 1986; Walker 1986).

Attribution of emission and absorption features in spectra of circumstellar shells around C-rich objects has been contested. Two leading proposals vie for identification of emission features: polycyclic aromatic hydrocarbons (here, PAHs), and hydrogenated amorphous carbon (here, HAC). Since R CrBs are H-deficient to different degrees, observation and analysis of their infrared emission features affords an opportunity to investigate the role of hydrogen in formation of the very large molecules or dust grains. Hydrogen deficiency and other circumstances peculiar to R CrB circumstellar shells may promote formation of other species.

With the advent of the *Infrared Space Observatory* (*ISO*, Kessler et al. 1996), there came an opportunity to obtain IR spectra of R CrB stars in and beyond the narrow windows open to ground-based observers. In this paper we discuss and interpret spectra of three R CrBs obtained with *ISO* over the wavelength region $2.5 - 45 \mu\text{m}$. At the H-rich end of the range is V854 Cen with a H

abundance only a factor of 100 below normal. We contrast V854 Cen with RY Sgr and R CrB, two stars with considerably less H than V854 Cen.

2. Observations

2.1. The Stars

Two stars were satisfactorily observed under our *ISO* program: RY Sgr and V854 Cen. Data for R CrB were retrieved from the *ISO* archive. Details of the observations including the parameters TDT and AOT are summarized in Table 1. At the time of observation, RY Sgr and R CrB were close to maximum light according to the AAVSO light curves. When observed V854 Cen appears to have been recovering from minimum light: the V magnitude according to Lawson et al. (1999) was about 7.7 or 0.5 magnitudes below maximum light.

2.2. Spectra - Reduction and Calibration

The spectra were obtained with the *Short-Wavelength Spectrometer* (*SWS*, de Graauw et al. 1996) on the *ISO* spacecraft. Spectra were recorded on separate 1×12 detector arrays. The complete spectrum was recorded as 12 different but overlapping spectral bands using different combinations of grating, aperture, and detector array. Our spectra of RY Sgr and V854 Cen were observed at the slowest scanning speed, and, therefore, the highest *SWS* resolving power of $R = \lambda/\Delta\lambda \sim 1000$. The archived R CrB spectrum had been taken at a higher speed and lower resolving power ($R \sim 400$).

Standard pipeline processing of the raw *SWS* data involves three principal steps: subtraction of the dark current, assignment of the wavelength scale, and the determination of the absolute fluxes. Dark current measurements were obtained just prior and after an observation. These were averaged and subtracted from the signal from the source. Wavelengths were calculated from the recorded grating positions and grating constants in the *SWS* database (Valentijn et al. 1996). Similarly, the raw signals were converted to absolute fluxes (in Janskys) using calibrations in the *SWS* database. Each of the 12 bands was reduced separately, and then combined to form a composite spectrum.

TABLE 1
THE SWS OBSERVATIONS

Star	Date ^a		V ^b	TDT	AOT	Speed
	UT	JD'				
V854 Cen	1996 Sep 9	50336	7.7	29701401	SWS01	4
R CrB	1998 Jan 15	50828	6.1	79200268	SWS01	2
RY Sgr	1997 Mar 25	50533	6.6	49500503	SWS01	4

^aJD = JD' + 240000

^bSee text

The output of the pipeline processing was a spectrum with discontinuities across the edges between adjacent bands. Additional reduction procedures were then applied using the interactive software *ISAP* (Roelfsema et al. 1993). Spectra in the 12 bands were grouped into four larger bands covering the wavelength regions: 2.35 - 4.1 μm , 4.0 - 12.0 μm , 12.0 - 28.0 μm , and 29.0 - 45.2 μm . A spectrum from a single detector was inspected for abrupt jumps in signal level, and excessive noise levels. Detectors so affected were dropped from the sample. A few bad data points were edited out. Following recommendations by Rita Loidl (private communication), we reduced the discontinuities by applying additive corrections relative to the signal from a reference detector within each AOT band selected based on two criteria: (1) it should provide the smallest discontinuity at the band edges, and the slope that is most consistent with other detectors, and (2) choice of the detector should not change the mean flux level. The first band (2.4 - 2.6 μm) was taken as the flux reference. We found that band 7 (7 - 12.0 μm) showed a significant offset between the scanning directions; the ‘down’ scan spectra were always of higher flux than the ‘up’ scan spectra. This is attributed to ‘memory effects’; the ‘down’ scans were rejected. This procedure provided spectra of higher signal to noise ratio than achieved in all of our earlier attempts to eliminate the discontinuities. Final spectra after rebinning to a resolving power of $R = 1000$ for RY Sgr and V854 Cen and $R=400$ for R CrB are shown in Figure 1 .




Fig. 1.— *ISO SWS* spectra of R CrB, RY Sgr, and V854 Cen (solid lines). The adopted photospheric continuum is shown by the dotted line in each panel. Assumed effective temperatures are 6900 K (R CrB), 7250 K (RY Sgr), and 6750 K (V854 Cen). The dashed line shows for R CrB and RY Sgr the blackbody continuum from the circumstellar dust for the temperatures given in the text. The sum of the dust and photospheric continua is represented by the crosses. In the case of V854 Cen, the very small contribution of the photospheric continuum is neglected and the crosses give the dust continuum. Excess emission over the fitted fluxes (crosses) is more clearly displayed in Figures 2 and 3.

2.3. Spectral Energy Distributions

Since the stars are of variable flux in the infrared, we should not expect the *ISO* fluxes to match exactly published fluxes. V854 Cen was caught by *ISO* in an unusually bright phase at IR wavelengths: for example, the *ISO* flux density at $10\ \mu\text{m}$ is 25 Jy while 15 Jy was found by *IRAS*. *IRAS* fluxes for RY Sgr are 60 - 70% higher than the *ISO* fluxes. R CrB was about 40% brighter as observed by *ISO* than by *IRAS*. Walker et al. (1996) observed R CrB with *ISO* when the star was at minimum light or 7 magnitudes below maximum light. The *ISOPHOT* instrument was used for photometry at 6 bandpasses from $60\ \mu\text{m}$ to $200\ \mu\text{m}$, and a low resolution spectrum was acquired from $2.5\ \mu\text{m}$ to $5\ \mu\text{m}$ and $5.8\ \mu\text{m}$ to $11.6\ \mu\text{m}$ at a resolving power of about 100. The *ISO* photometry seems to show that R CrB was a factor of about two brighter at minimum than when measured with *IRAS*. The $2.5\ \mu\text{m}$ to $11.6\ \mu\text{m}$ spectrum differs in shape but not average flux from ours: the *ISOPHOT* spectrum shows less flux than the *SWS* spectrum for wavelengths shorter than about $4\ \mu\text{m}$ and longer than about $8.5\ \mu\text{m}$ with differences largest at the wavelength limits of the *ISOPHOT* spectrum. Between $4\ \mu\text{m}$ and $8.5\ \mu\text{m}$, the *ISOPHOT* spectrum has the higher flux by up to about a factor of two. The *ISOPHOT* spectrum cannot be well fitted by a blackbody spectrum.

Infrared excesses are conveniently expressed using a fit of a blackbody spectrum after consideration of the photospheric spectrum. This is an adequate artifice here because we are primarily interested in the emission and absorption features. Predicted fluxes from model atmospheres show that the IR photospheric flux is essentially identical to that of a black body at the stellar effective temperature (Asplund et al. 1997a,b). We resolve the observed spectrum into photospheric and circumstellar components; the former does not exceed a 10% contribution at the short wavelength end of the *SWS* bandpass. Then, we fit the circumstellar spectrum with that of a blackbody. Figure 1 shows the photospheric and circumstellar components with the observed spectra. The blackbody's contribution is subtracted to show more clearly the emission and absorption features that may be present (see Figure 2, 3, and 4). Over the *SWS* spectrum, interstellar reddening, slight at B and V, is negligible.

For V854 Cen and RY Sgr, the infrared excess is well fitted by a single blackbody: temperatures of 1040 ± 20 and 820 ± 10 K are obtained for V854 Cen and RY Sgr, respectively. The former temperature is slightly hotter than the 900 K estimated by Lawson & Cottrell (1989) from *IRAS* measurements, but is equal to the excitation temperature of circumstellar C_2 molecules reported by Rao & Lambert (2000). Our temperature for RY Sgr fits the *IRAS* measurements; Walker (1985) derived a blackbody temperature of 800 ± 50 K from a fit to broad-band measurements out to $60\ \mu\text{m}$. A range of 600 K to 900 K was found by Feast et al. (1977) from ground-based photometry. To fit the R CrB *ISO* spectrum requires two blackbody components: one at $T=610\pm 60$ K provides an adequate fit at wavelengths longer than $5\ \mu\text{m}$, and the contribution from a hotter black body, $T = 1390\pm 270$ K, is needed to fit the shorter wavelengths. Walker (1985) obtained a temperature of 650 ± 50 K from *IRAS* broad-band fluxes from 12 to $100\ \mu\text{m}$. We note that the *IRAS* observations suggested a similar ($T = 680$ K) dust temperature (Rao & Nandy 1986). In each case, the blackbody simulation of the *SWS* spectrum fits the observations to a few per cent with small deviations largely arising from emission features.

Our adoption of a blackbody to represent the infrared fluxes is an artifice enabling us to display more clearly the emission features. Representation of an infrared flux distribution by a blackbody is tantamount to assuming that either the dusty clouds are isothermal and optically thick at all wavelengths or the clouds are optically thin with an absorption coefficient that is quasi-gray in the infrared. An attempt to model the clouds as optically thin was made using laboratory measurements of the non-gray absorption coefficient for amorphous carbon samples from Colangeli et al. (1995 - see also Bussoletti et al. 1987); the measurements for their BE sample were adopted. The inferred dust temperature is slightly lower, and the overall fit to the observed fluxes somewhat inferior to the results for the optically thick case. Presumably, the latter result is, in part, due to the failure of the adopted laboratory measured absorption coefficients to represent the actual circumstellar dust; the laboratory results depend on how the amorphous carbon is prepared. More refined models than either our simple optically thick




Fig. 2.— Difference spectra of R CrB, RY Sgr, and V854 Cen from 3 to 10 μm . These are the *ISO SWS* spectra after subtraction of the stellar and circumstellar continua, i.e., Figure 1’s solid line minus the fitted distribution represented by the crosses. Note the change of flux scale.




Fig. 3.— Difference spectra of R CrB, RY Sgr, and V854 Cen from 10 to 25 μm . These are the *ISO SWS* spectra after subtraction of the stellar and circumstellar continua, i.e., Figure 1’s solid line minus the fitted distribution represented by the crosses. Note the change of flux scale.

and thin approximations would presumably provide higher quality fits but our present purpose of investigating the narrow emission features is served adequately by adopting the blackbody fit to the spectrum.

Amorphous carbon particles show increased absorption between about 6 μm and 14 μm (see, for example, Colangeli et al.’s Figure 5, 1995) that appear to account for the broad emission seen from all three R CrB stars. In the optically thick case, one expects these features to appear in absorption on the grounds that the temperatures of dust grains on clouds’ outer edges will decrease towards the surface facing away from the star. Optically thin clouds will show emission bands associated with the increased absorption. Colangeli et al.’s measurements, and the temperatures found from the ‘optically thin’ fit to the infrared fluxes provide a rough fit to the broad emission features.

3. Emission and Absorption Features

3.1. Unidentified Infrared Emissions

The principal goal of this investigation was to search for spectral features. Such features are seen in the residual or difference spectra (Figures 2, and 3) obtained by subtracting the best-fitting blackbody spectrum from the observed spectrum. Emission features may be classified as ‘broad’ or ‘narrow’. Narrow features are seen only in V854 Cen. Broad features are seen in all three R CrB stars.

V854 Cen. The narrow emission features (Table 2) seen in V854 Cen’s spectrum correspond to some of the famous unidentified infrared features (UIR) seen in emission from many post-AGB stars, planetary nebulae, and H II regions. The 3.29 μm UIR is certainly present (Figure 2), the 3.4 μm UIR is possibly present, but the UIR at 3.51 μm is not seen above the noise. In the 5-10 μm interval, the UIR at 6.29 μm but not those at 5.6, 6.9, 7.3, 7.7, and 8.6 μm stand out above the noise and/or the smooth profile of the broad feature that extends from 6-9 μm . Three UIRs are seen in the 10-25 μm region (Figure 3): emission is clearly seen at 11.3, and 13.5, and possibly 14.6 μm . The latter two UIRs were discovered in *ISO* spectra of Red Rectangle (Waters et al. 1998).

The remaining emission features are broad. A feature appears at 3.95 μm but it occurs at the

border of two spectral bands. The $6.3 \mu\text{m}$ UIR feature appears at the short wavelength limit of a broad feature extending to about $9 \mu\text{m}$ with local peaks at $6.9 \mu\text{m}$ (possibly the intrusion of the $6.9 \mu\text{m}$ UIR) and $8.1 \mu\text{m}$. Clayton et al. (1995) observed V854 Cen from 8.5 to $8.8 \mu\text{m}$ at a resolving power of 1000 and found a featureless continuum, an observation consistent with our spectrum. An apparently broad feature at $12.6 \mu\text{m}$ is bracketed by UIRs at $11.3 \mu\text{m}$ and $13.5 \mu\text{m}$. Inspection of Buss et al.'s (1993) collection of $6 - 13 \mu\text{m}$ spectra of 'transition objects' (i.e., stars evolving from the AGB to the planetary nebula phase) shows a gross similarity between V854 Cen and the C-rich protoplanetary nebula *IRAS* 22272+5435. The spectrum of V854 Cen is definitely different from those of the planetary nebulae NGC 7027 and CPD - 56°8032.

RY Sgr. Two broad features are present. One extends from about $6 \mu\text{m}$ to $9 \mu\text{m}$, and the second from about $11 \mu\text{m}$ to $18 \mu\text{m}$. There is little evidence of structure within these features. Except for the presence of the $6.3 \mu\text{m}$ feature in V854 Cen, the appearance of these broad features is similar for V854 Cen and RY Sgr. Clayton et al. (1995) obtained a spectrum of the interval 8.2 to $8.8 \mu\text{m}$ at a resolution of 1000 and reported that it was featureless, a result consistent with our spectrum. The *IRAS LRS* spectrum shown by Clayton et al. provides a marginal detection of the 11.5 - $15 \mu\text{m}$ feature, and clear evidence for stronger emission extending from the $8 \mu\text{m}$ limit of the *LRS* spectrum to $9 \mu\text{m}$, which we identify with the 6 - $9 \mu\text{m}$ broad emission feature.

R CrB. Again, two broad emissions are present. We confirm the 6 to $9 \mu\text{m}$ feature found by Buss et al. (1993), and partially recorded on the *IRAS LRS* 8 - $22 \mu\text{m}$ spectrum (Clayton et al. 1995). An emission peak at $6.5 \mu\text{m}$ in the feature reported by Buss et al. is not confirmed. A second broad feature extends from 11.5 to about $15 \mu\text{m}$. Both features commence at the same wavelength in all three stars but span a shorter wavelength interval in R CrB than do the apparently similar features in V854 Cen and RY Sgr.

Absorption near $4.6 \mu\text{m}$ seen in V854 Cen, and RY Sgr is likely due to the fundamental vibration-rotation absorption lines from circumstellar CO molecules. Absorption at $7.2 \mu\text{m}$ in RY Sgr appears to be real; the feature is present in each of

the scans of this interval.

3.2. A Search for C_{60}

Goeres & Sedlmayr (1992) may have been the first to have considered the spherical cage molecule buckminsterfullerene, C_{60} , as a possible constituent of dust clouds around R CrB stars. Their theoretical model predicted low abundances of C_{60} molecules. This prediction could be circumvented if the carbon-rich gas contained H atoms. Formation of large C-containing molecules is facilitated in the presence of H-containing molecules such as acetylene (C_2H_2) and by photoerosion of hydrogenated amorphous carbon. This suggests that V854 Cen's dust clouds might harbor C_{60} molecules, even if RY Sgr and R CrB do not. Understanding molecule and dust formation in the outer atmosphere of a R CrB (or any star!) is far from complete. Therefore, a search for these very stable carbon cages is of interest.

Laboratory infrared spectroscopy of free C_{60} molecules has shown that the strongest of the 46 possible vibrational bands are at 7.0 , 8.4 , 17.4 , and $18.8 \mu\text{m}$ (Nemes et al. 1994, see also Krätschmer et al. 1990 and Frum et al. 1991). The given wavelengths are laboratory measurements extrapolated to a temperature of 0 K . Positions (and profiles) of these bands will vary with temperature of the gas as more and more levels of the rotational ladder are populated with increasing temperature. Nemes et al. estimate, for example, that the $8.4 \mu\text{m}$ will show its bandhead at $8.58 \mu\text{m}$ at 1000 K with a width of $0.1 \mu\text{m}$. Possibly, some molecules may be ionized - positively or negatively - in the dust clouds. Infrared spectroscopy of C_{60}^+ molecules in a rare gas matrix has provided measurements of vibrational bands of the ion C_{60}^+ at 7.1 and $7.5 \mu\text{m}$ (Fulara, Jakobi, & Maier 1993); expected bands at longer wavelengths were not investigated. The negative ion C_{60}^- had strong bands at 7.2 and $8.3 \mu\text{m}$. These measurements include a matrix-dependent shift but the bands of the gas phase ions will be close to the measured positions. Temperature dependence of the band positions and widths will occur, as they do for C_{60} , for the free cations and anions.

Clayton et al. (1995) searched unsuccessfully for the C_{60} $8.6 \mu\text{m}$ band on ground-based $R = 1000$ spectra of the interval $8.49 \mu\text{m}$ to $8.82 \mu\text{m}$ at a S/N ratio of about 100 for R CrB but somewhat

TABLE 2
V854 CEN - THE NARROW EMISSION FEATURES

Parameters ^a			
λ [μ m]	FWHM [μ m]	Flux [10^{-17} W cm ²]	UIR ^b
3.294(02)	0.052(02)	0.116(12)	Yes
6.298(10)	0.177(11)	0.378(70)	Yes
11.280(10)	0.443(11)	0.363(30)	Yes
13.490(17)	0.220(20)	0.100(20)	Yes

^aThe estimated uncertainty is given in parantheses following the entry, i.e.,
3.294(02) = 3.294 \pm 0.002

^bSee list of UIRs given by
<http://www.ipac.caltech.edu/iso/lws/unidentified.html>




Fig. 4.— Comparison of the difference spectrum of V854 Cen and the laboratory emission spectrum of hydrogenated amorphous carbon at 773 K from Scott et al. (1997).

lower for V854 Cen and RY Sgr; the wavelength of 8.6 μ m is that expected for C₆₀ absorption at 1000 K. We confirm that the feature is not present. (The S/N ratio at 8.5 μ m is 60 (V854 Cen), 70 (RY Sgr), and 80 (R CrB).) Our *SWS* spectra permit a search for the other strong bands. A feature appears near the 7.1 μ m band in V854 Cen but this is at the boundary of two *SWS* bands and, therefore, we doubt that it is a real emission feature. This wavelength is also close to that of bands measured for matrix-isolated C₆₀⁺ and C₆₀⁻. Other bands of these ions at 7.5 μ m for C₆₀⁺ and 8.3 μ m for C₆₀⁻ are not present in our spectra. At longer wavelengths (Figure 3), we do not detect the C₆₀ 17.5 and 19.0 μ m bands. An apparent emission feature at 19.2 μ m is an artefact.

3.3. The [Ne II] 12.8 μ m Line

Optical spectra of R CrBs in deep declines show a few broad emission lines with base widths of 400 - 600 km s⁻¹. Carriers of permitted broad lines include He I, Na I D, Ca II, and the C₂ molecule. Forbidden broad lines of [N II], [O II] and other atoms and ions have been noted. Detailed reports for V854 Cen (Rao & Lambert 1993) and R CrB (Rao et al. 1999) list and discuss the detected lines. Broad lines, if present, would be resolved on the *SWS* R = 1000 spectra. Broad lines, which are detectable only after a star has faded by several magnitudes, are not to be confused with sharp ‘chromospheric’ lines, primarily from neutral and singly-ionized metals, seen early in and throughout a decline. Sharp lines would not be resolved

on our spectra.

Gas emitting the optical broad forbidden lines should also emit the [Ne II] 12.8 μm line. If the gas is of low electron density, as indicated for V854 Cen by the intensity ratio of the red [S II] lines (Rao & Lambert 1993), the predicted flux of the [Ne II] line is about 0.3% that of the 6584 \AA [N II] line for equal abundances of the ions N^+ and Ne^+ . Considerations of elemental abundance and rough estimates of the ionization conditions suggest that the predicted flux of the [Ne II] line is several orders of magnitude smaller than the detection limit. Optical [O I] and [C I] lines observed in spectra of R CrB at minimum light indicate electron densities in excess of the critical densities for [N II] and [Ne II] lines. Then, the fluxes in the [N II] and [Ne II] lines are comparable for equal ionic densities, but, even in this more optimistic case, the flux of the 12.8 μm line is at last a factor of 100 smaller than the detection limit. We conclude that the absence of the 12.8 μm [Ne II] line is consistent with optical detections of forbidden lines.

4. Discussion

4.1. Hydrogenated Amorphous Carbon Dust

Surely, the most intriguing result from these *ISO* spectra is the presence of some UIRs in V854 Cen but not in RY Sgr and R CrB. One may suspect this difference is related to the fact that V854 Cen is less H-deficient than the other two stars. Then, those UIRs seen in V854 Cen likely arise from transitions involving H and C bonds in free molecules, large clusters, or grains.

Analyses of the UV extinction by R CrBs suggested that the dust was amorphous carbon (Hecht et al. 1984). Laboratory studies of absorption and emission infrared spectra of aggregates of sub-micron particles or thin films of partially hydrogenated amorphous carbon have been reported with a view to identifying the UIRs (Borghesi, Bussoletti, & Colangeli 1987; Scott & Duley 1996; Scott, Duley, & Jahani 1997). In many cases, the precise profile of the absorption/emission feature depends on the thermal history of the laboratory sample. Nonetheless, there is a correspondence between the narrow emission features seen in the spectrum of V854 Cen and the laboratory spectra of HAC. This is well illustrated in Figure 4 where

the emission spectrum of a thin HAC film at 773 K (Scott et al. 1997) is plotted with our spectrum of V854 Cen. The temperature of 773 K was the highest temperature investigated by Scott et al. A majority of the emission features seen in the laboratory spectrum are identifiable in V854 Cen's spectrum: absorption at 4.3 μm in V854 Cen presumably masks the emission seen in the laboratory spectrum at a similar wavelength, and the laboratory emissions at 7.1 μm and 12.3 μm are not seen as distinct peaks in the stellar spectrum. Apart from differences in the laboratory and stellar profiles of the 7-9 μm and 11-15 μm bands, there is an obvious difference in the relative strengths of these bands: the ratio of the shorter to the longer wavelength emission is much greater for the stellar than for the laboratory spectrum.

In the laboratory experiments, the intensity of the 3.4 μm feature, characteristic of aliphatic hydrocarbons, decreased as the temperature was raised from 425 K to 775 K and the intensity of the 3.29 μm feature, attributable to aromatic hydrocarbon and seen in V854 Cen, appeared and strengthened. This change was attributed by Scott et al. to the transformation of HAC from a polymer to a protographitic solid. In V854 Cen, the dominant emission is at about 3.3 μm not at 3.4 μm . The 6.2 μm (aromatic C-C ring vibration) and 11.3 μm (aromatic CH-bending mode) are also more intense at the higher temperatures. Emission corresponding to the CH bending modes is also apparent at 12.3 and 13.2 μm in the 773 K laboratory spectra, and in V854 Cen. In short, these and other comparisons of relative intensities suggest that a laboratory spectrum at a temperature higher than 773 K, perhaps at the temperature of 1040 K of the dust shell, would prove to be an even closer match to V854 Cen's narrow features. UIRs seen in astronomical sources at 7.7 and 8.6 μm are apparently absent from V854 Cen, and were also reported as absent by Borghesi et al. (1987) in their studies of absorption spectra of small HAC grains. The 7.7 μm feature was attributed to a CN-stretch of a NH_2 group which owing to a lack of N in the laboratory samples did not appear in laboratory spectra. Nitrogen is not especially abundant in V854 Cen and, therefore, absence of this 7.7 μm UIR is likely understandable.

RY Sgr and R CrB show broad emission fea-

tures similar to those seen from V854 Cen but the sharp features are exclusive to the latter star. Profiles of the 7-9 μm and 11-15 μm bands differ from star to star. Apart from the sharp features shown by V854 Cen, the spectra of V854 Cen and R CrB are the most similar. The 7-9 μm band appears resolvable into two bands with minimum flux at about 7.4-7.6 μm , and the emission in the longer wavelength band does not extend beyond about 15 μm . By contrast, RY Sgr's 7-9 μm band does not show the minimum, and emission in the longer wavelength band extends to about 18 μm . These broad emission bands would appear to be a property of dehydrogenated amorphous carbon (see Koike, Hasegawa, & Manabe 1980; Bussoletti et al. 1987; Colangeli et al. 1995).

Hydrogenated amorphous carbon accounts well for V854 Cen's narrow emission features. Absence of strong features at 7.7 and 8.6 μm that are characteristic of PAHs suggest the free molecules are not abundant in its circumstellar shell. Amorphous carbon formed in a hydrogen containing atmosphere is composed of randomly oriented linked and connected PAH clusters, and, therefore, the spectrum of HAC will resemble that of free PAHs with bands blurred, shifted, and blended by solid state effects. By an extension of this argument, it is not surprising that an overall match to the HAC spectrum is also found from laboratory spectrum of coals such as anthracite and semi-anthracite (Guillois et al. 1996; Papoular et al. 1996). We suppose it more probable that HAC rather than coals form in V854 Cen's circumstellar shell.

4.2. The Energy Budget

It is of interest to investigate the energetics of the dust clouds which we assume are heated solely by absorption of starlight. Circumstellar dust is distributed in clouds around the star. After full recovery from a decline, a star always returns to the same brightness. Therefore, it is plausible to suppose that at these times there is no cloud along the line of sight. Although the detailed comparisons are lacking, available comparisons of observed (corrected for interstellar extinction) and predicted stellar flux distribution about visible wavelengths (see Asplund et al.'s Figure 8 1997a for R CrB) suggest that circumstellar extinction by small particles is not present at maximum light; large particles producing gray extinc-

tion cannot be ruled out by this comparison. Gray extinction from a changing cloud of large particles would cause fluctuations in the brightness of a star at maximum light. If large dust particles are made in the cloud responsible for a decline, they cannot persist in the line of sight once spectroscopic manifestations of a decline (i.e., sharp emission lines) have disappeared in the recovery to maximum light.

Given these assumptions, there is a simple inequality that must be satisfied by the integrated infrared emission from the dust (f^d) and the integrated stellar flux (f^*), where f denotes flux received at the Earth. The inequality is $f^d < f^*$ and recognizes that (i) the dust clouds may be optically thin, and (ii) the clouds may not subtend 4π of solid angle at the star. The dust emission, f^d , is estimated as described above from the measured spectrum after subtraction of a small contribution from the star. Stellar emission, f^* , is estimated from published UBVRIJ magnitudes at maximum light with estimates added for the ultraviolet flux from *IUE* spectra and for flux beyond J.

With the *ISO* fluxes, we find $f^d/f^* \simeq 0.9$ for V854 Cen, and 0.4 for R CrB and RY Sgr. The larger ratio for V854 Cen may well result from the fact that we observed the star in the recovery from a deep decline when the infrared fluxes were enhanced due to a contribution from the recently formed dust cloud that caused the decline. This suggestion is consistent with Forrest's (1974) observations of R CrB showing that the infrared flux increased after minimum light with f^d/f^* increasing from 0.2 to 0.7, by his estimates. Noting too that the *IRAS* fluxes for RY Sgr were 60 - 70% higher than the *ISO* fluxes, we suggest $f^d \sim 0.5f^*$ for these three stars at maximum light between declines. Near-equality of f^d and f^* implies that the ensemble of clouds provides a covering factor approaching 4π , and that a majority of the clouds are optically thick or almost so. Given this result, it seems surprising that the stars are visible! Or to phrase this surprise in another way - are there R CrB stars that have eluded discovery because they are embedded in a very optically thick conglomeration of clouds?

If the dust clouds are optically thin, the mass in amorphous carbon can be estimated readily. If the mean absorption coefficient at 7 μm , the peak of the dust emission, is taken from Colangeli et al.

(1995), and the stellar distance derived from the stellar flux at about $1 \mu\text{m}$ and an assumed stellar radius of 100 solar radii, the dust masses (in solar masses) are 2×10^{-8} , 4×10^{-8} , and 3×10^{-8} for V854 Cen, R CrB, and RY Sgr, respectively. These are probably underestimates by a modest factor. At minimum light, a R CrB is approximately 7 magnitudes fainter than at maximum light. If the obscuring cloud covers the surface uniformly, an optical depth of about 6 at visual wavelengths exists. Given the approximately inverse wavelength dependence of the dust absorption coefficient, the optical depth of the cloud at $8 \mu\text{m}$ is about 0.4. This estimate for a fresh cloud is likely an overestimate for the typical cloud in the dust shell.

To build a more complete picture of the clouds, imposition of radiative equilibrium is required. Hartmann & Apruzese (1976) were able to fit Forrest's (1974) infrared photometry for infrared maximum and minimum (both corresponding to visual maximum) with dust distributed out to 300 to 500 stellar radii. Optical depth at infrared wavelengths for the assumed dust distributions is very small. The dust mass derived of about $5 \times 10^{-7} M_{\odot}$ is larger than our estimate but this is possibly due to differences in the adopted absorption coefficients. A recalculation of model circumstellar shells of HAC grains in radiative equilibrium would be of interest.

5. Concluding Remarks

Dust in a cloud is reluctant to betray its identity. Yet, the *ISO* 3 - 25 μm spectra of the ensemble of dust clouds around the stars V854 Cen, R CrB, and RY Sgr offer a few new clues. Sharp emission features coincident with certain of the UIRs are present only in V854 Cen's spectrum, and are a fair replica of emission seen in laboratory spectra of hydrogenated amorphous carbon. These associations are consistent with the fact that V854 Cen, although H-poor relative to normal stars, is H-rich by a factor of 1000 with respect to R CrB and RY Sgr. Spectra of all three stars show a double-peaked broad emission feature between $6 \mu\text{m}$ and $14 \mu\text{m}$ that corresponds to a feature in the extinction curve of amorphous carbon. In summary, amorphous carbon is a major contributor to the infrared emission from R

CrB and, in the case of V854 Cen, the amorphous carbon is hydrogenated.

We are especially grateful to Rita Loidl for advice on how to reduce the *SWS* spectra. We thank B. Gustafsson and M. Asplund for their interest in this project, and W. Duley for providing the laboratory spectrum illustrated in Figure 4. The *ISO* Spectral Analysis Package (*ISAP*) is a joint development by the *LWS* and *SWS* Instrument Teams and Data Centers with CESR, IAS, IPAC, MPE, RAL, and SRO as contributing institutions. This research was supported in part by NASA through grants NAG5-3348 and CITJ-961543.

REFERENCES

- Asplund, M., Gustafsson, B., Kiselman, D., & Eriksson, K. 1997a, *A&A*, 318, 521
- Asplund, M., Gustafsson, B., Kiselman, D., & Eriksson, K. 1997b, *A&A*, 323, 286
- Borghesi, A., Bussoletti, E., & Colangeli, L. 1987, *ApJ*, 314, 422
- Buss, R.H.Jr., Tielens, A.G.G.M., Cohen, M., Werner, M.W., Bregman, J.D., & Witteborn, F.C. 1993, *ApJ*, 415, 250
- Bussoletti, E., Colangeli, L., Borghesi, A. & Orofino, V. 1987, *A&AS*, 70, 257
- Clayton, G.C., Bjorkman, K.S., Nordsieck, K.H., Zellner, N.E.B., & Schulte-Ladbeck, R.E., 1997, *ApJ*, 476, 870
- Clayton, G.C., Kelly, D.M., Lacy, J.H., Little-Marenin, I.R., Feldman, P.A., & Bernath, P.F. 1995, *AJ*, 109, 2096
- Colangeli, L., Mennella, V., Palumbo, P., Rotundi, A., & Bussoletti, E. 1995, *A&AS*, 113, 561
- de Graauw, Th. et al. 1996, *A&A*, 315, L49
- Feast, M.W., Catchpole, R.M., Lloyd Evans, T., Robertson, B.S.C., Dean, J.F., & Bywater, R.A. 1977, *MNRAS*, 178, 415
- Feast, M.W., & Glass, I.S. 1973, *MNRAS*, 161, 293
- Forrest, W.J. 1974, thesis, Univ. of California, San Diego

- Forrest, W.J., Gillett, F.C., & Stein, W.A. 1972, *ApJ*, 178, L129
- Frum, C.I., Engleman, R. Jr., Hedderich, H.G., Bernath, P.F., Lamb, L.D., & Huffman, D.R. 1991, *Chem. Phys. Lett.*, 176, 504
- Fulara, J., Jakobi, M., & Maier, J.P. 1993, *Chem. Phys. Lett.*, 211, 227
- Gillett, F.C., Backman, D.E., Beichman, C., & Neugebauer, G. 1986, *ApJ*, 310, 842
- Glass, I.S. 1978, *MNRAS*, 185, 23
- Goeres, A., & Sedlmayr, E. 1992, *A&A*, 265, 216
- Guillois, O., Nenner, I., Papoular, R., & Reynaud, C. 1996, *ApJ*, 464, 810
- Hartmann, L., & Apruzese, J.P. 1976, *ApJ*, 203, 610
- Hecht, J.H., Holm, A.V., Donn, B., & Wu, C.-C. 1984, *ApJ*, 280, 228
- Kessler, M.F. et al. 1996, *A&A*, 315, L27
- Koike, C., Hasegawa, H., & Manabe, A. 1980, *Ap. Sp. Sci.*, 67, 495
- Krätschmer, W., Lamb, L.D., Fostiropoulos, K., & Huffman, D.R. 1990, *Nature*, 347, 354
- Lawson, W.A., & Cottrell, P.L. 1989, *MNRAS*, 240, 689
- Lawson, W.A., Maldoni, M.M., Clayton, G.C., Valencic, L., Jones, A.F., Kilkenny, D., van Wyk, F., Roberts, G., & Marang, F. 1999, *AJ*, 117, 3007
- Lee, T.A., & Feast, M.W. 1969, *ApJ*, 157, L173
- O’Keefe, J.A. 1939, *ApJ*, 90, 240
- Papoular, R., Conard, J., Guillois, O., Nenner, I., Reynaud, C., & Rouzaud, J.-N. 1996, *A&A*, 315, 222
- Rao, N.K., & Lambert, D.L. 1993, *AJ*, 105, 1915
- Rao, N.K., Lambert, D.L., Adams, M.T., Doss, D.R., Gonzalez, G., Hatzes, A.P., James, C.R., Johns-Krull, C.M., Luck, R.E., Pandey, G., Reinsch, K., Tomkin, J., & Woolf, V.M. 1999, *MNRAS*, 310, 717
- Rao, N.K., & Lambert, D.L. 2000, *MNRAS*, 313, L33
- Rao, N.K., & Nandy, K. 1986, *MNRAS*, 222, 357
- Rao, N.K., & Raveendran, A.V. 1993, *A&A*, 274, 330
- Roelfsema, P.R., Kester, D.J.M., Wesselius, P.R., Sym, N., Leech, K., Wieprecht, E. 1993 in *Astronomical Data Analysis Software and systems*, II, ed. R.J. Hanisch, R.J.V. Brissenden, & J. Barnes, *ASP Conf. Ser.* 52, 254
- Scott, A., & Duley, W.W. 1986, *ApJ* 472, L123
- Scott, A.D., Duley, W.W., & Jahani, H.R. 1997, *ApJ*, 490, L175
- Stanford, S.A., Clayton, G.C., Meade, M.R., Nordsieck, K.H., Whitney, B.A., Murison, M.A., Nook, M.A., & Anderson, C.M., 1988, *ApJ*, 325, L9
- Stein, W.A., Gaustad, J.E., Gillett, F.C., & Knacke, R.F. 1969, *ApJ*, 155, L3
- Valentijn, E.A. et al. 1996, *A&A*, 315, L60
- Walker, H.J. 1985, *A&A*, 152, 58
- Walker, H.J. 1986, in *Hydrogen Deficient Stars and Related Objects*, ed. K. Hunger, D. Schönberner & N.K. Rao (Dordrecht:Reidel), 407
- Walker, H.J., Heinrichsen, I., Richards, P.J., Klaas, U., & Rasmussen, I.L. 1996, *A&A*, 315, L249
- Waters, L. B. F. M., Cami, J., de Jong, T., Molster, F. J., van Loon, J. Th., Bouwman, J., de Koter, A., Waelkens, C., Van Winckel, H., Morris, P. W. 1998, *Nature*, 391, 868

This 2-column preprint was prepared with the AAS L^AT_EX macros v5.0.