Distant Cluster Hunting II: A Comparison of X-ray and Optical Cluster Detection Techniques and Catalogs from the ROX Survey

Megan Donahue^{1,5}, Caleb Scharf^{1,2}, Jennifer Mack¹, Paul Lee¹, Marc Postman¹, Piero Rosati³, Mark Dickinson^{1,5}, G. Mark Voit¹, John T. Stocke^{4,5}

ABSTRACT

We present and analyze the optical and X-ray catalogs of moderateredshift cluster candidates from the ROSAT Optical X-ray Survey, or ROXS. The survey covers the sky area contained in the fields of view of 23 deep archival ROSAT PSPC pointings, 4.8 square degrees. The crosscorrelated cluster catalogs were constructed by comparing two independent catalogs extracted from the optical and X-ray bandpasses, using a matched-filter technique for the optical data and a wavelet technique for the X-ray data. We cross-identified cluster candidates in each catalog. As reported in Paper I, the matched-filter technique found optical counterparts for at least 60% (26 out of 43) of the X-ray cluster candidates; the estimated redshifts from the matched filter algorithm agree with at least 7 of 11 spectroscopic confirmations ($\Delta z \lesssim 0.10$). The matched filter technique, with an imaging sensitivity of $m_I \sim 23$, identified approximately 3 times the number of candidates (155 candidates, 142 with a detection confidence > 3σ) found in the X-ray survey of nearly the same area. There are 57 X-ray candidates, 43 of which are unobscured by scattered light or bright stars in the optical images. Twenty-six of these have fairly secure optical counterparts. We find that the matched filter algorithm, when applied to images with galaxy flux sensitivies of

¹Space Telescope Science Institute, 3700 San Martin Drive, Baltimore, MD 21218, donahue@stsci.edu

²Columbia University, Columbia Astrophysics Lab, Mail Code 5247 Pupin Hall, 550 W. 120th St., New York, NY 10027

³European Southern Observatory, Karl-Schwarzschild-Str. 2, Garching, D-85748

⁴University of Colorado, CASA, CB 389, Boulder, CO 80309

⁵Visiting Astronomer, Kitt Peak National Observatory

 $m_I \sim 23$, is fairly well-matched to discovering $z \leq 1$ clusters detected by wavelets in ROSAT PSPC exposures of 8,000-60,000 seconds. The difference in the spurious fractions between the optical and X-ray (30% and 10% respectively) can not account for the difference in source number. In Paper I, we compared the optical and X-ray cluster luminosity functions and we found that the luminosity functions are consistent if the relationship between X-ray and optical luminosities is steep $(L_x \propto L_{opt}^{3-4})$. Here, in Paper II, we present the cluster catalogs and a numerical simulation of the ROXS. We also present color-magnitude plots for several of the cluster candidates, and examine the prominence of the red sequence in each. We find that the X-ray clusters in our survey do not all have a prominent red sequence. We conclude that while the red sequence may be a distinct feature in the color magnitude plots for virialized massive clusters, it may be less distinct in lower-mass clusters of galaxies at even moderate redshifts. Multiple, complementary methods of selecting and defining clusters may be essential, particularly at high redshift where all methods start to run into completeness limits, incomplete understanding of physical evolution, and projection effects.

Subject headings: catalogs, galaxies: clusters: general, dark matter, X-rays: galaxies: clusters

1. Introduction: Why Conduct an Optical - X-ray Survey for Clusters of Galaxies?

Clusters of galaxies are the most massive gravitationally-bound systems in the universe. Because they sample the high mass end of the mass function of collapsed systems, they can be used to determine cosmological parameters such as Ω_m (e.g. Donahue & Voit 1999). The largest clusters (~ $10^{15} M_{\odot}$), are the products of the collapse of matter from a very large volume of space ($r \sim 16h^{-1}(\Omega/0.2)^{-1/3}$ Mpc). Therefore they are thought to be "fair samples" of the universe – that the mass to light ratio or the baryonic mass fraction defined within the domain of a cluster of galaxies is representative of that ratio in the universe as a whole. They are purported to be "closed boxes" to star formation and evolutionary processes that occur within their domain. In this paper we present and analyze the catalogs from our joint optical-X-ray search for clusters of galaxies. We conducted this survey in order to provide a sample of clusters to test such assumptions about clusters of galaxies and

to investigate the impact of sample selection on studies of cluster evolution and the evolution of their member galaxies.

Understanding cosmological or galaxy evolution studies of clusters critically requires an understanding of the biases in any sample of clusters. For example, the evolution of the number density of systems with cluster-sized masses as a function of mass and redshift is a fundamental prediction of cosmological structure formation models. To know the number density of clusters, we must know the biases inherent in how we find them, preferably as a function of cluster mass. Furthermore, testing the "fair sample" hypothesis requires reliable and unbiased selection of the most massive clusters.

Ever since Abell (1958) and Zwicky (Herzog, Wild & Zwicky 1957; Zwicky 1961) began publishing catalogues of optically selected clusters of galaxies, the definition of a cluster and the definition of biases inherent in the cluster detection process have been lively topics of debate. In 1978, the launch of the first X-ray imaging telescope, the Einstein observatory, began a new era of cluster discovery, as clusters proved to be luminous (> 10^{42-45} erg s⁻¹), extended ($r \sim 1-5$ Mpc) X-ray sources, readily identified in the X-ray sky. The intracluster gas, in nearly hydrostatic equilibrium with the gravitational potential of the cluster, radiates optically thin thermal bremstrahlung and line radiation. X-ray selection of clusters is more robust against contamination along the line of sight than traditional optical methods since the richest clusters are relatively rare and since X-ray emissivity, which is proportional to the gas density squared, is far more sensitive to physical overdensities than is the projected number density of galaxies on the sky.

One cluster sample differs from another depending on how the clusters were detected. Optical selection of clusters using traditional methods looking for overdensities of galaxy counts (e.g. Abell 1958) was rife with contamination problems. However, modern methods such as the "matched filter" algorithm (Postman et al. 1996, P96 hereafter) provide automated, uniform detection of galaxy overdensities in deep optical images. The matched filter technique searches for local density enhancements in which galaxies follow a magnitude distribution characteristic of that expected for a cluster of galaxies. The results include statistically quantifiable estimates of cluster richness, redshift, and significance. The first X-ray selection method used to construct the Extended Medium Sensitivity Survey (EMSS; Gioia et al. 1990b) was biased somewhat towards selecting clusters with high central surface brightnesses. Now there are several algorithms optimized for detecting extended sources, including wavelets (Rosati et al. 1995) and Voronoi-Tesselation Percolation methods (Scharf et al. 1997).

A decade ago, optical and X-ray surveys apparently disagreed about how much clusters have evolved since $z \sim 0.5 - 1.0$. Optical surveys indicated very little evolution since $z \sim 0.5 - 1$ (Gunn, Hoessel & Oke 1986), but the accurate measurements of survey volumes and cluster properties required for quantitative assessment of this evolution were difficult to quantify in these first high-redshift cluster surveys and the volumes were small so uncertainties were large. X-ray studies suggested modest evolution (Gioia et al. 1990a). The most recently compiled X-ray samples of clusters over a range of redshifts out to $z \sim 0.8 - 1.2$ agree that the X-ray luminosity function for moderate luminosity clusters has not evolved significantly since $z \sim 0.8$ (Borgani et al. 1999; Nichol et al. 1999; Rosati et al. 1998, 2000; Jones et al. 1998), while the most luminous systems, contained in the EMSS, might have evolved somewhat (Henry et al. 1992; Nichol et al. 1997, Vikhlinin et al 1998, 2000; Gioia et al. 2001) or very little (Lewis et al. 2002). More recent optical surveys for distant clusters continue to find very little evidence for cluster number density evolution at moderate redshifts (Couch et al. 1991; P96). The explanation of what may seem like a persisting discrepancy is that if any evolution exists in the X-ray cluster population, it is only occuring in the highest luminosity systems which are also the most rare systems. The optical surveys of Couch et al. (1991) and P96 were too small and shallow to detect the putative evolution of the rarest systems.

While the most recent optical and X-ray results are now at least in statistical agreement on the question of evolution since z < 0.8, the question remains whether both techniques are selecting the same clusters. The fundamental quantity, from the viewpoint of comparison to cosmological simulations, is the cluster's mass. We do not know *a priori* whether optical luminosity or X-ray luminosity should be better correlated with a cluster's mass. The fundamental question, from the viewpoint of "fair sample" techniques of measuring universal ratios, is whether clusters are truly a "fair sample". For example, M/L ratios depend on the bandpass of the light and the star formation history of the constituent galaxies. If the gas fractions or the M/L ratios vary significantly from cluster to cluster they are obviously not representative of the universe as a whole.

X-ray selection is generally thought to be superior to optical selection. Observationally, the hot gas is a larger fraction of the cluster mass than the stellar mass, and the X-ray luminosity of a cluster is far easier to measure than its optical luminosity. For X-ray selected clusters, studies of gas fractions and cluster M/L ratios show that these quantitites are statistically constant (Evrard 1997; Arnaud & Evrard 1999; Carlberg et al. 1996). However, if X-ray selection biases the selection of the clusters, high-mass clusters of galaxies with low hot gas fractions (if they exist) would be omitted from such studies. Massive clusters, under the "fair sample" hypothesis, should have nearly identical baryon fractions, however they are discovered.

With the ROSAT Optical X-ray Survey (ROXS) for clusters of galaxies, we have endeavored to address such issues by obtaining optical images of complete 30' by 30' fields centered on positions of deep ROSAT PSPC pointings. In contrast to previous ROSAT PSPC serendipitous surveys such as those conducted by Rosati et al. (1995, 1998), Jones et al. (1998), Romer et al. (2000), and Vikhlinin et al. (1998), the ROXS includes optical imaging for the entire field of view of each X-ray pointing. The X-ray selection and optical selection of cluster candidates was then done independently of each other. We observed 23 ROSAT pointings for a total of nearly 5 square degrees in I band. For five of these fields we also obtained V-band imaging. In this paper (Paper II) we present the catalogs, survey windowing functions, data reduction and observation details, an analysis of detection likelihoods, as well as an expanded discussion and further analysis, including numerical simulations of the survey. In §2, we describe the X-ray field selection criteria and the optical observations. In §3, we present the optical cluster candidates catalog, crossidentification of clusters in the V and I bands. In §4, we present the X-ray cluster candidate catalogs and the X-ray/optical cross-identification procedure. In §5 we describe properties of the cluster candidates, including the distribution of observed properties of objects in the sample, the estimated richness vs. L_x , the V - I vs Icolor magnitude diagrams for the clusters identified in both the V and I bands. We discuss and summarize our results in §6 and §7 respectively.

For all derived quantities, we have used $H_0 = 75h_{75}$ km s⁻¹ Mpc⁻¹, and $q_0 = 0.5$.

2. Observing Strategy, Observations and Data Reduction

2.1. Field Selection and Observations

The sample of 23 target fields were chosen from a sample of archival X-ray observations. All ROSAT PSPC pointings with exposure times of more than 8,000 seconds and Galactic latitude of $|b_{II}| > 20$ degrees and declination $\delta > -20$ degrees were considered for optical imaging from Kitt Peak. We avoided fields with bright

stars ($m_V < 9$). Even so, bright stars outside the field occasionally scattered light into the field of view and stars with $m_I < 16$ had diffraction spikes. Pixels affected by bright stars and scattered light were masked in our subsequent analysis, and thus the total survey area is 4.8 square degrees, less than the nominal 5.75 square degrees that would have been covered if optical detection success were unaffected by stars. Table 1 summarizes the fields observed as well as the associated data.

We obtained the optical data during two observing runs at the Kitt Peak National Observatory 4-meter telescope during March 1996, and May 1997. We were awarded 4 observing runs for this project, but one (November 1995) was cancelled because the dome mechanism was broken and another had extremely poor weather (November 1996). Therefore, the entire survey was conducted in the northern spring sky. The KPNO 4-meter prime focus CCD camera T2KB has a 16' field of view (0.47" pixel⁻¹). We mosaiced 2×2 of these fields to created full 30' by 30' fields, each centered on a deep ROSAT pointing, overlapping each quadrant by ~ 1'.

A total of 900 seconds of exposure were obtained through the I-band filter for each ROSAT field quadrant. Our 5σ detection limit of I = 23 (Vega magnitude) was sufficient to detect cluster galaxies 2 magnitudes fainter than the typical unevolved first-ranked elliptical at z = 1 (Postman et al. 1998a). For five of the ROX fields, we also obtained V-band data, for a total exposure of 600 seconds for each quadrant. The data were obtained under near photometric conditions and low air masses (< 1.2, typically). When the sky was photometric, we obtained short exposures of the center of each field for calibration purposes, and thus all of our data were fluxcalibrated (see §3.1.1). The filters used were the V and I filters from the Harris set.

2.2. Data Reduction

The CCD images were first reduced with standard IRAF tools for bias level subtraction, frame trimming, and flat-field division. A cosmic ray rejection algorithm based on median filtering techniques was used to remove cosmic rays from the images. In the first observing run, each pointing was divided into two 450 second exposures because the I-band sky was so bright that the dynamic range of the A-to-D converter was swamped. We used the two images to reject cosmic rays. For our second run, the A-to-D converter had been updated to unsigned integers, and the cosmic-ray rejection made possible with two images was determined to be of little advantage compared to the overhead time and additional data-reduction complexity. Data obtained during subsequent telescope visits thus consisted of single 900-second observations. The two 450 second exposures for each quadrant obtained in earlier runs were coadded. We achieved a detection limit of 23 mags in I and 24 mags in V (Vega magnitudes).

To prepare for the galaxy catalog construction, the following processing steps as discussed in P96 were employed. A sky model for each (coadded) frame was created from the data to remove the remaining CCD signatures. We fit the median sky level as a function of row, and subtracted the appropriate sky components from each image row. These steps produced frames with extremely flat sky levels.

3. Catalogs

Here we describe the procedures and assumptions we used to create the galaxy catalog, the matched filter cluster catalog, the X-ray catalog, and the cross-identifications of probable detections of the same physical cluster system.

3.1. Optical Catalog

3.1.1. Galaxy Catalog Construction

We followed the procedures discussed in P96 to construct the galaxy catalog. A modified version of the Faint Object Classification and Analysis System (Jarvis & Tyson 1981, Valdes 1982) was used to detect, measure, and classify objects in the calibrated CCD images. The detection algorithm estimated a point spread function (PSF) for each frame, to compensate for seeing variations from exposure to exposure. To separate stars from galaxies, FOCAS classification parameters were chosen to cover PSF variations but to be robust to star-galaxy distinctions. Perfection in distinguishing galaxies from stars is not critical to our experiment since galaxies dominate the detected object counts in the magnitude range of interest (P96).

Astrometric calibration was performed by selecting unsaturated bright stars. The J2000 celestial positions of the stars were measured from the Digitized Sky Survey (the "Quick V" survey, epoch 1983) based on Palomar Observatory Schmidt plates (Lasker et al. 1990). An astrometric solution was computed based on a 6-term per coordinate polynomial for each field. Typical solution uncertainty is < 0.5". The absolute photometry for Johnson V and Cousins I fluxes was obtained using

Harris V and I-band observations of Landolt Standards fields PG1047+003, 1159-035, SA107, and PG1657+078 (Landolt 1992). For our figures and tables we use Johnson V and Cousins I. Those magnitudes are related to AB magnitudes using the following relations. V(Johnson) = V(AB) - 0.02 and I(Cousins) = I(AB) - 0.45.

3.1.2. Cluster Catalog Construction

The procedures in P96 were followed in constructing the cluster catalog. The algorithm based on the matched filter method, tuned to redshifts in the range 0.2 <z < 1.2, was used to generate cluster catalogs in $\Delta z = 0.1$ intervals. $H_0 = 75h_{75}$ km s^{-1} Mpc⁻¹ and $q_0 = 0.5$ were assumed for all quantities. Clusters were identified by searching for the local maxima within a moving box which was $1.667h_{75}^{-1}$ Mpc across. A candidate cluster was registered when the central pixel in the box was the local maximum and it lay above a prescribed threshold. We used the following cluster detection parameters and cluster assumptions to generate the cluster catalog. The slope of the radial filter profile was 1.40. A cluster core radius of $133.3h_{75}^{-1}$ kpc was assumed, and the smoothed map was sampled every 0.50 core radii. The cut-off radius for cluster detection was set to $1333h_{75}^{-1}$ kpc, with a detection box half-width of $1666.7h_{75}^{-1}$ kpc. The profile kernel halfwidth was 20 map pixels, defined by the 0.50 core radius for each map, the catalog scale was 1 arcsecond/unit, and the CCD scale was 0.45 arcsecond/pixel. For the luminosity function filter we assumed a Schechter function (Schechter 1976) with $M^* = -21.0 + 5 \log h_{75}$ in the V band and $M^* = -21.90 + 5 \log h_{75}$ in the I band with a slope of -1.10 for both bands. Each cluster candidate is assigned a central position, effective radius (corresponding to the area of detection equal to πr_{eff}^2), an estimated redshift based on the best-fit luminosity function, an average galaxy magnitude (I_{mag} or V_{mag} depending on the bandpass), and a detection confidence in units of sigma (σ). The algorithm also estimates an effective optical luminosity Λ_{cl} , which corresponds to the equivalent number of L^* galaxies in the cluster such that the optical cluster luminosity $L_{cl}/L_{\odot} =$ $\Lambda_{cl}L^*/L_{\odot} = \Lambda_{cl}10^{-0.4(M_{\odot}-M^*)}$ where M_{\odot} is the absolute magnitude of the Sun in the appropriate band (P96). The quantity Λ_{cl} is relatively insensitive to H_0 ; variations $\sim 5\%$ were measured if $h_{75} = 1.33$ rather than 1.0. For reference, a richness class 1 cluster at $z \leq 0.7$ had $\Lambda_{cl} = 30 - 65$ in the I-band (P96).

These parameters generated cluster catalogs that are directly comparable to cluster candidates in the Postman et al. (1998b) 16 square degree I-band survey (called "Deeprange"). The similarity in optical methods and instrumentation for the ROXS and the Deeprange surveys allowed us to compare the spurious rates of cluster identification and the reliability of the estimated redshifts with respect to spectroscopic redshifts for the ROXS from the larger Deeprange survey, in addition to the P96 survey.

Table 2 lists the cluster candidates for 23 fields observed. We have extracted the optical selection function as a function of redshift in Figure 1. These probabilities were computed following P96, assuming a K-correction appropriate to a galaxy population dominated by ellipticals.

3.2. V and I band Cross-Identifications

Here we describe the comparison of the cluster candidate catalogs resulting from the matched filter algorithm for the I-band data and the V-band data that we obtained for five of the ROXS fields. We find that 60-65% of the clusters identified in the I band are also detected in the V-band with similar estimated redshifts if we exclude all candidates with high redshift estimates (z > 1).

For five fields (Table 1), we obtained V-band images, with a source detection limit $V_{AB} \sim 24$ magnitudes within an exposure time of 600 seconds. We used the matched filter algorithm, adjusted for the V-band to provide appropriate estimated redshift estimates and Λ_{cl} , to provide an independent sample of optical candidates. In the five fields, we found 46 V-band candidates with a detection confidence of > 3σ and 33 I-band candidates with a > 3σ confidence.

When we cross-identify the candidates from the I-band and the V-band, we find that of the 33 I-band candidates, there are positional matches for 15 candidates with one or more V-band candidates (Table 3). All but one of these 15 candidates also match in estimated redshift ($\Delta z \leq 0.1$). Regions in the V-band images containing 6 I-band candidates were seriously affected by scattered light or bright stars – a more serious observational effect at V than at I. Therefore, a maximum of 27 of the I-band clusters could have been detected in V band. Of those 27, 4 were high redshift candidates ($z \geq 1$), leading to a V-band detection rate of z < 1 I-band candidates of 15/23 or 65% (Table 5). The failure to detect high-redshift I-band candidates in the V-band is not unexpected since the 4000-Angstrom break spectral feature would make the elliptical galaxies in such clusters, even if the clusters are real, very difficult to see in the V-band. Additionally, the matched filter algorithm reliability degrades rapidly for our I-band data at redshifts greater than ~ 1 because that is where the faint end of the luminosity function is severely truncated by the survey magnitude limit. In this regime, corrections for the missing optical light are only mildly reliable and the result is a significantly higher error in the derived Λ_{cl} value.

Correspondingly, of the 46 V-band candidates, 18 had one or more counterparts in the I-band (20 with a detection significance > 2.9σ), and only 2 candidates were undetectable because of stray light in the I-band image, for a detection fraction of 18/44 or 41%. This detection fraction has been diluted by many high redshift V-band candidates with no I-band counterparts (15 with z > 1). We suspect that all of the 17 z > 1 V-band candidates may be spurious. Of the two high redshift V-band candidates with plausible I-band counterparts, one has a loose association with a z = 1 I-band candidate (120436.7+280520, OC4 in the 1202+281 field) and the other (154923+212325, OC3 in the 3C324 field) has an estimated redshift of 0.7 from the I-band data, compared to the V-band estimate of 1.2.

We expect that the reliability of the matched filter method may become poor for the highest estimated redshifts, especially in the V-band for the following reasons. At the highest redshifts, the luminosity function filtering algorithm in the matched filter cluster detection method is only sensitive to the brightest galaxies in a cluster, especially for the V-band where K-correction effects are strongest. Moreover, the contrast of the most distant clusters against the foreground/background galaxy population becomes very weak, especially at bluer wavelengths due to the combination of K-correction effects and the steeper number counts of faint field galaxies (themselves potentially clustered) at bluer wavelengths.

We can therefore provide plausible explanations for many if not all of the failures to cross-correlate between I-band identifications and V-band identifications (Table 4) by a cluster candidate (either in I or V) with a high estimated redshift which may be a warning flag for unreliable or weak detections, or by obscuration or confusion with scattered light or spikes from bright stars.

Our resulting efficiency in cross-identifying clusters in either V-band or I-band is ~ 60-65%, for candidates with estimated redshifts of z < 1 and if we correct for the candidates which are detected through one filter but undetectable in the other because of scattered light. We present a summary of our numbers in Table 5.

We cannot explain the failure to detect 8 I-band cluster candidates of moderate redshift in V. Four of these candidates have possible V-band counterparts with redshifts from the V-band data similar to those derived from the I-band data, but with low detection significance or with centroids separated by > 3 - 4'. These possible counterparts are listed in Table 3. We also cannot explain the lack of an I-band counterpart for 11 V-band cluster candidates of moderate redshift (as in I band, 3 of these candidates have low significance counterparts or widely separated Iband counterparts with similar redshift estimates). We note that while the detection significance of these unmatched cluster candidates is similar to the rest of the sample (not much dynamic range), the estimated richnesses tend to be rather low Λ_{cl} 20–50, mostly \leq 30. We could be seeing incompleteness effects rather than spurious effects at lower Λ_{cl} , but it is possible that some of these unconfirmed cluster candidates are false.

If we count only the most secure cross-identifications, the I- and V-band cross identification statistics imply a spurious fraction of < 35% for matched filter candidates with estimated cluster redshifts of less than one – consistent with the estimates of the spurious fraction of 25-30% from the Deeprange survey (Postman et al 1998b), with simulations done in P96 and in Postman et al (in preparation), and with spectroscopic observations (for z < 0.6) in Postman et al. (in preparation). The spurious fraction for candidates with estimated redshifts larger than one is completely unknown since no candidate with $z \ge 1$ in one bandpass has a secure counterpart in the other bandpass.

Comparison of matched filter results from the I and V band do not address important sources of spurious cluster candidates, such as unbound aggregates of galaxies seen in a "pole-on" filament. Both the I- and V-band data would be contaminated by such structures. For this reason, comparision of the matched filter method with X-ray observations and other cluster detection methods, such as Sunyaev-Zel'dovich or weak-lensing observations and spectroscopy of member galaxies is critical. With ROXS, we make the direct comparison with the X-ray observations; in this paper, we provide the catalogs of optical and X-ray candidates for follow-up with other cluster hunting techniques.

4. X-ray Catalog

Here we describe the creation of the catalog of X-ray cluster candidates and the cross identification of X-ray candidates with optical candidates detected by the matched filter algorithm in the I-band images.

4.1. X-ray Source Detection and Upper Limits for Optical Sources

A wavelet-based technique, described by Rosati et al. (1995), was used to create an catalog of X-ray clusters of galaxy candidates for each field observed at Kitt Peak. Several of these fields overlapped with the orginal ROSAT Deep Cluster Survey (RDCS; Rosati et al. 1995, 1998) sample, and thus the X-ray cluster candidates in some of the fields already have been confirmed and have spectroscopic redshifts. The flux limits for ROXS are approximately the same as those of the RDCS $(F_x > 10^{-14} \text{ erg s}^{-1} \text{ cm}^{-2})$, but of course vary with exposure time in both surveys. Table 6 lists the 57 X-ray candidate clusters and their associated X-ray parameters for each ROSAT field, and the spectroscopic redshift, when available. The X-ray parameters (Table6) measured are a centroid position (RA and Dec J2000), the net number of X-ray photons in the source (Counts), the off-axis angle of the source in arcminutes (Theta), the FWHM of the source in arcseconds (FWHM), the confidence level of the extended nature of the source in σ (Sig-ext), and the X-ray flux and error in the 0.5-2.0 keV bandpass (F_x and eF_x). The comment field includes the spectroscopic redshift, if available. A notation of "d" indicates the source may be a double source. Occasionally, a source has a large flux error, reflecting not the confidence in the detection, but the uncertainty in measuring the flux from an extended, complex object.

A by-product of our analysis was a catalog of all the X-ray point sources in the field down to the flux limit of each ROSAT observation. These sources matched very nearly one-to-one to sources available in the WGACAT (White, Giommi, Angelini 1995), so we do not list them here. We will provide this list on request. In Table 2, we identify the optical candidates without an associated extended X-ray source but with an X-ray point source within 30-60".

For every optical cluster candidate, we estimated the X-ray detection threshold defined by an upper limit to the observed 0.5 - 2 keV flux as the flux corresponding to a 4σ fluctuation above the limiting surface brightness, within a r = 1' aperture, as a function of position in the ROSAT field of view. The variation of X-ray source detection efficiency, arising from the degradation of the Point Source Function (PSF), as a function of off-axis angle can be seen in Figure 2 where we plot the mean number of X-ray sources per unit area in radial bins. There is almost a factor of ten reduction in the source counts at the largest off-axis angles. This reduction corresponds to as much as a factor of 4-5 change in the flux limit, which is taken into account in our upper-limit estimates in Table 2. Using the known, deep, cumulative source number counts in the 0.5 - 2 keV band (e.g. Hasinger et al. 1998) we can map the trend seen in Figure 2 into flux-limit vs off-axis angle. A simple, approximate, linear fit to this yields a relationship of (in units of 10^{-14} erg s⁻¹ cm⁻² (0.5 - 2 keV)): $f_{lim} \simeq A + 0.078\theta$, where A is the flux limit at $\theta = 0'$ and θ is the off-axis angle in units of arcminutes. Since our X-ray detection algorithm provides us with an estimate of the background surface brightness we can normalize the above relationship with the flux of a 4σ , on-axis source in a given aperture, and for a given field. The detection thresholds F_{lim} in units of 10^{-14} erg s⁻¹ cm⁻² (0.5 - 2.0 keV) are reported in Table 2.

We convert the detection thresholds for the ROSAT fields into X-ray detection probabilities, as plotted in Figure 3, as a function of X-ray luminosity (L_x) and redshift, by assuming a canonical L_x -core radius relationship for the cluster surface brightness profiles (e.g. Jones et al. 1998). The cluster surface brightness profiles are assumed to be standard β -profiles with $\beta = 2/3$. The detection probability is computed as the net probability to detect a cluster with a given redshift and L_x in the entire ROXS coverage, based on exposure time, background surface brightness, and radial detection efficiency due to PSF variations discussed in the previous paragraph (Scharf et al. 1997; Rosati et al. 1995).

4.2. X-ray-Optical Cross-Identification

We cross-identified each X-ray candidate and I-band optical candidate by visually inspecting the overlap of the X-ray surface brightness contours and the optical contours of the detection thresholds for filtered optical maps at the nominal detected estimated redshift. Optical and X-ray contour shapes did not match in detail. Cluster candidates which overlapped significantly were identified as cross-identification candidates. We have noted the separation of the nominal optical and X-ray centroids in the tables. The centroid of the optical candidate could be shifted from the true position because of regions contaminated by bright stars, which must be masked from the data. The centroids of both X-ray and optical candidates are likely only to be good to $\pm 0.5'$; cluster diameters at any redshift are usually > 2'. Typical centroid separations are < 2'. An example I-band image quadrant with the X-ray and optical sources labelled is shown in Figure 4.

As reported in Paper I, of the 57 X-ray candidates, 43 would have been visible on our optical frames (that is, not obscured by bright stars or scattered light). Of the 43, 29 were visually identified with potential cluster candidates with centroid separations of $\leq 3'$ or with significant contour overlap, of which 26 are very secure (separations $\leq 2'$ or visual confirmation of X-ray and optical correspondence of somewhat complex structure). The other three less secure identifications are more questionable affiliations of sprawling optical galaxy systems or filaments with a more compact X-ray cluster candidate. One such X-ray candidate, a confirmed cluster at z = 0.167, MS1201+283 or Abell 1455, does not have a formal optical counterpart, but has up to 3 optical possible counterparts based on a visual inspection of the field. The optical field around this cluster was cut-up by bright stars, and the matched filter seems to have detected individual cluster candidates in the remnants.

The remaining 14 X-ray cluster candidates can be divided into 3 categories. Six candidates are bona fide, optically faint candidates, three of which have lowsignificance counterparts ($< 3\sigma$) that are interesting because of their high estimated redshift, but they are not counted as true X-ray/optical coincidences. The other two categories contain sources that have a lower probability of a true cluster: double sources and sources with very uncertain X-ray fluxes ($F_x/\sigma_{F_x} < 3$). There are two double sources without optical counterparts, and six sources with uncertain X-ray fluxes.

Only two of the optically-blank X-ray sources have yet been classified or confirmed at other wavelengths. One source (RXJ1256.9+4720) is tentatively associated with 3C280, the target of the original observation. Here we count this X-ray detection as an X-ray cluster candidate but not an optical cross-identification. It was the original target; this correspondence is the only example of the original target coinciding with a cluster candidate. The statistics are not skewed by the inclusion of this one very interesting cluster candidate. There was a 2σ detection at this position in the optical data. The estimated optical redshift (z = 1.0) was so remarkably close to that of 3C280's spectroscopic redshift (z = 0.997) that we quote the matchedfilter parameters of the optical candidate despite our extremely low confidence in its reality. We note that the 3C280 cluster only has one redshift in-hand, and a preliminary investigation of the Chandra image of this source shows X-ray structure associated with the radio source but no obvious cluster emission (M. Donahue, private communication.) The other candidate, RXJ1120.0+2115, was flagged as a double source; Rosati and his collaborators have found a cluster with z = 0.75 (P. Rosati, private communication).

Since the spurious fraction typical of the X-ray surveys is $\sim 10\%$ (Rosati et al. 1998; Vikhlinin et al. 1998), approximately 6 of the 57 sources are expected to be false – X-ray sources which are not really extended or are constellations of point-like X-ray sources. Some of these candidates may be bona fide, albeit optically-faint

clusters. Truly high redshift clusters are very faint in the I-band because of the 4000Å break in elliptical galaxy spectra. Near IR imaging is probably the best tool to reveal the presence of any high redshift clusters.

Eleven X-ray candidates have confirmed spectroscopic redshifts from Rosati's followup of clusters in the RDCS (Rosati et al., in preparation). We list the candidates in Table 7. All X-ray spectroscopic confirmations which were not obscured by bright stars had an optical counterpart with a detection confidence of $\geq 3\sigma$, except for the X-ray candidates associated with 3C280 at z = 1 and with RXJ1120.0+2115, which now has a confirmed redshift of 0.75 (Rosati, private communication).

The redshift estimated by the matched filter algorithm has proven to be surprisingly good for the clusters of galaxies, at least for the X-ray selected clusters (Table 7). For 7 out of 11 extended X-ray sources with spectroscopic redshifts, the discrepancies between the photometric and the spectroscopic redshift are less than $\Delta z = 0.1$, finer than our redshift grid. From Deeprange spectroscopic followup (Postman, private communication), the mean difference between spectroscopic and estimated redshifts for candidates with $0.3 \leq z_{est} \leq 0.4$ is 0.04 with an rms scatter of 0.07, based on ~ 25 clusters. The scatter for all Deeprange candidates out to $z \sim 1$ is closer to 0.15 (Postman, private communication).

Of the five X-ray cluster candidates contained in the fields with both V-band and I-band data, all five were detected in both the V-band and I-band.

5. Cluster Candidate Properties

In this section we discuss the cluster properties such as X-ray luminosity, estimated for the cluster candidates in our samples. We find that while the cluster candidates in the ROXS sample have X-ray and optical properties consistent with the range of properties in other cluster samples, the clusters are somewhat less X-ray luminous, optically poorer, and have much less prominent sequences of red galaxies in color-magnitude diagrams than the typical rich and massive clusters of galaxies found in all-sky surveys. This sample thus may include cluster candidates that may be missed in pure X-ray, optical, or color searches for clusters. We make what may seem a pedantic distinction here: These clusters may be missed by a search not because the sample is incomplete, but because their properties fall outside the boundary of the sample selection function.

5.1. X-ray Luminosities

Here we show that the cluster luminosities are typical, with estimated bolometric $L_x \sim 10^{43} - 10^{44} \text{ erg s}^{-1}$, and that the upper limits are all $L_x \geq 10^{42} \text{ erg s}^{-1}$. However, a couple of rich optical candidates are found with very low X-ray upper limits.

The X-ray luminosity for each cluster candidate with an optical counterpart was computed assuming the matched-filter estimate for the cluster redshift, the X-ray flux from the wavelet detection algorithm, and a self-consistent bolometric correction commensurate with the X-ray temperature T_x implied by the $L_x - T_x$ relation from Markevitch (1998). The $L_x - T_x$ relation is nearly constant with redshift (Donahue et al. 1999; Borgani et al. 2001). The optical cluster candidates which were not detected in the X-ray band all have X-ray upper limits from a r = 1' aperture (see § 4.1), ranging from $2 - 10 \times 10^{-14} \text{erg s}^{-1} \text{ cm}^{-2}$ (the bolometric corrections for the upper limits were computed assuming an X-ray temperature of kT = 4 keV and the estimated redshifts). A mean $N_H = 2.33 \times 10^{20} \text{cm}^{-2}$ column density for high Galactic latitudes was assumed for all luminosities and upper limits. In Figure 5, we have plotted the distribution of estimated X-ray luminosities of cross-identified clusters and the upper limits of the optical candidates without X-ray counterparts. Note that none of the clusters are extremely luminous $(> 6 \times 10^{44} h_{75}^{-2} \text{erg s}^{-1})$ and that none of the X-ray candidates has an X-ray luminosity upper limit significantly below a $2 \times 10^{42} h_{75}^{-2} \text{erg s}^{-1}$. Since this limit is approximately the luminosity of poor clusters, our X-ray data are not quite sensitive enough to rule out the presence of an X-ray group or low-luminosity X-ray cluster. However, a significant number of the optical candidates (54) have flux limits and estimated redshifts which imply X-ray cluster luminosities which are between $10^{42-43}h_{75}^{-2}$ erg s⁻¹, typical of luminosities of poor clusters of galaxies and even groups (Figure 5). Two such cluster candidates, with redshifts of 0.4 and 0.3 respectively, have $\Lambda_{cl} \geq 50$ (relatively rich) and upper limits on $L_x < 10^{43} h_{75}^{-2} \text{erg s}^{-1}$, clusters (OC1) 1118+2107 and (OC6) 1024+4707 (See Figure 6 for the I-band images of these candidates.) The optical detection significance for each of these two clusters was $\sim 5\sigma$. If these clusters turn out to be X-ray faint (L_x significantly lower than that predicted by the $L_x - T_x$ or $L_x - \sigma_v$ relations for clusters), that would argue that X-ray surveys could miss some relatively massive systems. Such a hypothesis would be straightforward to test with XMM or Chandra X-ray imaging observations, alongside ground-based galaxy spectra to confirm the redshift and establish a velocity dispersion.

5.2. X-ray Luminosity - Optical Richness

Here we show that the X-ray luminosities and optical richness of the ROXS clusters are in the expected range and ratios typical of clusters of galaxies; however, as expected, the ROXS clusters are on average fainter in X-rays than those clusters selected from a very large area X-ray survey at much shallower X-ray flux depths.

In Paper I (Donahue et al. 2001) we evaluated the relationship between the X-ray luminosity L_x and the estimate of optical luminosity Λ_{cl} and while we found a marginally statistically significant correlation between these two properties, independent of estimated redshift, we also demonstrated that this relationship has significant scatter beyond that indicated by the estimated observational uncertainties.

To further investigate the relationship between the X-ray and optical properties of the ROXS we have compared the X-ray luminosities or upper limits and optical richnesses with those obtained from the Abell clusters composing the X-ray Brightest Abell-type Cluster Survey (XBACS) ROSAT survey of Ebeling et al (1996). The XBACS survey is composed of the Abell clusters detected in the northern portion of the Rosat All-Sky Survey, and therefore contains many high luminosity clusters of galaxies. Since such clusters are extremely rare, they are not expected in much smaller surveys such as ours. However, the X-ray luminosities spanned in ROXS are also spanned in the XBACS coverage, albeit for lower-redshift clusters.

We have combined the 283 XBACS X-ray luminosities with the Abell richness number counts N_R from the Abell/ACO catalogue. For comparison to the ROXS, we have converted the matched filter richness parameter Λ_{cl} (approximately equivalent to the number of L^* galaxies in the system (P96)) to Abell richness counts N_R . However, as demonstrated in P96 the scatter between Λ_{cl} and N_R , (where N_R is defined as per Abell's galaxy number counts within a 1.0h⁻¹Mpc radius rather than Abell's 1.5h⁻¹Mpc) is large, albeit with a positive correlation. For our purposes we use a relatively crude conversion from Λ_{cl} to Abell counts; $N_R^{Abell} = (\Lambda_{cl} \pm 23)/0.72$, where the factor 0.72 converts galaxy counts from within a 1.0h⁻¹Mpc radius to a 1.5h⁻¹Mpc radius (P96).

In Figure 7 panel (a) the X-ray luminosity (0.1-2.4 keV, converted to $H_0 = 75$ km s⁻¹ Mpc⁻¹, $q_0 = 0.5$) is plotted against N_R^{Abell} for all XBACS clusters (open circles) and for all ROXS optical candidates with X-ray counterparts (filled circles). The uncertainties in the X-ray luminosities of ROXS clusters ranges from ~ 7% to ~ 80%. In Figure 7 panel (b) the XBACS clusters are again plotted, but now

with the upper limit on X-ray luminosities for all non-X-ray detected ROXS cluster candidates. The resulting relationship between Abell richness number counts and X-ray luminosity for ROXS plotted in Figure 7 has large scatter, a result consistent with our conclusions from Paper I (Donahue et al. 2001) and also with results from Borgani & Guzzo (2001), which we will discuss later.

It is also clear for both the X-ray detected and non-detected objects that the ROXS criteria tend to include more X-ray-poor systems than the Abell/ACO catalogue intersection with the XBACS. The XBACS half-sky survey has sufficient sky coverage and depth to find more luminous clusters, but it is not deep enough to detect proportionally as many X-ray faint clusters as are found in the ROXS survey. As we will also show in our red sequence analysis in § 5.5, the cluster candidates that we sample here are all likely to be somewhat lower in mass than their massive, X-ray luminous cousins found in surveys of larger sky area. The general form of the $L_X - N_R$ relationship of the ROXS cluster candidates is qualitatively similar to that found for the low- L_x clusters in the XBACS,

5.3. Cluster-Point Source Correlation

One goal of our survey was to see whether a criterion of requiring cluster X-ray candidates to have significant extent in the ROSAT PSPC image filtered out clusters with X-ray emission too compact to be unambiguously resolved. Such a criterion could screen out high redshift sources or sources with prominent cooling flows. Since we were imaging complete fields we could investigate whether X-ray point sources, ignored by some cluster surveys, could actually be high-redshift or compact clusters.

Of the 142 optical candidate clusters with a detection confidence of > 3σ , 27 have extended X-ray counterparts (one has two counterparts). Up to twenty-nine additional optical candidates have possibly interesting X-ray point sources either near the central core of galaxies (within 1'-2') or near a possible cluster member. When an X-ray point source is near an optical candidate, we have indicated so in the Comments column of Table 2. For example, in the field of MKN 78, OC2 has a point source very close to it, RXJ0741.7+6525, an active galaxy with a redshift of 1.65 (RIXOS F234_001, with Mg II and CIII emission seen by Puchnarewicz et al. 1997). The estimated redshift of the cluster, however, is 0.4. In the field of 10214+4724, OC10 (z = 0.3) is a compact optical candidate that is exactly coincident with the point source RXJ1025.4+4716; such exact alignments are worth investigating. However, most of the point sources with positional coincidence with optical cluster candidates were not exact; coincidental alignments are likely, given the size of a typical cluster candidate. The number of coincidences between point sources and optical cluster candidates in our survey can be estimated by assuming that a typical optical cluster candidate has a radius of ~ 100", and that the X-ray point source density at the 0.2-2.0 keV flux limit of $F_x \sim 10^{-14} \text{erg s}^{-1} \text{ cm}^{-2}$ is 100 sources deg⁻² (Rosati et al. 1998). With 142 optical candidates, the estimated number of chance coincidences is 34, comparable to the number that we identify. Therefore we do not detect a significant excess of X-ray point sources near optical cluster candidates.

In order to see whether the correspondence with point sources increased when we applied our judgement as to the reality of a cluster candidate, we visually inspected each optical candidate and assessed a subjective believability index to it, of "probable", "blend", or "unlikely". This subjective assessment showed that many of the optical candidates are likely to be blends with other optical candidates at different redshifts. Lacking a purely objective means of sorting out blends, we have listed all blend candidates with notations in the Comments column (Table 2). Some 57 out of the 142 optical candidates without X-ray counterparts passed this subjectivity test; 17 of these may have associated point sources. The fraction of optical candidates without extended X-ray counterparts but with possible X-ray point source counterparts did not increase significantly with this subjective assessment. Therefore, we suspect that most of these associations are likely to be random projections of background AGN with the optical candidates.

We note the fraction of optical candidates which have X-ray counterparts or which are deemed believable in a subjective assessment is a strong function of detection confidence (σ) and estimated redshift. The percentage of X-ray clusters plus "probable" optical clusters is 70% of the total at z = 0.2, dropping to 45% at z = 0.8, and 0 - 20% at z = 1.0 - 1.2 (Figure 8a and 10). The same fraction as a function of Λ_{cl} (Figure 8b) does not vary as strongly as it does with detection confidence (σ) and estimated redshift, presumably since Λ_{cl} is a measure of cluster richness (and richer clusters are in general more powerful X-ray sources and may subjectively look more "probable") but candidates with higher Λ_{cl} also tend to have higher estimated redshifts (and thus are harder to detect by X-ray methods and are less likely to be assessed as "probable" in a subjective review of the optical data.) However, only a few optically selected clusters with $\Lambda_{cl} < 30$ were considered "probable" (Figure 8b). Our statistical analysis in Paper I was only for cluster candidates with $\Lambda_{cl} > 30$. of > 3σ . The presence or absence of this high redshift candidate does not affect our analysis in this paper.

5.4. Redshift, Λ_{cl} , and Detection Confidence Distributions

Here we provide an analysis of the detection limitations of our dual band surveys. We compare the properties of the X-ray detected cluster candidates with those which had no X-ray counterparts; we find that the optical and X-ray surveys were well-matched in terms of depth and sensitivity.

As described previously and in Paper I, each optical cluster candidate was assigned an estimated redshift and a Λ_{cl} from the best-fit luminosity function by the matched filter algorithm. The detection efficiency of the matched filter, when confronted with the limitations placed by the depth of the I-band images and the spectrum of cluster galaxies, drops dramatically at z > 0.5 - 0.6 for clusters with $\Lambda_{cl} \leq 50$. Optical clusters with Λ_{cl} of 100 or more are detectable beyond $z \sim 1$ (Figure 1). This detection efficiency can be demonstrated in the distribution of estimated redshifts for our sample. The redshift distribution of all of the candidates is shown in Figure 8.

The estimated richness parameter from the matched filter algorithm, Λ_{cl} , may be corrected for aperture effects by multiplying the original Λ_{cl} by $(1 + z^9)^{0.7}$ (P96). We do not use this correction since it does not significantly correct the Λ_{cl} for cluster candidates at z < 1. For comparison, the distribution of cluster candidates with Λ_{cl} is plotted in Figure 8b. The distribution of redshift-corrected Λ_{cl} is plotted in Figure 9.

The relative depth of the X-ray and matched filter selection techniques and respective flux sensitivities can be assessed by looking at the distribution of the optical detection significance from the matched filter algorithm in Figure 10. If the optical detection threshold were not 3 but 4σ , nearly half of the optical/X-ray detections would be excluded. Therefore, in terms of depth, the detection constraints of the two surveys are compatible. In Figure 11, we plot the relationship between the detection significance and the estimated Λ_{cl} . This Figure demonstrates that Λ_{cl} and detection significance σ are not correlated.

In order to see how much the depth of the X-ray exposure affected the fraction of clusters detected in the X-ray (or in the optical), we divided the ROSAT exposures into two subsamples with roughly equal number of fields in each, one with $t_{exp} \leq$

25,000 seconds and the other with $t_{exp} > 25,000$ seconds. The fraction of optical sources which are detected in the X-ray was statistically equal in both samples, at 0.21 ± 0.05 and 0.24 ± 0.07 respectively. The fractions of X-ray cluster candidates within the optical coverage (not obscured by bright stars or scattered light) which were also detected in the optical in those two samples are 0.75 ± 0.23 and 0.68 ± 0.25 respectively. The uncertainties reflect only the Poisson statistics (\sqrt{N} uncertainties in each sample). We conclude that the exposure length of the X-ray observation did not make a statistically significant effect in the fractions of cross-identified sources, at least for the limited dynamic range of exposure times in our survey coverage.

Even for the deepest X-ray images, the optical survey methods failed to find fewer than 1-2 X-ray cluster candidates in a field, a number that could be further reduced by spurious or false X-ray clusters (~ 10% of the X-ray cluster candidates in our sample are probably spurious, being overlapping point sources). We suggest that the matched filter algorithm applied to images with flux sensitivies of $m_I \sim 23$ is fairly well-matched to discovering the $z \leq 1$ X-ray clusters detected by wavelet analysis of Rosat PSPC exposures between 8,000 and 60,000 seconds.

5.5. Galaxy Color Distributions - Red Sequence

The identification of a red sequence of old, evolved galaxies in a color- magnitude diagram has been proposed as another method to identify cluster candidates. The application of this method has resulted in the detection of high redshift clusters (e.g. Gladders & Yee 2000; Stanford et al. 1997). Here we examine the clusters detected in both I- and V-band to see how many of these candidates show a distinct red sequence and whether the clusters with X-ray counterparts showed strong red sequences. We reserve a direct application of the red sequence selection of clusters for a future paper. Utilizing the V and I band imaging, we determined the limiting magnitude for each field and then matched galaxies by their sky positions in the FOCAS catalogues. We then extracted magnitudes for each galaxy, measured in a fixed 4.7" radius aperture, and we calculated the V-I colors.

If there was a corresponding X-ray counterpart, we used the X-ray contours to define the boundary of the cluster, and we plotted colors for galaxies in this region. If no matching X-ray cluster existed, we compared the cluster centroid positions in V and I to determine the optical cluster location. We also compared the 3σ optical detection contours, since this threshold was the criteria for optical cluster detection. Often, however, this threshold encompassed too large an area (~ 15 to 50 square

arcminutes). When available, we also compared the areas defined by the 4 and 5σ contours to improve the location of the core of the optical cluster. Generally galaxy colors were measured in a sky area of ~ 5 square arcminutes or smaller, depending on the redshift of the optical cluster candidate.

In Figures 12-15, we present the galaxy color-magnitude diagrams for the background galaxies and the candidate clusters, along with the size of the cluster aperture, the estimated redshifts from the I (V) band data, the optical detection significance in I (V), and the corresponding value of Λ_{cl} in I (V). We overplot the color magnitude relations for 7 different redshifts between z = 0.1 - 0.7, corrected to the Landolt (1992) system (Johnson V and Cousins I) from the AB V and I magnitudes used in Gladders & Yee (2000). The darkest line represents the estimated redshift from the matched filter output for the V and I data. The uncertainties in I and V-I are relatively small for I < 22 - over a magnitude above our completeness limit, so we are confident that our photometry is sufficiently deep and accurate to show a red sequence at least for clusters with z < 0.7 - 0.8. However, visual inspection shows that very few of the clusters has a particularly strong red sequence - even the confirmed X-ray clusters have fewer than 10 galaxies near the expected red sequence lines.

To quantify statistically our visual assessment of the color-magnitude plots, we constructed a two-dimensional two-sample Kolmogorov Smirnov test (Press et al. 1997) in order to estimate the probability that the photometric V-I and I values of the possible cluster galaxies and the field galaxies were simply drawn from the same parent distribution. We list the results in Table 3, for all cluster candidates where analysis was possible. For each field we list the number of galaxies in the field population, typically ~ 2000 galaxies in each. The column N_{gals} lists the numbers of galaxies in the cluster candidate sample and the column P(KS) contains the probability that the V-I and I values in the cluster candidate sample were randomly drawn from the same parent population as the field galaxies, as plotted in the first panel for each of the 5 ROSAT fields. Only 7 out of the 16 I-band cluster candidates showed a KS probability lower than 2%. The K-S test is not a test of the presence of a red sequence; but a lack of a significant K-S result for over 50% of these candidates, including 50% of the candidates with X-ray counterparts, reflects the qualitative difficulty of discerning a red sequence in these data. One of the 5 X-ray candidates had only limited galaxy photometry data; 2 of the remaining 4 X-ray candidates had high and thus insignificant K-S probabilities distinguishing the colors and magnitudes of galaxies in the core from those of background galaxies. The result of the K-S test confirms our sense that not many of the cluster candidates

exhibit a strong red sequence on the color-magnitude plots.

This result suggests that for many of the ROXS candidates with both a V and an I detection, a red sequence is not particularly prominent, even for cluster candidates where both I and V-band counterparts exist nearby an X-ray cluster candidate. If these candidates are true clusters – and it seems unlikely that these triply-identified candidates are spurious – this result implies that searches for a prominent red sequence may miss clusters of galaxies at any given redshift, at least at the mass scales sampled by this survey. The red-sequence method and its creators do not claim to create a sample of clusters with galaxy populations representative of all clusters at a given redshift, since the method selects clusters by their old galaxy populations. (Furthermore, the red sequence method may not require a prominent red sequence.) Therefore the ROXS result may not be surprising since none of the ROXS candidates are particularly X-ray luminous. The observed 0.5-2.0 keV X-ray luminosities of the X-ray candidates with V and I counterparts lie between $3-70 \times 10^{42} h_{75}^{-2}$ erg s⁻¹, which places them in the category of richness class 0 clusters and even groups; we note however that the estimated Λ_{cl} for these cluster candidates are as high as 70-85, not particularly consistent with groups.

A prominent red sequence may preferentially exist in the most X-ray luminous clusters and groups, which are the most likely to be dominated by elliptical galaxies in their cores. X-ray luminous groups with central ellipticals and prominent elliptical galaxy populations are more X-ray luminous than their spiral-dominated counterparts (Mulchaey & Zabludoff 1998). Since the ROXS cluster candidates are not very X-ray luminous (whether they were detected in the X-ray or not), they might have a higher spiral component and are more like the Virgo cluster $(L_x \sim 10^{43} \text{erg s}^{-1})$ than they are like Coma ($L_x \sim 10^{44} \mathrm{erg \, s^{-1}}$). Stanford, Eisenhardt & Dickinson (1998; SED98) looked at various properties of the color-magnitude relation for early-type galaxies in high-redshift clusters, including the scatter of those properties, and found that for most of the SED98 clusters there was little variation in either the properties or their scatter. However, all of the SED98 clusters are more luminous than the ROXS clusters in our color-magnitude study. Therefore if the explanation for the ROXS results is that there is an X-ray luminosity threshold below which the colormagnitude properties break down, the SED98 study would not have seen it. A red sequence may be the most distinct at the highest masses and at the lowest redshifts, but at present it is uncertain at what mass and at what redshift the sequence may begin to be indistinct and difficult to see in clusters.

5.6. Simulations of ROX

In Paper I, we produced a Λ_{cl} function based on our data and, using the $L_x - \Lambda_{cl}$ relation, we compared that function to the X-ray luminosity function for clusters of galaxies. We found that a steep $L_x - \Lambda$ relation (steeper than the predicted $L_x \propto \Lambda_{cl}^2$) was needed to explain the observed X-ray and optical luminosity functions for the ROXS clusters. The proportionality, $L_x \propto \Lambda^2$ derives from the observed $L_x \propto T_x^3$ relation for clusters, an assumption of a constant M/L ratio for clusters, and $T_x \propto M^{2/3}$ from hydrodynamics simulations. We also demonstrated in Paper I that unless our observation errors are underestimated, the scatter in the observed $L_x - \Lambda$ relation is large and intrinsic.

We have performed a numerical simulation of the ROXS, accounting for all major selection and measurement effects. We confirm the basic empirical results, namely that the instrinsic cluster population must have both a moderately steep richness to X-ray-luminosity relationship and significant physical scatter between richness and X-ray luminosity in order to reproduce the survey results. In this section we briefly describe the simulation methodology and analysis.

Using Monte-Carlo techniques we first generate a large X-ray cluster population $(5 \times 10^4 \text{ members})$ occupying a volume from z = 0 to z = 2. We draw members according to the local X-ray luminosity function (0.1-2.4 keV) of Ebeling et al (1997), which we assume to be unevolving with z. Current constraints show at most weak (less than 10%) evolution in the cluster population to redshifts of at least z = 0.5 (cf. Gioia et al 2001; Lewis et al. 2002). Since ultimately very few clusters will be selected at z > 1 by the ROXS criteria we consider a non-evolving X-ray luminosity function to be a valid assumption for our first-order goals.

Assuming a optical to X-ray luminosity relationship of the form $\Lambda = AL_x^{\alpha}$ and an instrinsic scatter expressed by a normally distributed random variable ΔA where the amplitude is either fixed or a fractional value of A, we assign a Λ to each cluster. Via a Monte-Carlo approach we then sample this cluster population and apply *both* the X-ray selection criteria and optical selection criteria of the ROXS until a total of N clusters have been "observed". The X-ray detection criteria are slightly complicated by the known dependency on the combination of cluster angular extent and net flux. Using an empirical relationship between cluster luminosity and X-ray core radius (assuming a standard King profile for cluster density) of: $r_c = 0.125(L_x h^{-2}/1.25 \times 10^{44} \text{ erg s}^{-1})^{0.2}$ Mpc h⁻¹ (Jones et al 1998), we derive the angular size of the X-ray cluster. The X-ray selection functions for the simulation are essentially the same as the X-ray selection functions in Figure 3 in § 4, except for the simplifying notion that all clusters were observed on-axis by the ROSAT PSPC. Cluster luminosities are converted to the X-ray detection band (0.5-2 keV) assuming a constant 4 keV thermal spectrum, which is a reasonable approximation for our observed sample.

Clusters are entirely discarded which fall outside of the 0.2 < z < 1.2 range of the ROXS. Clusters which are only detected in either the X-ray or optical, but not both, are flagged. Those detected only in the optical are then assigned an X-ray flux upper-limit (and X-ray luminosity upper limit) based on the mean X-ray flux limit of the ROXS (~ 10^{-14} erg s⁻¹ cm⁻² 0.5-2 keV). Finally, the Λ_{cl} for each cluster is assigned an observational error as a function of redshift, based on the known error distributions of the matched-filter algorithm, which correspond to ~ 10% errors for z < 0.7 and ~ 50% errors for z > 1 (P96).

We are therefore left with four datasets: joint X-ray and optical detected clusters, optically detected clusters with X-ray upper limits, X-ray detected but not optically detected clusters, and entirely undetected clusters.

In Figure 16 we plot the results for a simulated dataset where $L_x \propto \Lambda^2$ (the fiducial relationship expected under basic assumptions) and that the $\Lambda - L_x$ scatter is zero. Observational errors are still included in Λ , so some spread is still seen. The total number of systems is arbitrary, but chosen here for clarity of presentation. It is immediately apparent that this particular simulation does not qualitatively agree with the observations, both in the shallowness of the relation and in the scatter between L_x and Λ . The relative numbers of joint, only optical, or only X-ray detections are also in strong disagreement with observations. We observe approximately 4 cluster candidates in the ROXS which are only detected optically for every 1 cluster which is jointly detected, and no more than 3% of all candidates are solely X-ray candidates. In the simulation of Figure 16 the mean numbers of detections (over 10 runs) yields a straight 1:1 ratio between jointly detected and optically detected clusters, at odds with the real ROXS, although the fraction of the purely X-ray detected numbers are consistent and are at about the 3% level. Adding a significant intrinsic scatter ($1\sigma = 28\%$ of the normalization A) to the $L_x - \Lambda$ relationship does not alter these results.

In contrast, in Figure 17 we plot the results for $L_x \propto \Lambda^3$ and an intrinsic scatter of $1\sigma = 39\%$. It is apparent that there is much better qualitative agreement between this simulation and the ROXS results. Furthermore, the relative numbers of X-ray vs. optical cluster detections are now in much better agreement. The simulation yields a 4 to 1 ratio for optical detections vs joint detections, and a pure X-ray detection rate of 1-2%.

Interestingly, a steeper relationship $(L_x \propto \Lambda^4)$ appears to be less successful at reproducing the ROXS observations, even when we adjust the allowed scatter. Typically, the number of optical-only detections are overproduced by at least a factor of 2 in such tests.

We note here that the simulations are seriously limited in their ability to provide a faithful reproduction of the actual ROXS data. This limitation is for a variety of reasons. Without much greater complexity we cannot mimic the details of the X-ray selection, namely the variation in detection sensitivity as a function of off-axis position in the ROSAT fields, and the detailed variations in field exposure times and backgrounds. We are also reliant on the Λ selection functions, which are known to suffer increasing uncertainties with z. The basic findings are however clear. The $L_x - \Lambda$ relationship must be steeper than Λ^2 , and the intrinsic scatter between L_x and Λ must be large - these are *not* due to systematics in the observations.

A somewhat more quantitative evaluation of the simulation results is summarized in Figure 18. We have applied a 2D K-S test (Press et al 1997) to the ROXS joint detections (X-ray and optically detected clusters) and the equivalent simulation outputs for a range of parameters in the $\Lambda = AL_x^{\alpha}$ relation. Specifically, we have chosen a fixed instrinsic fractional dispersion in this relation such that the 1σ width is 100% when the normalization A is 50 and approximately 30% when A = 175. We have then varied A and set α to 1/2, 1/3 and 1/4. In Figure 18 (upper panel) we plot the resultant 2D K-S probabilities. These should be considered only in relation to each other rather than as absolute indicators of a 'goodness-of-fit'. The higher the probability the more significant the agreement between distributions. It is apparent that when $\alpha = 1/2$ the agreement between the ROXS and the simulations is never very good, but does seem to peak around A = 100. In constrast, for $\alpha \leq 1/3$, the agreement appears best for low A, comparable to $\alpha = 1/2$ for $a \simeq 100$ and significantly worse for A > 120. Variations as large as 50% in the intrinsic scatter used in the simulations do not significantly alter these results.

However, as described above, visual inspection of the data in the $L_x - \Lambda$ panel reveals that the 2D K-S test may be somewhat misleading, and in fact neither the low- α results at low A or the $\alpha = 1/2$ results at $a \sim 100$ would be classed as very good fits 'by-eye'. For example, when $\alpha = 1/2$ the simulated and observed distributions overlap only at the lower L_x end, which boosts the K-S probability relative to $\alpha = 1/3$, although the latter distribution is better aligned, its overlap is smaller. The situation is considerably clearer when we also take into account the number ratios of joint detections versus optical only detections (as discussed above). In Figure 18 (lower panel) we plot the ratio of the number of joint detections to the number of optical-only detections for the 3 cases described above. We also plot the actual ROXS ratio (dashed line). It is immediately apparent that only the $\alpha = 1/3 - 1/4$ curves intersect the observations for reasonable A. Weighing the 2D K-S results by this additional constraint we conclude that the simulations are in best agreement with the data when $\alpha \simeq 1/3 - 1/4$ (since the KS probabilities are overall higher) and when $A \simeq 100 - 150$. $L_x \propto \Lambda^2$ is therefore excluded as a plausible intrinsic relationship.

6. Discussion

The optical matched-filter selection technique works well to find candidate clusters of galaxies, but the spurious fraction is high at $\sim 30\%$, and it is demonstrably not 100% complete - it misses clusters at a rate of 10-20% at least. The selection window for a matched filter cluster sample is more difficult to quantify because of its sensitivity to the spectral energy distribution of the galaxies for which it is searching.

On the other hand, the X-ray selection technique, while generating fewer spurious cluster candidates (~ 10%) and reliably finding the (apparently) more massive clusters, misses some high Λ_{cl} cluster candidates. We do not know yet whether these are true massive clusters or whether they are fortuitous projections of less massive systems. Redshifts and X-ray observations are required to determine the nature of these candidates. Zabludoff & Mulchaey (1998) and Mulchaey & Zabludoff (1998) showed in their sample of 12 optically selected groups that groups without detected X-ray emission tend to be lower velocity dispersion systems with few or no elliptical galaxies. The undetected groups in the Zabludoff & Mulchaev sample may not even be bound. Some X-ray observations of optically-selected, high redshift clusters have revealed such clusters to be "underluminous" in the X-ray (Castander et al 1994.) The high-significance, optically rich ROXS candidates without X-ray counterparts are prime targets for observational tests whether X-ray selection indeed selects on the basis of cluster mass. The correlation of the presence of a dense, X-ray emitting intracluster media and the presence of a significant elliptical galaxy population is another testable hypothesis with further observations of the ROXS sample. With the limited color information we have in hand, we were not able to show that X-ray cluster candidates at the moderate X-ray luminosity levels available in our sample

are more likely to have red elliptical sequences. But we were only able to make this test for a small number of cluster candidates, given the available photometry.

A related result from ROXS is that the cluster richness N_R is not well-correlated with cluster X-ray luminosity, a result which is supplementary to our result in Paper I regarding cluster X-ray luminosity and the Λ_{cl} richness parameter. The clusters in our sample are on average poorer and less massive than those found in a halfsky survey (e.g. XBACS), owing to the smaller sky area and greater depth of our survey. For $\Lambda > 30$, Donahue et al. (2001) suggested that the joint redshift and Λ_{cl} distribution arises from the population of X-ray clusters with a steep dependence between L_x and Λ_{cl} . Equivalently, we show here that we can also reproduce this joint distribution with a large scatter between L_x and Λ_{cl} in Monte-Carlo simulations of the ROXS.

A correlation between cluster X-ray luminosity and cluster richness has been suspected at least since Jones & Forman (1978) plotted cluster richness class against X-ray luminosity for nearby (z < 0.07) clusters. But even in their sample, a 3×10^{44} erg s⁻¹ cluster is equally as likely to have a cluster richness of 0 as it is 2 or 3. Only the most luminous clusters ($L_x > 10^{45}$ erg s⁻¹) had cluster richnesses reliably of 2 or 3. The XBACS clusters (Figure 7) show this correlation and very large scatter with cluster richness. However, richness has long been a suspect observational parameter for X-ray astronomers (e.g. Mushotzky et al 1978), since the measurements rely on number counts that only become more difficult to ascertain at higher redshift with the associated higher contamination levels. The matched filter method provides a possibly more robust and objective means of estimating a cluster candidate's luminosity in the form of Λ_{cl} , yet, this measure too is not strongly correlated with cluster X-ray luminosity.

The lack of strong correlation between X-ray and optical luminosity, and the large scatter of those two quantities, are confirmed by the ROX Survey. A review by Borgani & Guzzo (2001) showed that the velocity dispersions and cluster optical luminosities of an optically-selected sample of clusters are not as well-correlated as the velocity dispersions and the optical luminosities of a subsample of clusters selected for their X-ray fluxes from the original sample. The velocity dispersions of these X-ray selected clusters are even better correlated with their X-ray luminosities, suggesting that X-ray luminosity is better correlated with cluster mass than is optical luminosity. The ROX Survey has no such independent test of correlation with mass; however, followup observations of this sample in the X-ray and the optical will further test the Borgani & Guzzo suggestion at higher redshifts.

The ROXS, since it is a uniformly selected sample at both X-ray and optical wavelengths, has examples of both X-ray candidates without optical counterparts and optical candidates without X-ray counterparts. Both subsamples probe the extremes of the L_x/L_{opt} distribution. Explaining the extremes will go a long way towards explaining why and how the emissivity of the intracluster gas is related to the light emitted by the stars in the cluster. It is particularly important to see if there is a "third parameter" such as galaxy formation efficiency, evolutionary stage of the cluster, or galaxy morphology populations, that creates the large spread in the L_x versus richness relationship for clusters, or if the large spread is due to projection effects significantly impacting the optical selection technique. One possible clue is the existence of a difference in the conclusions based on cluster contents from an optically-selected, rather heterogenous sample of clusters (the MORPHS sample, in Smail et al 1997) and those found in an X-ray selected sample of clusters by Ellingson et al (2001). The X-ray selected clusters show cores that exhibit very little evolution between z = 0.5 and the present, whereas the MORPHS clusters show evidence that the S0 population may be turning into ellipticals during that same time. It is possible that the presence of a well-developed intracluster medium is ubiquitous in a massive cluster, but in less massive clusters, the galaxy populations are still evolving in the cluster cores. Uniform morphological studies of a cluster sample diverse in X-ray and optical properties are needed to test such a statement, and such a test could be accomplished with a sample such as the ROXS.

Rosati et al. (1995, 1998) selected clusters for the RDCS not only for their X-ray emission, but used an extent criteria as well. There was some worry that this selection criteria may have missed the most compact X-ray clusters which remain unresolved by the ROSAT PSPC, particularly those at high redshift. However, our survey does not reveal an obvious population of high redshift cluster candidates with X-ray point sources. The correspondence between an optical cluster and an X-ray point source is not more than we would expect by chance - therefore we do not believe we have found a population of cluster sources which would go undetected by the ROSAT cluster surveys which select for extent, a result consistent with the findings of the Wide Angle ROSAT Pointed Survey (WARPS; Jones et al 1998), a ROSAT serendipitous search which did not filter for extent. We have some examples of optical clusters with X-ray point source counterparts which could be AGN in those clusters. Approximately 25 of our candidates have X-ray point sources within 1'-2' of the cluster centroid; optical identification of these sources is required to ascertain whether the candidate and the point source are physically related.

Probably the most surprising result is that the ROXS cluster candidates do not,

as a rule, show distinct or strong red sequences in the color-magnitude diagrams of their galaxy populations. Only about 50% of the optical cluster candidates detected in both I- and V-band show galaxy colors distinct from those of the field population. And only $\sim 50\%$ of the X-ray clusters so identified show such sequences. It is possible that the red sequence is ubiquitous in the most massive, virialized clusters. But perhaps at some unknown threshold in mass or level of virialization or luminosity, the red sequence ceases to be distinct and prominent. We plan to use the red sequence detection method (Gladders & Yee 2000), which does not rely on an obvious red sequence to select cluster candidates, to see what cluster candidates it finds in the ROXS galaxy photometry data.

The ROXS sample identifies several problems that warrant further investigation:

- A red sequence is not prominent in all X-ray selected clusters. At what X-ray luminosity (or equivalently at what mass) does the red sequence cease to be prominent? Is the presence of a red sequence dependent on mass?
- The estimated optical luminosity and richness of a cluster are not at all wellcorrelated with the X-ray luminosity. There are at least three unsolved issues regarding the correlation of the optical luminosity, X-ray luminosity, and cluster virial mass:
 - 1. Is the lack of correlation representative of the difficulties of estimating the optical luminosity or is it representative of an intrinsic variation or scatter in the M/L ratios in clusters?
 - 2. If the estimated optical or X-ray luminosity, and therefore the detectability of a cluster, has enormous scatter with respect to the mass of the cluster, selection by mass using optical or X-ray light may be very difficult. Knowing the scatter with respect to cluster mass with both optical and X-ray luminosity is important. Individual studies show good correspondence of both quantities with velocity dispersion (e.g. Girardi et al. 2000 and Mahdavi & Geller 2001 for L_B and L_x respectively), but a comparative analysis of the masses and luminosities of a well-chosen, homogeneous sample has not been done yet.
 - 3. If there is such a large intrinsic variation in the optical M/L for clusters, what evolutionary process controls this variation? Can galaxy formation efficiency or evolutionary history be so radically different from one cluster environment to another to produce this effect? Does M/L correlate with any other property of a cluster of galaxies?

• While galaxies and hot gas may trace the gravitational potentials of the most massive clusters, mergers and other physics may disrupt that correspondence. True optical condensations of galaxies with low levels of X-ray emission may be galaxies which have been temporarily separated from the hot intracluster gas while undergoing a merger event; galaxies experience the merger as a non-collisional fluid while the gas experiences hydrodynamic effects of shocks and cooling. An observational example of such a merger is Abell 754 (Zabludoff & Zaritsky 1995). Studies of the ROXS "X-ray poor" cluster candidates may reveal their true nature, whether they are mergers, minor systems embedded in filaments, or mere projection effects.

Because the ROXS has greater depth and smaller sky area than surveys like the EMSS or XBACS, the ROXS clusters must be relatively low-mass clusters compared to typical clusters in those surveys. The low-mass end of the cluster mass function may be where the physics of the intracluster medium collides with the physics of star formation and galactic winds. The energy injected by stellar processes is closer to the specific energy per particle in the gravitationally bound gas of a low mass cluster of galaxies. Mergers may be more common or at least more significant in the lowmass clusters – when the relative velocities of the galaxies are not much greater than the internal velocities of the galaxies, collisions and interactions are more likely to induce star formation. The least massive clusters may be the most recent members to the cluster hierarchy, whose galaxies are most recently accreted from the field. The optical and X-ray properties of such clusters may thus still be correlated, but only very weakly and with high scatter. Such recent formation may explain the weakness or lack of a red sequence, which may only dominate in clusters where the oldest ellipticals formed preferentially in high density regions at an early time. The other inference possible from ROXS is that at some point below a given mass on the mass scale of clusters, some treasured assumptions about clusters, their M/Lconstancy, the invariance of the baryon fraction in clusters, and their ubiquitous red galaxy content may break down.

7. Summary

No single cluster selection method is perfect. Multiple, complementary methods of selecting clusters may be essential for cluster studies, particularly at high redshift where all methods start to run into completeness limits, incomplete understanding of physical evolution, and projection effects. Understanding the biases inherent in any cluster selection method is essential to extracting conclusions about cluster evolution or the evolution of galaxies in clusters from any cluster sample. The ROSAT Optical X-ray Survey (ROXS) for clusters of galaxies provides a well-matched, head-to-head comparison of two of the most popular methods for finding clusters of galaxies: a matched filter in the optical and extended source detection algorithms such as the wavelet transform in the X-ray. Both algorithms do seem to agree on the location of at least some of the cluster candidates in our sky coverage; however, the overlap was far from perfect.

We find that the correlation is weak between the X-ray and optical luminosity or richness of a cluster, and it has considerable intrinsic scatter. Some of this scatter may arise from underestimating our observational uncertainties, but the scatter is too large for it to be accounted for by pure observational uncertainties in L_x or Λ_{cl} (Paper I). We confirm these results in a numerical model of the ROXS. While we did not find any bias that would affect the conclusions on the evolution of clusters of galaxies, we did find evidence for possible bias that could affect conclusions regarding galaxy evolution in clusters. X-ray and optical cluster detection techniques sample different cluster populations, a statement which should worry those who use cluster samples to make claims about galaxy evolution.

We find that the red-sequence does not appear to be very strong in our cluster candidates, regardless of whether they have X-ray counterparts. We plan to do a proper red sequence detection with the ROXS galaxy photometry and position data in an upcoming paper.

All cluster methods have their merits; the cluster community should consider, however, the effect of cluster selection and the measurement of cluster properties on their conclusions. The ROXS has provided a number of follow-up avenues to assess in detail the real differences between cluster candidates which were detected in the X-ray and those which were not. Observations to obtain the weak lensing signatures, velocity dispersions, and/or ICM temperatures would provide the most fundamental property of a cluster from a cosmologist's perspective - its virial mass. Confirmation of at least a subsample of the optical clusters with the faintest X-ray limits would more strongly test our working hypothesis that we have no need to invoke the presence a bi-modal population of X-ray faint and X-ray luminous clusters of galaxies at the mass scales of $10^{14} - 10^{15} M_{\odot}$. Clusters whose X-ray/optical luminosities lie outside the normal distribution could reveal clusters in early evolutionary states or clusters embedded in unusual projection with respect to filaments and other large scale structure, or they may represent a fundamental challenge to our assumptions about cluster physics and formation.

The authors would like to acknowledge the support of the NASA LTSA grant NAG5-3257 and a STScI DDRF grant 82208. This research has made use of data obtained from the High Energy Astrophysics Science Archive Research Center (HEASARC), provided by NASA's Goddard Flight Center, in particular the ROSAT observation catalogs and data, and the NASA/IPAC Extragalactic Database (NED) which is operated by the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration. JTS acknowledges support of a NASA/Astrophysical Data Products grant NAG5-6936.

REFERENCES

- Abell, G. O. 1958, ApJS, 3, 211.
- Arnaud, M., et al. 1992, A&A, 254, 49.
- Arnaud, M. & Evrard, A. E. 1999, MNRAS, 305, 631.
- Borgani, S., Rosati, P., Tozzi, P., Norman, C. 1999, ApJ, 517, 40.
- Borgani, S., et al. 2001, ApJ, in press, (astro-ph/0106428).
- Borgani, S. & Guzzo, L. 2001, Nature, 409, 39.
- Carlberg, R. G., Yee, H. K. C., Ellingson, E., Abraham, R., Gravel, P., Morris, S., Pritchet, C. J. 1996, 462, 32.
- Castander, F. J., Ellis, R. S., Frenk, C. S., Dressler, A., Gunn, J. E. 1994, ApJ, L79.
- Couch, W. J., Ellis, R. S., MacLaren, I., Malin, D. 1991, MNRAS, 249, 606.
- Donahue, M. & Voit, G. M. 1999, ApJ, 523, L137.
- Donahue, M., Voit, G. M., Scharf, C. A., Gioia, I. M., Mullis, C. R., Hughes, J. P., Stocke, J. T. 1999, ApJ, 527, 525.
- Donahue, M., Mack, J., Scharf, C., Lee, P., Postman, M., Rosati, P., Dickinson, M., Voit, G. M., Stocke, J. T. 2001, ApJ, 552, L93 (Paper 1).
- Ebeling, H. et al. 1996, MNRAS, 281, 799. Erratum 1996, MNRAS, 283, 1103.
- Ebeling, H. et al., 1997, ApJ, 479, L101.
- Ellingson, E., Lin, H., Yee, H. K. C., Carlberg, R. C., 2001, ApJ, 547, 609.
- Evrard, A. E. 1997, MNRAS, 292, 289.
- Gioia, I. M., et al., 1990a, ApJS, 72, 567.
- Gioia, I. M., et al., 1990b, ApJ, 357, L36.
- Gioia, I. M., et al., 2001, ApJ, 553, L103.
- Girardi, M., Borgani, S., Giuricin, G., Mardirossian, F., Mezzetti, M. 2000, ApJ, 530, 62.
- Gladders, M. D. & Yee, H. K. C. 2000, AJ, 120, 2148.
- Gunn, J. E., Hoessel, J. G. & Oke, J. B. 1986, ApJ, 306, 30.
- Hasinger, Burg, R., Giacconi, R., Schmidt, M., Trumper, J., 1998, A&A 329, 482
- Henry, J. P. et al. 1992, ApJ, 386, 408.
- Herzog, E., Wild, P., Zwicky, F. 1957, PASP, 69, 409.

- Jarvis, J.F., & Tyson, J.A. 1981, AJ, 86, 476.
- Jones, L. R., Scharf, C., Ebeling, H., Perlman, E., Wegner, G., Malkan, M., Horner, D. 1998, ApJ, 495, 100.
- Jones, C. & Forman, W. 1978, ApJ, 224, 1.
- Landolt, A. U. 1992, AJ, 104, 340.
- Lasker, B. M., Sturch, C. R., McLean, B. J., Russell, J. L., Jenkner, H. & Shara, M. M. 1990, AJ, 99, 2016.
- Lewis, A. D., Stocke, J. T., Ellingson, E., Gaidos, E. J., 2002, ApJ, in press (astro-ph/0110156).
- Mahdavi, A. & Geller, M. J. 2001, ApJ, 554, L129.
- Markevitch, M. 1998, ApJ, 504, 27.
- Mulchaey, J. S. & Zabludoff, A. I. 1998, ApJ, 496, 73.
- Mushotzky, R. F., Serlemitsos, P. J., Boldt, E. A., Holt, S. S., Smith, B. W. 1978, ApJ, 225, 21.
- Nichol, R. C., et al. 1999, ApJ, 521, L21.
- Nichol, R. C., et al. 1997, ApJ, 481, 644.
- Postman, M., Lubin, L.M., Gunn, J.E., Oke, J.B., Hoessel, J.G., Schneider, D.P., Christensen, J.A. 1996, AJ, 111, 615 (P96).
- Postman, M., Lubin, L. M., Oke, J. B., 1998a, AJ, 116, 560.
- Postman, M., Lauer, T. R., Szapudi, I., Oergerle, W. 1998b, ApJ, 506, 33.
- Press, W. H., Teukolksy, S. A., Vettering, W. T., Flannery, B. P. "Numerical Recipes in C", 1997 printing, Cambridge University Press.
- Puchnarewicz, E. M., et al. 1997, MNRAS, 291, 177.
- Romer, A. K., et al. 2000, ApJS, 126, 209.
- Rosati, P. et al. 1995, ApJ, 445, L11.
- Rosati, P. et al. 1998, ApJ, 492, L21.
- Rosati, P. et al. 2000, in Large Scale Structure in the X-ray Universe, Santorini, Greece, eds. Plionis, M. & Georgantopoulos, I., atlantisciences (Paris), p.13.
- Scharf, C. A., Jones, L. R., Ebeling, H., Perlman, E., Malkan, M., Wegner, G. 1997, ApJ, 477, 79.
- Scharf, C. A., Donahue, M., Voit, G. M., Rosati, P., Postman, M. 2000, ApJ, 528, L73.

- Schechter, P. 1976, ApJ, 203, 297.
- Smail, I. et al. 1997, ApJS, 110, 213.
- Stanford, S. A., Elston, R., Eisenhardt, P. R., Spinrad, H., Stern, D., Dey, A. 1997, AJ, 114, 2232.
- Stanford, S. A., Eisenhardt, P. R., & Dickinson, M. 1998, ApJ, 492, 461.
- Ulrich-Demoulin, M.-H. & Molendi, S. 1996, ApJ, 457, 77.
- Valdes, F. 1982, Proc, SPIE, 331, 465.
- Vikhlinin, A. et al. 1998, ApJ, 502, 558.
- Vikhlinin, A. et al. 2000, in Large-Scale Structure in the X-ray Universe, eds. M. Plionis & I. Georgantopoulos, atlantisciences (Paris), p. 31.
- White, N., Giommi, P., Angelini, L. 1995, http://lheawww.gsfc.nasa.gov/users/white/wgacat/wgacat
- Zabludoff, A. & Mulchaey, J. S. 1998, ApJ, 496, 39.
- Zabludoff, A. & Zaritsky, D. 1995, ApJ, 447, L21.
- Zwicky, F., et al, 1961, Catalogue of Galaxies and of Clusters of Galaxies, California Institute of Technology, vol. 1.

This preprint was prepared with the AAS LATEX macros v5.0.



Fig. 1.— The optical matched filter detection probabilities in the I-band as a function of redshift for systems of richness $\Lambda = 30, 50, 100, 200, 300$ (in order of increasing line weight). The probabilities were determined using the methods of P96.



Fig. 2.— Mean number of extended and pointlike X-ray sources per unit area as a function of distance from the center of the ROSAT field.



Fig. 3.— The X-ray detection probability in the 0.5-2 keV band as a function of redshift for intrinsic X-ray luminosities (0.5-2 keV) of 0.01, 0.1, 1.0, 5.0 and 10 $\times 10^{44}$ erg s⁻¹ ($H_0 = 75$, $q_0 = 0.5$) (in order of increasing line weight, from left to right). The detection probability is plotted as the net probability over all 23 ROXS fields, based on exposure time, background surface brightness, and radial detection efficiency due to PSF variations for a cluster with a given X-ray luminosity and redshift and a standard radial surface profile (Jones et al. 1998) and $\beta = 2/3$ (Scharf et al. 1997; Rosati et al. 1995).

Zel'dovich Pancake : NE quad



Fig. 4.— The rich northeast quadrant of the Zel'dovich Pancake field. X-ray point source detections are plotted as small circles. The large circles are extended X-ray sources. The optical cluster candidates are identified with crosses and single contours. In this quadrant there are 3 optical cluster candidates, two of which match an X-ray cluster candidate.



Fig. 5.— The black line shows the distribution of the upper limits to the bolometric X-ray luminosities of the cluster candidates which were not detected in the X-ray. The dashed lines show the distribution of the optical cluster candidates which were cross-identified with X-ray sources. The lower luminosity candidates and optical candidates with low luminosity upper limits lie at low estimated redshifts $z \sim 0.2 - 0.3$.



Fig. 6.— Two optical candidates with high detection significance from the matched filter algorithm (> 4σ) but no associated X-ray flux, OC6 1024+4707 from the 1024+4707 field (z = 0.4) and OC1 1118+2107 in the 1116+215 field (z = 0.3).



Fig. 7.— X-ray luminosities vs. optical richness for XBACS clusters (open circles, both panels) and ROXS optical cluster candidates with X-ray counterparts (filled circles, left panel) and the upper limits on the X-ray luminosity for ROXS optical candidates without X-ray counterparts (filled triangles, right panel). The size of the error bars for the ROXS optical richness reflects the uncertainty in the conversion from Λ_{cl} , the matched filter optical luminosity parameter, to N_R , optical richness.



Fig. 8.— The redshift (left) and Λ_{cl} (right) histograms of all 155 optically selected cluster candidates, of which 3 have $< 2.9\sigma$. The dotted line indicates those clusters with X-ray counterparts. The dashed line indicates the histogram of candidates with X-ray counterparts and optical candidates deemed "probable" in a subjective assessment.



Fig. 9.— The distribution of aperture-corrected $\Lambda_{cl}(1+z^9)^{0.7}$ (correction from P96), with the same line codes as Figure 8. The majority of the optically selected cluster candidates with corrected $\Lambda_{cl} > 100$ have estimated z > 1, and are not probable clusters, according to our subjective assessment.



Fig. 10.— The distribution of the optical detection confidence in units of σ for all of the optical candidates is plotted in a solid line. The same distribution for the X-ray/optical cross-identifications is plotted with a dotted line. Those clusters which were additionally identified as "likely" clusters are shown in a dashed line. Note that the majority of the X-ray/optical clusters lie in the bin with detection significance $3-4\sigma$.



Fig. 11.— The optical detection confidence is plotted here against Λ_{cl} , an estimate of the cluster's optical luminosity. The two quantities are not strongly correlated.



Fig. 12.— V-I vs. I distributions for mutual V and I cluster candidates in 1202+281 ROSAT field and for 1 of the 2 candidates in the 3C356 ROSAT field. The solid line(s) plot the sequence at the estimated redshift(s) of the cluster candidates. The dashed lines show the red sequences predicted for z = 0.1 - 0.7 in increments of $\Delta z = 0.1$. The upper left label indicates the field, the quadrant, the estimated redshift from the I(V). The upper right label lists the detection confidence (sig) in I(V), and Λ_{cl} (lamb) from the I(V) data. Lower right label lists the cluster core size.



Fig. 13.— V-I vs. I distributions for mutual V and I cluster candidates in the 3C356 (1 of 2) and 3C324 ROSAT fields. Plots are annotated as in Figure 12.



Fig. 14.— V-I vs. I distributions for mutual V and I cluster candidates in the Zel'dovich Pancake and the 0902+34 ROSAT fields. Plots are annotated as in Figure 12.



Fig. 15.— V-I vs. I distributions for 3 mutual V and I cluster candidates in Zel'dovich Pancake ROSAT field. Plots are annotated as in Figure 12.



Fig. 16.— L_x vs. Λ is plotted for the three cluster detection datasets in a ROXS simulation with $L_x \propto \Lambda^2$, normalized to approximate the observed relationship. Leftmost panel plots clusters *jointly* detected in the X-ray and optical, point size is proportional to cluster redshift (0.2 < z < 1.2). Solid curve is the $L_x \propto \Lambda^2$ relationship used to generate the clusters, dashed curve is an $L_x \propto \Lambda^4$ relationship. Center plot is of X-ray upper limits for clusters only detected optically. Note that since these are 4σ upper limits to L_x and the clusters are typically at higher z for a given Λ than the detected systems, these points occupy a region above the instrinsic curve. Rightmost panel plots those clusters which are X-ray detected but not optically detected (1 object in this simulation). L_x is in units 10⁴⁴ erg s⁻¹.



Fig. 17.— As for Figure 16, but simulation now uses $L_x \propto \Lambda^3$ and an instrinsic scatter with $1\sigma = 39\%$ of the amplitude.



2D K-S Probability of match

1

0.1

40

60

80

Fig. 18.— Summary of 2D K-S test and simulation results. In the upper panel the 2D K-S probabilities (of a significant match between the ROXS joint X-ray/optical detection Λ , L_x data and the simulation joint detections) are plotted as a function of the normalization A in the simulation $\Lambda = AL_x^{\alpha}$ relationship. Simulation data is taken from single Monte Carlo runs which generate at least 50 joint detection data points, scatter between runs is small compared to the true variations in these curves. Lightest curve is for $\alpha = 1/2$, medium is $\alpha = 1/3$, and the heaviest is $\alpha = 1/4$. Lower panel shows the number ratio of jointly detected clusters to those only detected optically, curves are as for the upper panel. The dashed horizontal line denotes the observed ratio for the ROXS.

100

120

Normalization (A)

140

160

180

200

ROSAT Field	ROSAT Target	ROSAT Exptime	RA	Dec	Area	KPNO Date	
DD700110		(sec)	J2000	J2000	deg ²	I-Band	V-Band
RP700112	MKN 78	19844	07:42:40	65:10:52	0.17518	03/18/96	
RP700326N00	0902 + 34	14355	09:05:31	34:07:48	0.24414	03/17/96	05/06/97
RP900327	Zeldovich Pancake	55611	$09{:}06{:}52$	33:40:16	0.24161	03/16/96	03/18/96
WP700540	10214 + 4724	19359	10:24:33	47:09:04	0.24401	03/17/96	
WP201243N00	PG1034+001	18111	10:37:05	-00:08:21	0.19623	05/09/97	
WP700228	1116 + 215	24585	11:19:09	21:19:16	0.24343	03/17/96	
WP201367M01	PG1159-035	32219	12:04:25	-03:40:16	0.23667	03/18/96	
WP700232	1202 + 281	27859	12:04:43	27:54:04	0.23531	03/18/96	05/06/97
RP700864A01	3C 270.1	19290	12:20:33	33:43:16	0.19836	05/06/97	
RP600242A01	Giovanelli-Haynes Cl	24830	12:27:43	01:36:04	0.18036	05/08/97	
RP700073	3C 280	48051	12:56:57	47:20:28	0.16501	03/17/96	
RP700216A00	B2 1308+32	8316	13:10:29	32:20:59	0.19679	05/09/97	
RP700117	3CR294	22692	14:06:43	34:11:27	0.19156	05/07/97	
WP700248	1411 + 442	25319	14:13:48	44:00:04	0.23088	03/16/96	
RP700122	Q1413+1143	28066	14:15:45	11:29:46	0.24770	03/18/96	
RP800401A01	4C23.37	11811	14:15:58	23:07:12	0.23037	03/16/96	
RP700257N00	MKN 841	16842	15:04:02	10:26:24	0.18221	05/09/97	
RP701373N00	3C 324	15434	15:49:50	21:25:52	0.22442	03/17/96	05/06/97
RP800239N00	CL1603	28763	16:04:29	43:13:12	0.18933	05/08/97	
RP300021N00	MS1603.6+2600	24506	16:05:46	25:51:36	0.19376	05/08/97	
WP170154	WFC/PSPC Calibr	37490	16:29:32	78:04:52	0.18426	05/09/97	
701457n00	HS1700+6416	27408	17:01:00	64:11:59	0.18889	05/08/97	

Table 1. ROSAT fields

- 55 - Table 1—Continued

RP800395N00 3C 356 18563 17:24:19 50:57:36 0.22243 03/18/96 05/06/97

_

Optical ID	RA	Dec	Sigma	Lambda	z	I-mag	Radius (")	Theta (')	$^{\mathrm{F}_{lim}1'}_{(10^{-14})}$	Xray Match	Comment
						RP7001	12 - MKN	78			
OC1 0740+6504 OC2 0741+6524	$\begin{array}{c} 07 \ 40 \ 57.25 \\ 07 \ 41 \ 44.06 \end{array}$	$65 \ 04 \ 00.9 \\ 65 \ 24 \ 42.9$	$3.42 \\ 3.25$	$31.17 \\ 38.69$	$0.30 \\ 0.40$	$19.46 \\ 20.29$	$138 \\ 162$	$12.85 \\ 15.11$	$1.9 \\ 2.1$		PtSource within 30"
OC2a 0742+6519 OC3 0743+6458	$\begin{array}{c} 07 \ 42 \ 14.19 \\ 07 \ 43 \ 45.82 \end{array}$	$\begin{array}{c} 65 19 43.7 \\ 64 58 14.9 \end{array}$	$3.31 \\ 5.45$	$30.13 \\ 64.88$	$\begin{array}{c} 0.30\\ 0.40\end{array}$	$\begin{array}{c} 19.60\\ 20.13\end{array}$	170 162	$9.35 \\ 14.31$	$1.7 \\ 2.1$	RXJ0743.7+6457	Cluster within 1'
						RP700326	N00 - 0902	2+34			
OC1 0904+3411 OC2 0905+3359 OC3 0905+3421 OC4 0905+3421	09 04 23.88 09 05 10.18 09 05 11.11	34 11 04.9 33 59 37.6 34 21 53.0 24 02 17 5	4.25 2.94 5.14 2.58	43.38 25.04 34.79	0.40 0.30 0.20 1.00	20.13 19.24 18.39	70 59 116	14.27 9.24 14.67 6.40	2.5 2.1 2.5		Edge,Compact Compact,Low Sig Edge,Pt Source within 1' Pland 0.052
OC5 0905+3407 OC5 0906+3417 OC7 0906+3405 OC8 0906+3359	$\begin{array}{c} 03 & 03 & 03 & 03 \\ 09 & 05 & 57.01 \\ 09 & 06 & 17.59 \\ 09 & 06 & 28.25 \\ 09 & 06 & 31.74 \\ 00 & 06 & 02 \\ \end{array}$	$34\ 05\ 11.3$ $34\ 07\ 00.6$ $34\ 17\ 16.0$ $34\ 05\ 11.9$ $33\ 59\ 51.1$	3.35 3.39 5.40 4.05	48.56 28.87 36.56 70.13	0.60 0.30 0.20 0.70	21.28 19.24 18.39 21.65	101 177 79 275 165	5.44 13.50 12.13 14.89	1.8 1.8 2.4 2.3	RXJ0906.3+3417	Blend OC7 and OC4? Cluster within 15" PtSource within 2.0' PtSource within 1.5', blend OC7?
0C9 0906+3413	09 06 32.70	34 13 17.4	3.35	89.67	1.10	22.71	50	13.88	2.5		
					RP	900327 - 2	Lel'dovicn	Pancake			
OC1 0906+3354 OC2 0907+3330 OC3 0907+3343	$\begin{array}{c} 09 \ 06 \ 12.55 \\ 09 \ 07 \ 18.64 \\ 09 \ 07 \ 27.69 \\ \end{array}$	$\begin{array}{c} 33 \ 54 \ 21.9 \\ 33 \ 30 \ 44.5 \\ 33 \ 43 \ 02.5 \end{array}$	$4.32 \\ 5.23 \\ 4.59$	25.13 85.19 74.79	$0.20 \\ 0.50 \\ 0.50$	18.10 20.71 20.85	227 174 268	$16.45 \\ 10.88 \\ 7.80$	1.7 1.3 1.0	RXJ0907.3+3330 RXJ0907.4+3342	Cluster within 30" Cluster within 30"
OC4 0907 + 3355 OC5 0907 + 3350	$09\ 07\ 47.16$ $09\ 07\ 58.29$	33 55 00.4 33 50 09.6	$3.50 \\ 3.09$	28.21 67.27	$0.30 \\ 0.70$	19.47 21.57	68 135	$18.62 \\ 16.86$	1.9	RXJ0907.8+3351	Edge,Compact,Near cut area Cluster within 2',PtSource within 15"
						WP700540	0 - 10214+	4724			
OC1 1023+4722 OC2 1024+4657 OC3 1024+4721 OC4 1024+4714 OC5 1024+4721	$\begin{array}{c} 10 \ 23 \ 18.89 \\ 10 \ 24 \ 06.16 \\ 10 \ 24 \ 08.26 \\ 10 \ 24 \ 19.52 \\ 10 \ 24 \ 42.64 \end{array}$	$\begin{array}{c} 47 \ 22 \ 59.8 \\ 46 \ 57 \ 05.7 \\ 47 \ 21 \ 02.4 \\ 47 \ 14 \ 25.6 \\ 47 \ 21 \ 41.7 \end{array}$	3.20 3.09 3.53 3.45 3.02	37.61 22.08 52.37 56.14 77.59	0.40 0.20 0.50 0.60 1.20	$20.18 \\18.26 \\20.72 \\21.13 \\22.85$	$74 \\ 55 \\ 115 \\ 103 \\ 56$	$18.86 \\ 12.80 \\ 12.78 \\ 5.93 \\ 12.79$	2.5 2.0 2.0 1.5 2.0		Edge Compact
OC6 1024+4707 OC7 1024+4656 OC8 1025+4701	$\begin{array}{c} 10 \ 24 \ 43.91 \\ 10 \ 24 \ 52.51 \\ 10 \ 25 \ 01 \ 77 \end{array}$	$\begin{array}{c} 47 \ 07 \ 21.1 \\ 46 \ 56 \ 56.2 \\ 47 \ 01 \ 53 \ 7 \end{array}$	4.76 4.01 2.96	55.85 59.42 76.12	$0.40 \\ 0.50 \\ 1.20$	20.14 20.69 22.91	108 176 58	2.41 12.49 8.57	1.2 2.0 1.7		Blend with OC 11 Low Sig
OC9 1025+4722 OC9a 1025+4723 OC10 1025+4716 OC11 1025+4701	$\begin{array}{c} 10 & 25 & 07.14 \\ 10 & 25 & 07.14 \\ 10 & 25 & 26.37 \\ 10 & 25 & 22.73 \\ 10 & 25 & 35.85 \end{array}$	47 22 58.8 47 23 16.4 47 16 13.9 47 01 23.7	3.44 3.25 3.63 7.35	$ \begin{array}{r} 40.46 \\ 52.92 \\ 30.16 \\ 52.92 \end{array} $	$\begin{array}{c} 0.40 \\ 0.60 \\ 0.30 \\ 0.20 \end{array}$	20.12 21.33 19.04 18.25	71 93 39 356	15.09 16.84 11.03 13.05	2.2 2.3 1.9 2.0	RXJ1025.4+4703	Near Edge; Blend with 9A? Near Edge; Blend with 9?; PtSrc Compact,PtSource within 30" Cluster within 3'; Blend OC7
					W	P201243N	00 - PG10	34+001			
OC1 1036+0005	10 36 36.01	00 05 24.9	4.30	64.68	0.80	21.78	137	7.79	1.2		Near Edge,PtSource within 1' Blend 1a, 1b
OC1a 1036+0002 OC1b 1037+0005 OC2 1036-0018 OC3 1037-0007	$\begin{array}{c} 10 \ 36 \ 50.70 \\ 10 \ 37 \ 03.35 \\ 10 \ 36 \ 43.81 \\ 10 \ 37 \ 22.54 \\ 10 \ 37 \ 27 \ 57 \end{array}$	00 02 36.2 00 05 37.1 -00 18 52.9 -00 07 59.0	3.86 3.27 4.18 3.34 2.02	$\begin{array}{c} 43.35 \\ 52.58 \\ 29.47 \\ 37.53 \\ 16.60 \end{array}$	$0.50 \\ 0.90 \\ 0.30 \\ 0.50 \\ 0.90 \\ $	20.69 22.17 19.26 20.55	131 89 98 119	6.78 2.81 11.72 4.45 8.47	$1.2 \\ 0.9 \\ 1.6 \\ 1.0 \\ 1.2 \\ 0.9 $		Near Edge, blend 1, 1b Near Edge; blend 1a, 1 Near Edge Blend with OC3a
OC3a 1037-0010 OC4 1038-0008	$\begin{array}{c} 10 \ 37 \ 37.70 \\ 10 \ 38 \ 03.24 \end{array}$	$-00\ 10\ 25.0$ $-00\ 08\ 22.4$	$3.93 \\ 3.11$	16.60 34.88	$0.20 \\ 0.50$	18.27 20.66	200 67	$8.47 \\ 14.61$	1.3 1.8		PtSource within 1'; Blend with OC3

Table 2. Optical Cluster Candidates

 $^{-57}$

Table	2	Continued
10010	-	Commuou

	I					WP	700228 -	-1116+2	15		
OC1 1118+2107	11 18 09.36	$21 \ 07 \ 04.7$	5.26	52.96	0.30	19.42	122	18.56	2.8		Near Edge
OC2 1118+2126	$11\ 18\ 29.45$	$21 \ 26 \ 48.0$	2.93	56.79	1.00	22.54	39	12.04	2.3		Low Sig
OC3 1119+2116	$11 \ 19 \ 18.80$	$21 \ 16 \ 54.9$	3.80	46.36	0.50	20.79	184	3.13	1.6	RXJ1119.2+2117	Cluster within 30", Blend OC3a
OC3a 1119+2118	$11 \ 19 \ 31.88$	$21 \ 18 \ 54.8$	4.09	41.21	0.30	19.46	219	5.20	1.8		Blend OC3, OC5; PtSrc
OC4 1119+2107	$11 \ 19 \ 26.77$	$21 \ 07 \ 08.9$	6.07	46.53	0.20	18.18	189	12.70	2.4	RXJ1119.4+2106	Cluster within 30"
OC5 1119+2124	$11 \ 19 \ 35.04$	$21 \ 24 \ 19.2$	3.38	44.92	0.70	21.63	76	7.83	2.0		PtSource within 1'; Blend OC6, 3a
OC6 1119+2127	11 19 35.95	21 27 18.2	3.89	43.00	0.40	20.14	192	10.16	2.2	RXJ1119.7+2126	Cluster within 1.5',PtSource within 1' Blend OC5
OC7 1119+2131 OC8 1119+2124	11 19 55.78 11 19 59.29	$21 \ 31 \ 36.6$ $21 \ 24 \ 34.1$	$3.36 \\ 3.13$	$33.80 \\ 74.57$	$0.30 \\ 1.10$	$19.48 \\ 22.72$	158 76	$16.41 \\ 12.75$	$2.7 \\ 2.4$		
						WP20	01367 -	PG1159-	035		I
0.01 1902 0244	10 02 07 66	02 44 95 5	2.44	42.10	0.40	20.24	00	15.07	17		Neer Edge Difference within 20"
0C1 1203-0344	12 03 27.00	-03 44 20.0 03 50 34 2	3.44	43.18	0.40	20.24 21.57	00 70	17.66	1.7		Near Edge PtSource within 1'
0C3 1203-0330	12 03 20.17	-03 30 34.3	3.04	60.49	0.70	21.07	19	15.89	1.9		Compact
OC4 1203-0333	12 03 29.23	-03 42 45 8	3.84	64 10	0.80	21.69	137	6 72	1.0		Blend with OC4a, OC4b
OC4 1204-0042	12 04 00.10	-03 42 12 5	3.26	86.45	0.00	221.10	119	4 40	0.9		PtSource within 30": Blend OC4 OC4b
OC4b 1204-0348	12 04 14 18	-03 48 16 5	3.85	131.82	1 10	22.20	174	8 47	1.2		Blend with OC4 OC4a
OC5 1204-0351	12 04 21.80	-03 51 15.3	4.90	28.06	0.20	18.38	229	11.00	1.4	RXJ1204.3-0350	Cluster within 30"
OC6 1204-0330	$12 \ 04 \ 41.21$	-03 30 22.3	3.38	32.56	0.30	19.33	100	10.70	1.4		
OC7 1204-0338	12 04 45.34	-03 38 17.0	3.16	30.45	0.30	19.54	99	5.28	1.0		
OC8 1204-0330	$12 \ 04 \ 55.19$	-03 30 52.9	2.10	55.10	0.90	22.21	11	_	_	?RXJ1205.0-0332	Cluster within 2' (note low sigma)
						WP7	700232 -	1202+2	81		
OC1 1203+2758	12 03 47.81	27 58 36.5	3.92	56.48	0.50	20.72	91	13.07	1.7	?RXJ1204.1+2807	A1455?
OC2 1203+2743	$12 \ 03 \ 53.23$	$27 \ 43 \ 54.8$	3.25	39.62	0.40	20.13	130	14.97	1.9		
OC3 1204 + 2744	$12 \ 04 \ 26.02$	$27 \ 44 \ 12.8$	3.16	89.91	1.00	22.52	53	10.50	1.5		
OC4 1204 + 2805	$12 \ 04 \ 36.71$	$28 \ 05 \ 19.9$	3.35	95.30	1.00	22.51	117	11.42	1.6	?RXJ1204.1+2807	A1455?
OC5 1204 + 2801	$12 \ 04 \ 38.01$	$28 \ 01 \ 52.0$	3.96	28.01	0.20	18.25	145	7.95	1.3	?RXJ1204.1+2807	A1455?
$OC6\ 1204 + 2757$	$12 \ 04 \ 53.71$	27 57 59.1	3.21	39.02	0.40	20.32	92	4.61	1.0		
OC7 1204 + 2753	$12 \ 04 \ 56.99$	27 53 04.8	3.18	90.66	1.00	22.48	49	3.18	0.9		
$OC8 \ 1205 + 2805$	$12\ 05\ 18.49$	28 05 17.1	3.19	31.43	0.30	19.51	73	13.71	1.8		Sprawling
OC9 1205+2740	12 05 43.83	27 40 30.2	4.15	50.47	0.40	20.05	130	19.04	2.2		Near Edge; PtSource 1'
						RP70	0864A0	1 - 3C27	0.1		
OC1 1219+3331	$12 \ 19 \ 35.76$	$33 \ 31 \ 44.0$	3.44	30.79	0.30	19.21	65	16.41	1.9		Edge; PtSource
OC2 1219+3345	12 19 41.34	$33 \ 45 \ 20.6$	4.95	26.82	0.20	18.33	257	11.07	1.5		Blend OC2a
OC2a 1219+3343	$12 \ 19 \ 57.94$	$33 \ 43 \ 54.2$	3.67	44.68	0.40	20.20	189	7.45	1.2		Blend OC2
OC3 1220 + 3334	12 20 08.42	33 34 02.7	2.64	91.13	1.20	22.86	34	-	_	?RXJ1220.0+3334	Cluster within 1.5' (note low sigma)
OC4 1221+3344	12 21 11.30	33 44 31.3	3.20	28.66	0.30	19.47	85	7.95	1.3	RXJ1220.8+3343 RXJ1220.9+3343	Both Xray clusters within 4', PtSource within 15
OC5 1221+3352	$12 \ 21 \ 35.57$	$33 \ 52 \ 28.4$	3.61	124.68	1.20	22.87	84	15.86	1.9		Edge

Table 2—Continued

OC1 1227+0143 OC1a 1228+0142 OC2 1227+0136	$\begin{array}{c} 12 \ 27 \ 48.26 \\ 12 \ 28 \ 07.53 \\ 12 \ 27 \ 46.77 \end{array}$	$\begin{array}{c} 01 \ 43 \ 38.8 \\ 01 \ 42 \ 36.5 \\ 01 \ 36 \ 01.0 \end{array}$	2.96 3.79 2.91	$34.13 \\ 65.44 \\ 61.06$	0.30 0.50 0.80	19.47 20.76 21.86	$ \begin{array}{r} 114 \\ 246 \\ 42 \\ 42 \end{array} $	$7.76 \\ 9.01 \\ 0.94$	$1.7 \\ 1.8 \\ 1.2$	RXJ1227.8+0143	Cluster within 45";Blend OC1a Blend OC1, OC4; PtSources Compact,Low Sig	
OC3 1228 + 0136 OC4 1228 + 0136	$\begin{array}{c} 12 \ 28 \ 08.53 \\ 12 \ 28 \ 11.07 \end{array}$	$01 \ 48 \ 17.5$ $01 \ 36 \ 08.7$	$3.37 \\ 3.13$	$63.64 \\ 19.81$	$\begin{array}{c} 0.70 \\ 0.20 \end{array}$	$21.56 \\ 18.66$	$110 \\ 193$	$13.85 \\ 7.01$	$\frac{2.2}{1.6}$		Edge Blend OC1, OC4	
							RP700	0073 - 3C	280			
OC1 1257+4719	12 57 03.89	47 19 09.2	2.02	76.15	1.00	22.52	9	-	-	?RXJ1256.9+4720	Cluster within 1.5'(note low sigma)	
OC2 1257 + 4709 OC3 1257 + 4708	12 57 13.31 12 57 53.29	47 09 01.3 47 08 39.3	6.04	42.15	0.80 0.20	18.48	325	$11.09 \\ 15.09$	$1.5 \\ 1.7$		rtSource within 1.5	
						RP'	700216A	00 - B2	1308 + 3	2		
OC1 1309+3218	13 09 54.66 12 10 04 20	$32 \ 18 \ 25.4$	3.86	90.60	0.90	22.10	140	7.51	1.3	DV 11200 0 1 2000	Blend with OC2	-0.945), Pland OC1
OC2 1310+3221 OC3 1310+3214	$13 \ 10 \ 04.30$ $13 \ 10 \ 44.83$	$32 \ 21 \ 17.8$ $32 \ 14 \ 04.3$	$4.48 \\ 4.11$	44.68 22.97	$0.30 \\ 0.20$	19.47 18.39	$124 \\ 117$	$5.03 \\ 7.77$	$1.1 \\ 1.3$	RAJ1309.9+3222	Cluster within 2 (MS1308.8+3244,z=	(0.245); Blend OCI
OC4 1311+3228 OC4a 1311+3228	$\begin{array}{c} 13 \ 11 \ 12.72 \\ 13 \ 11 \ 28.36 \end{array}$	$32 \ 28 \ 47.3 \\ 32 \ 28 \ 52.9$	$5.28 \\ 5.14$	$29.49 \\ 66.19$	$0.20 \\ 0.40$	$18.62 \\ 20.14$	$274 \\ 180$	$12.23 \\ 14.97$	$1.6 \\ 1.8$	RXJ1313.2+3229	Cluster within 30"; Blend OC4a PtSource within 1'; Blend with OC4	
	Į						RP700)117 - 3C	294			I
OC1 1406+3401	14 06 05.35	$34\ 01\ 57.5$	3.22	43.47	0.60	21.05	63	12.23	1.9		PtSrc	90
OC2 1406 + 3411 OC3 1406 + 3403	$14\ 06\ 07.62$ $14\ 06\ 30.63$	$34 11 01.2 \\ 34 03 05.3$	3.30 3.34	8.10 19.97	0.20 0.30	18.86 19.47	110	7.33 8.68	1.5 1.6		PtSource within 1.5	
OC4 1406 + 3407	$14\ 06\ 36.45$	$34 \ 07 \ 33.3$	3.04	34.56	0.50	20.56	64	4.06	1.2		Compact	
OC5 1406+3402	14 06 59.14	34 02 04.2	3.09	72.07	0.90	22.13	67 76	9.89	1.7		C	
OC6 1407 + 3420 OC7 1407 + 3415	$14\ 07\ 41.45$ $14\ 07\ 41\ 81$	$34 \ 20 \ 45.9$ $34 \ 15 \ 24 \ 4$	3.90 5.86	8.97 66.54	0.20	18.77	144	15.28	2.1	BX 11407 $6+3415$	Compact; PtSrc Cluster within 30" PtSource within 1!	5'
OC8 1407+3400	14 07 44.94	34 00 17.3	3.18	19.03	0.30	19.47	35	16.96	2.2	10101101.0 0410	Compact,Near Edge	5
						I	NP7002	48 - 141	1 + 442			
$OC1 \ 1412 + 4347$	$14\ 12\ 50.50$	$43 \ 47 \ 57.6$	3.89	106.22	1.00	22.48	176	15.90	2.2		Blend OC3	
OC2 1412+4353	14 13 02.49	43 53 16.8	3.29	56.36	0.60	21.10	169	10.60	1.7		Blend with OC2a	
OC2a 1413 + 4347 OC2 1412 + 4250	14 13 26.52	43 47 20.9	3.75	64.29 60.71	0.60	20.98	150	13.23	2.0		Blend with OC2 Blend OC1	
OC3 1413 + 4359 OC4 1413 + 4351	14 13 12.83 14 13 30.58	43 59 10.9	4.83	37.77	0.30	19.19	318	9.08	1.4		Bielid OC1	
OC5 1414+4347	$14 \ 14 \ 12.61$	43 47 23.8	3.45	68.96	0.70	21.44	84	13.36	2.0			
OC6 1415+4348	$14\ 15\ 05.58$	$43 \ 48 \ 16.0$	3.21	25.13	0.30	19.27	104	18.26	2.3		Near Edge, PtSource within 1'	
OC7 1415 + 4352	$14\ 15\ 09.62$	$43 \ 52 \ 13.3$	2.98	101.76	1.20	22.87	39	16.64	2.2		Edge,Low Sig	
OC8 1415+4409	14 15 10.84	44 09 34.5	3.07	24.02	0.30	18.96	34	17.68	2.3		Compact,Off Edge	
				NO 15	0.00	R	P700122	2 - Q1413	3+1143			
0C1 1414 + 1123 0C2 1414 + 1118	14 14 50.97	11 23 35.9 11 18 22 2	6.25 2.07	59.40 87.49	0.30	19.09	264 122	14.71 16.76	1.9	KAJ1415.2+1119	Cluster within 7', PtSource within 15" Low Sig	
OC3 1414+1143	14 14 53.24 14 14 58.61	11 43 54.4	4.09	38.91	0.30	19.14	155	18.28	2.0 2.1		Near Edge	
OC4 1415+1140	14 15 21.90	11 40 06.4	4.47	33.13	0.20	18.22	196	11.92	1.6		Blend with OC4a	
OC4a 1415+1141	$14\ 15\ 30.56$	$11 \ 41 \ 28.2$	3.57	47.88	0.50	20.88	176	12.33	1.7		Blend with OC4	
OC5 1416 + 1141	$14 \ 16 \ 41.07$	$11 \ 41 \ 06.5$	3.19	30.29	0.30	19.32	102	17.74	2.1			

RP600242A01 - Giovanelli-Haynes Cloud

 $^{\circ}59$

							RP8004	401A01 -	4C23.3	7		
OC1 1415+2259 OC2 1415+2317 OC3 1415+2307 OC3a 1415+2311 OC4 1417+2255	$\begin{array}{c} 14 \ 15 \ 14.04 \\ 14 \ 15 \ 40.08 \\ 14 \ 15 \ 55.85 \\ 14 \ 15 \ 26.72 \\ 14 \ 17 \ 01.06 \end{array}$	$\begin{array}{c} 22 \ 59 \ 36.9 \\ 23 \ 17 \ 08.5 \\ 23 \ 07 \ 26.1 \\ 23 \ 11 \ 40.9 \\ 22 \ 55 \ 41.8 \end{array}$	5.02 3.96 4.76 3.86 3.26	$\begin{array}{r} 44.39 \\ 39.08 \\ 42.09 \\ 58.86 \\ 25.66 \end{array}$	$0.30 \\ 0.40 \\ 0.30 \\ 0.50 \\ 0.20$	$19.17 \\ 20.16 \\ 19.25 \\ 20.62 \\ 18.23$	$240 \\ 105 \\ 215 \\ 120 \\ 68$	12.46 10.67 0.35 8.28 18.70	2.1 1.9 1.1 1.7 2.6	RXJ1415.8+2316 RXJ1415.9+2307	PtSource within 30" Cluster within 2.5',PtSource with Cluster within 30" PtSource within 30" Near Edge	hin 1'
						1	RP7002	257N00 -	MKN84	41		
$\begin{array}{c} {\rm OC1} \ 1503{+}1014 \\ {\rm OC2} \ 1503{+}1021 \\ {\rm OC2a} \ 1503{+}1018 \\ {\rm OC2b} \ 1503{+}1021 \\ {\rm OC3} \ 1504{+}1022 \end{array}$	$\begin{array}{c} 15 \ 03 \ 16.51 \\ 15 \ 03 \ 33.48 \\ 15 \ 03 \ 33.92 \\ 15 \ 03 \ 45.45 \\ 15 \ 04 \ 00.49 \end{array}$	10 14 16.9 10 21 32.5 10 18 39.7 10 21 35.4 10 22 13.3	3.60 3.82 3.60 4.12 2.98	$\begin{array}{r} 46.43 \\ 23.28 \\ 66.46 \\ 44.77 \\ 67.95 \end{array}$	$0.40 \\ 0.20 \\ 0.60 \\ 0.30 \\ 0.80$	20.24 18.67 21.13 19.51 21.93	159 290 149 217 73	$16.48 \\ 8.52 \\ 10.36 \\ 6.29 \\ 4.19$	3.3 2.7 2.8 2.5 2.3		Edge Blend with OC2a, OC2b PtSource within 30"; Blend with Blend with OC2, OC2a Low Sig	оС2, ОС2ь
							RP701	1373N00	- 3C324	L		
OC1 1548+2113 OC2 1548+2127	$\begin{array}{c} 15 \ 48 \ 50.06 \\ 15 \ 48 \ 55.88 \end{array}$	$\begin{array}{cccccccccccccccccccccccccccccccccccc$	3.39 3.89	$39.86 \\ 38.51$	$\begin{array}{c} 0.50\\ 0.40\end{array}$	$\begin{array}{c} 20.76\\ 20.14 \end{array}$	106 171	$\begin{array}{c} 18.42 \\ 12.78 \end{array}$	$2.5 \\ 2.1$	RXJ1548.8+2126 Blend w/ OC1?.OC2a, OC2b	Blend with OC2/2a/2b? Cluster within 1.5',PtSource with	hin 1'
OC2a 1549+2119 OC2b 1549+2122 OC3 1549+2123	15 49 02.22 15 49 02.37 15 49 32.13	$\begin{array}{cccccccccccccccccccccccccccccccccccc$	$5.60 \\ 3.88 \\ 3.85$	$41.59 \\ 31.53 \\ 70.35$	$0.20 \\ 0.30 \\ 0.70$	$18.02 \\ 19.15 \\ 21.55$	304 198 81	$12.68 \\ 11.68 \\ 4.88$	$2.1 \\ 2.0 \\ 1.5$,,,	PtSource within 1'; Blend with C Blend with OC1?,OC2,OC2a	DC1?,OC2,OC2
$\begin{array}{c} \text{OC4 } 1549 + 2129 \\ \text{OC5 } 1550 + 2112 \\ \text{OC6 } 1550 + 2121 \end{array}$	$\begin{array}{c} 15 \ 49 \ 55.45 \\ 15 \ 50 \ 02.46 \\ 15 \ 50 \ 10.33 \end{array}$	$\begin{array}{cccccccccccccccccccccccccccccccccccc$	$2.94 \\ 3.18 \\ 2.96$	$23.95 \\ 23.64 \\ 54.04$	$0.30 \\ 0.20 \\ 0.70$	$19.28 \\ 17.65 \\ 21.49$		$3.65 \\ 13.47 \\ 6.51$	$1.4 \\ 2.2 \\ 1.6$		Low Sig,Compact Near Edge; PtSrc Low Sig	I
$OC7 1550+2125 \\ OC8 1550+2134$	$\begin{array}{c} 15 50 13.35 \\ 15 50 44.69 \end{array}$	$\begin{array}{cccccccccccccccccccccccccccccccccccc$	$3.64 \\ 3.69$	$ \begin{array}{r} 113.13 \\ 36.51 \end{array} $	$\begin{array}{c} 1.10 \\ 0.40 \end{array}$	$\begin{array}{c} 22.71 \\ 20.08 \end{array}$	87 148	$5.36 \\ 15.38$	$1.5 \\ 2.3$		-	
							RP800	239N00 -	CL160	3		
OC1 1603+4323 OC2 1604+4313 OC3 1604+4310 OC4 1604+4324	$\begin{array}{c} 16 \ 03 \ 29.85 \\ 16 \ 04 \ 11.46 \\ 16 \ 04 \ 39.05 \\ 16 \ 04 \ 51.73 \end{array}$	$\begin{array}{c} 43 \ 23 \ 52.2 \\ 43 \ 13 \ 53.8 \\ 43 \ 10 \ 19.7 \\ 43 \ 24 \ 16.4 \end{array}$	$3.92 \\ 3.25 \\ 4.38 \\ 5.58$	$71.12 \\ 45.90 \\ 35.31 \\ 44.93$	$0.60 \\ 0.50 \\ 0.30 \\ 0.30$	21.14 20.85 19.34 19.38	140 121 172 175	15.02 3.11 3.50 11.88	$1.8 \\ 0.9 \\ 0.9 \\ 1.6$		PtSource within 1'	
						RP3	300021N	100 - MS	1603.6+	-2600		
$\begin{array}{c} \text{OC1 } 1604 + 2558 \\ \text{OC2 } 1604 + 2554 \\ \text{OC3 } 1605 + 2541 \\ \text{OC4 } 1605 + 2553 \\ \text{OC4 } 1605 + 2553 \end{array}$	$\begin{array}{c} 16 \ 04 \ 55.58 \\ 16 \ 04 \ 56.65 \\ 16 \ 05 \ 18.39 \\ 16 \ 05 \ 38.20 \\ \end{array}$	$\begin{array}{cccccccccccccccccccccccccccccccccccc$	2.99 3.54 3.86 4.84	23.00 34.93 67.98 26.15	0.40 0.60 0.90 0.20	20.26 21.21 22.35 18.69	63 77 117 292	$13.3 \\ 11.22 \\ 11.70 \\ 2.08 \\ 0.01$	1.9 1.7 1.8 1.0	?RXJ1605.0+2552 RXJ1605.5+2553	Compact,Low Sig Cluster within 2.5',PtSource with PtSource within 15"; QSO w/z= Cluster within 1.5'	hin 45" 1.07
$\begin{array}{c} \text{OC4a } 1605 + 2554 \\ \text{OC5 } 1606 + 2602 \\ \text{OC6 } 1606 + 2556 \end{array}$	$ \begin{array}{c} 16 \ 05 \ 58.12 \\ 16 \ 06 \ 27.81 \\ 16 \ 06 \ 40.05 \\ \end{array} $	$\begin{array}{c} 25 \ 54 \ 12.4 \\ 26 \ 02 \ 35.1 \\ 25 \ 56 \ 22.8 \end{array}$	$3.88 \\ 3.65 \\ 3.85$	$30.15 \\ 94.53 \\ 37.92$	$0.40 \\ 1.10 \\ 0.60$	20.02 22.75 21.22	131 107 140	3.94 14.6 13.27	1.2 2.0 1.9	CAL	PtSource within 15"	
	1					WE	170154	4 - WFC/	PSPC	CAL		

Table 2—Continued

Table	2	Continue	d
10010	_	Contract	Ċ.

OC1 1626+7817	16 26 20.25	$78\ 17\ 01.5$	3.81	55.28	0.50	20.64	118	15.66	2.1	RXJ1626.3+7816	Cluster within 15", Blend with OC1a
OC1a 1625+7814	$16\ 25\ 54.51$	$78 \ 14 \ 39.2$	4.81	44.17	0.30	19.14	126	14.86	2.0		Blend with OC1
$OC2 \ 1626 + 7759$	$16\ 26\ 52.23$	77 59 05.3	3.18	29.19	0.30	19.27	83	10.13	1.7	?RXJ1627.1+7756	Cluster within 3', PtSource within 45"
$OC3 \ 1627 + 7753$	$16\ 27\ 46.31$	77 53 38.6	3.06	51.74	0.60	21.24	75	12.48	1.8		PtSource within 30"
OC4 1629+7811	$16\ 29\ 39.77$	$78 \ 11 \ 12.1$	3.12	28.70	0.30	19.43	127	6.41	1.4		PtSource within 15"; Blend OC4a, OC4b
OC4a 1629+7803	$16\ 29\ 41.64$	$78 \ 03 \ 30.0$	3.81	73.76	0.70	21.48	182	1.37	0.9		Blend with OC4, OC4b
OC4b 1629+7806	16 29 53.81	$78 \ 06 \ 36.5$	3.94	49.58	0.40	20.19	206	2.10	1.0		Blend with OC4, OC4a
$OC5 \ 1632 + 7806$	$16 \ 32 \ 23.64$	78 06 17.1	3.37	48.96	0.50	20.68	130	8.92	1.6		PtSource within 1'
$OC6 \ 1633 + 7808$	$16 \ 33 \ 39.75$	$78 \ 08 \ 35.8$	3.89	35.76	0.30	19.23	100	13.23	1.9		Near Edge, PtSource within 30"
-	RP800395N00 - 3C356										
OC1 1723+5059	17 23 14.13	50 59 25.1	3.36	38.02	0.50	20.75	61	10.37	1.6		Compact
OC2 1723+5108	$17 \ 23 \ 24.79$	$51 \ 08 \ 34.9$	4.49	41.06	0.20	18.03	191	13.89	1.9		PtSource within 1'
OC3 1725+5059	$17 \ 25 \ 05.24$	50 59 00.1	2.94	55.58	0.70	21.59	70	7.41	1.4		PtSource within 1.5',Low Sig
OC4 1725 + 5055	$17\ 25\ 25.99$	$50 \ 55 \ 21.9$	3.93	28.99	0.30	19.34	91	10.79	1.6		
					WF	701457N	00 - HS	51700 + 64	16		
OC1 1700+6413	17 00 41.94	$64 \ 13 \ 16.9$	3.83	91.04	0.30	19.29	209	2.35	0.8	RXJ1700.6+6413	Cluster $(z=0.22)$
OC2 1701+6412	17 01 30.22	$64 \ 12 \ 05.4$	3.15	99.89	0.70	21.51	109	3.29	0.8	RXJ1701.3+6414	Cluster (z=0.45)
OC3 1701+6421	17 01 48.77	$64 \ 21 \ 03.0$	3.99	62.42	0.20	18.20	260	10.49	1.4	RXJ1701.6+6421	Cluster
OC4 1702+6420	$17 \ 02 \ 13.58$	$64 \ 20 \ 07.3$	3.34	79.25	0.30	19.26	177	11.39	1.5	RXJ1702.2+6420	Cluster $(z=0.23)$

-61 -

I-band ID	$\Lambda_{cl}(I)$	z(I)	RA (J2000)	Dec (J2000)	Sigma	$\Lambda_{cl}(V)$	$\mathbf{z}(\mathbf{V})$	V-mag	Radius	${N_{gals}}^1$	$P(KS)^2$	
				0902+34 (1849 §	$_{\rm galaxies})^3$							
OC2 090510.2+335938 ^a	25.04	0.30	09 05 11.36	33 59 30.6	5.45	29.87	0.30	20.66	141	22	3e-4	
OC5 090557.0+340701	48.56	0.60	$09 \ 05 \ 44.01$	34 08 59.4	3.83	50.72	0.50	22.45	163	27	0.25	
OC6 090617.6+341716	28.87	0.30	$09 \ 06 \ 14.26$	$34 \ 18 \ 15.3$	5.01	27.45	0.30	20.58	124	12	0.015	
			$09 \ 06 \ 13.16$	$34 \ 16 \ 25.5$	3.62	18.44	0.20	19.44	138			
$OC7 090628.2 + 340512^{b}$	36.56	0.20	$09 \ 06 \ 14.70$	$33 \ 55 \ 57.1$	4.95	27.11	0.30	20.34	114			
			$09 \ 06 \ 14.76$	$34 \ 08 \ 45.4$	3.25	17.82	0.30	20.73	108			
				Pancake (1820 g	galaxies)							
OC1 090612.5+335422	25.13	0.20	09 06 22.21	33 52 14.2	3.93	32.10	0.30	20.45	134	29	9e-5	
OC2 090718.6+333044	85.19	0.50	$09 \ 07 \ 35.41$	$33 \ 32 \ 59.4$	3.26	26.62	0.30	20.88	183	16	0.20 ^e	
OC3 090727.7+334302	74.79	0.50	$09 \ 07 \ 23.35$	33 38 36.8	3.60	72.12	0.50	22.45	175	14	0.02	
OC5 090758.3+335010	67.27	0.70	$09 \ 07 \ 44.66$	$33 \ 48 \ 25.6$	4.39	87.85	0.50	22.57	161	10	0.39	
1202+281 (2397 galaxies)												
OC1 120347.8+275836	56.48	0.50	$12 \ 03 \ 53.36$	27 53 14.5	5.51	71.72	0.40	21.64	161	43	0.15	
			$12 \ 03 \ 44.70$	27 59 32.9	4.86	41.40	0.20	19.10	217			
OC2 120353.2+274355	39.62	0.40	$12 \ 04 \ 00.27$	$27 \ 43 \ 47.3$	3.36	43.74	0.40	21.93	113	37	0.068	
$OC4 \ 120436.7 + 280520^{\circ}$	95.30	1.00	$12 \ 04 \ 19.52$	$28 \ 07 \ 07.2$	4.41	155.55	1.10	24.15	95			
OC5 120438.0+280152	28.01	0.20	$12 \ 04 \ 36.76$	$28 \ 01 \ 38.8$	4.11	34.99	0.20	19.28	179	23	0.017	
OC7 120457.0+275305	90.66	1.00	$12 \ 04 \ 55.47$	27 53 35.6	*2.67	100.32	0.90	23.99	67			
OC9 120543.8+274030 ^C	50.47	0.40	$12 \ 05 \ 26.85$	$27 \ 44 \ 4.9$	3.49	29.69	0.20	19.51	104			
				3C324 (2145 ga	alaxies)							
OC2 154855.9+212722	38.51	0.40	15 48 55.83	21 28 29.5	*2.85	31.81	0.40	21.53	83	13	0.17	
OC2a 154902.2+211953	41.59	0.20	$15 \ 49 \ 1.13$	$21 \ 21 \ 16.6$	3.61	29.57	0.30	20.64	146	81	0.011	
OC2b 154902.4+212226	31.53	0.30										
$OC3 154932.1 + 212325^{d}$	70.35	0.70	$15 \ 49 \ 31.97$	$21 \ 23 \ 28.4$	4.23	477.83	1.20	24.11	136	29	0.179	
OC8 155044.7+213436	36.51	0.40	15 50 34.25	$21 \ 31 \ 53.1$	3.61	29.53	0.30	20.56	192	28	0.012	
				3C356 (1591 ga	alaxies)							
OC2 172324.8+510835	41.06	0.20	17 23 19.59	51 09 13.5	4.56	22.23	0.20	19.43	91	48	0.32	
OC4 172526.0+505522	28.99	0.30	$17\ 25\ 35.94$	$50\ 57\ 34.6$	4.08	19.66	0.20	19.46	100	9	0.14	

 1 X-ray flux units for the upper limits are 10^{-14} erg s⁻¹ cm⁻². Upper limits were computed for all optical candidates as a function of off-axis position in the ROSAT field and of ROSAT exposure time, regardless the presence of an X-ray counterpart.

 $^1N_{gals}$ is the number of galaxies near the centroid of the optical candidate used for the 2-d K-S test.

 2 The KS probability between 0 and 1 that the galaxies were randomly selected from the same parent population as the rest of the galaxies in the same field. Typically ~ 2000 galaxies were measured in the same field.

 3 Name of original target. (Number of galaxies in the optical field used as the parent population sample.)

^aSignificance in I-band is 2.94 σ , formally below a 3σ threshold.

 $^{\rm b}$ The center of the I band candidate was significantly obscured by scattered light in the V-band image, but the matched filter algorithm found a couple of cluster candidates in the unaffected regions of the V image.

 c These two cluster candidates overlap at the 3σ contour level, but the centroids differ significantly. These systems, if a true match, are filamentary and unlikely to be virialized.

 $^{\rm d}{\rm Note}$ significant difference in estimated redshifts between the I and V data.

^eThe joint photometry for this candidate was significantly distant from the X-ray core because of scattered light limitations.

Note. — This table lists the matched-filter parameters derived from the V-band for cluster candidates cross-identified with cluster candidates in the I-band. The units for the matched-filter quantities are the same as in Table 2, although Λ_{cl} and V-mag refer to V-band quantities. Note though that Λ_{cl} from the I and V bands should be similar, nevertheless. The **boldfaced** entries in column 1 are cluster candidates with X-ray counterparts (Table 6).

Band name	RA	Dec	Sigma	Lambda	\mathbf{z}	V-mag	Radius	Comment
				0	00124			
				05	902+34			
Ι	$09 \ 04 \ 23.88$	$34 \ 11 \ 04.9$	4.25	43.38	0.40	20.13	70	off V frame
Ι	$09 \ 05 \ 11.11$	$34 \ 21 \ 53.0$	5.14	34.79	0.20	18.39	116	off V frame
Ι	$09 \ 05 \ 52.95$	$34 \ 03 \ 17.5$	3.58	88.98	1.00	22.51	101	"high z"
Ι	$09 \ 06 \ 31.74$	33 59 51.1	4.05	70.13	0.70	21.65	165	cut from V frame
I	$09 \ 06 \ 32.70$	$34 \ 13 \ 17.4$	3.35	89.67	1.10	22.71	50	cut from V frame
V	09 04 33.06	34 15 30.1	4.07	22.32	0.30	20.25	79	<i>"</i>
V	09 04 54.98	34 02 33.6	3.64	248.79	1.20	24.15	87	"high z"
V	09 05 19.40	34 20 04.8	4.99	313.70	1.10	24.12	162	"high z" "Li-L -"
v	09 05 22.27	34 11 47.3	3.43	215.58	1.10	24.12	80	nign z
				Р	ancake			
Ι	$09 \ 07 \ 47.16$	$33 \ 55 \ 00.4$	3.50	28.21	0.30	19.47	68	off V frame
V	$09 \ 05 \ 40.86$	$33 \ 54 \ 05.0$	3.47	188.25	1.20	24.16	57	Very low sig $(2.1, z = 1.1)$ I candidate
V	$09 \ 05 \ 48.45$	$33 \ 32 \ 11.6$	3.86	17.20	0.20	19.03	97	
V	09 06 11.17	33 43 41.5	3.19	173.15	1.20	24.14	60	high z
V	09 06 41.30	33 27 41.1	3.50	155.25	1.00	24.07	78	high z
V	09 06 52.22	33 34 29.3	3.43	185.96	1.20	24.14	76	high z
V	09 06 57.94	33 54 04.2	4.58	248.20	1.20	24.15	75	high z
V	09 06 59.65	33 47 48.3	3.40	187.81	1.20	24.14	88 140	nign z
v	09 07 10.45	33 30 23.2	4.30	35.09	0.30	20.51	149	l
				12	02+281			
Ι	$12 \ 04 \ 26.02$	$27 \ 44 \ 12.8$	3.16	89.91	1.00	22.52	53	"high z"
Ι	$12 \ 04 \ 53.71$	$27 \ 57 \ 59.1$	3.21	39.02	0.40	20.32	92	
Ι	$12 \ 05 \ 18.49$	$28 \ 05 \ 17.1$	3.19	31.43	0.30	19.51	73	
V	$12 \ 03 \ 48.00$	$28 \ 06 \ 55.5$	2.97	111.49	0.90	23.94	48	"high z"
V	12 04 33.52	27 51 24.4	4.03	142.08	1.10	24.15	69	"high z"
V	12 05 16.06	28 06 42.6	4.97	175.41	1.10	24.15	113	"high z"
V	12 05 18.25	27 46 51.8	4.67	164.63	1.10	24.16	79	"high z"
				:	3C324			
I	$15 \ 48 \ 50.06$	$21 \ 13 \ 54.3$	3.39	39.86	0.50	20.76	106	
I	15 50 02.46	$21 \ 12 \ 37.5$	3.18	23.64	0.20	17.65	136	off V frame
Ι	$15 \ 50 \ 13.35$	$21 \ 25 \ 21.6$	3.64	113.13	1.10	22.71	87	"high z"
V	$15 \ 49 \ 01.37$	$21 \ 35 \ 35.8$	3.44	12.68	0.20	19.20	59	
V	$15 \ 49 \ 13.73$	$21 \ 32 \ 04.1$	3.51	396.85	1.20	24.08	60	"high z"
V	$15 \ 49 \ 33.91$	$21 \ 33 \ 20.6$	3.23	117.30	0.80	23.79	64	Very low sig $(2.3, z = 0.7)$ I candidate
V	15 50 03.67	21 19 01.6	3.29	371.34	1.20	24.08	57	cut out, "high z"
V	15 50 46.75	21 15 57.0	4.99	47.04	0.20	19.49	84	Very low sig $(2.6, z = 0.2)$ l candidate
				:	3C356			
I	17 23 14.13	50 59 25.1	3.36	38.02	0.50	20.75	61	
V	$17 \ 23 \ 00.86$	$50 \ 57 \ 51.2$	4.57	272.56	1.20	24.16	121	"high z"
V	$17 \ 23 \ 45.28$	$51 \ 02 \ 55.6$	3.75	45.64	0.40	21.53	98	
V	17 24 36.12	50 59 19.8	3.73	45.37	0.40	21.58	74	cut l region
V	$17 \ 24 \ 44.99$	$51 \ 05 \ 14.1$	6.00	357.66	1.20	24.16	135	"high z"

Table 4. Unmatched Cluster Candidates in V and I

Table 5: Summary of Cross-Identification Statistics for I- and V-band Matched Filter Candidates

	I-band	V-band
Total number of candidates detected in 5 fields $(> 3\sigma)$	33	46
Candidates unavailable in other band (scattered light)	6	2
"High redshift" candidates undetected in other band	4	15
Net viable candidates in 5 fields ¹	23	29
Candidates with one or more counterparts in other $band^2$	16	18
Cross ID efficiency	70%	62%

¹V and I counterparts were counted if and only if both candidates had detection confidence of $> 3\sigma$ and the centroids were closer than $\sim 3 - 4'$ in both RA and Dec. Overlapping matched filter candidates with centroids this close have contours which overlap significantly.

 $^{^{2}}$ The net number of viable candidates is the difference between the total number of candidates and the sum of the candidates unavailable in the other band and the "high redshift" candidates.

Xray ID	RA J2000	Dec J2000	Counts	Theta (')	FWHM (")	Sig-ext (σ)	Fx^1	eFx^1	Comment	Optical Match	Comment
							RP7001	12			
RXJ0743.7+6457	$07 \ 43 \ 45.01$	+64 57 19.4	52.6	15.2	90.2	5.4	3.51	1.60		OC3 0743+6458	z=0.4, within 1'
						F	RP700326	N00			
RXJ0906.3+3417	$09 \ 06 \ 18.73$	$+34 \ 17 \ 24.8$	17.4	13.8	67.2	2.0	1.50	0.95		OC6 0906+3417	z=0.3, within 15"
							RP9003	27			
RXJ0907.3+3330 RXJ0907.4+3342 RXJ0907.8+3351	09 07 18.26 09 07 27.10 09 07 51.76	+33 30 28.1 +33 42 32.0 +33 51 26.6	$61.3 \\ 32.9 \\ 74.6$	$11.2 \\ 7.6 \\ 16.6$	$54.1 \\ 73.9 \\ 74.2$	2.2 5.9 2.5	$2.84 \\ 0.94 \\ 4.27$	$0.46 \\ 0.35 \\ 0.50$	(0.470?)	OC2 0907+3330 OC3 0907+3343 OC5 0907+3350	z=0.5, within 30" z=0.5, within 30" z=0.7, within 2'
							WP7005	40			
RXJ1025.4+4703 RXJ1025.8+4709	$\begin{array}{cccccccccccccccccccccccccccccccccccc$	$^{+47\ 03\ 44.4}_{+47\ 09\ 01.1}$	$30.6 \\ 53.6$	$\begin{array}{c} 10.3 \\ 13.2 \end{array}$	$\begin{array}{c} 69.4 \\ 90.7 \end{array}$	$4.6 \\ 6.3$	$1.84 \\ 3.21$	$\begin{array}{c} 0.38\\ 1.31 \end{array}$	d	OC11 1025+4701	z=0.2, within 3' O. Half of area excluded
						v	VP201243	N00			
							WP7002	28			
RXJ1118.9+2117 RXJ1119.2+2117 RXJ1119.4+2106 RXJ1119.7+2126 RXJ1120.0+2115	11 18 59.90 11 19 16.73 11 19 25.41 11 19 43.21 11 20 02.45	$\begin{array}{r} +21 \ 17 \ 56.3 \\ +21 \ 17 \ 32.3 \\ +21 \ 06 \ 44.3 \\ +21 \ 26 \ 36.9 \\ +21 \ 15 \ 10.2 \end{array}$	$93.7 \\ 142.5 \\ 162.7 \\ 93.3 \\ 32.5$	2.6 2.4 13.1 10.8 13.0	53.8 73.5 60.0 55.0 63.3	$ \begin{array}{r} 4.0 \\ 6.6 \\ 2.3 \\ 2.5 \\ 2.8 \end{array} $	$\begin{array}{c} 4.55 \\ 6.92 \\ 8.38 \\ 4.68 \\ 1.68 \end{array}$	$\begin{array}{r} 4.98 \\ ?21.00 \\ 0.89 \\ 0.69 \\ 0.64 \end{array}$	(0.176) (0.061)	OC3 1119+2116 OC4 1119+2107 OC6 1119+2127	z=0.5,within 30" z=0.2,within 30" z=0.4,within 1.5',(z-spec=f.g.galaxy)
						W	VP201367	M01			
RXJ1204.3-0350 RXJ1205.0-0332	$\begin{array}{c} 12 \ 04 \ 22.58 \\ 12 \ 05 \ 02.23 \end{array}$	-03 50 53.9 -03 32 25.1	$207.1 \\ 54.0$	$\begin{array}{c} 10.7 \\ 12.1 \end{array}$	$\begin{array}{c} 74.2 \\ 60.3 \end{array}$	$5.1 \\ 2.8$	$9.83 \\ 2.69$	$\begin{array}{c} 1.16 \\ 0.55 \end{array}$	(0.261)	OC5 1204-0351 ?OC8 1204-0330	z=0.2,within 30" z=0.9,within 2',(opt sigma=2.1)
							WP7002	32			
RXJ1204.1+2807 RXJ1205.2+2752	$\begin{array}{c} 12 \ 04 \ 03.15 \\ 12 \ 05 \ 16.06 \end{array}$	$^{+28\ 07\ 03.1}_{+27\ 52\ 48.9}$	$740.2 \\ 64.5$	$15.7 \\ 7.4$	$79.9 \\ 65.3$	$3.8 \\ 4.8$	32.83 2.79	$0.75 \\ 0.31$		OC1,4,5 ³	Area excluded; MS1201+2823/A1455 $(z=0.167)$
						F	RP700864	A01			
RXJ1220.0+3334 RXJ1220.8+3343 RXJ1220.9+3343	$ \begin{array}{r} 12 & 20 & 01.31 \\ 12 & 20 & 52.82 \\ 12 & 20 & 54.48 \end{array} $	+33 34 46.4+33 43 50.6+33 43 52.2	49.2 8.2 33.9	10.8 4.1 4.4	53.0 88.4 84.2	2.2 8.5 7.8	$3.05 \\ 0.50 \\ 2.06$	$0.29 \\ 0.45 \\ 1.18$	d d d	?OC3 1220+3334 OC4 1221+3344 OC4 1221+3344	z=1.2,within 1.5',(opt sigma=2.6) z=0.3,within 4' same Xray cluster as above?

Table 6. X-ray Clusters of Galaxies

Xray ID	RA J2000	Dec J2000	Counts	Theta (')	FWHM (")	$\begin{array}{c} \text{Sig-ext} \\ (\sigma) \end{array}$	Fx^1	eFx^1	Comment	Optical Match	Comment	
RP600242A01												
RXJ1227.8+0143 RXJ1228.5+0134	$\begin{array}{cccccccccccccccccccccccccccccccccccc$	$+01 \ 43 \ 37.9 \\ +01 \ 34 \ 42.7$	$39.3 \\ 39.0$	7.9 11.8	$87.8 \\ 66.1$	$7.7 \\ 3.6$	$\begin{array}{c} 1.88 \\ 1.91 \end{array}$	$0.79 \\ 0.38$		OC1 1227+0143	z=0.3,within 45" Most of area excluded	
RP700073												
RXJ1255.6+4712 RXJ1256.6+4715 RXJ1256.8+4727 RXJ1256.9+4720 RXJ1257.0+4738 RXJ1257.3+4729 RXJ1257.6+4737 RXJ1257.7+4723	$\begin{array}{c} 12 \ 55 \ 36.25 \\ 12 \ 56 \ 38.56 \\ 12 \ 56 \ 53.16 \\ 12 \ 56 \ 57.02 \\ 12 \ 57 \ 05.17 \\ 12 \ 57 \ 22.20 \\ 12 \ 57 \ 36.97 \\ 12 \ 57 \ 43.50 \end{array}$	$\begin{array}{r} +47 \ 12 \ 02.5 \\ +47 \ 15 \ 29.8 \\ +47 \ 27 \ 23.5 \\ +47 \ 20 \ 46.7 \\ +47 \ 38 \ 17.1 \\ +47 \ 29 \ 56.1 \\ +47 \ 37 \ 06.3 \\ +47 \ 23 \ 21.3 \end{array}$	$186.5 \\ 213.5 \\ 40.8 \\ 53.3 \\ 223.8 \\ 77.8 \\ 153.4 \\ 64.6 \\$	$16.1 \\ 5.9 \\ 7.0 \\ 0.3 \\ 17.9 \\ 10.4 \\ 18.0 \\ 8.4$	89.576.764.255.3113.869.0130.953.4	$\begin{array}{c} 4.8 \\ 6.6 \\ 4.8 \\ 4.2 \\ 7.1 \\ 4.5 \\ 9.4 \\ 3.0 \end{array}$	$\begin{array}{c} 4.60 \\ 5.46 \\ 1.01 \\ 1.29 \\ 7.02 \\ 1.97 \\ 4.67 \\ 1.56 \end{array}$	$\begin{array}{c} 0.62 \\ 0.83 \\ 0.67 \\ 0.27 \\ 0.86 \\ 0.73 \\ 0.92 \\ 0.22 \end{array}$	(0.410) $(0.997)^2$?OC1 1257+4719	Off frame Optical area excluded Optical area excluded z=1.0,within 1.5',(opt sigma=2.0) Off frame Half of area excluded	
						RP70	0216A00					
RXJ1309.9+3222 RXJ1310.5+3217 RXJ1313.2+3229	$\begin{array}{c} 13 09 55.79 \\ 13 10 35.95 \\ 13 11 12.44 \end{array}$	$\begin{array}{r} +32 \ 22 \ 23.0 \\ +32 \ 17 \ 36.0 \\ +32 \ 29 \ 07.6 \end{array}$	$29.0 \\ 17.5 \\ 178.4$	$7.1 \\ 3.7 \\ 12.3$	$76.5 \\ 54.9 \\ 66.7$	$6.4 \\ 4.0 \\ 3.5$	4.06 2.44 24.95	$1.00 \\ 0.63 \\ 1.75$		OC2 1310+3221 OC4 1311+3228	z=0.3,within 2' (MS1308.8+3244, $z=0.245$) Star spike; area excluded z=0.2,within 30"	- 99
RP700117												
RXJ1407.6+3415	$14\ 07\ 39.76$	$+34 \ 15 \ 11.0$	29.1	12.3	58.6	2.1	1.53	0.37		OC7 1407+3415	z=0.5, within 30"	
						WP	700248					
RXJ1412.6+4359 RXJ1413.5+4411	$\begin{array}{c} 14 \ 12 \ 36.30 \\ 14 \ 13 \ 30.94 \end{array}$	$^{+43}_{+44} \begin{array}{c} 59 \\ 02.9 \\ +44 \\ 11 \\ 44.2 \end{array}$	$39.4 \\ 32.6$	$\begin{array}{c} 12.9 \\ 12.1 \end{array}$	$76.9 \\ 81.1$	$4.6 \\ 5.5$	$\begin{array}{c} 1.86 \\ 1.51 \end{array}$	$0.89 \\ 0.61$			Optical area excluded Optical area excluded	
						RP	700122					
RXJ1415.2+1119	$14 \ 15 \ 15.77$	$+11 \ 19 \ 32.6$	75.2	12.5	75.7	4.6	3.18	0.63		?OC1 1414+1123	z=0.3, within 7'	
RP800401A01												
$\begin{array}{c} \text{RXJ1415.8+2316} \\ \text{RXJ1415.9+2307} \\ \text{RXJ1416.3+2309} \\ \text{RXJ1416.4+2315} \\ \text{RXJ1416.4+2302} \end{array}$	$\begin{array}{c} 14 \ 15 \ 50.54 \\ 14 \ 15 \ 57.16 \\ 14 \ 16 \ 22.51 \\ 14 \ 16 \ 26.88 \\ 14 \ 16 \ 28.28 \end{array}$	$\begin{array}{r} +23 \ 16 \ 10.9 \\ +23 \ 07 \ 37.4 \\ +23 \ 09 \ 59.6 \\ +23 \ 15 \ 37.2 \\ +23 \ 02 \ 31.6 \end{array}$	$ 19.0 \\ 99.1 \\ 28.1 \\ 499.0 \\ 15.8 $	$9.1 \\ 0.4 \\ 6.4 \\ 10.8 \\ 8.5$	$ \begin{array}{r} 64.2 \\ 67.9 \\ 66.5 \\ 109.6 \\ 55.7 \\ \end{array} $	4.2 5.9 5.2 9.8 3.3	2.01 10.20 2.90 51.34 1.67	$ 1.55 \\ 0.78 \\ 0.69 \\ 1.53 \\ 1.16 $		OC2 1415+2317 OC3 1415+2307	z=0.4,within 2.5' z=0.3,within 30" Optical area excluded	

Table 6—Continued

RP700257N00

Xray ID	RA J2000	Dec J2000	Counts	Theta (')	FWHM (")	$\begin{array}{c} \text{Sig-ext} \\ (\sigma) \end{array}$	Fx^1	eFx^1	Comment	Optical Match	Comment	
						RP70	1373N00					
RXJ1548.8+2126	$15 \ 48 \ 52.57$	$+21 \ 26 \ 06.1$	61.9	13.4	110.8	8.9	5.08	2.42	d	OC2 1548+2127	z=0.4, within 1.5'	
						RP80	0239N00					
RXJ1603.6+4316	$16\ 03\ 39.01$	$+43 \ 16 \ 19.1$	53.7	9.6	56.1	3.0	2.17	0.36				
						RP30	0021N00					
RXJ1605.0+2552	$16\ 05\ 04.66$	$+25\ 52\ 47.4$	32.9	9.3	52.8	2.7	1.76	0.68	d	?OC2 1604+2554	z=0.6, within 2.5', Cluster center excluded	
RXJ1605.5+2553	$16 \ 05 \ 30.99$	$+25\ 53\ 16.7$	40.0	3.8	74.8	6.5	2.06	1.10	d	OC4 1605+2553	z=0.2, within 1.5'	
RXJ1605.6 + 2548	$16 \ 05 \ 41.71$	+25 48 27.7	39.6	3.2	75.7	6.8	2.04	1.05				
RXJ1606.1 + 2558	$16\ 06\ 10.83$	$+25\ 58\ 42.8$	22.0	9.1	55.8	3.1	1.17	0.69				
						WP	170154					
RXJ1626.3+7816	$16\ 26\ 23.25$	$+78 \ 16 \ 59.6$	303.9	15.5	87.0	4.8	10.63	0.62	(0.580)	OC1 1626+7817	z=0.5, within 15"	19
RXJ1627.1+7756	$16\ 27\ 10.77$	+77 56 06.8	102.7	11.4	74.1	4.8	3.47	0.49	d	?OC2 1626+7759	z=0.3, within 3'	
RXJ1629.7+7757	$16\ 29\ 45.43$	+77 57 59.5	122.2	6.9	78.6	6.7	4.12	0.64	d			
RXJ1629.9+7819	$16\ 29\ 56.33$	+78 19 18.6	92.2	14.5	98.6	6.8	3.11	0.75			Off frame	
RXJ1630.2+7815	$16 \ 30 \ 12.72$	$+78 \ 15 \ 35.3$	44.1	10.9	59.9	3.1	1.54	0.57			Optical area excluded	
						RP80	0395N00					
						WP70	1457N00					
RXJ1700.6+6413	17 00 40.58	$+64 \ 13 \ 04.5$	551.7	2.3	44.6	2.8	24.54	0.44	(0.22)	OC1 1700+6413	z=0.3	
RXJ1701.3+6414	$17 \ 01 \ 21.31$	+64 14 16.5	428.3	3.2	53.0	3.8	19.05	0.44	(0.45)	OC2 1701+6412	z=0.7	
RXJ1702.0+6407	$17 \ 02 \ 00.94$	$+64 \ 07 \ 39.0$	35.0	8.0	58.2	3.7	1.56	0.25				
RXJ1701.6+6421	$17 \ 01 \ 42.96$	$+64 \ 21 \ 20.2$	42.7	10.4	83.5	6.4	1.89	0.75		OC3 1701+6421	z=0.2	
RXJ1702.2+6420	$17 \ 02 \ 10.14$	$+64 \ 20 \ 00.3$	36.1	11.1	60.7	3.2	1.65	0.37	(0.23)	OC4 1702 + 6420	z=0.3	

Table 6—Continued

 1 Flux units are 10^{-14} erg s⁻¹ cm⁻².

²The extended X-ray source RXJ1256.9+4720 is likely to be associated with the radio galaxy 3C280, the target of the original observation. The X-rays could be from the galaxy or a cluster surrounding the galaxy.

 3 This source (Abell 1455) was obscured by a nearby F3 star in the I-band data, but 3 optical candidates were identified nearby. The matched filter algorithm may have picked up the outskirts of this cluster.

Table 7: Spectroscopic vs. Matched Filter Estimated Redshifts

Xray ID	Optical ID	Spec z	Est z	Comments
RXJ0907.4+3342	OC3 0907+3343	0.470	0.5	
RXJ1119.4+2106	OC4 1119+2107	0.176	0.2	
RXJ1119.7+2126	OC6 1119+2127	0.061	0.4	foreground galaxy
RXJ1204.1+2807	OC1 1203+2758	0.167	0.5	$MS1201 + 2823/A1455^{1}$
	OC4 1204+2805	0.167	1.0	$MS1201 + 2823/A1455^{1}$
	$OC5\ 1204 + 2801$	0.167	0.2	$MS1201 + 2823/A1455^1$
RXJ1204.3-0350	OC5 1204-0351	0.261	0.2	
RXJ1256.9+4720	OC1 1257+4719	0.997	1.0	Optical 2.0σ ; $3C280$
RXJ1309.9+3222	OC2 1310+3221	0.245	0.3	EMSS1308.8+3244
RXJ1626.3+7816	OC1 1626+7817	0.580	0.5	
RXJ1700.7+6413	OC1 1700+6413	0.220	0.3	
RXJ1701.4+6414	OC2 1701+6412	0.45	0.7	
RXJ1702.2+6420	OC4 1702+6420	0.23	0.3	

 $^{^{1}}$ The X-ray cluster was obscured by the diffraction spike of an F3 star; the matched filter algorithm found 3 candidates in the vicinity, none close enough for an official match.