

Dust-to-gas ratio and star formation history of blue compact dwarf galaxies

Hiroyuki Hirashita^{1,*}, Yuka Y. Tajiri², and Hideyuki Kamaya^{2,**}

¹ Osservatorio Astrofisico di Arcetri, Largo E. Fermi, 5, 50125 Firenze, Italy
e-mail: irasita@arcetri.astro.it

² Department of Astronomy, Faculty of Science, Kyoto University, Sakyo-ku, Kyoto 606-8502, Japan
e-mail: tajiri, kamaya@kustastro.kyoto-u.ac.jp

Received 25 July 2001 / accepted 29 March 2002

Abstract. This paper investigates the origin of the observed large variety in dust-to-gas ratio, \mathcal{D} , among blue compact dwarf galaxies (BCDs). By applying our chemical evolution model, we find that the dust destruction can largely suppress the dust-to-gas ratio when the metallicity of a BCD reaches $12 + \log(O/H) \sim 8$, i.e., a typical metallicity level of BCDs. We also show that dust-to-gas ratio is largely varied owing to the change of dust destruction efficiency that has two effects: (i) a significant contribution of Type Ia supernovae to total supernova rate; (ii) variation of gas mass contained in a star-forming region. While mass loss from BCDs was previously thought to be the major cause for the variance of \mathcal{D} , we suggest that the other two effects are also important. We finally discuss the intermittent star formation history, which naturally explains the large dispersion of dust-to-gas ratio among BCDs.

Key words. ISM: dust, extinction — galaxies: dwarf — galaxies: evolution — galaxies: ISM — stars: formation

1. Introduction

Dust grains absorb stellar ultraviolet–optical light and emit far-infrared (FIR) light, thereby affecting the spectral energy distributions of galaxies (e.g., Takagi et al. 1999). Since the spectral energy distribution is frequently analysed to infer the star formation history (SFH) that provides us with a key to understand the evolutionary history of galaxies, the research on the origin of the interstellar dust is important to this issue. A key quantity concerning grains is the dust-to-gas mass ratio, \mathcal{D} . Oort & van de Hulst (1946) have observationally shown a strong correlation between the densities of gas and dust. This indicates that dust traces dense environments, which should be rich in heavy elements. Thus, it is worth examining the evolution of \mathcal{D} in the context of the chemical evolution of galaxies.

The condensation of heavy elements is an important process for the formation of grains. One of the environments where condensation takes place is the atmosphere

of a cool giant star (Hoyle & Wickramasinghe 1963). High-dispersion spectroscopic observation of C_2 in post AGB stars may indicate the condensation process on C_2 as a dust kernel (Crawford & Barlow 2000; Kameswara Rao & Lambert 2000). Another environment for condensation is a supernova (SN). Dwek & Scalo (1980) have shown that SNe can be the dominant source of dust grains. This is partly because of a rich metal content in SNe. Indeed, a significant amount of dust is observed within hot SN remnants (Dwek et al. 1983; Moseley et al. 1989; Kozasa et al. 1989) and theoretical work by Todini & Ferrara (2001) has explained some principal features of dust formation in SN 1987A. Hirashita (1999a) has shown that observational dust amounts suggest that $\sim 10\%$ (the fraction in mass) of the heavy elements ejected from stars condense into dust grains. Since his model prediction reproduces the observed trend between dust-to-gas ratio and metallicity of dwarf galaxies, we adopt this fraction for dust condensation.

The processes of dust destruction should also be considered. As shown in Dwek & Scalo (1980), dust grains are not only made from heavy elements but also destroyed in SN shocks (see also McKee 1989; Jones et al. 1996) in a cycle of the birth and death of stars. In short, \mathcal{D} of a galaxy reflects its SFH via the regulation of dust formation and destruction. To investigate what determines the value of \mathcal{D} , investigation of a star-forming galaxy is the most inter-

Send offprint requests to: H. Hirashita

* Research Fellow of the Japan Society for the Promotion of Science.

** Visiting Academics at Department of Physics, University of Oxford, Keble Road, Oxford OX1 3RH, UK (1st March – 31st December 2001)

esting since SNe are expected to affect most largely their dust amount.

One of the observational features which give us a key to understand the regulation of \mathcal{D} is the variance of \mathcal{D} itself (e.g., Lisenfeld & Ferrara 1998, hereafter LF98). In this paper we examine the variance of \mathcal{D} among galaxies. According to LF98, there is a large variance in \mathcal{D} for a sample of blue compact dwarf galaxies (BCDs). By definition, galaxies categorised as BCD more or less show active star-forming activity. Therefore, dust formation and destruction are expected to occur in BCDs. Since grains are composed of heavy elements, application of the theory of galactic chemical evolution is useful and interesting. Then, we particularly consider the variance of \mathcal{D} in BCDs by using the chemical evolution model developed by LF98 and Hirashita (1999b, hereafter H99).

LF98 applied a chemical evolution model to explain \mathcal{D} of dwarf galaxies including BCDs. They have suggested that if the dust-to-gas ratio in outflow (galactic wind) is different from that in the interstellar medium (ISM), the large dispersion of \mathcal{D} of dwarf galaxies can be explained. In this scenario, a significant outflow of gas is indispensable to explain the observed variance of dust-to-gas ratio of BCDs. However, is the variance of \mathcal{D} determined simply by mass outflow? In this paper, after further investigation in the framework of LF98, we answer this question and point out that other factors are also important to explain the value and variance of \mathcal{D} .

Indeed, a recent analysis by Tajiri & Kamaya (2002) has suggested that outflow is not so efficient for BCDs. They estimated the current momentum supply from SNe by using $H\alpha$ luminosity, and concluded that the supplied momentum is not sufficient to blow away the $H\text{ I}$ envelopes surrounding star-forming regions and that BCDs do not currently suffer significant mass loss. Legrand et al. (2001) also suggested that low density halos around BCDs can be an obstacle for the ISM to escape from the galaxies themselves. Thus, it is worth examining mechanisms other than mass outflow. Since Tajiri & Kamaya (2002) adopted a sample in Sage et al. (1992), we also use the sample. Moreover, the sample in LF98 is also included because this paper is an extended study of LF98.

This paper is organised as follows. First, in Sect. 2 we explain the model that describes the evolution of dust content in a galaxy. In Sect. 3, we consider dust destruction, which is the most important process in this paper. Then, in Sect. 4, model predictions are presented in comparison with observations. In Sect. 5, we discuss the results and propose a physical mechanism that can explain the observations. Finally, we summarise the contents of this paper.

2. Model description

In order to consider the dust formation and destruction, we analyse the dust-to-gas ratio along with the chemical evolution model by H99. The model is based on Eales & Edmunds (1996), LF98 and Dwek (1998). In our model,

we do not need to model any SFH. This has an advantage in considering the dust-to-gas ratio of BCDs since the SFH of a dwarf galaxy is generally complex (e.g., Grebel 2001) and is difficult to model. We focus especially on dust destruction, because LF98 have not fully considered it. As we see later, our model is a powerful tool to know the metallicity level where dust destruction becomes effective enough to suppress the dust-to-gas ratio.

2.1. Brief review of the model

In order to investigate dust content in a galaxy, H99 has established a set of model equations describing dust formation and destruction processes. In H99, a galaxy is treated as one zone to focus on the quantities averaged over the whole galaxy. The galaxy is assumed to be a closed system; that is, mass inflow and outflow are not considered. If the metallicity of the infalling material is zero or much lower than that of the ISM in the galaxy, the relation between dust-to-gas ratio and metallicity, with which we will be concerned in this paper, is not altered by infall (Edmunds 2001; Hirashita 2001). This is because the infall dilutes both metallicity and dust-to-gas ratio at almost the same rate. Our model does not include the effect of outflow, and this is different to LF98, in which outflow is essential to explain the observed variance of the dust-to-gas ratio in BCDs. Since Tajiri & Kamaya (2002) and Legrand et al. (2001) have suggested that outflow is not efficient for BCDs, it is worth examining a case of no outflow. Indeed, we present another clear possibility to explain the large scatter of \mathcal{D} among BCDs later.

The model equations in H99 (see the paper for details; see also LF98) describe the evolution of total gas mass (M_g), the total mass of metals (both in gas and dust phases) labeled as i (M_i ; $i = \text{O, C, Fe, etc.}$), and the mass of metal i in a dust phase ($M_{d,i}$). We neglect dust growth in clouds, since Hirashita (1999a) has shown that this process in low-metallicity systems such as dwarf galaxies is much less efficient than the formation of dust around stars. Then, we adopt an instantaneous recycling approximation as in LF98 and H99 according to the formalism in Tinsley (1980): Stars less massive than m_t (present turn-off mass set to be $1 M_\odot$) live forever and the others die instantaneously.

2.2. Solution of the model

Dust-to-gas ratio and metallicity of galaxies are observationally known to correlate with each other (e.g., Issa et al. 1990). This relation has recently been used as a test for chemical evolution models including dust formation and destruction (LF98; H99; Hirashita 1999a; Edmunds 2001). The model by H99 reduces the following differential equation:

$$\mathcal{Y}_i \frac{d\mathcal{D}_i}{dX_i} = f_{\text{in},i}(\mathcal{R}X_i + \mathcal{Y}_i) - (\mathcal{R} + \beta_{\text{SN}})\mathcal{D}_i, \quad (1)$$

where \mathcal{D}_i is the mass fraction of an element i locked up in dust (i.e., $\mathcal{D}_i \equiv M_{d,i}/M_g$); X_i is the mass fraction of an element i (i.e., $X_i \equiv M_i/M_g$); $f_{in,i}$ quantifies what fraction of an element i ejected from stars condenses into dust grains; β_{SN} is “dust destruction efficiency,” which we explain later; \mathcal{R} is the fraction of stellar mass subsequently returned to the interstellar space, and \mathcal{Y}_i is the mass fraction of an element i newly produced and ejected by stars.¹ The definition of β_{SN} is as follows:

$$\beta_{\text{SN}} = \frac{M_g}{\tau_{\text{SN}}\psi}, \quad (2)$$

where τ_{SN} is the timescale of dust destruction by SN shocks (Eq. 5), and ψ is the star formation rate (SFR). Hereafter, β_{SN} is called “dust destruction efficiency”, because it is inversely proportional to the timescale of dust destruction (τ_{SN}). This is the most important parameter in this paper.

When all the quantities except X_i and \mathcal{D}_i are constant in time, the analytical solution obtained by LF98 is applicable. With our notations, it is rewritten as

$$\mathcal{D}_i(X_i) = \frac{b}{a}X_i + (1 - e^{-aX_i}) \left(\frac{c}{a} - \frac{b}{a^2} \right), \quad (3)$$

where $a \equiv (\mathcal{R} + \beta_{\text{SN}})/\mathcal{Y}_i$, $b \equiv f_{in,i}\mathcal{R}/\mathcal{Y}_i$, and $c \equiv f_{in,i}$.

Here, we select oxygen as a traced element (i.e., $i = \text{O}$) according to LF98, because (i) most of the oxygen is produced by massive stars (Type II SNe and their progenitors), (ii) oxygen is one of the main constituents of dust grains, and (iii) the common tracer for the metal abundance in BCDs is an oxygen emission line. The first item (i) means that an instantaneous recycling approximation may be reasonable for the investigation of oxygen abundances, since the generation of oxygen is a massive-star-weighted phenomenon. In other words, results are insensitive to the value of m_t . Following H99, we adopt $(\mathcal{R}, \mathcal{Y}_\text{O}) = (0.32, 7.2 \times 10^{-3})$, which are consistent with the relation between dust-to-gas ratio and metallicity of nearby galaxies including our BCD sample.

3. Destruction efficiency of dust

In this paper, we discuss the variance of dust-to-gas ratio of a BCD sample in terms of the variation of β_{SN} defined in Eq. (2). We estimate β_{SN} based on McKee (1989) and LF98, while we also address their differences.

We assume that gas is divided into two components: gas in the star-forming region and that in the H I envelope. Such an envelope is generally observed around a star-forming region of a BCD (e.g., van Zee et al. 1998). We denote the gas mass fraction in the star-forming region as X_{SF} and that in the H I envelope as X_{HI} (i.e., $X_{\text{SF}} + X_{\text{HI}} = 1$). We distinguish the two regions for the comparison with the *IRAS* sample (Sect. 4.1). *IRAS* FIR

¹ $\mathcal{R} = R$ and $\mathcal{Y}_i = y(1 - R)$ for the notation in LF98. These two parameters are calculated in the formalism of the instantaneous recycling approximation.

bands are sensitive to dust hotter than about 25 K. Such “warm” dust exists in star-forming regions, not in H I envelopes. Calzetti et al. (1995) have also shown by using the *IRAS* sample of actively star-forming galaxies that 70% of the FIR flux comes from such a warm component of dust. Thus, dust mass derived from the *IRAS* observation of a BCD is considered to trace the dust in the star-forming region. This suggests that it is useful for us to consider dust contained in star-forming regions as long as we are interested in the comparison of our result with the *IRAS* observations.

Thus, we estimate τ_{SN} and β_{SN} in star-forming regions. Gas mass accelerated to a velocity of v_s by a SN, $M_s(v_s)$, is estimated as

$$M_s(v_s) = 6800 \frac{E_{51}}{v_{s7}^2} M_\odot, \quad (4)$$

where E_{51} is energy released by a SN in units of 10^{51} erg and v_{s7} is v_s in units of 10^7 cm s⁻¹ (McKee 1989). In swept ISM, dust is not fully destroyed. Thus, the fraction of destroyed dust, ϵ (~ 0.1 ; McKee 1989), should be multiplied. Since we are interested in mass swept by multiple SNe in a star-forming region, SN rate must be considered, which is denoted as γ . Then, τ_{SN} is expressed as

$$\tau_{\text{SN}} = \frac{M_g X_{\text{SF}}}{\epsilon \gamma M_s (100 \text{ km s}^{-1})}. \quad (5)$$

Here, we substitute v_s with the threshold velocity for dust destruction (100 km s^{-1} ; McKee 1989). Comparing equations (2) and (5), we obtain

$$\beta_{\text{SN}} = \epsilon M_s (100 \text{ km s}^{-1}) \frac{\gamma}{\psi} \frac{1}{X_{\text{SF}}}. \quad (6)$$

We should consider both Type Ia and II SNe in estimating γ . Type II SNe occur “soon” after a star formation, because the progenitors of Type II SNe have lifetimes much shorter than the age of the universe. On the other hand, Type Ia SNe, whose progenitors are low-mass stars, occur about 1 Gyr after a star formation. Thus, γ and β_{SN} should be sensitive to the SFH. We express γ as

$$\gamma = \gamma_{\text{Ia}} + \gamma_{\text{II}}, \quad (7)$$

where γ_{Ia} and γ_{II} are the rate of Type Ia and Type II SNe, respectively. We note that LF98 estimated β_{SN} to be 5. But, for example, the fraction between γ_{Ia} and γ_{II} changes according to SFH of BCDs. In the next section, we examine the relation between dust-to-gas ratio and metallicity for various β_{SN} , whose probable range is constrained.

4. Sample and result

4.1. Observational samples

We compare the model calculation in Eq. (3) with the observed values of dust-to-gas ratio and metallicity of BCDs. We adopt the data in Table 2 of LF98 and Table 1 of Sage et al. (1992). The latter sample has been adopted by Tajiri

Table 1. Data of Blue Compact Dwarf Sample

Object	D (Mpc)	S_{60} (Jy)	S_{100} (Jy)	$\log M_{\text{HI}}$ [M_{\odot}]	$\log M_{\text{d}}^{\text{IRAS}}$ [M_{\odot}]	12 + (O/H)	ref. ^a
II Zw 40	9.1	6.6	5.8	8.30	4.55	8.15	S92
Haro 2	20.3	4.8	5.5	8.68	5.39	8.4	S92
Haro 3	13.7	5.2	6.7	8.76	5.21	8.3	S92
UM 439	12.6	0.39	1.2	8.23	4.96	7.98	S92
UM 462	11.9	0.99	1.1	8.15	4.21	7.89	S92
UM 465	13.2	0.99	1.3	7.71	4.48	8.9	S92
UM 533	10.4	0.51	0.54	7.76	3.75	8.10	S92
UM 448	6.00	4.14	4.32	7.54	4.16	8.08	LF98
IC 3258	21.2	0.490	0.970	8.55	5.03	8.44	LF98
Mrk 7	42.3	0.480	0.970	9.56	5.64	8.54	LF98
Mrk 33	21.6	4.68	5.30	8.77	5.41	8.40	LF98
Mrk 35	14.5	4.95	6.74	8.73	5.29	8.30	LF98
Mrk 450	12.1	0.480	0.820	7.77	4.37	8.21	LF98
NGC 4670	12.1	2.63	4.47	8.52	5.10	8.30	LF98
NGC 4861	12.9	1.97	2.26	9.13	4.60	8.08	LF98
II Zw 70	17.6	0.710	1.24	8.56	4.89	8.11	LF98

^a S92 — Sage et al. (1992); LF98 — Lisenfeld & Ferrara (1998)

& Kamaya (2002). We select BCDs whose 21-cm H I emission and FIR dust emission are both detected. The data are summarised in Table 1.

The observational dust-to-gas ratio in a star-forming region should be

$$\mathcal{D}^{\text{obs}} \equiv M_{\text{d}}^{\text{IRAS}} / (M_{\text{HI}} X_{\text{SF}}), \quad (8)$$

where $M_{\text{d}}^{\text{IRAS}}$ and M_{HI} are dust mass determined from the *IRAS* observation and the total H I gas determined from the 21-cm observations. M_{HI} traces H I gas in both the star-forming region and the envelope of a galaxy. On the other hand, $M_{\text{d}}^{\text{IRAS}}$ traces the amount of dust in the star-forming region, because the *IRAS* is sensitive to high-temperature ($\gtrsim 25$ K) dust. In the following two paragraphs, we describe the two quantities further.

LF98 divided the observed H I mass by a factor of 2 to obtain the gas mass in only the star-forming region. In other words, LF98 assumed that X_{SF} is 0.5. However, X_{SF} is hardly constrained observationally. In this paper we assume $X_{\text{SF}} = 1$ for the observational sample to obtain the first result. As we discuss in Sect. 5, the variation of X_{SF} also contributes to the variation of β_{SN} . Therefore, X_{SF} affects our analysis in the following two ways:

1. Observationally, the estimate of dust-to-gas ratio in a star-forming region is affected by the value of X_{SF} .
2. Theoretically, the value of X_{SF} affects the value of β_{SN} , thus changing the dust-to-gas ratio.

It is difficult to discuss the former quantitatively, because we need a high-resolution spatial map of H I distribution and a reasonable observational definition of a star-forming region on the map. However, we can discuss the latter because the effect of X_{SF} on β_{SN} is well determined from Eq. (6).

The observational dust mass, $M_{\text{d}}^{\text{IRAS}}$, is derived from the luminosity densities at wavelengths of 60 μm and 100 μm observed by *IRAS* using Eq. (4) of LF98. The *IRAS* bands are insensitive to the cold ($\lesssim 20$ K) dust that lies out of star-forming regions. Moreover, since the dust in a star-forming region suffers destruction by SN shocks, we should take into account an efficient destruction in the star-forming region. Therefore, in order to discuss the *IRAS* sample and the selective dust destruction in star-forming regions, we need to define β_{SN} as the *dust destruction efficiency* (Sect. 3) in star-forming regions as we have done in Eq. (6).

Finally, we hypothesise that the fraction of oxygen contained in dust grains is constant for all the sample BCDs. Following H99, we assume the Galactic composition of the grains:

$$\mathcal{D} = 2.2\mathcal{D}_{\text{O}}. \quad (9)$$

4.2. Conclusion from our model

In Figure 1, we show analytical results calculated according to Eqs. (3) and (9) with $i = \text{O}$ for various values of β_{SN} as solid line ($\beta_{\text{SN}} = 1$), dotted line ($\beta_{\text{SN}} = 5$; the case of LF98) and dashed line ($\beta_{\text{SN}} = 25$), respectively. The black and gray squares indicate the observational samples in Sage et al. (1992) and LF98, respectively. The number ratio of oxygen atoms to the hydrogen atoms is denoted as (O/H). We convert the mass fraction of oxygen, X_{O} , to (O/H) for the model prediction, assuming $\log X_{\text{O}} + 10.80 = 12 + \log(\text{O}/\text{H})$. In this figure, $f_{\text{in},\text{O}} = 0.1$ is assumed to concentrate on the variation in β_{SN} . LF98 have shown a large variety of $f_{\text{in},\text{O}}$ to explain the large scatter of the relation between \mathcal{D} and X_{O} in their Figure




Fig. 1. Relation between dust-to-gas ratio and metallicity. The solid, dotted, and dashed lines show the model results with the dust destruction efficiencies (Eq. 2) $\beta_{\text{SN}} = 1, 5,$ and $25,$ respectively. The black and gray squares show the data for the blue compact dwarf galaxies in Sage et al. (1992) and in Lisenfeld & Ferrara (1998), respectively.

7. Since Hirashita (1999a) has shown that $f_{\text{in,O}} \sim 0.1$ reproduces the observed trend between dust-to-gas ratio and metallicity, it is worth considering the effect of β_{SN} by setting $f_{\text{in,O}} = 0.1$.

From Figure 1, we see that the variance in dust-to-gas ratio is reproduced by an order-of-magnitude variation in β_{SN} (1–25 here). We note that LF98’s value ($\beta_{\text{SN}} = 5$) is within this range. Even if outflow does not efficiently occur in BCDs, we can explain the variance of the dust-to-gas ratio of BCDs with the various “destruction efficiency” of dust, β_{SN} , although we can never reject the importance of outflow in the framework of this paper. The important point is that we have demonstrated that dust destruction by SNe can play an important role in producing the variance of \mathcal{D} . The result also indicates that the difference in dust-to-gas ratio among the three lines becomes clear at an oxygen abundance, $12 + \log(\text{O}/\text{H}) \sim 8$. Thus, we conclude

1. that the dust destruction by SNe can vary the dust-to-gas ratio when the metallicity, $12 + \log(\text{O}/\text{H})$, reaches about 8 ($\sim 10\%$ of the solar metallicity), and
2. that the large variation of the dust-to-gas ratio among BCDs is explained by the variation of dust destruction efficiency, β_{SN} .

5. Discussion

5.1. Scenario

We propose a scenario for large variation of β_{SN} for each BCD along with intermittent SFH. The time variability of SFH on a short (< 1 Gyr) timescale has been suggested observationally by Searle & Sargent (1972) and

theoretically by Gerola et al. (1980). Nonlinear processes in the ISM may also cause an intermittent star formation (Ikeuchi 1988; Kamaya & Takeuchi 1997 and references therein). Thus, it is pertinent to consider the intermittent SFH of BCDs.

As shown in the following, an intermittent star formation history leads to the time variation of β_{SN} , because γ/ψ depends on time. We consider an intermittent SFH: a starburst whose SFR is ψ_{burst} and an inter-starburst epoch whose SFR is ψ_{inter} . We assume that $\psi_{\text{burst}} = 100\psi_{\text{inter}}$, for example. Such a two-orders-of-magnitude variation in SFR is proposed theoretically by Gerola et al. (1980) and Kamaya & Takeuchi (1997). While the starburst is going on, we expect that almost all the SNe are Type II ($\gamma_{\text{burst}} \sim \gamma_{\text{II}}$, where γ_{burst} is a typical SN rate in the bursting epoch). However, in the inter-burst epoch, Type Ia SNe can be dominated ($\gamma_{\text{inter}} \sim \gamma_{\text{Ia}}$, where γ_{inter} is a typical SN rate in the inter-burst). According to the model by Bradamante et al. (1998), a given stellar population releases energy in the form of Type II and Ia SNe with a ratio of 5:1 (Bradamante et al. 1998; see their Fig. 9. This value is essentially determined by the initial mass function (IMF), and they assumed the Salpeter IMF with a stellar mass range from 0.1 to $100 M_{\odot}$)². Since they assumed the same energy between Type Ia and II SNe, this means that the number ratio between Type Ia and II SNe is 5:1. Therefore, we expect that $\gamma_{\text{inter}} \sim \gamma_{\text{burst}}/5$. The intermittent star formation finally predicts that $\gamma_{\text{inter}}/\psi_{\text{inter}} \sim 20\gamma_{\text{burst}}/\psi_{\text{burst}}$. This means that an intermittent SFH can cause a 20-times variation in $\beta_{\text{SN}} \propto \gamma/\psi$ during a single star formation cycle.

Furthermore, Fig. 1 shows that the value of β_{SN} has little effect on the relation between dust-to-gas ratio and metallicity for $12 + \log(\text{O}/\text{H}) < 8$. Therefore, until the metallicity level becomes $12 + \log(\text{O}/\text{H}) \sim 8$, the relation between dust-to-gas ratio and metallicity evolves in the same way whatever the value of β_{SN} might be. On the contrary, the relation is largely affected by β_{SN} if $12 + \log(\text{O}/\text{H}) > 8$. Then, we study the response of the relation between dust-to-gas ratio and metallicity to the change of β_{SN} at $12 + \log(\text{O}/\text{H}) \sim 8$, as we are interested in the intermittent SFH.

First, we shall estimate a typical metallicity increment during a single star formation epoch of the intermittent SFH. The metallicity increase during an episode of star formation, ΔZ , can be estimated by $\Delta Z \sim yM_{*}/M_{\text{g}}$, where M_{*} is the mass of stars formed in the episode, and y is a chemical yield. If the IMF is similar to that of the Galaxy, $y \sim Z_{\odot}$ (i.e., $\Delta Z \rightarrow Z_{\odot}$ for $M_{*} \rightarrow M_{\text{g}}$). We estimate M_{*} by multiplying observed SFR with a duration of a star formation activity. Assuming that the SFR is $0.1 M_{\odot} \text{ yr}^{-1}$ and that the duration is 10^7 yr (Legrand et al. 2001), we obtain $M_{*} \sim 10^6 M_{\odot}$. With typical gas mass $M_{\text{g}} \sim 10^7 M_{\odot}$, we obtain $\Delta Z \sim 0.1Z_{\odot}$. This

² We assumed the Salpeter IMF with the range from 0.1 – $120 M_{\odot}$, but the difference in the higher mass cut has little influence on the SN rate.




Fig. 2. Relation between dust-to-gas ratio and metallicity. The three dotted lines show the model results same as Fig. 1. The squares are the data points same as Fig. 1. The two solid lines represents the result of the calculations which change β_{SN} (from 5 to 1 and from 5 to 25 for upper and lower lines, respectively).

corresponds, for example, to the metallicity increase from $12 + \log(\text{O}/\text{H}) = 8.0$ to 8.2 . The model by Bradamante et al. (1998) also indicates that one episode of star formation can result in such a metallicity increment.

As shown above, the effect of intermittence can be examined by changing β_{SN} . In order to examine the effect of time variation of β_{SN} , thus, we calculate the relation between dust-to-gas ratio and metallicity in the following two cases:

1. $\beta_{\text{SN}} = 5$ for $12 + \log(\text{O}/\text{H}) \leq 8$ and $\beta_{\text{SN}} = 25$ for $12 + \log(\text{O}/\text{H}) > 8$
2. $\beta_{\text{SN}} = 5$ for $12 + \log(\text{O}/\text{H}) \leq 8$ and $\beta_{\text{SN}} = 1$ for $12 + \log(\text{O}/\text{H}) > 8$

In Figure 2, we show the result of the two calculations (two solid lines). The lower and upper branches represent the cases 1 and 2, respectively. The three dotted lines show the results as in Fig. 1. We see that when the metallicity $12 + \log(\text{O}/\text{H})$ increases from 8.0 to 8.2 , the line rapidly converges to the two dotted lines, which represent the result for constant β_{SN} (1 and 25, respectively). This rapid convergence supports the idea that the dust-to-gas ratio varies widely in response to the time-evolution of β_{SN} in an episode of star formation at the metallicity level of $12 + \log(\text{O}/\text{H}) \sim 8$. Thus, it is possible to determine the variance of the dust-to-gas ratio among BCDs from the time variation of β_{SN} .

5.2. Age difference

If the age of BCDs varies, the present turn-off mass of stars is changed. As a result, the returned fraction of gas (\mathcal{R}), the metal yield (\mathcal{Y}_i), and the dust supply from stars

($f_{\text{in},i}$) are effectively different among BCDs. The effect of varying \mathcal{R} and \mathcal{Y}_i on the relation between dust-to-gas ratio and metallicity has been examined by H99.³ However, the resulting relation is less sensitive to the two parameters than to β_{SN} . The dependence of $f_{\text{in},i}$ on the turn-off mass can be important since it largely affects the dust amount in low-metallicity systems (LF98; Hirashita 1999a).

Thus, if the BCD sample proves to have a large age variation, we should reconsider the variation in dust-to-gas ratio with a time-dependent formulation. Indeed, we cannot reject the possibility that some BCDs are much younger than the cosmic age. A metal-poor BCD SBS 0335–052 may be younger than 5×10^6 yr (Vanzi et al. 2000). Recently, Hirashita, Hunt, & Ferrara (2002) have succeeded in explaining the dust amount of SBS 0335–052 with a time-dependent formulation applicable to young galaxies.

5.3. Future observations

The time variation of X_{SF} also leads to the time dependence of β_{SF} (Eq. 6), although X_{SF} was assumed as unity (Sect. 4.1). If the various X_{SF} for the BCD sample is interpreted to reflect the time evolution of X_{SF} in each BCD, we can suggest that the gas mass in a star-forming region should change temporally because of the mass exchange between the star-forming region and the envelope. Such a mass exchange during episodic star formation activity in BCDs is indeed suggested by e.g., Saitō et al. (2000). The temporal change of X_{SF} is also possible if the ISM in the star-forming region is consumed for star formation and locked in stellar remnants like white dwarfs, neutron stars, and black holes.

In order to constrain X_{SF} , we need to observe H I emission or FIR emission with angular resolution fine enough. The present typical angular resolution of $1'$ corresponds to 2.9 kpc in physical size if a galaxy lies at a typical distance of 10 Mpc. Future large space FIR telescopes such as *Herschel*⁴ (e.g., Pilbratt 2000) or *SPICA*⁵ (e.g., Nakagawa et al. 2000) will resolve the star-forming regions of the BCDs. For example, the Japanese future infrared satellite *SPICA* will have a diameter larger than 3.5 m. If the diffraction limit is achieved, the angular resolution becomes $6''$ at the wavelength of $100 \mu\text{m}$. This corresponds to 290 pc at the distance of 10 Mpc, and is comparable to or smaller than the half-light radius of a typical BCD (Marlowe et al. 1999).

6. Summary

In this paper, we have analysed the relation between dust-to-gas ratio and metallicity of BCDs by using our chemical

³ See Fig. 3 of the paper. Although H99 examined “different IMF”, this difference only affects the values of \mathcal{R} and \mathcal{Y}_i . Therefore, his result can be used to see the effect of varying \mathcal{R} and \mathcal{Y}_i .

⁴ <http://astro.estec.esa.nl/First/>

⁵ <http://www.ir.isas.ac.jp/SPICA/index.html>

evolution model. We have focused on the dust destruction process, because this process was not investigated in LF98. We have shown that the dust destruction significantly affects the dust-to-gas ratio when the metallicity is larger than $12 + \log(O/H) \sim 8$. The intermittent SFH can explain the large variety of dust-to-gas ratio among BCDs through the “dust destruction efficiency” of β_{SN} .

Acknowledgements. We thank the anonymous referee for invaluable comments and suggestions which improved this paper very much. We thank A. Ferrara and K. Yoshikawa for useful discussions on dwarf galaxies and excellent environments for our study. We are also grateful to S. Mineshige and J. Silk for continuous encouragement. H. H. was supported by the Research Fellowship of the Japan Society for the Promotion of Science for Young Scientists. We fully utilised the NASA’s Astrophysics Data System Abstract Service (ADS).

References

- Bradamante, F., Matteucci, F., & D’Ercole, A. 1998, *A&A*, 337, 338
- Calzetti, D., Bohlin, R. C., Kinney, A. L., Storchi-Bergmann, T., & Heckman, T. M. 1995, *ApJ*, 443, 136
- Crawford, I. A., & Barlow, M. J. 2000, *MNRAS*, 311, 370
- Dwek, E. 1998, *ApJ*, 501, 643
- Dwek, E., & Scalo, J. M. 1980, *ApJ*, 239, 193
- Dwek, E., et al. 1983, *ApJ*, 274, 168
- Eales, S. A., & Edmunds, M. G. 1996, *MNRAS*, 280, 1167
- Edmunds, M. G. 2001, *MNRAS*, 328, 223
- Gerola, H., Seiden, P. E., & Schulman, L. S. 1980, *ApJ*, 242, 517
- Grebel, E. K. 2001, *Dwarf Galaxies and their Environment*, ed. K. S. de Boer, R.-J. Dettmar, & U. Klein (Shaker Verlag, Bonn), 45
- Hirashita, H. 1999a, *ApJ*, 522, 220
- Hirashita, H. 1999b, *ApJ*, 510, L99 (H99)
- Hirashita, H. 2001, PhD thesis, Department of Astronomy, Kyoto University
- Hirashita, H., Hunt, L. K., & Ferrara, A. 2002, *MNRAS*, 330, L19
- Hoyle, F., & Wickramasinghe, N. C., 1963, *MNRAS*, 126, 401
- Ikeuchi, S. 1988, *Fundam. Cosmic Phys.*, 12, 255
- Issa, M. R., MacLaren, I., & Wolfendale, A. W. 1990, *A&A*, 236, 237
- Jones, A. P., Tielens, A. G. G. M., & Hollenbach, D. J. 1996, *ApJ*, 469, 740
- Kamaya, H., & Takeuchi, T. T. 1997, *PASJ*, 49, 271
- Kameswara Rao, M., & Lambert, D. L., 2000, *MNRAS*, 313, L33
- Kozasa, T., Hasegawa, H., & Nomoto, K. 1989, *ApJ*, 344, 325
- Legrand F., Tenorio-Tagle G., Silich S., Kunth D., & Cervino M. 2001, *ApJ*, 560, 630
- Lisenfeld, U., & Ferrara, A. 1998, *ApJ*, 496, 145 (LF98)
- Marlowe, A. T., Meurer, G. R., & Heckman, T. M. 1999, *ApJ*, 522, 183
- McKee, C. F. 1989, in *IAU Symp. 135, Interstellar Dust*, ed. L. J. Allamandola & A. G. G. M. Tielens (Dordrecht: Kluwer), 431
- Moseley, S. H., Dwek, E., Glaccum, W., Greham, J. R., & Loewenstein, R. F. 1989, *Nature*, 340, 697
- Nakagawa, T., et al. 2000, *The Institute of Space and Astronautical Science Report SP No. 14, Mid- and Far-Infrared Astronomy and Future Space Missions*, ed. T. Matsumoto & H. Shibai, (ISAS, Sagamihara), 189
- Oort, J. H., & van de Hulst, H. C., 1946, *Bull. Astr. Inst. Netherlands*, 10, 187
- Pilbratt, G. L. 2000, *The Institute of Space and Astronautical Science Report SP No. 14, Mid- and Far-Infrared Astronomy and Future Space Missions*, ed. T. Matsumoto & H. Shibai, (ISAS, Sagamihara), 211
- Sage, L. J., Salzer, J. J., Loose, H.-H., & Henkel, C. 1992, *A&A*, 265, 19
- Saitō, M., Kamaya, H., & Tomita, A. 2000, *ASP Conf. Ser. 218, Mapping the Hidden Universe: The Universe Behind the Milky Way*, *ASP Conf. Ser. 218*, (ASP, San Francisco), 305
- Searle, L., & Sargent, W. L. W. 1972, *ApJ*, 173, 25
- Takagi, T., Arimoto, N., & Vansevicius, V. 1999, *ApJ*, 523, 107
- Tajiri, Y. Y., & Kamaya, H. 2002, *A&A*, in press
- Tinsley, B. M. 1980, *Fundam. Cosmic Phys.*, 5, 287
- Todini, P., & Ferrara, A. 2001, *MNRAS*, 325, 726
- Searle, L., & Sargent, W. L. W. 1972, *ApJ*, 173, 25
- van Zee, L., Skillman, E. D., & Salzer, J. J. 1998, *AJ*, 116, 1186
- Vanzi, L., Hunt, L. K., Thuan, T. X., & Izotov, Y. I. 2000, *A&A*, 363, 493