## Tracking K-essence

Takeshi Chiba Department of Physics, Kyoto University, Kyoto 606-8502, Japan (November 2, 2018)

# Abstract

We derive a condition for converging a common evolutionary track for kessence (a scalar field dark energy with non-canonical kinetic terms). For the Lagrangian density  $V(\phi)W(X)$  with  $X = \dot{\phi}^2/2$ , we find tracker solutions with  $w_{\phi} < w_B$  exist if  $\Gamma \equiv V''V/(V')^2 > 3/2$ . Here  $w_{\phi}(w_B)$  is the equation-ofstate of the scalar field (background radiation/matter). Our condition may be useful for examining the existence of the attractor-like behavior in cosmology with k-essence (for example, rolling tachyon).

PACS numbers: 98.80.Cq ; 98.80.Es ; 95.35.+d

Typeset using REVTEX

#### I. INTRODUCTION

Scalar fields play important role in cosmology. For example, inflation in the early universe may be caused by inflatons, or an ultra-light scalar field with mass  $\sim 10^{-33}$  eV (or quintessence) may cause the universe to accelerate recently.

In the case of inflation models, the conditions for existence of inflationary solutions are conveniently described in terms of the slow-roll parameters without solving the equation of motion directly:  $(V'/V)^2/2\kappa^2 < 1$  $(V'/V)^2/2\kappa^2 < 1$  and  $V''/\kappa^2 V < 1$  with  $\kappa^2 = 8\pi G$  [1].

Similarly, some quintessence models admit so called tracker fields which have attractorlike solutions in the sense that a very wide range of initial conditions rapidly converge to a common cosmic evolutionary track[[2\]](#page-6-0). It may be particularly useful to express the condition for the existence of tracking solutions in terms of a simple condition of  $V(\phi)$  without having to solve the equation of motion directly. For quintessence, the condition for the existence of trackersolutions with  $w_{\phi} < w_B$  is  $\Gamma \equiv V''V/(\overline{(V')^2} > 1$  $\Gamma \equiv V''V/(\overline{(V')^2} > 1$  $\Gamma \equiv V''V/(\overline{(V')^2} > 1$  [2]. Here  $w_{\phi}(w_B)$  is the equation-ofstate of quintessence (background radiation or matter).

Usually the quintessence field (or inflaton) is modeled by a scalar field with a canonical kinetic term and a potential term. However, as shown in[[3](#page-6-0),[4\]](#page-6-0), a scalar field with solely kinetic terms can (even without potential terms), albeit they are non-canonical, mimic such a (canonical) quintessence/inflaton field.<sup>1</sup> However, the condition for the existence of tracker solutions for k-essence is not known to date.

In view of recent interest in scalar field cosmology with non-canonical kinetic terms [\[6](#page-6-0)–[8\]](#page-6-0), in this paper we shall express the condition for tracker solutions in terms of a function of the Lagrangian density.

#### II. TRACKING K-ESSENCE

### A. Basics of K-essence

The action of K-essence minimally coupled with gravity is

$$
S = \int d^4x \sqrt{-g} \left( \frac{1}{2\kappa^2} R + p(\phi, X) \right),\tag{1}
$$

where  $\kappa^2 = 8\pi G$  and  $X = -\nabla^{\mu}\phi\nabla_{\mu}\phi/2$ . The pressure of the scalar field  $\phi$ ,  $p_{\phi}$ , is given by  $p(\phi, X)$  itself and the energy density  $\rho_{\phi}$  is given by  $\rho_{\phi} = 2X \partial p/\partial X - p$  [\[3,4](#page-6-0)].

The field equations in flat Friedmann-Robertson-Walker spacetime are

$$
H^{2} := \left(\frac{\dot{a}}{a}\right)^{2} = \frac{\kappa^{2}}{3}(\rho_{B} + \rho_{\phi}) = \frac{\kappa^{2}}{3}\left(\rho_{B} + 2X\frac{\partial p}{\partial X} - p\right),\tag{2}
$$

$$
\frac{\ddot{a}}{a} = -\frac{\kappa^2}{6} (\rho_B + 3p_B + \rho_\phi + 3p_\phi) =: -\frac{\kappa^2}{6} ((1 + 3w_B)\rho_B + (1 + 3w_\phi)\rho_\phi),\tag{3}
$$

<sup>&</sup>lt;sup>1</sup>Wenamed such quintessence as "kinetic" quintessence in [[4](#page-6-0)]. However, later the name "k-essence" was coined by the authors of[[5](#page-6-0)] and got popularity. So in this paper we reluctantly use the term k-essence for such kinetically driven quintessence.

$$
\ddot{\phi} \left( \frac{\partial p}{\partial X} + \dot{\phi}^2 \frac{\partial^2 p}{\partial X^2} \right) + 3H \frac{\partial p}{\partial X} \dot{\phi} + \frac{\partial^2 p}{\partial X \partial \phi} \dot{\phi}^2 - \frac{\partial p}{\partial \phi} = 0, \tag{4}
$$

<span id="page-2-0"></span>where  $\rho_B$  and  $p_B$  are the energy density and the pressure of the background matter and/or radiation, respectively.

In this paper, as a first step toward more general case, we shall derive a tracker condition for a k-essence with the following factorized form of  $p(\phi, X)$ :

$$
p(\phi, X) = V(\phi)W(X).
$$
\n(5)

This form of the Lagrangian is suggested for that of tachyon by using the boundary string field theory [\[9](#page-6-0),[10](#page-6-0)]. Moreover, any Lagrangian containing only  $\dot{\phi}^2$  and  $\dot{\phi}^4$  terms can be recast in the factorized form after field redefinition.

The equation of motion of the scalar field is then written as

$$
\ddot{\phi}(W_X + 2XW_{XX}) + 3HW_X \dot{\phi} + (2XW_X - W)\frac{V'}{V} = 0,
$$
\n(6)

where  $W_X = dW/dX$ .

## B. Tracker Equation

We can express the equation of motion of  $\phi$  in alternative form which may be useful for the following analysis:

$$
\frac{V'}{V^{3/2}} = \mp \frac{\kappa}{2} \sqrt{\frac{(1 + w_{\phi})W_X}{3\Omega_{\phi}}} (6 + Ay'), \qquad (7)
$$

$$
A = \frac{(XW_X - W)(2XW_{XX} + W_X)}{XW_X^2 - WW_X - XWW_{XX}} = \frac{1 - w_{\phi}}{c_s^2 - w_{\phi}},
$$
\n(8)

where  $y = (1 + w_{\phi})/(1 - w_{\phi})$  and  $y' = d \ln y/d \ln a$ , and minus(plus) sign corresponds to  $\dot{\phi} > 0 \, (< 0),$  respectively.  $c_s^2$  $s<sub>s</sub><sup>2</sup>$  is the speed of sound of k-essence defined by [\[11\]](#page-6-0)

$$
c_s^2 = \frac{p_X}{p_X + 2Xp_{XX}}.\tag{9}
$$

Note that for quintessence with a canonical kinetic term,  $c_s^2 = 1$ . For a tracker solution  $(w_{\phi} \simeq \text{const.})$ , we obtain a relation:

$$
\frac{1}{\sqrt{\Omega_{\phi}}} = \pm \frac{1}{\kappa \sqrt{3(1 + w_{\phi})W_X}} \frac{V'}{V^{3/2}},\tag{10}
$$

which might be called the k-essential counterpart of the tracker condition[[2\]](#page-6-0).

Similar to [\[2](#page-6-0)], we define a dimensionless function  $\Gamma$  by  $\Gamma = VV''/V'^2$ . After taking the time derivative of Eq.(7), we obtain

<span id="page-3-0"></span>
$$
\Gamma - \frac{3}{2} = -\frac{1}{(1+w_{\phi})(6+Ay')} \left[ 3(w_{\phi}-w_B)(1-\Omega_{\phi}) + \frac{(1-w_{\phi})^2}{2(c_s^2-w_{\phi})} y' + \frac{2(1-w_{\phi})(c_s^2-w_{\phi})y'' + 2(\dot{w}_{\phi}(1-c_s^2) - (c_s^2)^2(1-w_{\phi}))y'/H}{(6+Ay')(c_s^2-w_{\phi})^2} \right],
$$
\n(11)

where  $y'' = d^2 \ln y/d \ln a^2$ . Eq.(11) might be called the k-essential counterpart of the tracker equation. Therefore for the tracker solution (assuming  $\Gamma \simeq$  const. and  $\Omega_{\phi} \ll 1^2$ ) we can write  $w_{\phi}$  in terms of Γ:

$$
w_{\phi} = \frac{w_B - 2(\Gamma - 3/2)}{2(\Gamma - 3/2) + 1} \simeq \text{const.}
$$
 (12)

## C. Tracking Condition

Convergence toward the tracker solution. We now examine the stability of the tracker solution. Consider a solution which is perturbed from the tracker solution with  $w_0$  (Eq.(12)) by an amount  $\delta$ , then the tracker equation Eq.(11) is expanded to lowest order in  $\delta$  to obtain

$$
2\delta'' + 3(1 + w_B - 2w_0)\delta' + 9(c_s^2 - w_0)(1 + w_B)\delta = 0,
$$
\n(13)

where the prime means  $d/d \ln a$ . The solution of this equation is

$$
\delta \propto a^{\gamma} \tag{14}
$$

$$
\gamma = -\frac{3}{4}(1+w_B - 2w_0) \pm \frac{3}{4}\sqrt{(1+w_B - 2w_0)^2 - 8(c_s^2 - w_0)(1+w_B)}.
$$
\n(15)

In order for the real part of  $\gamma$  to be negative so that  $\delta$  decays exponentially and the solution approaches the tracker solution, it is required that

$$
w_0 < \frac{1+w_B}{2} \qquad \text{and} \qquad w_0 < c_s^2,\tag{16}
$$

where  $c_s^2 \geq 0$  is assumed for stability against perturbation. Note that for the canonical quintessence  $(c_s^2 = 1)$  the second requirement in Eq.(16) is automatically satisfied.

From Eq.(12), the above requirements are written in terms of  $\Gamma$ :

$$
\Gamma > \frac{3}{2} - \frac{c_s^2 - w_B}{2(c_s^2 + 1)} \quad \text{and} \quad \Gamma > \frac{3}{2} - \frac{1 - w_B}{6 + 2w_B} \tag{17}
$$

$$
\text{or} \quad \Gamma < 1. \tag{18}
$$

<sup>2</sup>This assumption is implicit in [\[2](#page-6-0)].

Tracking behavior with  $w_{\phi} < w_B$ . In this case,  $\Omega_{\phi}$  increases with increase of time. Then according to Eq.([10\)](#page-2-0),  $|V'/V^{3/2}\sqrt{W_X}|$  decreases for a tracker solution. On the other hand, using the equation of motion, we find that

$$
\left(\frac{V'}{V^{3/2}\sqrt{W_X}}\right) = \frac{V'\dot{\phi}}{V^{5/2}\sqrt{W_X}}\left(\Gamma - \frac{3}{2}\right). \tag{19}
$$

Hence,  $|V'/V^{3/2}\sqrt{W_X}|$  decreases if  $\Gamma > 3/2$ . Therefore, tracking behavior with  $w_{\phi} < w_B$ occurs if  $\Gamma > 3/2$  and nearly constant. From the stability analysis, this tracker solution is stable if  $\Gamma > \max(3/2, 3/2 - (c_s^2 - w_B)/2(c_s^2 + 1))$ . In particular for  $w_B = 0$ , the condition is simply  $\Gamma > 3/2$ .

Tracking behavior with  $w_{\phi} > w_B$ . This is possible for  $1 < \Gamma < 3/2$  and nearly constant. However, from the stability analysis, we find that  $w_{\phi} < \min((1 + w_B)/2, c_s^2)$  is additionally required, which demands  $\Gamma - 3/2 > \max(-(c_s^2 - w_B)/2(c_s^2 + 1), -(1 - w_B)/(6 + 2w_B)).$ 

Tracking behavior with  $w_{\phi} < -1$ . This is possible for  $\Gamma < 1$  and nearly constant. From the stability analysis, we find that this solution is stable.

### D. Examples

As an example, we consider the universe consisting of matter/radiation and a k-essential scalarfield with a power-law model,  $V \propto \phi^{-\alpha}$ , studied in [[4](#page-6-0)]. Since  $\Gamma = (\alpha + 1)/\alpha$ , if  $0 < \alpha < 2$ , then tracking behavior with  $w_{\phi} < w_B$  occurs, while if  $\alpha < 0$ , tracking behavior with  $w_{\phi} < -1$  occurs. From Eq.([12](#page-3-0)), in this case we have for the tracker solution

$$
w_{\phi} = (1 + w_B)\frac{\alpha}{2} - 1,\tag{20}
$$

which indeed coincides with the equation-of-state of the attractor solution studied in [\[4\]](#page-6-0) and indeed the solution is stable [\[4](#page-6-0)].

Another example is the universe consisting of matter/radiation and the rolling tachyon with  $W(X) = -\sqrt{1-2X}$ . The exact classical potential of it has been computed in [\[9](#page-6-0)] and is given by

$$
V(T) = V_0 \left( 1 + T/T_0 \right) \exp(-T/T_0), \tag{21}
$$

where T is the tachyon field and  $V_0$  is the tension of some unstable bosonic D-brane and  $T_0$ is a constant of the order of the string scale. In this case we have

$$
\Gamma = 1 - (T/T_0)^{-2}
$$
 (22)

and Γ becomes nearly constant if  $T/T_0 \gg 1$ . In the limit of large  $T/T_0$ , we have  $\Gamma = 1$  and obtain  $w_{\phi} = -[w_B(1 - \Omega_{\phi}) + 1]/\Omega_{\phi}$  if we do not neglect  $\Omega_{\phi}$  in Eq.[\(12](#page-3-0)). Since the rolling tachyon respects the weak energy condition,  $w_{\phi} \geq -1$ , this implies  $\Omega_{\phi} \geq 1$  during the radiation dominated epoch (the same result holds for the matter dominated epoch), which would be incompatible with the success of the Big-Bang Nucleosynthesis. Thus the above potential does not admit viable tracker solutions, which implies the need of fine-tuning to account for  $\Omega_{\phi} \sim 1$  today in agreement with the recent analysis [\[8](#page-6-0)].

#### III. SUMMARY

We have derived a (sufficient) condition for the existence of tracker solutions for the system of matter/radiation and a scalar field with Lagrangian density of the form Eq.([5](#page-2-0)). Our results are summarized as follows:

- Tracking behavior with  $w_{\phi} < w_B$  occurs if  $\Gamma > \max(3/2, 3/2 + (w_B c_s^2))$  $\binom{2}{s}/2(c_s^2+1)$ and nearly constant. In particular for  $w_B = 0$ , the condition is simply  $\Gamma > 3/2$  and nearly constant.
- Tracking behabior with  $w_{\phi} > w_B$  occurs if  $3/2 > \Gamma > 3/2 + \max(-(c_s^2 w_B)/2(c_s^2 +$ 1),  $-(1 - w_B)/(6 + 2w_B)$  and nearly constant.
- Tracking behavior with  $w_{\phi} < -1$  occurs if  $\Gamma < 1$  and nearly constant.

Interestingly, thanks to the factorizable ansatz  $Eq.(5)$  $Eq.(5)$  $Eq.(5)$ , the conditions are similar to those of canonical quintessence.

## ACKNOWLEDGMENTS

We would like to thank Masahide Yamaguchi for useful comments. This work was supported in part by a Grant-in-Aid for Scientific Research (No.13740154) from the Japan Society for the Promotion of Science and by a Grant-in-Aid for Scientific Research on Priority Areas (No.14047212) from the Ministry of Education, Science, Sports and Culture, Japan.

## REFERENCES

- <span id="page-6-0"></span>[1] J.E. Lidsey, A.R. Liddle, E.W. Kolb, E.J. Copeland, T. Barreiro, M. Abney, Rev. Mod. Phys. 69, 373 (1997).
- [2] P.J. Steinhardt, L. Wang and I. Zlatev, Phys. Rev. D 59, 123504 (1999).
- [3] C. Armendáriz-Picón, T. Damour and V. Mukhanov, Phys. Lett.  $\mathbf{B458}$ , 209 (1999).
- [4] T. Chiba, T. Okabe and M. Yamaguchi, Phys. Rev. D 62, 023511 (2000).
- [5] C. Armendariz-Picon, V. Mukhanov, and P.J. Steinhardt, Phys. Rev. Lett. 85, 4438 (2000).
- [6] A. Sen, [hep-th/0203211;](http://arxiv.org/abs/hep-th/0203211) [hep-th/0203265.](http://arxiv.org/abs/hep-th/0203265)
- [7] G.W. Gibbons, [hep-th/0204008](http://arxiv.org/abs/hep-th/0204008); M. Fairbain and M.H.G. Tytgat, [hep-th/0204070;](http://arxiv.org/abs/hep-th/0204070) S. Mukohyama, [hep-th/0204084;](http://arxiv.org/abs/hep-th/0204084) T. Padmanabhan, [hep-th/0204150](http://arxiv.org/abs/hep-th/0204150).
- [8] A. Frolov, L. Kofman and A. Starobinsky, [hep-th/0204187](http://arxiv.org/abs/hep-th/0204187); X. Li, J. Hao and D. Liu, [hep-th/0204252](http://arxiv.org/abs/hep-th/0204252); G. Shiu and I Wasserman, [hep-th/0205003](http://arxiv.org/abs/hep-th/0205003); L. Kofman and A. Linde, [hep-th/0205121](http://arxiv.org/abs/hep-th/0205121).
- [9] D. Kutasov, M. Marino and G. Moore, JHEP 0010 (2000) 045.
- [10] S. Sugimoto and S. Terashima, [hep-th/0205085.](http://arxiv.org/abs/hep-th/0205085)
- [11] J. Garriga and V.F. Mukhanov, Phys. Lett. B458, 219 (1999).