

On the Spectrum of Magnetohydrodynamic Turbulence

Stanislav Boldyrev

Department of Astronomy and Astrophysics, University of Chicago, 5640 South Ellis Avenue, Chicago, IL 60637

`boldyrev@uchicago.edu`

ABSTRACT

We propose a phenomenological model for incompressible magnetohydrodynamic turbulence. We argue that nonlinear-wave interaction weakens as the energy cascade proceeds to small scales, however, the anisotropy of fluctuations along the large-scale magnetic field increases, which makes turbulence strong at all scales. To explain the weakening of the interaction, we propose that small-scale fluctuations of the velocity and magnetic fields become increasingly dynamically aligned as their scale decreases, so that turbulent “eddies” become locally anisotropic in the plane perpendicular to the large-scale magnetic field. In the limit of weak anisotropy, that is, weak large-scale magnetic field, our model reproduces the Goldreich-Sridhar spectrum, while the limit of strong anisotropy, that is, strong large-scale magnetic field, corresponds to the Iroshnikov-Kraichnan scaling of the spectrum. This is in good agreement with recent numerical results.

Subject headings: MHD–turbulence

1. Introduction.

Magnetohydrodynamic (MHD) turbulence is a state of a randomly stirred conducting fluid in the limit of very small fluid viscosity and resistivity; it plays an essential role in a variety of astrophysical systems, from planets and stars, to interstellar and intergalactic media (see, e.g., Biskamp 2003). Despite more than 35 years of analytical, numerical, and observational investigations, the spectrum of MHD turbulence remains a subject of controversy. The standard results and recent developments in the theory of MHD turbulence can be found in many excellent texts (see, e.g., Iroshnikov 1963; Kraichnan 1965; Goldreich &

Sridhar 1995, 1997; Ng & Bhattacharjee 1996; Cho & Vishniac 2000; Biskamp & Müller 2000; Maron & Goldreich 2001; Cho, Lazarian & Vishniac 2002; Galtier, et al. 2000, 2002; Müller, Biskamp & Grappin 2003; Biskamp 2003; Kulsrud 2005). In the present Letter we first analyze the approaches of Iroshnikov (1963) and Kraichnan (1965) and Goldreich & Sridhar (1995) and point out the discrepancies of these theories with numerical results. Then, we propose a new model for the turbulent MHD cascade, which is free of these discrepancies, and which is in good agreement with recent high-resolution numerical findings of Maron & Goldreich (2001), Müller, Biskamp & Grappin (2003), and Padoan, et al. (2004). Our results are summarized in Conclusions.

The Iroshnikov-Kraichnan energy cascade. The MHD equations describing evolution of the fluid velocity field, $\mathbf{v}(\mathbf{r}, t)$, and fluctuations of the magnetic field, $\mathbf{b}(\mathbf{r}, t)$, have an especially simple form when expressed in the Elsässer variables, $\mathbf{z} = \mathbf{v} - \mathbf{b}$, and $\mathbf{w} = \mathbf{v} + \mathbf{b}$:

$$\partial_t \mathbf{z} + (\mathbf{V}_A \cdot \nabla) \mathbf{z} + (\mathbf{w} \cdot \nabla) \mathbf{z} = -\nabla P, \quad (1)$$

$$\partial_t \mathbf{w} - (\mathbf{V}_A \cdot \nabla) \mathbf{w} + (\mathbf{z} \cdot \nabla) \mathbf{w} = -\nabla P, \quad (2)$$

where the pressure P is determined from the incompressibility condition, $\nabla \cdot \mathbf{z} = 0$ or $\nabla \cdot \mathbf{w} = 0$. In equations (1) and (2), $\mathbf{V}_A = \mathbf{B}_0 / \sqrt{4\pi\rho}$ is the Alfvén velocity, ρ is the fluid density, \mathbf{B}_0 is the large-scale external magnetic field, and we have neglected small viscosity and resistivity. Note that for $\mathbf{w} = 0$, any function $\mathbf{z} = \mathbf{f}(\mathbf{r} - \mathbf{V}_A t)$ is a solution of the system (eqs. 1,2); analogously, for $\mathbf{z} = 0$, the solution is $\mathbf{w} = \mathbf{g}(\mathbf{r} + \mathbf{V}_A t)$, where the function \mathbf{g} is arbitrary.

Iroshnikov and Kraichnan used this fact to propose that the interacting Alfvén-wave packets (or “eddies”) are those propagating in *opposite* directions along the large-scale magnetic-field lines (Iroshnikov 1963; Kraichnan 1965). From that, they deduced that the energy transfer time from wave packets of size λ to smaller ones is increased compared with the simple dimensional estimate $\tau(\lambda) \sim \lambda / \delta v_\lambda$. (We denote by δv_λ and δb_λ the velocity and magnetic-field fluctuations in the “eddy.” In the Alfvén wave, $\delta v_\lambda \sim \delta b_\lambda$.) Indeed, consider two wave packets of size λ propagating in the opposite directions along a magnetic-field line with the Alfvén velocities. Assuming that the “eddies” are decorrelated at the field-parallel scale λ , one can estimate from equations (1) and (2) that during one collision the distortion of the “eddy” is $\Delta \delta v_\lambda \sim (\delta v_\lambda^2 / \lambda)(\lambda / V_A)$. The distortions add up randomly, therefore, the “eddy” will be distorted relatively strongly after $N \sim (\delta v_\lambda / \Delta \delta v_\lambda)^2 \sim (V_A / \delta v_\lambda)^2$ collisions. The energy transfer time is, therefore,

$$\tau_{IK}(\lambda) \sim N \lambda / V_A \sim \lambda / \delta v_\lambda (V_A / \delta v_\lambda). \quad (3)$$

It is important that in the Iroshnikov-Kraichnan interpretation, an “eddy” has to ex-

perience *many* uncorrelated interactions with oppositely moving “eddies” before its energy is transferred to a smaller scale. Moreover, turbulence becomes progressively *weaker* as the energy cascade proceeds toward smaller scales. The requirement of constant energy flux over scales $\delta v_\lambda^2/\tau_{IK}(\lambda) = \text{const}$ immediately leads to the scaling of fluid fluctuations $\delta v_\lambda \propto \lambda^{1/4}$, which resulted in the energy spectrum (the Iroshnikov-Kraichnan spectrum),

$$E_{IK}(k) = |\delta v_k|^2 k^2 \propto k^{-3/2}. \quad (4)$$

Iroshnikov and Kraichnan did not consider anisotropy of the spectrum, so $E_{IK}(k)$ was assumed to be three-dimensional and isotropic.

Anisotropy: The Goldreich-Sridhar energy cascade. Over the years, isotropy of the MHD spectrum in a strong external magnetic field seemed to contradict analytical and numerical findings, (see, e. g., Biskamp 2003). A theory of anisotropic MHD turbulence was proposed by Goldreich & Sridhar (1995). They argued that “eddies” are strongly anisotropic; they are elongated along the large-scale magnetic field lines. As a consequence, the time of relatively strong “eddy” distortion (the energy transfer time) is on the order of a crossing time required for two oppositely moving “eddies” to pass each other.

Suppose that the “eddy” has a transverse (to the large-scale field) size λ . Then, its field-parallel size l can be found from the so-called critical-balance condition, proposed by Goldreich & Sridhar (1995). This condition has two explanations that are equivalent in the Goldreich-Sridhar picture. First, the critical balance can be understood as a formal balance of the second and third terms in equations (1) and (2): $V_A/l \sim \delta b_\lambda/\lambda$. Second, it follows from the geometric distortion of magnetic-field lines in the turbulent “eddy.” Indeed, the “eddy” displaces the lines in their perpendicular direction by a distance $\xi \sim \delta b_\lambda l/V_A$, and this displacement equals the perpendicular “eddy” size λ .

The critical balance condition is the same for all scales, so, contrary to the Iroshnikov-Kraichnan picture, the turbulence strength does not change with the scale. The energy transfer time predicted in the Goldreich-Sridhar theory is

$$\tau_{GS}(\lambda) \sim l/V_A \sim \lambda/\delta v_\lambda. \quad (5)$$

Assuming that the energy cascade is independent of the scale, $\delta v_\lambda^2/\tau_{GS}(\lambda) = \text{const}$, one obtains the scaling of fluid fluctuations, $\delta v_\lambda \propto \lambda^{1/3}$. The corresponding energy spectrum is

$$E_{GS}(k_\perp) = |\delta v_{k_\perp}|^2 k_\perp \propto k_\perp^{-5/3}. \quad (6)$$

This spectrum coincides with the Kolmogorov spectrum of incompressible non-magnetized fluid turbulence, as it should since the energy transfer time coincides with the “eddy” turnover time, $\tau(\lambda) \sim \lambda/\delta v_\lambda$, in both approaches. The anisotropy of fluctuations is described by

the condition that follows from the critical balance, $l \propto \lambda^{2/3}$. One can therefore write that the fluctuations scale with the field-parallel size of the “eddy” as $\delta v_l \propto l^{1/2}$.

The Goldreich-Sridhar picture, however, does not fully agree with numerical simulations. As was recently discovered by Müller, Biskamp & Grappin (2003), the anisotropic spectrum depends on the strength of the external magnetic field. Denote $\gamma = B_0^2 / \langle \rho \delta v_L^2 \rangle$, where δv_L is the velocity field at the outer scale of turbulence, L . It was found that the *field-perpendicular* scaling of fluctuations changed from the Goldreich-Sridhar form to the Iroshnikov-Kraichnan form as the field increased from $\gamma \ll 1$ to $\gamma \gg 1$. A similar result for $\gamma \gg 1$ was obtained earlier by Maron & Goldreich (2001). These intriguing numerical findings motivated our interest in the problem. In the next section, we propose a phenomenological model of MHD turbulence, which agrees well with available numerical results, for any strength of the external field. In the limiting case of a weak external field, our model reproduces the anisotropic spectrum of Goldreich & Sridhar. In the other limiting case of a very strong external field, anisotropy of the spectrum is *stronger*; however, the *field-perpendicular* spectrum formally coincides with the spectrum predicted in the Iroshnikov-Kraichnan model.

2. A model for MHD turbulence.

To begin with, we make a certain assumption about reduction of the nonlinear interaction, which is not present in either the Iroshnikov-Kraichnan or Goldreich-Sridhar picture. We postpone the justification of this assumption until the end of this section, when we obtain the corresponding solution for the turbulent spectra and compare it with numerical simulations.

Reduction of nonlinear interaction. Let us assume that the nonlinear interaction of the counter-propagating fluctuations is reduced by a factor $(\delta v_\lambda / V_A)^\alpha$, where α is some undetermined exponent, $0 \leq \alpha \leq 1$. In other words, we assume that the nonlinearity in equations (1) and (2) is “depleted,” so that the interaction terms are of order

$$(\mathbf{w} \cdot \nabla) \mathbf{z} \sim (\mathbf{z} \cdot \nabla) \mathbf{w} \sim (\delta v_\lambda^2 / \lambda) (\delta v_\lambda / V_A)^\alpha. \quad (7)$$

Thus, the fluid fluctuations at the transverse distance λ become decorrelated on the time-scale

$$\tau_N(\lambda) \sim (\lambda / \delta v_\lambda) (V_A / \delta v_\lambda)^\alpha. \quad (8)$$

Their decorrelation length along the magnetic field line can be found from the causality principle. For $\delta v_\lambda < V_A$, the perturbation cannot propagate along the field line faster than

V_A ; therefore, the correlation length along the field line is on the order of $l \sim V_A \tau_N(\lambda)$. This condition is analogous to the critical balance condition of the previous section in that it satisfies the same balance between the linear and nonlinear interaction terms in equations (1) and (2). However, its geometric meaning is different, and will be discussed below.

We see that the interaction strength (eq. 7) decreases for smaller scales; however, the field-parallel “eddy” size l is adjusted in such a way that the energy transfer to the smaller scales always takes one “eddy” crossing time. Note that contrary to the Iroshnikov-Kraichnan formalism, and similar to the Goldreich-Sridhar approach, in our picture turbulence is strong and fluctuations are highly anisotropic. Now, we require that the energy cascade be constant over scales $\delta v_\lambda^2 / \tau_N(\lambda) = \text{const}$. We derive $\delta v_\lambda \propto \lambda^{1/(3+\alpha)}$, and the anisotropy condition reads $l \propto \lambda^{2/(3+\alpha)}$. The corresponding energy spectrum is given by

$$E(k_\perp) = |\delta v_{k_\perp}|^2 k_\perp \propto k_\perp^{-(5+\alpha)/(3+\alpha)}. \quad (9)$$

One can formally define the corresponding longitudinal spectrum of fluctuations, from the condition $E(k_\perp) dk_\perp = E(k_\parallel) dk_\parallel$, with $k_\perp \propto k_\parallel^{(3+\alpha)/2}$. The answer is $E(k_\parallel) \propto k_\parallel^{-2}$; note that it is independent of α . Therefore, the scaling of fluid fluctuations with respect to the field-parallel distance l is always $\delta v_l \propto l^{1/2}$.

Comparison with numerical simulations. Obviously, our result with $\alpha = 0$ corresponds to the Goldreich-Sridhar scaling, while $\alpha = 1$ produces the Iroshnikov-Kraichnan scaling. Simulations by Müller, Biskamp & Grappin (2003) for a range of large-scale external magnetic fields have shown that the scaling of the second-order structure function with respect to the field-perpendicular scale λ changes from the Goldreich-Sridhar value in the case of a weak external field, $\gamma \ll 1$, to the Iroshnikov-Kraichnan value in the case of a strong field, $\gamma \gg 1$. At the same time, the scaling of the second-order structure function with respect to the field-parallel distance l does not change much and stays close to $\langle (\delta v_l)^2 \rangle \propto l^{0.9}$. This result is consistent with our prediction, $(\delta v_l)^2 \propto l$. The same scalings for the case of a strong external magnetic field, were earlier observed in simulations by Maron & Goldreich (2001). In these simulations, it was also found that the scaling of the energy-cascade time was $\tau_E \propto \lambda^{1/2}$. This contradicts the Goldreich-Sridhar picture (see the discussion in § 6.1.3 of Maron & Goldreich 2001), but coincides with our formula $\tau_N(\lambda) \propto \lambda^{(1+\alpha)/(3+\alpha)}$, for $\alpha = 1$.

It is important to note that when the perpendicular Iroshnikov-Kraichnan scaling is formally reproduced in our model, the turbulent fluctuations are strong and anisotropic, with $l \propto \lambda^{1/2}$. Moreover, the turbulence possessing the Iroshnikov-Kraichnan spectrum is more anisotropic than the turbulence in the Goldreich-Sridhar model, which is not surprising, since, as just discussed, the Iroshnikov-Kraichnan spectrum corresponds to a much stronger external magnetic field.

Scale-dependent dynamic alignment. We next explain a possible physical origin for the proposed reduction of the non-linear interaction (7). A hint toward the explanation can be obtained from geometric considerations. The displacement of magnetic field lines in the direction perpendicular to the large-scale magnetic field, produced by the wave packet with $l \propto \lambda^{2/(3+\alpha)}$ and $\delta b_\lambda \propto \lambda^{1/(3+\alpha)}$, is given by $\xi \propto \delta b_\lambda l \propto \lambda^{3/(3+\alpha)}$. We thus obtain that the transverse displacement of magnetic field lines in the shear-wave packet is on the order of ξ at distances $\lambda \ll \xi$. Therefore, this packet should be highly anisotropic in the plane *perpendicular* to the large-scale magnetic field. In this plane, it is elongated in the direction of the field fluctuations. This may be consistent with the numerically supported picture that dissipative structures in MHD turbulence are micro-current sheets (Biskamp 1993, 2003; Biskamp & Müller 2000; Maron & Goldreich 2001). For comparison, the Goldreich-Sridhar scaling, $\alpha = 0$, would correspond to filament-like dissipative structures. In Figure 1 we sketch the shapes of the “eddies” in the Goldreich-Sridhar picture and in our picture. In our model, the anisotropy of the “eddies” is weakest for $\alpha = 0$, and therefore it is natural to identify this limit with zero large-scale magnetic field. In the other limit, $\alpha = 1$, the anisotropy of the “eddies” is strongest, therefore, this case should be identified with a strong magnetic field, $\gamma \gg 1$.¹

Let us now explain how the envisioned anisotropy of fluctuations in the field-perpendicular plane reduces the strength of the nonlinear-wave interaction. For this purpose, consider the third terms in the MHD equations (1,2), say, $(\mathbf{w} \cdot \nabla) \mathbf{z}$. Since both \mathbf{w} and \mathbf{z} are divergence-free and perpendicular to the large-scale magnetic field, this term is proportional to the angle θ_λ between the directions of \mathbf{w}_λ and \mathbf{z}_λ (if this angle is small). But we just established that this angle is $\theta_\lambda \sim \lambda/\xi \propto \lambda^{\alpha/(3+\alpha)}$. Quite remarkably, we have reproduced the reduction factor $(\delta v_\lambda/V_A)^\alpha$ in equation (7). This demonstrates that our initial assumption is self-consistent. Moreover, a detailed analysis of equations (1) and (2) suggests that the alignment $\theta_\lambda \propto \delta v_\lambda^\alpha$ is indeed consistent with the MHD dynamics, when θ_λ is small. In this limit we can obtain the following approximate evolution equations, $\partial_t \delta v_\lambda \sim \delta v_\lambda^2 \theta_\lambda / \lambda$ and $\partial_t \theta_\lambda \sim \delta v_\lambda \theta_\lambda^2 / \lambda$. The alignment exponent α would be determined by numeric coefficients in these equations, which cannot be obtained from the dimensional analysis.

As an important analogy, we note that *decaying* MHD turbulence approaches the so-called Alfvénic state, where either \mathbf{w} or \mathbf{z} is zero depending on the initial conditions (see e.g., Grappin, et al 1982; Grappin, Pouquet, & L  orat 1983). Based on our analysis, we propose that driven turbulence behaves in a similar manner, although \mathbf{b} does not become

¹As is shown in (Grappin, et al 1982) with the aid of the EDQNM closure, strong large-scale magnetic field leads to the Alfv  nic decorrelation time in aligned turbulence. In our model, this means that α cannot exceed 1.

exactly equal to $\pm\mathbf{v}$. Rather, magnetic fluctuations $\delta\mathbf{b}_\lambda$ tend to align their direction, but not their magnitude, with that of the velocity fluctuations, $\pm\delta\mathbf{v}_\lambda$. As a result, the fluctuations $\delta\mathbf{w}_\lambda$ and $\delta\mathbf{z}_\lambda$ are of the same order, and the directions of $\delta\mathbf{w}_\lambda$ and $\pm\delta\mathbf{z}_\lambda$ are aligned within the angle $\theta_\lambda \sim \lambda/\xi \propto \lambda^{\alpha/(3+\alpha)}$. The degree of the alignment increases progressively as the scale decreases. Such scale-dependent dynamic alignment (and the associated depletion of nonlinearity) can in principle be checked numerically, although the numerical analysis may be complicated by the rather slow dependence of the alignment angle θ_λ on the scale. There is, however, a numerical indication that MHD turbulence indeed has a tendency to create correlated regions of polarized fluctuations (Maron & Goldreich 2001).

On the non-universality of the turbulent spectrum. Our analysis points to an interesting possibility that MHD turbulence is non-universal in that it depends on the large-scale magnetic field. We may further speculate that, in principle, other large-scale conditions may affect the scaling properties of turbulence. For example, the dynamic alignment may be sensitive to the level of cross-helicity fluctuations; an analogous result is known for the case of *decaying* turbulence, (see, e.g., Grappin, et al 1982; Grappin, Pouquet, & L  orat 1983). An alternative possibility is that the spectrum is *universal* and has the Iroshnikov-Kraichnan scaling but, for $\gamma \ll 1$, the dynamic alignment with $\alpha = 1$ is established rather slowly as the scale decreases, and the resolution of numerical simulations is not large enough to reach the universal regime.

3. Conclusions.

To conclude, we summarize our main results:

1. Our consideration is motivated by the recent numerical observations (Maron & Goldreich 2001; M  ller, Biskamp & Grappin 2003) that incompressible MHD turbulence is not completely described by either the Iroshnikov-Kraichnan or the Goldreich-Sridhar model. The scaling of velocity fluctuations was found in these papers to depend on the strength of the large-scale external magnetic field: the Iroshnikov-Kraichnan scaling appeared in the limit of strong external magnetic field, while the Goldreich-Sridhar scaling appeared in the limit of weak field.

2. To explain these numerical findings, we propose that turbulent fluctuations become increasingly dynamically aligned as the energy cascade proceeds to smaller scales. Velocity fluctuations $\delta\mathbf{v}_\lambda$ tend to align their direction with that of magnetic field fluctuations, $\pm\delta\mathbf{b}_\lambda$, and the smaller the scale λ , the stronger the alignment. The dynamic alignment leads to reduction of the nonlinear-wave interaction (so-called depletion of nonlinearity).

3. As a result of point 2, fluctuating “eddies” are three-dimensionally anisotropic. The “eddies,” whose smallest scale is λ , have the scale $\xi \propto \lambda^{3/(3+\alpha)}$ in the direction of the shear (the direction of the magnetic field line distortion), and the scale $l \propto \lambda^{2/(3+\alpha)}$ in the direction of the large-scale magnetic field, as is sketched in Figure 1. The scaling and anisotropy of fluctuations are described by a single parameter α , which depends on the strength of the external magnetic field, and which is determined in our work from comparison with numerical simulations by Maron & Goldreich (2001) and Müller, Biskamp & Grappin (2003).

4. The energy distribution is given by

$$E(k_{\perp}) \propto k_{\perp}^{-(5+\alpha)/(3+\alpha)}.$$

This coincides with the Goldreich-Sridhar spectrum in the limit of weak anisotropy ($\alpha = 0$), and with the Iroshnikov-Kraichnan spectrum in the limit of strong anisotropy ($\alpha = 1$). As the external magnetic field increases from $\gamma \ll 1$ to $\gamma \gg 1$, the anisotropy of turbulent fluctuations increases, and the parameter α changes from $\alpha = 0$ to $\alpha = 1$. According to point 3, the corresponding scalings of the fluctuations with respect to their field-perpendicular and field-parallel dimensions are $\delta v_{\lambda} \propto \lambda^{1/(3+\alpha)}$, $\delta v_{\xi} \propto \xi^{1/3}$, and $\delta v_l \propto l^{1/2}$.

5. The smallest-scale “eddies” in our turbulent cascade ($\lambda \rightarrow 0$ for $\alpha \neq 0$) have a sheet like morphology, in agreement with micro current-sheet dissipative structures, numerically observed in MHD turbulence (Biskamp & Müller 2000). In the case of zero external magnetic field, $\alpha = 0$, the dissipative structures are filaments, which also agrees with numerical simulations (Padoan, et al. 2004).

6. Previous attempts to explain the numerically observed Iroshnikov-Kraichnan scaling (and the associated cascade-time increase (3)), in anisotropic sub-Alfvénic MHD turbulence ($\gamma \gg 1$) essentially invoked intermittency effects, (see, e.g., § 6.6.4 in Maron & Goldreich 2001; Müller, Biskamp & Grappin 2003). The intermittency effects are essential for explaining higher order statistics of MHD turbulence, that is, scaling of higher order structure functions of \mathbf{w} and \mathbf{z} . However, they usually provide only small corrections to the scaling of the second-order structure functions, and of the energy spectra. Such effects are not addressed in the present Letter; our derivation is based on the idea of scale-dependent dynamic alignment, which does not require intermittency.

I am grateful to Fausto Cattaneo and Samuel Vainshtein for many discussions of MHD turbulence, to Jason Maron, the referee of this Letter, for important comments on the results of Maron & Goldreich (2001), to Wolf-Christian Müller for discussion of the results of Müller, Biskamp & Grappin (2003), and to Dmitri Uzdensky for helpful comments on the physics and the style of the paper. This work was supported by the NSF Center for Magnetic Self-

Organization in Laboratory and Astrophysical Plasmas at the University of Chicago. The hospitality of the Aspen Center for Physics, where part of this work was done, is gratefully acknowledged.

REFERENCES

- Biskamp, D. *Magnetohydrodynamic Turbulence*. (Cambridge University Press, Cambridge, 2003).
- Biskamp, D. *Nonlinear Magnetohydrodynamics*. (Cambridge University Press, Cambridge, 1993).
- Biskamp, D. & Müller, W.-C., *Phys. Plasmas* **7** (2000) 4889.
- Cho, J. & Vishniac, E., *ApJ*, **539** (2000) 273.
- Cho, J., Lazarian, A., & Vishniac, E., *ApJ* **564** (2002) 291.
- Galtier, S., Nazarenko, S. V., Newell, A. C., & Pouquet, A., *J. Plasma Physics* **63** (2000) 447.
- Galtier, S., Nazarenko, S. V., Newell, A. C., & Pouquet, A., *ApJ* **564** (2002) L49.
- Goldreich, P. & Sridhar, S., *ApJ* **438** (1995) 763.
- Goldreich, P. & Sridhar, S., *ApJ* **485** (1997) 680.
- Grappin, R., Frisch, U., Léorat, J., & Pouquet, A., *Astron. Astrophys.* **105** (1982) 6.
- Grappin, R., Pouquet, A., & Léorat, J., *Astron. Astrophys.* **126** (1983) 51.
- Iroshnikov, P. S., *AZh*, **40** (1963) 742; *Sov. Astron.*, **7** (1964) 566.
- Kraichnan, R. H., *Phys. Fluids*, **8** (1965) 1385.
- Kulsrud, R. M., *Plasma Physics for Astrophysics*. (Princeton University Press, 2005).
- Maron, J., & Goldreich, P., *ApJ* **554** (2001) 1175.
- Müller, W.-C., Biskamp, D., & Grappin, R., *Phys. Rev. E* **67** (2003) 066302.
- Ng, C. S., & Bhattacharjee, A., *ApJ* **465** (1996) 845.
- Padoan, P., Jimenez, R., Nordlund, Å., & Boldyrev, S., *Phys. Rev. Lett.* **92** (2004) 191102.




Fig. 1.— A sketch of the shapes of turbulent fluctuations, “eddies,” in the Goldreich-Sridhar picture (left), and in our picture (right). The large-scale magnetic field is denoted by \mathbf{B} . As the turbulent cascade proceeds toward the smallest, dissipative scales, $\lambda \rightarrow 0$, the Goldreich-Sridhar “eddy” assumes the shape of a filament, while the “eddy” predicted in our model turns into a current sheet.