

# New constraints on MOND from galaxy clusters

Etienne Pointecouteau<sup>1\*</sup> and Joseph Silk<sup>1†</sup>

<sup>1</sup>*Astrophysics, University of Oxford, Keble Road, Oxford OX1 3RH*

## ABSTRACT

We revisit the application of Modified Newtonian Dynamics (MOND) to galaxy clusters. We confront the high quality X-ray data for eight clusters of galaxies observed by the *XMM-Newton* satellite with the predictions of MOND. We obtain a ratio of the MOND dynamical mass to the baryonic mass of  $M_m/M_b = 4.94 \pm 0.50$  in the outer parts (i.e  $r \sim 0.5 R_{vir}$ ), in the concordance cosmological model where the predicted asymptotic ratio, if any baryons are present, is  $7.7^{+1.4}_{-1.1}$  (at  $r \geq 0.3 R_{vir}$ ). We confirm that the MOND paradigm lowers the discrepancy between the binding mass and the baryonic mass in clusters by a factor of  $\sim 1.6$  at about half the virial radius. However, at this radius about 80% of the mass is still missing, and as pointed out by Sanders (2003), this necessitates a component of dark baryons or neutrinos in the cluster core. Concerning the neutrino hypothesis, application of the new data requires a minimum neutrino mass of  $m_\nu > 1.74 \pm 0.34$  eV to fill this gap. The corresponding  $2\sigma$  lower limit of  $m_\nu > 1.06$  eV is marginally inconsistent with the current constraints from the cluster number counts, and from the CMB and large scale structure data. MOND must invoke neutrinos to represent the main component that account for the missing mass problem in clusters.

**Key words:** Cosmology: observations – Galaxies: clusters: general

## 1 INTRODUCTION

The missing mass problem in clusters of galaxies arises from the comparison of the observed baryonic mass with the observed dynamic mass. The baryonic mass is mainly due to the hot intracluster gas that is well observed in X-rays via its free-free emission. The current status of the observed gas fraction in clusters gives a fairly well constrained value of about 12% (see Grego et al. 2001; Allen et al. 2003, for instance). Taking into account the stellar mass, this makes the discrepancy between the observed dynamic mass and the observed baryonic mass larger than a factor of 7.

The dark matter (DM hereafter) hypothesis appears to provide a seductive explanation of this problem. A new component of non-baryonic matter, insensitive to all interactions but gravitation, is introduced to fill the gap between the baryonic matter and the binding mass. While cosmological evidence is accumulating in favour of this scenario (see for instance Freedman et al. 2001; Spergel et al. 2003; Tegmark et al. 2004), it is disconcerting that the nature of the non-baryonic dark matter is completely unknown. Of course there are many candidates of varying degrees of detectability and plausibility (eg, review by Bertone et al. 2004)

As an alternative to dark matter scenarios, Milgrom (1983c) proposed a modification of the Newtonian dynamics effective at galactic and extra-galactic scales. This modified Newtonian dynamics (MOND hereafter) has been notably successful in explaining the discrepancy between rotation and luminosity curves in spiral galaxies (Milgrom 1983a,b), and claims other phenomenological successes (see Sanders & McGaugh 2002, for a review). Given that there is now a relativistic, Lorentz invariant generalisation of MOND that can incorporate both gravitational lensing and cosmology (Bekenstein 2004), it is timely to reexamine one of the few admitted failures of MOND. The discrepancy between the baryonic mass and the dynamical mass in clusters of galaxies is perhaps foremost among the issues that MOND has yet to convincingly address.

The first confrontation of X-ray observations of clusters with MOND (Gerbal et al. 1992) emphasised, despite some minor controversy (Milgrom 1993; Gerbal et al. 1993), the difficulties faced by MOND in passing the cluster test. The problem was revisited by Sanders (1994, 1999) and ended in a remaining discrepancy of a factor of 2-3 between the baryonic and the MOND masses. More recently, Aguirre et al. (2001) discussed observational evidence for three clusters for which the observed discrepancy is about 1-5 within 1 Mpc and is boosted to a factor  $\sim 10$  within the central 200 kpc, further weakening the reliability of the MOND paradigm. However Sanders (2003) responded with an update of his

\* E-mail: etienne@astro.ox.ac.uk (EP)

† E-mail: silk@astro.ox.ac.uk (JS)

earlier work, introducing an added *ad hoc* dark component at the cluster centre to reduce the discrepancy to only a factor of 1-3 overall in the cluster.

Some other tests have also been carried out using gravitational lensing data. They have also pointed out the difficulties faced by MOND at the cluster scale (Gavazzi 2002; Clowe et al. 2004).

In this paper, we test the MOND prediction in terms of dynamical mass with respect to the observed baryonic mass, by basing our work on high quality X-ray data recently obtained from observations with the *XMM-Newton* satellite for ten nearby relaxed clusters (Pointecouteau et al. 2005; Arnaud et al. 2005). Section 2 presents the context and the formalism within which the mass of clusters is derived in the MOND case. In section 3 we present the data and the working framework we adopt to compute the observed MOND masses. The comparison of the MOND mass with the Newtonian dynamic mass and the baryonic mass is developed in Sec. 4. The results are discussed with respect to previous studies in Sec. 5. Our new result is that, in addition to the known problem in the cluster cores, there is a significant discrepancy in the outer cluster, where an additional *ad hoc* dark component is needed to rescue MOND. We investigate the neutrino hypothesis for such a possibility.

Unless mentioned otherwise, we choose to work in a concordance model, using as cosmological parameters:  $\Omega = 1$ ,  $\Omega_m = 0.3$ ,  $\Omega_\Lambda = 0.7$  and  $H_0 = 70$  km/Mpc/s (referred as LCDM).

## 2 MOND DYNAMICAL MASSES

For a spherical system in hydrostatic equilibrium, the density and the temperature distributions are connected via the equation of hydrostatic equilibrium, so that the dynamical mass of a spherical system can be expressed as :

$$M_d(r) = -\frac{kT r}{G\mu m_p} \left[ \frac{d \ln n_g}{d \ln r} + \frac{d \ln T}{d \ln r} \right] \quad (1)$$

where  $G$  is the gravitational constant,  $m_p$  is the proton mass and  $\mu = 0.609$ .  $M_d(r)$  is the dynamical mass enclosed within a sphere of radius  $r$ . Here,  $n_g(r)$  and  $T(r)$  are the density and temperature radial distributions.

In the framework of modified Newtonian dynamics, the gravitational acceleration  $a$  is linked to the Newtonian acceleration  $g$  as follows:

$$a f(a/a_0) = g \quad (2)$$

where  $a_0$  is assumed to be a universal constant of acceleration in MOND. Its value as determined from the rotation curves of galaxies is about  $10^{-8}$  cm/s<sup>2</sup>. In the following, we adopt the value of  $0.8 \times 10^{-8}$  used by Sanders (1999) (hereafter S99 – referring to Begeman et al. 1991). The transition between the Newtonian and the MOND regimes is expressed through the functional  $f(x)$ ,  $x$  being  $a/a_0$ , that is also derived from the application of MOND to the rotation and luminosity curve of galaxies:

$$f(x) = \frac{1}{\sqrt{1+x^{-2}}} \quad (3)$$

We now quote Eq. 9 from S99, that gives the MOND

gravitational acceleration  $a$ , quantity that decreases with increasing radius:

$$a(r) = -\frac{kT}{\mu m_p r} \left[ \frac{d \ln n_g}{d \ln r} + \frac{d \ln T}{d \ln r} \right] \quad (4)$$

Eq. 8 from S99 relates the dynamical MOND mass,  $M_m$ , to the dynamical Newtonian mass,  $M_d$ :

$$M_m(r) = \frac{M_d(r)}{\sqrt{1+(a_0/a(r))^2}} \quad (5)$$

Further details of the MOND formalism can be found, for instance, in S99, Sanders & McGaugh (2002) and Sanders (2003).

According to Eq. 1 and Eq. 4, we can express the ratio of the Newtonian dynamical mass to the MOND mass as:

$$\frac{M_d}{M_m}(r) = \sqrt{1 + \left( \frac{a_0}{G} \frac{r^2}{M_d} \right)^2} \quad (6)$$

In the following we refer to the Newtonian dynamical mass,  $M_d(r)$ , as the dynamical mass, and to the dynamical mass in modified Newtonian dynamics,  $M_m$ , as the MOND mass.


## 3 WORKING FRAMEWORK

### 3.1 The X-ray data

Recently, Pointecouteau et al. (2005, hereafter PAP05) have published the observed mass profiles for a set of ten nearby ( $z \leq 0.15$ ) and relaxed galaxy clusters observed with the *XMM-Newton* satellite. These clusters, chosen for their regular X-ray morphologies, have dynamical masses determined through the hypothesis of spherical symmetry and the use of Eq. 1. They cover a large temperature range of [2 – 9] keV, corresponding to a dynamical mass range of  $[1.2 - 12.0] \times 10^{14} M_\odot$  enclosed within a radius  $R_{200}$  (i.e a radius encompassing 200 times the critical density of the Universe at the cluster redshift), and a range of radii between 0.01 to 0.7 in units of  $R/R_{200}$ .

From the initial sample of ten clusters, we kept only eight for the present work: A1991, A2717, A2597, A1068, A478, A1413, PKS0745, A2204. We excluded A1983 and MKW9. First they are the only two clusters of the sample that are not observed up to  $0.5 R_{200}$ . Indeed, we need to keep to the observed radial range of each cluster to avoid any extrapolation, and to derive reliable observational constraints. Meanwhile MWK9 has also the most disturbed morphology of the 10 clusters (Pratt & Arnaud 2005). Both also exhibit unexpectedly low gas fractions that may turn them into outliers in terms of the average gas fraction in clusters (Grego et al. 2001; Allen et al. 2003). As our sample is quite small, excluding those two clusters may avoid any poissonian bias on the computation of the intrinsic dispersion for the ratio of the MOND mass to the baryonic mass. However, at the scale of our sample, this also may slightly bias the results in favour of MOND predictions, for which low gas fraction systems will be difficult to explain (as it increases the discrepancy between the observed baryonic mass and the dynamical mass).

We used the observed density and temperature profiles derived by PAP05 to compute the observed ratios of the dynamical mass to the MOND mass and of the MOND mass to the baryonic mass.



**Figure 1.** Observed scaled profiles for the ratio of the dynamical mass to the MOND mass. The average profile over the sample is overplotted (solid black line) together with its  $1\sigma$  dispersion (dashed lines). The triangles mark the estimate from the self-similar hypothesis for each cluster (see Sect. 3.2).

### 3.2 Scaled MOND profiles

In hierarchical structure formation, the virialized part of clusters at a given redshift is encompassed within a sphere with a radius corresponding to a fixed density contrast ( $\sim 200$ ) with respect to the critical density of the Universe.  $R_{200}$  is thus considered to be the virial radius of the cluster (see Arnaud et al. 2004, and references therein).

Thus to cross-compare the internal structure of clusters, one has to look at quantities scaled according to  $R_{200}$ . As an extension, to compare the MOND and the dynamical masses as a function of radius, we choose to scale both quantities according to the virial radius (i.e.  $R_{200}$ ). For each cluster, we used the value of  $R_{200}$ , reported by PAP05 for a LCDM cosmology (see their table 2), obtained from the fit of NFW profiles (Navarro et al. 1997) over the observed mass profiles. In the framework of MOND structure formation, Knebe & Gibson (2004) have shown that the most massive galactic halos formed in their numerical simulations could be fitted by a NFW profile. We could thus fit each MOND mass profile with a NFW profile to derive the corresponding characteristic scaling radius  $R_{200}^M$ . However, for a given cluster, the values of  $R_{200}$  and  $R_{200}^M$  are likely to be different. Scaling  $M_d(r)$  and  $M_m(r)$  respectively with  $R_{200}$  and  $R_{200}^M$  will lead us to compare these two profiles at different physical scales. For our purpose, working with scaled radii is required by the use of a sample of clusters with different masses, but is not required to compare the MOND and dynamical masses for a given cluster. A single scaling has to be adopted for each cluster. We thus adopt  $R_{200}$  as a scaling radius for both dynamical and MOND profiles.

### 3.3 The baryonic mass in clusters

The observational constraints on the gas fraction in clusters strongly favour an asymptotic value of about 10-12% beyond  $0.3 R_{200}$  (Grego et al. 2001; Allen et al. 2003), in a LCDM cosmology. For the sample used here, the average gas fraction at a density contrast of  $\delta = 1000$  (i.e. at

$r = 0.47 \pm 0.02 R_{200}$  – see APP05) is  $\langle f_{gas} \rangle = 0.11 \pm 0.02$ . We adopted this value in the following, as the average asymptotic value for the gas fraction.

We now can further investigate the issue of the stellar mass. For the stellar mass, S99 used a correlation derived from the early work of David et al. (1990) between the luminosity and the gas masses in clusters leading to:  $f_\star = M_\star/M_{gas} \simeq 0.7(kT/\text{keV})^{-1}h^{3/2}$ . We used this estimate when working in CDM cosmology, making use of the spectroscopic temperature measured between  $[0.1, 0.5] R_{200}$  (see APP05). Nevertheless, more recent work allows us to revise this crude estimate of  $f_\star$ . For example, Lin et al. (2003) derived a variation of  $f_\star$  with the total mass, and thus with the temperature. From their equation 10 and the values of  $M_{500}$  derived by APP05 (see their table 1), we obtain an average value of  $\langle f_\star \rangle = 0.12 \pm 0.03$  (values ranging from 0.1 to 0.17). In related work, Voevodkin & Vikhlinin (2004) obtained consistent, but slightly higher, values, and concluded that the stellar mass represents about 15% of the gas mass. We adopted this value of  $f_\star = 15\%$  for our sample. The baryon fraction will then be  $f_b = 0.13 \pm 0.02$ . In order to cope with the baryonic mass in clusters, the MOND mass has then to be  $7.7_{-1.0}^{+1.4}$  times lower than the dynamical mass once the baryonic mass reaches its asymptotic value (i.e.  $r > 0.3 R_{200}$ ).

## 4 MATCHING THE BARYON CONTENTS OF CLUSTERS

In an initial approach, taking into account the self-similar nature of the cluster population, we can express the ratio  $M_d/M_m$  as a function of  $kT$ . We make use of the scaling relations between the mass and the temperature:  $M = A(z) T_\delta^\alpha$ , and between the mass and the radius  $M = 4\pi/3 \delta \rho(z) R_\delta^3$ . Then Eq. 6 can be rewritten as follows:

$$\frac{M_d}{M_m}(r) = \sqrt{1 + B(z) T_\delta^{-2\alpha}} \quad (7)$$

where  $B(z) = (3a_0/4\pi\delta\rho_c(z))A(z)^{-1}$ , and  $T_\delta$  is the temperature at density contrast  $\delta$ .

Using the  $R - T$  relation published by APP05 at  $\delta = 1000$  (a density contrast corresponding to observed radii for all eight clusters) we estimated from the above equation  $M_d/M_m = 1.63 \pm 0.29$ , which leads to  $M_m/M_b \simeq 4.7$ .

We can reverse the computation and estimate the value of  $a_0$  needed to reach an equality between the MOND mass  $M_m$  and baryonic mass  $M_b$ . We obtain  $a_0 = (4.75 \pm 1.24) \times 10^{-8} \text{ cm/s}^2$  at  $\delta = 1000$  using the eight clusters.

From the mass profiles derived by PAP05 in a LCDM cosmology, we computed the observed ratio  $M_d/M_m$  from Eq. 6 for the eight clusters of the sample. The profiles (scaled to  $R_{200}$ ) are shown in Fig. 1. The estimates from the self-similar hypothesis are shown for  $\delta = 1000$  as the observed triangles, and match the computed ratio very well.

With the previously given notations and Eq. 6, we can express the ratio of the MOND mass to the baryonic mass

as:

$$\frac{M_m}{M_b}(r) = \left[ f_{gas}(1 + f_*) \sqrt{1 + \left( \frac{a_0}{G} \frac{r^2}{M_d} \right)^2} \right]^{-1} \quad (8)$$

In Table 1, we report the average baryon fraction values and the average  $M_m/M_b$  and  $M_d/M_m$  ratios for the measured clusters at a given radius. The computation were done at the following radii: the physical radius of 750 kpc in order to directly compare with S99, and the radii corresponding to the density contrasts of  $\delta = 1000$  and  $\delta = 15000$  (i.e.  $0.47 \pm 0.02 R_{200}$  and  $0.10 \pm 0.01 R_{200}$  average over the sample). Those two radii mark the boundaries of the radial range over which the observational constraints are especially well tied down (see PAP05, APP05).

From Fig. 1, it is obvious that the ratio  $M_d/M_m$  decreases, thus the discrepancy between the MOND and the baryonic mass increases, towards the centre of clusters as the acceleration returns to the Newtonian regime. At  $\sim 0.1 R_{200}$ , the ratio is  $1.08 \pm 0.06$ . The discrepancy also increases for massive (i.e. hot) systems. Indeed if we only consider the 5 hot clusters<sup>1</sup> we obtain at  $\delta = 1000$  a ratio of  $M_d/M_m = 1.43 \pm 0.08$ , an even smaller value. The corresponding value of  $a_0$  to match the baryonic mass has to be  $a_0 = (5.58 \pm 0.63) \times 10^{-8} \text{ cm/s}^2$ .

## 5 DISCUSSION

### 5.1 MOND as a stand alone solution

For a direct comparison with S99, it is interesting to performed the computation at the radius of 750 kpc with  $\Omega_m = 1$  and  $H_0 = 50 \text{ km/s/Mpc}$ . For the five clusters observed beyond 750 kpc (i.e. the 5 hot clusters), the average gas fraction is then  $0.19 \pm 0.02$  and the derived ratio of the MOND mass to the baryonic mass is  $M_m/M_b = 3.84 \pm 0.33$ . For the same cosmological setup, S99 claimed that the MOND context reduces the discrepancy with the baryonic mass down to an “acceptable” factor of  $\sim 2$ . However, our value of 3.8 is significantly higher, and does not even agree with a factor of 2 within a  $3\sigma$  limit.

As we are working with a sample of nearby clusters, the most important cosmological parameter is the Hubble constant. Indeed, the baryon fraction in clusters scales with  $h^{-3/2}$ , thus decreases with increasing values of  $H_0$ . Current observational evidence strongly favours a flat Universe with a low matter density  $\Omega_m \sim 0.3$ , a component due to dark energy which can be represented by a positive cosmological constant  $\Lambda$ , and a high value for the Hubble constant ( $H_0 \sim 70 \text{ km/s/Mpc}$ ) (see Freedman et al. 2001; Spergel et al. 2003; Tegmark et al. 2004; Seljak & et al. 2005, for instance). In such a universe (i.e. LCDM)  $M_d/M_m$  decreases to  $1.63 \pm 0.31$  ( $1.08 \pm 0.06$ ) at  $\sim 0.5 R_{200}$  ( $\sim 0.1 R_{200}$ ), so that the discrepancy is increased between the MOND dynamic mass and the baryonic mass. The corresponding ratio of the MOND mass to the baryonic mass is now  $4.94 \pm 0.50$  ( $10.6 \pm 3.77$ ). This is more than a factor of

**Table 1.** Ratios of the dynamical mass to the MOND mass and of the MOND mass to the baryonic mass.

Radius	$f_b$	$M_d/M_m$	$M_m/M_b$	$N_c^a$
$R = 750 \text{ kpc}$	$0.13 \pm 0.01$	$1.34 \pm 0.14$	$5.61 \pm 0.55$	5
$\delta = 15000$	$0.09 \pm 0.03$	$1.08 \pm 0.06$	$10.6 \pm 3.77$	8
$\delta = 1000$	$0.13 \pm 0.02$	$1.63 \pm 0.31$	$4.94 \pm 0.50$	8
$\delta = 1000^b$	$0.14 \pm 0.02$	$1.43 \pm 0.08$	$5.10 \pm 0.56$	5

(a) Number of clusters used. (b) Ratio for the hot clusters only (i.e.  $kT > 3.5$ ).

two above the value derived by S99. The evidence is confirmed if we only consider the hot systems. Indeed, for clusters with ( $kT > 3.5 \text{ keV}$ ), we derive  $M_d/M_m = 1.43 \pm 0.08$  at  $\sim 0.5 R_{200}$ , which makes the ratio of the MOND mass to the baryonic mass  $\sim 5.10 \pm 0.56$ . Thus in all cases, within a  $3\sigma$  (i.e. 99% confidence) the ratio  $M_m/M_b$  in a MOND cosmology will be greater than 3.4, making MOND unable to fully overcome the missing mass problem in clusters.

The case of A1413 is even more eloquent. Indeed, this cluster has been observed beyond  $R_{500}$  (i.e. a physical radius of 1129 kpc corresponding to  $\sim 0.7 R_{200}$  – Pratt & Arnaud 2002), a radius that conservatively is often taken as the outer bound of the virialized part in clusters. Moreover, for this cluster the *XMM-Newton* results agrees very well with the results derived from the *Chandra* observations, especially in term of the shape of the temperature profile (Vikhlinin et al. 2005). At the radius of  $R_{500}$ , the measured gas fraction is  $f_{gas} = 0.15 \pm 0.01$ , which makes the baryon fraction  $0.17 \pm 0.01$  (see Sect. 3.3). We derived a corresponding ratio  $M_m/M_b = 3.6 \pm 0.7$ . Such an observed discrepancy measured within a radius closing the virial radius puts a very tight constraint on the MOND paradigm.

In fact, S99 relies on hypotheses that together might have biased its results down towards an optimistic value: (i) the use of a  $\beta$ -model to describe the gas distribution down to 750 kpc contributes to overestimate the gas mass at large radii. Moreover the spatial resolution of the current X-ray data has ruled out the  $\beta$ -model as a fitted representation of the observed X-ray surface brightness profiles (especially in nearby clusters – see PAP05 for instance); (ii) The use of physical radius as a working radius could have induced biases as clusters of very different masses are then compared at various different density contrasts; (iii) The isothermality of the intra-cluster medium was forced by the single over-all temperature measurements available for the considered sample. However, in contrast we have made use of accurately measured temperature profiles.

If we consider the problem in terms of the value of the characteristic MOND acceleration, we derive values of  $a_0$  that are 5 times larger (and still 3.5 times larger within the  $3\sigma$  limit) than the value derived from the rotation curves of galaxies, and are thus unacceptable values.

### 5.2 Adding a dark component

A last alternative to rescue MOND is to invoke a non-luminous component at the centre of clusters, as suggested recently by Sanders (2003) (S03 hereafter). This author in-

<sup>1</sup> i.e. A1413, A478, A2204, PKS0745, A1068

roduced this dark component on an *ad hoc* basis to explain the huge observed discrepancy within the cluster centre (Aguirre et al. 2001). Indeed, the baryon fraction is lower in the inner parts, so that the ratio to the MOND mass increases. With respect to our sample, if such a component was to exist it would have to account for 81% ( $\pm 4\%$ ) of the total mass at  $\sim 0.5 R_{200}$ . In other terms, this means that MOND just reduces the missing mass problem in clusters by about 20% (at half the virial radius) compare to a standard DM scenario, but does not solve it. Therefore, some exotic dark component has to be added to fill the remaining gap of 80%. This component will become the main explanation for the missing mass, and will thus draw a cosmological setup that will be closer to a mixed DM scenario than to a dominating MONDian scenario.

It is highly unlikely that 80% of the missing mass in cluster can be due to hidden baryons. Indeed, if atomic or molecular hydrogen exist within the intra-cluster medium, it will have to face its thermal conditions. Such a component could only account for a small fraction of the thermal baryons. The relativistic populations are also known to be a minor component in terms of mass, as otherwise they will be expected to produce a strong hard X-ray signal, and stronger radio emissions than those currently observed.

Nevertheless, we can follow the suggestion by S03 to consider, as potential candidates for this dark component, massive neutrinos that aggregate at cluster scale. In the MOND case, according to S03, the needed neutrino mass per neutrino flavour needed for MOND to cope with the observed baryonic mass is about 2 eV. Further assuming a constant density sphere for his dark component and taking into account the phase space density limit for neutrinos, S03 derived an upper limit for the neutrino density after their collapse and accretion within structure of:  $\rho_\nu \leq (4.8 \times 10^{-24})(m_\nu/2\text{eV})^4(T_{\text{keV}})^{3/2} \text{ kg m}^{-3}$ . We make use of this limit on the neutrino density, and we use the spectroscopic temperatures measured for each cluster of our sample between 0.1 and 0.5  $R_{200}$  (see PAP05). From our eight clusters, it is possible to compute the needed neutrino mass to equate the missing mass at a given radius (i.e.  $M_m(r) - M_b(r)$ ) with the contribution of the massive neutrino to the cluster total mass. S03 hypothesis of the neutrino accretion mainly concerned the central parts of clusters. In our study, to explain the  $\sim 80\%$  of missing mass in MOND, we add to extend the radius of the neutrino sphere down to  $R_{1000}$ . The minimum neutrino mass then required is  $m_\nu > 1.74 \pm 0.34 \text{ eV}$ . This is a strongly constraining value for the neutrino mass, which makes the lower bound for the neutrino mass becomes  $\sim 1.06 \text{ eV}$ , within a  $2\sigma$  limit (i.e. 95% confidence).

To date particle physics experiments (single and double beta-decay, neutrino oscillations measurements) lead to a wide range of upper limit for the neutrino mass going from  $\sim 0.8 \text{ eV}$  to  $\sim 3 \text{ eV}$  (Fogli et al. 2004). Those constraints directly apply to MOND, and they are consistent with our lower limit taking into account their large variation. If to date the constraints derived from astrophysical data seems to be tighter, they may be considered as irrelevant in a MOND framework (i.e.  $\Omega_\nu h^2 = \sum m_{\nu_i}/94\text{eV}$ ) as they are derived within the framework of a standard cosmological DM model for structure formations. Nevertheless, within the cosmological MOND+neutrino setup we end up

with, MOND plays a minor role, where the neutrinos are the major component to explain the missing mass in the Universe. We can thus reconsider the astrophysical constraints in this context. As the number of clusters is linked to the matter content of the Universe, thus to  $\Omega_m$  and  $\Omega_\nu$ , the cluster number counts can be used to give an upper limit on the neutrino mass (Kahniashvili et al. 2005; Elgarøy & Lahav 2005). With the following cosmological setup:  $\Omega_m = 0.3$ ,  $h = 0.7$ ,  $n = 1$  and  $0.7 < \sigma_8 < 1.1$  (to account for the variation in  $\sigma_8$  determinations – see for instance Tegmark et al. (2004); Seljak et al. (2005)), current cluster number counts lead to an  $2\sigma$  upper limit of  $m_\nu < 0.8 \text{ eV}$  (Fukugita et al. 2000; Allen et al. 2003). A value marginally incompatible with the lower bound was derived here. This inconsistency is even stronger with the  $2\sigma$  upper limit of  $\sim 0.3 \text{ eV}$  derived from the combination of the WMAP, 2dFGRS and SDSS data (Spergel et al. 2003; Tegmark et al. 2004; Seljak & et al. 2005; Elgarøy & Lahav 2005). However, one has to keep in mind that there is quite a large dispersion between those upper limits (Elgarøy & Lahav 2005), that may just leave enough room for the MOND+neutrino hypothesis to stand as a reliable paradigm.

## 6 CONCLUSION

We have revisited the case of the galaxy cluster scale in the framework of the MOND paradigm. We based our study on a set of consistent and recent high quality X-ray data obtained with the last generation of X-ray satellites (*XMM-Newton*). For eight nearby and relaxed clusters, we confirmed that MOND alone is not able to explain the missing mass problem in clusters of galaxies. Almost 80% of the mass is still missing at half the virial radius of clusters in a MOND Universe. Thus MOND is not the main solution to the missing mass problem at cluster scales.

Indeed, it undeniably requires extra exotic DM to survive. In this context we have investigated the hypothesis of massive neutrinos distributed as a sphere of constant density up to half the virial radius of galaxy clusters. Under this hypothesis, we derived a very tight observational lower bound for the neutrino mass,  $m_\nu > 1.06 \text{ eV}$  (95% confidence), which is marginally inconsistent with the constraints from the cluster number counts and from the CMB and LSS constraints. The consistency with the current direct experiments to measure the neutrino mass is spoiled by their associated large uncertainties. The perspectives of dedicated CMB and Large-scale structure experiments in coming years will bring definitive constraints on  $m_\nu$  in the astrophysical context, and should definitively settle the case of the MOND paradigm.

Note that recently Skordis et al. (2005) have been studying the formation of structure in the relativistic MOND framework (i.e. the Bekenstein theory – see Bekenstein (2004)). To reproduce the observed angular power spectrum of the CMB, those authors appeal to massive neutrinos with  $m_\nu \simeq 2 \text{ eV}$ . So unless the main fraction of the baryonic content of clusters remains hidden from the current observations, in any case a large neutrino mass (or another exotic massive candidate) is needed to promote MOND as a reliable paradigm. As such, massive neutrinos would then become a major component of the Universe mat-

ter content. This will turn the cosmological framework more into a mixed DM cosmology than into a MONDian cosmology.

## ACKNOWLEDGEMENTS

We thank the anonymous referee for her/his fruitful comments. We are grateful to Monique Arnaud, Mathieu Langer, Adi Nusser and Gabriel Pratt for useful discussions and comments. EP also thanks Monique Arnaud for software issues and Adi Nusser for enlightenments on the basis of MOND. EP acknowledges the financial support of the Leverhulme Trust (UK).

## REFERENCES

- Aguirre A., Schaye J., Quataert E., 2001, *ApJ*, 561, 550  
 Allen S. W., Schmidt R. W., Fabian A. C., Ebeling H., 2003, *MNRAS*, 342, 287  
 Arnaud M., Pointecouteau E., Pratt G. W., 2005, *A&A* accepted (astro-ph/0502210)  
 Arnaud M., Pratt G. W., Pointecouteau E., 2004, *Memorie della Societa Astronomica Italiana*, 75, 529  
 Begeman K. G., Broeils A. H., Sanders R. H., 1991, *MNRAS*, 249, 523  
 Bekenstein J. D., 2004, *Phys. Rev. D*, 70, 083509  
 Bertone G., Hooper D., Silk J., 2004, *Phys. Rep.*, 405, 279  
 Clowe D., Gonzalez A., Markevitch M., 2004, *ApJ*, 604, 596  
 David L. P., Arnaud K. A., Forman W., Jones C., 1990, *ApJ*, 356, 32  
 Elgarøy Ø., Lahav O., 2003, *Journal of Cosmology and Astro-Particle Physics*, 4, 4  
 Elgarøy Ø., Lahav O., 2005, *New Journal of Physics*, 7, 61  
 Fogli G. L., Lisi E., Marrone A., Melchiorri A., Palazzo A., Serra P., Silk J., 2004, *Phys. Rev. D*, 70, 113003  
 Freedman W. L., et al. , Madore B. F., Gibson B. K., Ferrarese L., Kelson D. D., Sakai S., Mould J. R., Kennicutt R. C., Ford H. C., Graham J. A., Huchra J. P., Hughes S. M. G., Illingworth G. D., Macri L. M., Stetson P. B., 2001, *ApJ*, 553, 47  
 Fukugita M., Liu G., Sugiyama N., 2000, *Physical Review Letters*, 84, 1082  
 Gavazzi R., 2002, *New Astronomy Review*, 46, 783  
 Gerbal D., Durret F., Lachieze-Rey M., Lima-Neto G., 1992, *A&A*, 262, 395  
 Gerbal D., Durret F., Lachieze-Rey M., Lima-Neto G., 1993, *A&A*, 273, L9+  
 Grego L., et al. , 2001, *ApJ*, 552, 2  
 Knebe A., Gibson B. K., 2004, *MNRAS*, 347, 1055  
 Kahnashvili T., von Toerne E., Arhipova N. A., Ratra B., 2005, *Phys. Rev. D*, 71, 125009  
 Lin Y., Mohr J. J., Stanford S. A., 2003, *ApJ*, 591, 749  
 Milgrom M., 1983a, *ApJ*, 270, 371  
 Milgrom M., 1983b, *ApJ*, 270, 384  
 Milgrom M., 1983c, *ApJ*, 270, 365  
 Milgrom M., 1993, *A&A*, 273, L5+  
 Navarro J. F., Frenk C. S., White S. D. M., 1997, *ApJ*, 490, 493  
 Pointecouteau E., Arnaud M., Pratt G. W., 2005, *A&A*, 435, 1  
 Pratt G. W., Arnaud M., 2002, *A&A*, 394, 375  
 Pratt G. W., Arnaud M., 2005, *A&A*, 429, 791  
 Sanders R. H., 1994, *A&A*, 284, L31  
 Sanders R. H., 1999, *ApJ (Lett.)*, 512, L23  
 Sanders R. H., 2003, *MNRAS*, 342, 901  
 Sanders R. H., McGaugh S. S., 2002, *Ann. Rev. Astron. Ap.*, 40, 263  
 Seljak U., et al. 2005, *Phys. Rev. D*, 71, 103515  
 Seljak U., et al., 2005b, *Phys. Rev. D*, 71, 043511  
 Skordis C., et al., 2005, *MNRAS* submitted (astro-ph/0505519)  
 Spergel D. N., et al. , 2003, *ApJ Suppl.*, 148, 175  
 Tegmark M., et al. , 2004, *Phys. Rev. D*, 69, 103501  
 Vikhlinin A., et al., 2005, astro-ph/0412306  
 Voevodkin A., Vikhlinin A., 2004, *ApJ*, 601, 610