ROTATION OF 10 Be STARS THROUGH FOURIER TRANSFORM ANALYSIS

NATAŠA GAVRILOVIĆ

¹ Astronomical Observatory, Volgina 7, 11000 Belgrade, Yugoslavia E−mail ngavrilovic@aob.aob.bg.ac.yu

Abstract. Here we determine the projected rotational velocity of 10 Be stars using Fourier Transform Method. Also, we discuss the gravity darkening and extend of deviation from solid body rotation for our sample of stars. We found that 7 of considered stars are affected by strong gravity darkening or/and solar differential rotation.

1. INTRODUCTION

Since Be stars are fast rotators, our knowledge about their fundamental parameters, in particular the projected rotational velocity, is subject to considerable uncertainty. Their gravity and temperature are aspect angle-dependent and do not have the straightforward meaning as they have in slow rotators (Slettebak 1976,1982; Brown & Verschneren 1997; Brown & Verschueren 1997; Halbedel 1996).

Projected rotational velocity $V_e \sin(i)$ for ten stars in our sample of Be stars has been determined using spectra from CDS data base¹. In order to determine the projected rotational velocity, Fourier Transform Method-FTM (Jankov 1995; Jankov et al. 2000) has been applied to each individual spectrum of our sample of stars, since the position of the first minimum is determined by the rotational broadening, and does not depend on other broadening mechanism. Reiners (2003) showed that the ratio of the first two minimum positions q_2/q_1 of the Fourier Transform can be also a reliable parameter to conclude the amount of gravity darkening or differential rotation. A value of $q_2/q_1 < 1.72$ is a direct indication for a gravity darkening or a solar-like differential rotation law, while $q_2/q_1 > 1.83$ indicates anti-solar differential rotation. The measurement of q_2/q_1 can be used without any modeling of line profiles.

2. RESULTS AND DISCUSSION

Using FTM and values for limb-darkening coefficient given by Claret (2000), we obtained projected rotational velocity, and also ratio of the first two minimum positions for 10 Be stars in our sample. In Table 1. are given values for q_2/q_1 , $V_e \sin(i)$ and (the last column) the determination of $V_e \sin(i)$ for considered stars by Chauville et al. (2001). As one can see from Table 1, our calculated values for $V_e \sin(i)$ are in good agreement with those determined by Chauville et al. (2001). Only three stars from the group have value $q_2/q_1 > 1.72$, so it can be concluded that the stars HD 75311 and HD 120324 have solid-body rotation and the mass concentration in the center of the star. The third one, HD 201733 has anti-solar differential rotation. Our analysis for the rest of the stars in the sample revealed a small q_2/q_1 ratio (less than 1.72) which could be interpreted as gravity darkening or/and solar differential rotation.

¹http://cdsweb.u-strasbg.fr/cgi-bin/Cat?J/A%2bA/378/861

HD	Wavelenght[nm]	$V_e \sin(i)[km/s]$	q_2/q_1	$V_e \sin(i)[km/s]^*$
10144	HeI 438.81	217	1.62	235
56139	HeI 438.81	88	1.16	85
57219	HeI 438.81	90	1.17	80
75311	HeI 438.81	264	1.74	268
77320	HeI 438.81	338	1.49	345
189687	HeI 438.81	211	1.27	200
201733	HeI 438.81	351	2.22	340
210129	HeI 438.81	141	1.70	130
120324	HeI 438.81	156	1.74	124
224544	HeI 438.81	239	1.36	260

Table 1: Determined values of $V_e \sin(i)$ in [km/s] and q_2/q_1 for the sample of 10 Be stars. In the last column (denoted with *) are values given by Chauville et al.(2001)




Figure 1: Flux line profile of the HeI 438.8 nm line of HD 10144 (top) and its Fourier transform (bottom). The Fourier frequency was reduced to velocity units, so that the first minimum of the Fourier transform of a rotational profile points to the projected rotational velocity of the star.

In the case of Achernar (HD 10144), no evidence has been found for differential rotation in high quality spectroscopy of the star (e.g. Gray 1977; Howarth & Smit 2001). It means that our result implies a strong gravity darkening which could be expected taking into account the extremely oblate shape of the star (Domiciano de Souza et al. 2003).

References

Brown, A.G.A., & Verschueren, W. 1997, Astron. Astrophys., 319, 811

Chauville, J., Zorec, J., Ballereau, D. et al. 2001, Astron. Astrophys., 378, 861 (Tables are only available in full in electronic form at CDS via

http://vizier.u-strasbg.fr/viz-bin/VizieR

Claret, A. 2000, Astron. Astrophys., 363, 1081

Domiciano de Souza, A., Kervella, P., Jankov, S., Abe, L., Vakili, F., et al. 2003, Astron. Astrophys., 407L, 47

Gray, D. F. 1977 ApJ, **211**, 198

Halbedel, E.M. 1996, *PASP*, **108**, 833 Howarth, I.D., Smith, K.C., 2001 *MNRAS*, **327**, 353

Jankov S.,1995, Publ. Obs. Astron.Belgrade, 50, 75

Jankov S., Janot-Pacheco, E., Leister, N.V., 2000, ApJ, 540, 535

Reiners, A. 2003, Astron. Astrophys., 408, 707

Slettebak, A., Collins II, G.W., & Traux, R. 1992, ApJS, 81, 335

Steele, I.A., Negueruela, I., & Clark, J.S. 1999, A & AS, 137, 147