

# CALIBRATING THE SOLAR DYNAMO: MAGNETIC ACTIVITY CYCLES OF SOUTHERN SUN-LIKE STARS

T. S. Metcalfe<sup>1</sup>, T. J. Henry<sup>2</sup>, M. Knölker<sup>1</sup>, and D. R. Soderblom<sup>3</sup>

<sup>1</sup>High Altitude Observatory, NCAR, P.O. Box 3000, Boulder, CO 80307 USA (travis@ucar.edu, knoe@ucar.edu)

<sup>2</sup>Department of Physics and Astronomy, Georgia State University, Atlanta, GA 30302 USA (thenry@chara.gsu.edu)

<sup>3</sup>Space Telescope Science Institute, 3700 San Martin Drive, Baltimore, MD 21218 USA (drs@stsci.edu)

## ABSTRACT

The solar magnetic activity cycle is responsible for periodic episodes of severe space weather, which can perturb satellite orbits, interfere with communications systems, and bring down power grids. Much progress has recently been made in forecasting the strength and timing of this 11-year cycle, using a predictive flux-transport dynamo model [1, 2]. We can strengthen the foundation of this model by extending it to match observations of similar magnetic activity cycles in other Sun-like stars, which exhibit variations in their Ca II H and K emission on time scales from 2.5 to 25 years [3]. This broad range of cycle periods is thought to reflect differences in the rotational properties and the depth of the surface convection zone for stars with various masses and ages. Asteroseismology is now yielding direct measurements of these quantities for individual stars, but the most promising asteroseismic targets are in the southern sky ( $\alpha$  Cen A,  $\alpha$  Cen B,  $\beta$  Hyi), while the existing activity cycle survey is confined to the north. We are initiating a long-term survey of Ca II H and K emission for a sample of 92 southern Sun-like stars to measure their magnetic activity cycles and rotational properties, which will ultimately provide independent tests of solar dynamo models.

Key words: magnetic fields; stars: activity, chromospheres, interiors, oscillations; surveys.

## 1. CONTEXT

Astronomers have been making telescopic observations of sunspots since the time of Galileo, gradually building a historical record showing a periodic rise and fall in the number of sunspots every 11 years. We now know that sunspots are regions with an enhanced local magnetic field, so this 11-year cycle actually traces a variation in surface magnetism. Attempts to understand this behavior theoretically often invoke a combination of differential rotation, convection, and meridional flow to modulate the field through a magnetic dynamo. Significant progress in this area quickly followed after helioseismology allowed




Figure 1. A flux-transport dynamo model correctly reproduces the relative amplitudes of the past 8 solar cycles, and then predicts the strength and timing of the next cycle (from [2], reproduced with permission).

the detailed characterization of the Sun's interior structure and dynamics over the past decade (e.g., the convection zone depth and meridional flow).

This has gradually led to the development of a predictive flux-transport dynamo model, which has recently been used to forecast the timing [1] and strength [2] of the upcoming solar magnetic cycle. By calibrating the relationship between sunspot area and surface magnetic flux using modern measurements, historical sunspot data going back more than 125 years could be used to test the model. After initializing the dynamo with the first several cycles, the model correctly reproduced the relative amplitudes of the subsequent 8 cycles, and was then used to predict the characteristics of future cycles (see Fig. 1). This is not merely of academic interest, since the magnetic maximum of the Sun is accompanied by severe space weather events. Advance knowledge of the timing and overall




Figure 2. The Ca II HK index of the Sun observed by the Mount Wilson survey (adapted from [3]).

strength of the coming cycle allows us to prepare for the adverse consequences, which include the destabilization of orbiting satellites, interference with communications systems, and the disruption of power grids.

While the success of this dynamo model in reproducing the solar cycle is very encouraging, there will soon be new opportunities for independent tests using the magnetic activity cycles of other Sun-like stars. The properties of the dynamo—most notably the cycle period—are sensitive to specific physical circumstances, such as the meridional flow, the degree of latitudinal differential rotation and the depth of the convection zone. By matching the dynamo model to the single set of circumstances represented by the Sun, we risk the possibility of fine tuning, which could ultimately limit our comprehension of the underlying physical mechanisms. If we exploit the magnetic activity cycles observed in a wide variety of Sun-like stars, we can extend the calibration to dozens of unique circumstances and ensure that the physical basis of the model is more broadly applicable. Specifically, observations of stellar activity cycle periods and differential rotation profiles (from this proposed survey, and its existing northern counterpart) as well as stellar convection zone depths (from ground- and space-based asteroseismology) will allow us to test the dynamo model by inverting the implied stellar meridional flow, and comparing it to the solar case.

Although we can rarely observe spots on other Sun-like stars directly, these areas of concentrated magnetic field produce strong emission in the Ca II H (396.8 nm) and K (393.4 nm) spectral lines. The intensity of the emission scales with the amount of non-thermal heating in the chromosphere, making these lines a useful spectroscopic proxy for the strength of, and fractional area covered by, magnetic fields [4]. Disk-integrated time-series measurements of Ca II H and K emission in the Sun easily reveal the 11-year cycle (see Fig. 2). Similar measurements for a large sample of northern Sun-like stars began at the Mount Wilson Observatory in 1966 [5], and continue to this day [6].

The Mount Wilson Ca HK survey, as it is known, has revealed magnetic activity variations in Sun-like stars with cycle periods ranging from 2.5 to more than 25 years. The frequent time sampling for 111 stars since 1980 also allows the measurement of rotation periods, from the re-

peated signature of individual active regions rotating into view and then out again. In some cases the rotation periods appear to drift over longer timescales, due to the migration of active regions to lower latitudes which are rotating faster (the same mechanism that is responsible for the “Butterfly Diagram” on the Sun). Thus, it is sometimes possible to measure or place limits on the degree of latitudinal differential rotation in other Sun-like stars. Early analysis of the Mount Wilson data revealed an empirical correlation between the mean level of magnetic activity and the rotation period normalized by the convective timescale [7]. The data also established a relation between the rotation rate and the period of the observed activity cycle, which generally supports a dynamo interpretation [8].

It was several decades before similar observations were conducted for a large sample of Sun-like stars in the southern sky [9]. These data confirmed the bimodal distribution in mean stellar activity *levels* first seen in the northern sample [10], but they could say nothing about activity *cycles* because the data were confined to a single epoch of observation. Another decade has passed, and there is still no time-domain survey of Ca HK emission in southern Sun-like stars. This is particularly disappointing in light of the fact that the most promising candidates for asteroseismology, which will soon allow the direct measurement of stellar convection zone depths (see Fig. 3), are stars in the southern sky—most notably the brightest solar-type stars  $\alpha$  Cen A,  $\alpha$  Cen B, and  $\beta$  Hyi which has long been studied as a “future Sun”.

Solar-like oscillations have already been detected in these stars [11, 12, 13], but tests of solar dynamo models will not be possible until their magnetic activity cycle periods are established. There is some evidence from sparse x-ray observations of  $\alpha$  Cen A that it may exhibit an activity cycle as short as 3.5 years [14], despite the fact that it is only 10% more massive than the Sun and just slightly older. A firm measurement of the cycle period in  $\alpha$  Cen A, particularly when combined with forthcoming asteroseismic data, could provide a very stringent second test of the flux-transport dynamo model.

## 2. PROJECT GOALS

We are proposing to initiate a long-term survey of Ca II H and K emission for a sample of southern Sun-like stars, establishing a southern counterpart of the Mount Wilson Ca HK survey.

During at least the first two years, we will monitor the sample with sufficient frequency (several times per week) to measure the rotation periods, and to place limits on the degree of latitudinal differential rotation. For those stars with the shortest activity cycles, we will either directly measure or provide firm upper limits on the cycle period. By comparing our observations to those from the single-epoch survey in 1996, we will also establish interesting limits on the cycle period for those stars with the slowest




Figure 3. The expected oscillation signal from the edge of the convective envelope of a  $1.1 M_{\odot}$  star with (dotted) and without (solid) overshoot (adapted from [15]). Data with this level of precision will soon emerge from space-based asteroseismology missions such as CoRoT and Kepler.

variations in activity. These initial results, particularly the analysis of  $\alpha$  Cen A, will provide a compelling justification for the continuation of the survey through external funding.

Our longer-term goal, which extends beyond the two-year period covered by our proposal, is to position NCAR as a leader in the observation and interpretation of magnetic activity cycles, complementing the role of the Mount Wilson Observatory in the north. We will constitute a direct link to facilitate the use of magnetic activity cycles and asteroseismic data to calibrate solar dynamo models, bridging a gap between the solar and stellar communities. At the very least, we plan to measure the activity cycle periods and rotational properties for a complete sample of 92 southern Sun-like stars, but we also hope eventually to upgrade the instrumentation and enlarge the sample as resources allow.

### 3. METHODOLOGY

The single-epoch survey of Henry et al. (1996) contained a total of 1016 observations of 815 individual stars with visual magnitudes between 0.0 and about 9.0, which were observed using the *RC Spec* instrument on the CTIO 1.5m telescope. The full survey was completed in 11 nights of observing time with typical integration times between 1-10 minutes, implying an average data acquisition rate of about 90 observations per night. Several sub-samples were defined, including the “Best & Brightest” (B) and “Nearby” (N) samples, which together contain 92 indi-

vidual stars with visual magnitudes between 0.0 and 7.9, and B–V colors that are approximately solar (see Table 1). All three of the most promising southern asteroseismic targets ( $\alpha$  Cen A,  $\alpha$  Cen B, and  $\beta$  Hyi) are included in this combined (B+N) sample. Since it is a full magnitude brighter than the complete sample, integration times will often be shorter—but the overhead of slewing the telescope between more widely dispersed targets will require more time between observations. Thus, we can expect a data acquisition rate roughly comparable to that achieved during the 1996 survey.

The CTIO 1.5m telescope is now operated by a consortium of about a dozen partners, known as SMARTS<sup>1</sup> (Small and Moderate Aperture Research Telescope System). This consortium runs the telescope in queue mode, with observations collected by a trained technician and made available for download by the principal investigator. Every 4-6 weeks the technician cycles between two available instruments (*RC Spec* and a broad-band infrared imager). In practical terms, this means that *RC Spec* is available for an average of 26 weeks per year. It is important to note that SMARTS operates the *only* southern telescope run in queue mode with an aperture and instrument that are appropriate for this project. Aside from a dedicated survey telescope like the one at Mount Wilson, SMARTS is the only option that makes a time-domain survey like this one feasible.

The SMARTS consortium is currently seeking a new partner for 2007, who can contribute at the level of \$50,000 per year in return for the equivalent of 40 nights

<sup>1</sup><http://www.astro.yale.edu/smarts/>

Table 1. Characteristics of the 92 stars in the survey sample.

HD	RA(2000)	Dec(2000)	V	B-V	$\log R'_{HK}$	HD	RA(2000)	Dec(2000)	V	B-V	$\log R'_{HK}$
1273	00 16 53.9	-52 39 04	6.840	0.655		128620	14 39 36.5	-60 50 02	0.010	0.710	-5.002
1581	00 20 04.3	-64 52 29	4.226	0.576	-4.839	128621	14 39 35.1	-60 50 14	1.350	0.900	-4.923
2151	00 25 45.1	-77 15 15	2.820	0.618	-4.996	130948	14 50 15.8	+23 54 43	5.863	0.576	
3443	00 37 20.7	-24 46 02	5.572	0.715	-4.893	131156	14 51 23.4	+19 06 02	4.700	0.730	-4.323
4308	00 44 39.3	-65 38 58	6.546	0.655		131977	14 57 28.0	-21 24 56	5.723	1.024	-4.484
4628	00 48 23.0	+05 16 50	5.742	0.890	-4.854	136352	15 21 48.1	-48 19 03	5.652	0.639	
7570	01 15 11.1	-45 31 54	4.959	0.571		136466	15 22 36.5	-47 55 16	7.685	0.722	
10360	01 39 47.7	-56 11 34	5.900	0.800	-4.753	140538	15 44 01.8	+02 30 55	5.865	0.684	
10361	01 39 47.2	-56 11 44	5.800	0.860	-4.881	140901	15 47 29.1	-37 54 59	6.010	0.715	
10476	01 42 29.8	+20 16 07	5.242	0.836	-4.930	141004	15 46 26.6	+07 21 11	4.422	0.604	
10700	01 44 04.1	-15 56 15	3.495	0.727	-4.962	147513	16 24 01.3	-39 11 35	5.385	0.625	
14412	02 18 58.5	-25 56 44	6.343	0.724		147584	16 28 28.1	-70 05 04	4.900	0.555	
16160	02 36 04.9	+06 53 13	5.791	0.918	-4.847	153631	17 01 10.8	-13 34 02	7.124	0.608	
17051	02 42 33.5	-50 48 01	5.400	0.561		155885	17 15 20.9	-26 36 09	5.110	0.860	-4.569
20407	03 15 06.4	-45 39 53	6.762	0.586		155886	17 15 20.9	-26 36 09	5.070	0.850	-4.577
20766	03 17 46.2	-62 34 31	5.529	0.641	-4.646	156026	17 16 13.4	-26 32 46	6.327	1.144	-4.447
20794	03 19 55.7	-43 04 11	4.260	0.711	-4.977	156274	17 19 03.8	-46 38 10	5.330	0.770	-4.941
20807	03 18 12.8	-62 30 23	5.239	0.600	-4.787	156384	17 18 57.2	-34 59 23	5.910	1.082	-4.721
21411	03 26 11.1	-30 37 04	7.877	0.716		158614	17 30 23.8	-01 03 47	5.314	0.715	-4.972
22049	03 32 55.8	-09 27 30	3.726	0.881	-4.469	160691	17 44 08.7	-51 50 03	5.127	0.694	
26965	04 15 16.3	-07 39 10	4.426	0.820		163840	17 57 14.3	+23 59 45	6.301	0.642	
30003	04 40 17.8	-58 56 38	6.529	0.677		165185	18 06 23.7	-36 01 11	5.900	0.615	
30495	04 47 36.3	-16 56 04	5.491	0.632		165341	18 05 27.3	+02 30 00	4.026	0.860	-4.637
32778	05 02 17.1	-56 04 50	7.023	0.636		165401	18 05 37.5	+04 39 26	6.800	0.610	
37655	05 38 01.9	-42 57 49	7.434	0.600		165499	18 10 26.2	-62 00 08	5.473	0.592	
43834	06 10 14.5	-74 45 11	5.080	0.714		172051	18 38 53.4	-21 03 07	5.858	0.673	
53705	07 03 57.3	-43 36 29	5.559	0.624		177565	19 06 52.5	-37 48 38	6.154	0.705	
59468	07 27 25.5	-51 24 09	6.726	0.694		178428	19 07 57.3	+16 51 12	6.086	0.702	
63077	07 45 35.0	-34 10 21	5.363	0.589		189567	20 05 32.8	-67 19 15	6.078	0.648	
65907	07 57 46.9	-60 18 11	5.595	0.573		190067	20 02 34.2	+15 35 31	7.166	0.714	
67458	08 07 00.5	-29 24 11	6.798	0.600		190248	20 08 43.6	-66 10 55	3.554	0.751	-4.999
71334	08 25 49.5	-29 55 50	7.797	0.643		190406	20 04 06.2	+17 04 13	5.788	0.600	
73524	08 37 20.0	-40 08 52	6.534	0.598		191408	20 11 11.9	-36 06 04	5.315	0.868	-4.988
76151	08 54 17.9	-05 26 04	6.000	0.661	-4.691	194640	20 27 44.2	-30 52 04	6.612	0.724	
88742	10 13 24.7	-33 01 54	6.377	0.592		196761	20 40 11.8	-23 46 26	6.363	0.719	
98231	11 18 10.9	+31 31 45	3.786	0.606		199288	20 57 40.1	-44 07 46	6.516	0.587	
102365	11 46 31.1	-40 30 01	4.892	0.664		201091	21 06 53.9	+38 44 58	5.200	1.069	-4.300
102438	11 47 15.8	-30 17 11	6.481	0.681		202940	21 19 45.6	-26 21 10	6.558	0.737	-4.870
114260	13 09 42.5	-22 11 33	7.356	0.718		206860	21 44 31.3	+14 46 19	5.945	0.587	
114613	13 12 03.2	-37 48 11	4.849	0.693		207129	21 48 15.8	-47 18 13	5.579	0.601	
115383	13 16 46.5	+09 25 27	5.209	0.585		209100	22 03 21.7	-56 47 10	4.688	1.056	-4.559
115617	13 18 24.3	-18 18 40	4.739	0.709	-4.963	211415	22 18 15.6	-53 37 37	5.363	0.614	
120690	13 51 20.3	-24 23 25	6.435	0.703		212330	22 24 56.4	-57 47 51	5.310	0.665	
122742	14 03 32.4	+10 47 12	6.288	0.733		214953	22 42 36.9	-47 12 39	5.988	0.584	
126053	14 23 15.3	+01 14 30	6.268	0.639		216803	22 56 24.1	-31 33 56	6.482	1.094	-4.272
126525	14 27 33.0	-51 55 59	7.829	0.682		217014	22 57 28.0	+20 46 08	5.469	0.666	

of observing time. Operating in queue mode, this allocation will allow us to obtain exposures for the entire available B+N sample (around 40 stars visible on any given date) three times each week whenever *RC Spec* is on the telescope. Over time, this sampling rate has proven sufficient to determine the rotation periods for many of the stars in the Mount Wilson survey [7], and in some cases allows a measurement or limit on the degree of latitudinal differential rotation.

#### 4. DISSEMINATION

The results of our survey, including raw and calibrated time-series measurements of the Ca HK emission for the entire B+N sample, will be made available without restriction through the project web-site. Updates will be made approximately once every two months, after each epoch of observation with *RC Spec* through the SMARTS queue scheduling system, which cycles between two available instruments at 4–6 week intervals. At the end of the first two years, we will publish the initial results of the survey in a refereed journal to advertise the availability of the data to the entire community. We will also promote the inclusion of the data in all appropriate future Virtual Observatory projects.

#### ACKNOWLEDGMENTS

The National Center for Atmospheric Research is a federally funded research and development center sponsored by the National Science Foundation.

#### REFERENCES

- [1] M. Dikpati. Solar magnetic fields and the dynamo theory. *Advances in Space Research*, 35:322–328, 2005.
- [2] M. Dikpati, G. de Toma, and P. A. Gilman. Predicting the strength of solar cycle 24 using a flux-transport dynamo-based tool. *Geophys. Res. Lett.*, 33:5102–5105, March 2006.
- [3] S. L. Baliunas, R. A. Donahue, W. H. Soon, J. H. Horne, J. Frazer, L. Woodard-Eklund, M. Bradford, L. M. Rao, O. C. Wilson, Q. Zhang, W. Bennett, J. Briggs, S. M. Carroll, D. K. Duncan, D. Figueroa, H. H. Lanning, T. Misch, J. Mueller, R. W. Noyes, D. Poppe, A. C. Porter, C. R. Robinson, J. Russell, J. C. Shelton, T. Soyumer, A. H. Vaughan, and J. H. Whitney. Chromospheric variations in main-sequence stars. *ApJ*, 438:269–287, January 1995.
- [4] R. B. Leighton. Observations of Solar Magnetic Fields in Plage Regions. *ApJ*, 130:366–380, September 1959.
- [5] O. C. Wilson. Chromospheric variations in main-sequence stars. *ApJ*, 226:379–396, December 1978.
- [6] S. Baliunas, P. Frick, D. Moss, E. Popova, D. Sokoloff, and W. Soon. Anharmonic and standing dynamo waves: theory and observation of stellar magnetic activity. *MNRAS*, 365:181–190, January 2006.
- [7] R. W. Noyes, L. W. Hartmann, S. L. Baliunas, D. K. Duncan, and A. H. Vaughan. Rotation, convection, and magnetic activity in lower main-sequence stars. *ApJ*, 279:763–777, April 1984.
- [8] R. W. Noyes, N. O. Weiss, and A. H. Vaughan. The relation between stellar rotation rate and activity cycle periods. *ApJ*, 287:769–773, December 1984.
- [9] T. J. Henry, D. R. Soderblom, R. A. Donahue, and S. L. Baliunas. A Survey of Ca II H and K Chromospheric Emission in Southern Solar-Type Stars. *AJ*, 111:439–465, January 1996.
- [10] A. H. Vaughan and G. W. Preston. A survey of chromospheric CA II H and K emission in field stars of the solar neighborhood. *PASP*, 92:385–391, August 1980.
- [11] T. R. Bedding, H. Kjeldsen, R. P. Butler, C. McCarthy, G. W. Marcy, S. J. O’Toole, C. G. Tinney, and J. T. Wright. Oscillation Frequencies and Mode Lifetimes in  $\alpha$  Centauri A. *ApJ*, 614:380–385, October 2004.
- [12] H. Kjeldsen, T. R. Bedding, R. P. Butler, J. Christensen-Dalsgaard, L. L. Kiss, C. McCarthy, G. W. Marcy, C. G. Tinney, and J. T. Wright. Solar-like Oscillations in  $\alpha$  Centauri B. *ApJ*, 635:1281–1290, December 2005.
- [13] T. R. Bedding, R. P. Butler, H. Kjeldsen, I. K. Baldry, S. J. O’Toole, C. G. Tinney, G. W. Marcy, F. Kienzle, and F. Carrier. Evidence for Solar-like Oscillations in  $\beta$  Hydri. *ApJ*, 549:L105–L108, March 2001.
- [14] J. Robrade, J. H. M. M. Schmitt, and F. Favata. X-rays from  $\alpha$  Centauri - The darkening of the solar twin. *A&A*, 442:315–321, October 2005.
- [15] M. J. P. F. G. Monteiro, J. Christensen-Dalsgaard, and M. J. Thompson. Seismic study of stellar convective regions: the base of the convective envelope in low-mass stars. *MNRAS*, 316:165–172, July 2000.