

## THE ORIGIN AND EVOLUTION OF THE MASS-METALLICITY RELATIONSHIP FOR GALAXIES: RESULTS FROM COSMOLOGICAL N-BODY SIMULATIONS

A. M. BROOKS<sup>1,2</sup>, F. GOVERNATO<sup>1</sup>, C. M. BOOTH<sup>3</sup>, B. WILLMAN<sup>4,5</sup>, J. P. GARDNER<sup>6</sup>, J. WADSLEY<sup>7</sup>, G. STINSON<sup>1</sup>, T. QUINN<sup>1</sup>

*Draft version March 13, 2018*

### ABSTRACT

We examine the origin and evolution of the mass-metallicity relationship (MZR,  $M_*-Z$ ) for galaxies using high resolution cosmological SPH + N-Body simulations that include a physically motivated description of supernovae feedback and subsequent metal enrichment. We discriminate between two sources that may contribute to the origin of the MZR: 1) metal and baryon loss due to gas outflow, or 2) inefficient star formation at the lowest galaxy masses. Our simulated galaxies reproduce the observed MZR in shape and normalization both at  $z=0$  and  $z=2$ . We find that baryon loss occurs due to UV heating before star formation turns on in galaxies with  $M_{bar} < 10^8 M_\odot$ , but that some gas loss due to supernovae induced winds is required to subsequently reproduce the low effective chemical yield observed in low mass galaxies. Despite this, we show that low star formation efficiencies, regulated by supernovae feedback, are primarily responsible for the lower metallicities of low mass galaxies and the overall  $M_*-Z$  trend. We find that the shape of the MZR is relatively constant with redshift, but that its normalization increases with time. Simulations with no energy feedback from supernovae overproduce metals at low galaxy masses by rapidly transforming a large fraction of their gas into stars. Despite the fact that our low mass galaxies have lost a majority of their baryons, they are still the most gas rich objects in our simulations due to their low star formation efficiencies.

*Subject headings:* galaxies: evolution — galaxies: formation — methods: N-Body simulations

### 1. INTRODUCTION

The observed trend that the metallicity of nearby galaxies decreases with galaxy mass has been well established in the literature (e.g., Tremonti et al. 2004; Lee et al. 2006). This trend has also recently been observed at high redshifts (Savaglio et al. 2005; Erb et al. 2006, hereafter S05 and E06, respectively). The gas metallicity of a galaxy depends on reprocessed material from its stars and subsequent inflow and outflow of gas. The mass-metallicity relationship (MZR) thus provides insight into two physical mechanisms important to the evolution of galaxies: star formation efficiency and gas inflow/outflow.

In a “closed box” system (no inflow or outflow of material), metallicity  $Z$  is given by

$$Z = y \ln f_{gas}^{-1}, \quad (1)$$

where  $y$  is the nucleosynthetic metal yield produced by stars and  $f_{gas}$  is the gas mass fraction ( $M_{gas}/[M_{gas} + M_{stars}]$ ) (e.g., Tinsley 1980). In general, low mass galaxies are observed to have higher gas fractions than high mass galaxies (e.g., Geha et al. 2006; West 2005). This observation can easily explain the origin of the

$M_*-Z$  relation (MZR) in the context of the closed box model. In this scenario, low mass galaxies are gas rich because they are inefficient at turning gas into stars, likely due to their low surface densities (Kennicutt 1998; Martin & Kennicutt 2001; Verde et al. 2002; Dalcanton et al. 2004).

Alternatively, SNe feedback might expel gas in galactic winds. In this scenario the loss of metals becomes progressively more important at lower masses due to shallower halo potential wells (for a review see Veilleux et al. 2005) and leads to the observed  $M_*-Z$  trend. However, the importance of supernova feedback as a function of galaxy mass has been investigated by various groups, with results that are still debated (Veilleux et al. 2005; Geha et al. 2006; Dalcanton 2006, E06).

Recently, studies have investigated how the MZR evolves with cosmic time (S05, E06, Davé et al. 2006; De Rossi et al. 2006; Kobayashi et al. 2006; Tassis et al. 2006). Fully cosmological numerical experiments are one of the best tools to infer the relative importance of star formation (SF) and feedback processes in a realistic setting (e.g., Okamoto et al. 2005; Robertson et al. 2006; Governato et al. 2006, hereafter G06). In this Letter, we investigate the origin and evolution of the MZR in a concordance  $\Lambda$ CDM cosmological setting with N-body + SPH simulations which include a physically motivated treatment of SF and SN feedback. These simulations allow us to study the formation history of galaxies over range of masses, and thus quantify the importance of feedback and blowout versus low SF efficiencies over their lifetimes.

### 2. THE SIMULATIONS

The simulations used in this study are the culmination of an effort to create realistic disk galaxies. As

arXiv:astro-ph/0609620v3 22 Dec 2006

<sup>1</sup> Astronomy Department, University of Washington, Box 351580, Seattle, WA, 98195-1580

<sup>2</sup> e-mail address: abrooks@astro.washington.edu

<sup>3</sup> Department of Physics, Institute for Computational Cosmology, University of Durham, South Road, Durham, DH1 3LE, UK

<sup>4</sup> Harvard-Smithsonian Center for Astrophysics, 60 Garden Street, Cambridge, MA, 02138

<sup>5</sup> Clay Fellow

<sup>6</sup> Department of Physics & Astronomy, University of Pittsburgh, 100 Allen Hall, 3941 O’Hara Street, Pittsburgh, PA, 15260

<sup>7</sup> Department of Physics and Astronomy, McMaster University, Hamilton, Ontario, L8S 4M1, Canada

discussed in G06, at  $z=0$  disk galaxies in our simulations fall on the Tully-Fisher (TF) and baryonic TF relations (Giovanelli et al. 1997; McGaugh 2005) and the age-mass relation (MacArthur et al. 2004). This is an important improvement over previous works, and is made possible due to the inclusion of a simple but physically motivated recipe (full details in Stinson et al. 2006) to describe SF and the effects of subsequent SNe feedback. Our adopted SN feedback and cosmic UV background (Haardt & Madau 1996) also drastically reduce the number of galaxy satellites containing a significant stellar population, making many of them “dark” (Quinn et al. 1996; Moore et al. 1999). Thus, the SN and SF efficiency parameters were adopted as in G06, with an IMF from Kroupa et al. (1993), with no additional tuning of the parameters to affect metals. Metal enrichment from both SN Ia&II is followed based on the prescription of Raiteri et al. (1996), who adopt yields from Thielemann et al. (1986) for SN Ia, and Weaver & Woosley (1993) for SN II.

We selected four “field” regions ( $d\rho/\rho \sim 0.1$ ) from a low resolution, dark matter (DM) simulation run in GASOLINE (Wadsley et al. 2004) using a concordance, flat,  $\Lambda$ -dominated cosmology:  $\Omega_0 = 0.3$ ,  $\Lambda=0.7$ ,  $h = 0.7$ ,  $\sigma_8 = 0.9$ , shape parameter  $\Gamma = 0.21$ , and  $\Omega_b = 0.039$  (Perlmutter et al. 1997; Efstathiou et al. 2002). These field regions were resimulated at higher resolution using the volume renormalization technique (Katz & White 1993), which allows us to resolve fine structure while capturing the effect of large scale torques. The field regions are centered around galaxies with peak rotation velocity  $V_{rot} \sim 150$  and 275 km/s, with each high resolution region having a comoving volume of  $\sim 100$ – $1000$  Mpc<sup>3</sup>. Two of the main galaxies are described in detail in G06, but for this study they have been rerun at higher resolution. At  $z=0$ , these runs yield a sample of 31 high resolution disk and dwarf galaxies over a range of peak velocities  $30 \text{ km/s} < V_{rot} < 275 \text{ km/s}$ , or total halo masses from  $3.4 \times 10^9 M_\odot$  to  $1.1 \times 10^{12} M_\odot$ . Seven of these 31 galaxies are satellites of the primaries. Galaxies and their parent halos were identified using AHF<sup>8</sup> (Knebe et al. 2001; Gill et al. 2004). Our highest (lowest) resolution runs have particle masses of  $7.6 \times 10^5 M_\odot$  ( $4.9 \times 10^6 M_\odot$ ),  $1.3 \times 10^5 M_\odot$  ( $8.5 \times 10^5 M_\odot$ ), and  $3.8 \times 10^4 M_\odot$  ( $2.5 \times 10^5 M_\odot$ ) for DM, gas, and stars, respectively, and a force resolution of 0.15–0.3 kpc.

### 3. THE ORIGIN OF THE MZR

The MZR for 31 high resolution galaxies in our simulations is shown in Fig. 1, at both  $z=0$  and  $z=2$ . The solid curved line is the empirical fit for  $> 53,000$  galaxies in SDSS from Tremonti et al. (2004, hereafter T04). It has been shifted down by 0.26 dex as found by E06 to correct for the fact that the method used by T04 is known to systematically find O abundances up to 0.3 dex larger than other methods (T04, S05, E06 Lee et al. 2006). The galaxies in our sample at  $z=0$  match the empirical fit of T04 extremely well, reproducing the observed trend over almost 4 orders of magnitude in stellar mass. We note that our lowest mass galaxies, however, have a lower O

abundance than that found for dwarf irregular galaxies by Lee et al. (2006).

Fig. 1 also shows that the MZR for our simulated galaxies at  $z=2$  (open diamonds) is in excellent agreement with high redshift data. The error bars overplotted on Fig. 1 display the observational  $z=2$  results from E06. We have identified in the simulations the  $z=0$  galaxies corresponding to those plotted at  $z=2$ . The dotted lines in Fig. 1 connect a few of these galaxies, which are representative of how all the galaxies have evolved in the  $M_*$ - $Z$  plane with time.

To make a meaningful comparison with the T04 sample, Fig. 1 uses only cool gas particles (with temperature  $< 40,000$  K) within the radius of each galaxy that encloses  $2/3$  of the stellar mass. The median “observed” diameter for our galaxies is 5 kpc, similar to that found by T04. At  $z=2$  we included all cold gas in each galaxy. Even if we include all cool gas at  $z=0$ , the  $M_*$ - $Z$  trend is unchanged up to  $10^{10} M_\odot$ , showing that the evolution seen in Fig. 1 is not an effect of our measurements.

A few runs were repeated at lower resolutions ( $1/2$  and  $1/20$  of the mass resolution). Low resolution runs undergo less SF, thus decreasing both  $M_*$  and the total amount of O produced. Our results from each run begin to significantly diverge from each other ( $\Delta M_* > 20\%$ ,  $\Delta 12+\log(\text{O}/\text{H}) > 0.2$  dex) if the number of DM particles in a halo drops below a few thousand and/or mass and force resolution degrade too much. Note that this low resolution effect can lead to an artificially steeper  $M_*$ - $Z$  trend, making it essential that results converge. Informed by these resolution tests, we have only included galaxies with more than 3500 DM particles in our sample (at high and low redshift), ensuring that their stellar and metal content have converged.

Having shown that our simulations reproduce the observed MZR at both  $z=0$  and  $z=2$ , we will next examine the origin of this relation. We focus on the importance of preferential metal loss versus low SF efficiency at low masses in producing the MZR.

#### 3.1. The Role of Mass Loss

An observable quantity commonly used to investigate mass loss from galaxies is the effective yield,  $y_{eff}$ . From Eq. 1, we can define the effective yield,

$$y_{eff} = Z / (\ln f_{gas}^{-1}). \quad (2)$$

In a closed box system,  $y_{eff}$  is equal to the true stellar nucleosynthetic yield, but it can be inferred from Eq. 2 that  $y_{eff}$  may deviate from the true yield due to inflow or outflow of gas. The top panel of Fig. 2 shows  $y_{eff}$  for our galaxies, derived again using inner cold gas abundances, versus the total baryonic mass of our galaxies at  $z=0$  (black circles). The effective yields reproduce the observed trend from T04 (solid line) over most of our mass range. Our two lowest mass galaxies have  $y_{eff}$  below that found by Lee et al. (2006), likely because these two have undergone additional tidal stripping of gas.

The black circles in the bottom panel of Fig. 2 show the baryonic to dark mass ratio (B/DM) of our galaxies, normalized by the cosmic ratio (the seven satellite galaxies have been excluded from this panel). In our simulations, large deviations below  $B/\text{DM} = 1$  can only be produced by mass loss. Our lowest mass galaxies have lost up to

<sup>8</sup> AMIGA’s Halo Finder, available for download at <http://www.aip.de/People/aknebe/AMIGA>

98% of their baryonic mass.

Using the halos defined by AHF (our halo finder), we identify a given halo and its gas particles back to  $z=3$ . Once all the gas particles that have ever belonged to a galaxy back to  $z=3$  are identified, those particles that are no longer within the virial radius at  $z=0$  are considered to be gas “lost” from the galaxy. The grey points in both panels of Fig. 2 represent this information, for all gas (remaining + lost).

When we include all of the gas lost since  $z = 3$ , we find that our low mass galaxies still have B/DM that fall significantly below the cosmic ratio. The bottom panel of Fig. 2 show that our lowest mass galaxies have lost 50 - 90% of their baryons prior to  $z = 3$ . We verify that a UV only simulation (SNe feedback turned off) removes up to 80% of baryons from our lowest mass galaxies, but that these galaxies evolve as closed boxes once SF commences. We thus find that SNe induced mass loss can account for up to 20% of baryon loss from galaxies with  $M_{bar} < 1 \times 10^8 M_{\odot}$ . Additional gas loss can occur via tidal stripping.

Thus, Fig. 2 clearly demonstrates that gas has been preferentially lost from our lowest mass galaxies, and that this gas loss is responsible for the observed  $y_{eff}$ - $M_{bar}$  trend. However, when all gas (remaining + lost) is used to rederive the MZR (Fig. 1), the observed trend that lower mass galaxies have lower metallicities still exists, and it cannot be mass loss alone that causes the MZR.

### 3.2. Star Formation Efficiencies

As explained above, the mass loss experienced by our lowest mass galaxies cannot account for the overall  $M_*$ - $Z$  trend seen in Fig. 1. We now investigate whether SF efficiencies can explain the origin of this trend. If lower mass galaxies convert gas into stars less efficiently than higher mass galaxies, then they will require a much longer period of time to enrich the remaining gas to the same metallicity as higher mass galaxies. This effect can lead to the observed MZR, with or without mass loss as an additional contributor.

We verified that all of our galaxies follow the Schmidt-Kennicutt law for SF (Kennicutt 1998; Martin & Kennicutt 2001). Our galaxies span a factor of 250 in gas surface densities, yielding a SFR that varies by over three decades from our lowest to highest mass galaxies. Our lowest mass galaxies are thus incredibly inefficient at SF compared to our high mass galaxies. The consumption rate ( $SFR/M_{coldgas}$ ) of our lowest mass galaxies is about two decades lower than our highest mass galaxies. That is, while our highest mass galaxies will use up all their gas in 1 Gyr or less, our lowest mass galaxies will take up to 60 Gyr if they continue SF at their current rate.

Fig. 3 shows the gas mass fraction,  $f_{gas}$ , versus rotational velocity for our galaxies (satellite galaxies excluded). As found by recent observations of isolated galaxies (e.g. West 2005; Geha et al. 2006), lower mass galaxies have increasing gas fractions, implying that lower mass galaxies have been less efficient at making stars. Note that while the lowest mass galaxies in our sample have lost more than 90% of their baryons (Fig. 2), they remain gas rich due to their low SF efficiencies (with the exception of two that have been tidally stripped of

much of their gas).

We reran our simulations without SNe feedback. In our models, without this feedback the MZR is flat because low mass galaxies experience unrealistically high SF rates, resulting in an overproduction of metals. This difference with the runs that do include SNe demonstrates that energy injection by SNe into the ISM regulates SF efficiency, independent of whether SNe also lead to mass loss. SNe feedback thus indirectly affects the shape of the MZR by modulating the gas surface densities, and hence the SFR, with galaxy mass.

## 4. THE EVOLUTION OF THE MZR

Fig. 3 shows that our simulated galaxies at  $z=2$  are less evolved than their  $z=0$  counterparts, with  $z=2$  galaxies being much more gas rich at a given rotational velocity than at  $z=0$ . By  $z=0$ , they have consumed more of their gas, increasing both their stellar mass and O abundance; hence the evolution of the MZR as shown in Fig. 1.

Both S05 and E06 examined the MZR at  $z \sim 0.7$  and  $z \sim 2$ , respectively, and drew conclusions similar to ours. As pointed out by S05 and Pettini (2006), at a given metallicity the high  $z$  data are about an order of magnitude more massive than the  $z=0$  trend (Fig. 1). This suggests that more massive galaxies enrich at a faster rate than lower mass galaxies. This is supported by our results in section 3.2. Both S05 and E06 conclude that the evolution of the MZR is best described by a model in which low mass galaxies are inefficient at SF relative to high mass galaxies for a similar formation epoch. Thus the period of SF is lengthened in lower mass galaxies and they remain more gas rich at a given redshift.

We find that a shift downward by 0.3 dex in  $\Delta \log(O/H)$  from fit shown in Fig. 1 is an excellent fit to our  $z=2$  data, as was also found by E06. A shift in  $\Delta \log(M_*)$  is much too steep to match our simulations. Thus, we find that the slope of the MZR remains relatively constant with redshift, but the normalization increases with time.

## 5. CONCLUSIONS

We have presented results from high resolution simulations of field galaxies in a  $\Lambda$ CDM context. The simulations are carried to  $z=0$  and include SF and a physically motivated description of the effects of SNe feedback and metal enrichment. We have shown in previous work (G06) that this approach leads to disk galaxies that fall on the TF and baryonic TF relations and reproduce the age-mass relation of disk galaxies and abundance of galaxy satellites. Here we present a natural extension of this analysis, showing that the simulated galaxies are also in excellent agreement with the observed MZR both at  $z=0$  and  $z=2$ .

Simulated galaxies match both the slope and normalization of the MZR over four orders of magnitude in stellar mass. We find that the slope of the MZR is roughly constant with redshift, but that its normalization increases with time. Such evolution agrees with the high- $z$  observations of E06, and with early theoretical predictions (Davé et al. 2006; De Rossi et al. 2006). Furthermore, the mass scales of the turnovers in our simulated  $Z$ ,  $y_{eff}$ , B/DM, and  $f_{gas}$  versus mass relations are in good agreement with a range of previous observational and theoretical studies that find critical transitions at  $\sim 120$  km/s (e.g., Dalcanton et al. 2004; Kereš et al.

2005; Dekel & Birnboim 2006).

Galaxies with  $M_{bar} < 10^8 M_{\odot}$  ( $V_{rot} \sim 50$  km/s), lose a significant amount of their baryons due to combined heating from the cosmic UV field and SN feedback. Baryon loss due to SN feedback results in a  $y_{eff}-M_{bar}$  trend in agreement with the results from the SDSS (T04). However, including gas that has been lost from galaxies does not alter the O abundances significantly, and does not change the overall shape of the MZR, confirming that the lower O abundances in our low mass galaxies are primarily due to their low SF efficiencies rather than directly to blowouts. Despite strong baryon loss from their halos, we find that our cold gas fractions at low galaxy masses are in qualitative agreement with recent observational samples (Geha et al. 2006).

SN feedback plays a crucial role in lowering the SF efficiency in low mass galaxies and originating the turnover in the MZR at the observed scales. Without energy injection from SNe to regulate SF, gas that remains in galaxies rapidly cools, forms stars, and increases the O abundance, even by  $z=2$  (see G06 for additional shortcomings of the no feedback case). With feedback from

SNe turned off, small galaxies produce too many metals too early at the expense of their cold gas content, producing  $M_{*}-Z$  and  $y_{eff}-M_{bar}$  relations too flat compared to observations, in agreement with some early theoretical predictions (De Rossi et al. 2006; Dalcanton 2006). SN feedback prevents low mass galaxies from enriching their gas to a high metallicity at all redshifts. In our runs, star formation in small field galaxies occurs over extended periods of time until the present, in agreement with observations (Dolphin et al. 2005).

We would like to thank C. Brook, H. Lee, E. Skillman, K. Venn, J. Dalcanton, A. Kravtsov, and A. West for helpful conversations during this project. Support for this work was provided by the Spitzer Space Telescope Theoretical Research Program. FG acknowledges support from NSF grant AST-0098557. Simulations were run at the Pittsburgh Supercomputing Center, SDSC, and Cineca. AB, GS, and TQ were supported by NSF ITR grant PHY-0205413.

#### REFERENCES

- Dalcanton, J. J. 2006, astro-ph/0608590  
Dalcanton, J. J., Yoachim, P., & Bernstein, R. A. 2004, ApJ, 608, 189  
Davé, R., Finlator, K., & Oppenheimer, B. D. 2006, astro-ph/0608537  
De Rossi, M. E., Tissera, P. B., & Scannapieco, C. 2006, astro-ph/0609243  
Dekel, A., & Birnboim, Y. 2006, MNRAS, 368, 2  
Dolphin, A. E., Weisz, D. R., Skillman, E. D., & Holtzman, J. A. 2005, astro-ph/0506430  
Efstathiou, G. et al. 2002, MNRAS, 330, L29  
Erb, D. K., Shapley, A. E., Pettini, M., Steidel, C. C., Reddy, N. A., & Adelberger, K. L. 2006, ApJ, 644, 813  
Geha, M., Blanton, M. R., Masjedi, M., & West, A. A. 2006, astro-ph/0608295  
Gill, S. P. D., Knebe, A., & Gibson, B. K. 2004, MNRAS, 351, 399  
Giovannelli, R., Haynes, M. P., Herter, T., Vogt, N. P., da Costa, L. N., Freudling, W., Salzer, J. J., & Wegner, G. 1997, AJ, 113, 53  
Governato, F., Willman, B., Mayer, L., Brooks, A., Stinson, G., Valenzuela, O., Wadsley, J., & Quinn, T. 2006, MNRAS, in press, astro-ph/0602351  
Haardt, F., & Madau, P. 1996, ApJ, 461, 20  
Katz, N., & White, S. D. M. 1993, ApJ, 412, 455  
Kennicutt, Jr., R. C. 1998, ApJ, 498, 541  
Kereš, D., Katz, N., Weinberg, D. H., & Davé, R. 2005, MNRAS, 363, 2  
Knebe, A., Green, A., & Binney, J. 2001, MNRAS, 325, 845  
Kobayashi, C., Springel, V., & White, S. D. M. 2006, astro-ph/0604107  
Kroupa, P., Tout, C. A., & Gilmore, G. 1993, MNRAS, 262, 545  
Lee, H., Skillman, E. D., Cannon, J. M., Jackson, D. C., Gehrz, R. D., Polomski, E. F., & Woodward, C. E. 2006, ApJ, 647, 970  
MacArthur, L. A., Courteau, S., Bell, E., & Holtzman, J. A. 2004, ApJS, 152, 175  
Martin, C. L., & Kennicutt, Jr., R. C. 2001, ApJ, 555, 301  
McGaugh, S. S. 2005, ApJ, 632, 859  
Moore, B., Ghigna, S., Governato, F., Lake, G., Quinn, T., Stadel, J., & Tozzi, P. 1999, ApJ, 524, L19  
Okamoto, T., Eke, V. R., Frenk, C. S., & Jenkins, A. 2005, MNRAS, 363, 1299  
Perlmutter, S. et al. 1997, ApJ, 483, 565  
Pettini, M. 2006, astro-ph/0603066  
Quinn, T. and Katz, N., & Efstathiou, G. 1996, MNRAS, 278, L49  
Raiteri, C. M., Villata, M., & Navarro, J. F. 1996, A&A, 315, 105  
Robertson, B., Bullock, J. S., Cox, T. J., Di Matteo, T., Hernquist, L., Springel, V., & Yoshida, N. 2006, ApJ, 645, 986  
Savaglio, S. et al. 2005, ApJ, 635, 260  
Stinson, G., Seth, A., Katz, N., Wadsley, J., Governato, F., & Quinn, T. 2006, MNRAS, in press, astro-ph/0602350  
Tassis, K., Kravtsov, A. V., & Gnedin, N. Y. 2006, astro-ph/0609763  
Thielemann, F.-K., Nomoto, K., & Yokoi, K. 1986, A&A, 158, 17  
Tinsley, B. M. 1980, Fundamentals of Cosmic Physics, 5, 287  
Tremonti, C. A. et al. 2004, ApJ, 613, 898  
Veilleux, S., Cecil, G., & Bland-Hawthorn, J. 2005, ARA&A, 43, 769  
Verde, L., Oh, S. P., & Jimenez, R. 2002, MNRAS, 336, 541  
Wadsley, J. W., Stadel, J., & Quinn, T. 2004, New Astronomy, 9, 137  
Weaver, T. A., & Woosley, S. E. 1993, Phys. Rep., 227, 65  
West, A. A. 2005, Ph.D. Thesis




FIG. 1.— The mass-metallicity relation of our simulated galaxies at  $z=0$  (solid points) and  $z=2$  (open diamonds). The solid curved line is the observational fit to  $>53,000$  galaxies in SDSS from T04, shifted down by  $-0.26$  dex as found by E06. Error bars show the observational mass-metallicity relation at  $z=2$  from E06. Dotted lines connect some of the  $z=0$  galaxies to their progenitors at  $z=2$ , showing how galaxies evolve in the  $M_*$ -Z plane with time.




FIG. 2.— *Top Panel:* Effective yield vs total baryonic mass for our galaxies at  $z=0$  (black circles). The solid curved line is the observational fit to  $>53,000$  galaxies from T04. *Bottom Panel:* Baryonic to dark matter mass ratio (relative to the cosmic abundance) vs total baryonic mass. Satellite galaxies excluded. In both panels, grey circles show what the value would be if no gas loss from the galaxies had occurred.




FIG. 3.— The gas mass fraction vs  $V_{rot}$  for our galaxies at  $z=0$  and  $z=2$  (satellite galaxies excluded). Faint dotted lines connect a few of the galaxies at  $z=2$  to their evolved counterpart at  $z=0$ .