

DEEP *HST* PHOTOMETRY OF NGC 6388: AGE AND HORIZONTAL BRANCH LUMINOSITY*

M. CATELAN¹, PETER B. STETSON^{2,3,4}, BARTON J. PRITZL⁵, HORACE A. SMITH⁶, KAREN KINEMUCHI⁷, ANDREW C. LAYDEN⁸, ALLEN V. SWEIGART⁹, AND R. M. RICH¹⁰

ApJ Letters, in press

ABSTRACT

Using the *Hubble Space Telescope*, we have obtained the first color-magnitude diagram (CMD) to reach the main-sequence turnoff of the Galactic globular cluster NGC 6388. From a comparison between the cluster CMD and 47 Tucanae's, we find that the bulk of the stars in these two clusters have nearly the same age and chemical composition. On the other hand, our results indicate that the blue horizontal branch and RR Lyrae components in NGC 6388 are intrinsically overluminous, which must be due to one or more, still undetermined, non-canonical second parameter(s) affecting a relatively minor fraction of the stars in NGC 6388.

Subject headings: stars: horizontal-branch — stars: variables: other — globular clusters: individual (47 Tucanae, NGC 6388, NGC 6441) — globular clusters: general

1. INTRODUCTION

NGC 6388 and NGC 6441 are two of the most intriguing Galactic globular clusters (GC's). The integrated-light study by Rich, Minniti, & Liebert (1993) revealed a strong far-UV flux for these metal-rich ($[Fe/H] \simeq -0.60$ and -0.53 , respectively; Harris 1996) bulge GC's. The far-UV flux in resolved GC's is dominated by hot horizontal branch (HB) stars (e.g., Dorman, O'Connell, & Rood 1995), especially when rare UV-bright post-asymptotic giant branch stars are not present. Accordingly, the most likely explanation for the far-UV flux in NGC 6388 and NGC 6441 was immediately recognized to be hot HB stars. However, while the UV-upturn phenomenon in elliptical galaxies is often attributed to blue HB stars (e.g., Dorman et al. 1995), no resolved metal-rich GC had been known with a blue HB morphology. The tendency for metal-rich GC's to have red HB's while metal-poor GC's have predominantly blue HB's reflects the "first parameter" of HB morphology. As a consequence, the presence of blue HB stars in NGC 6388 and NGC 6441 would represent an example of the so-called "second-parameter (2ndP) phenomenon."

*BASED ON OBSERVATIONS WITH THE NASA/ESA *HUBBLE SPACE TELESCOPE*, OBTAINED AT THE SPACE TELESCOPE SCIENCE INSTITUTE, WHICH IS OPERATED BY THE ASSOCIATION OF UNIVERSITIES FOR RESEARCH IN ASTRONOMY (AURA), INC., UNDER NASA CONTRACT NAS 5-26555, AND ON OBSERVATIONS RETRIEVED FROM THE ESO ST-ECF ARCHIVE.

¹ Pontificia Universidad Católica de Chile, Departamento de Astronomía y Astrofísica, Av. Vicuña Mackenna 4860, 782-0436 Macul, Santiago, Chile; e-mail: mcatelan@astro.puc.cl

² Dominion Astrophysical Observatory, Herzberg Institute of Astrophysics, National Research Council, 5071 West Saanich Road, Victoria, BC V9E 2E7, Canada; e-mail: Peter.Stetson@nrc.gc.ca

³ Guest Investigator of the UK Astronomy Data Centre.

⁴ Guest User, Canadian Astronomy Data Centre, which is operated by the Herzberg Institute of Astrophysics, National Research Council of Canada.

⁵ Macalester College, 1600 Grand Avenue, Saint Paul, MN 55105

⁶ Dept. of Physics and Astronomy, Michigan State University, East Lansing, MI 48824

⁷ Dept. of Physics and Astronomy, University of Wyoming, Laramie, WY 82071

⁸ Department of Physics and Astronomy, Bowling Green State University, 104 Overman Hall, Bowling Green, OH 43403

⁹ NASA Goddard Space Flight Center, Exploration of the Universe Division, Code 667, Greenbelt, MD 20771

¹⁰ Division of Astronomy, Department of Physics and Astronomy, UCLA, Los Angeles, CA 90095

The presence of blue HB stars extending almost as faint in V as the turnoff (TO) point in both NGC 6388 and NGC 6441 was confirmed by Rich et al. (1997), who presented *Hubble Space Telescope* (*HST*) photometry for both these GC's from the survey by Piotto et al. (2002). A remarkable feature of the published diagrams is the presence of a *strongly sloped* HB at colors where other GC's have a much more nearly "horizontal" HB. As emphasized by Sweigart & Catelan (1998, hereafter SC98), such a sloped HB cannot be simply the result of an older age or of enhanced mass loss along the red giant branch (RGB): while these are able to move a star horizontally along the HB, neither is able to increase the luminosity of a blue HB star compared to the red HB or RR Lyrae stars. Likewise, while strong differential reddening might explain the sloping HB of a red HB cluster, it obviously cannot produce RR Lyrae and blue HB stars. SC98 conclude therefore that *non-canonical* 2ndP candidates must be at play in NGC 6388 and NGC 6441.

However, these conclusions were challenged by Raimondo et al. (2002, hereafter R02), who computed models with non-standard values of the chemical abundance and mixing length parameter. Some of their models did reveal sloped HB's, but only as a consequence of an anomalously *faint red HB* (in V), together with a blue HB having a V -band luminosity consistent with the canonical models (see, e.g., their Fig. 2).

Additional insights are provided by stellar variability and spectroscopic studies. Silbermann et al. (1994), Layden et al. (1999), Pritzl et al. (2000, 2001, 2002, 2003), and Corwin et al. (2006) have shown that the RR Lyrae variable stars in these GC's, which occupy the normally "horizontal" part of the HB, have much longer periods than field RR Lyrae of similar metallicity, thus strongly suggesting that they are intrinsically more luminous (SC98). Moreover, theoretical calculations by Pritzl et al. (2002) have shown that, contrary to the suggestions by Ree et al. (2002), the RR Lyrae components in both clusters cannot be explained in terms of evolution away from a position on the blue HB—and neither can the sloping nature of the HB be reproduced in this way. On the other hand, the first spectroscopic measurements of the gravities of blue HB stars in both NGC 6388 and NGC 6441 (Moehler, Sweigart, & Catelan 1999) revealed surface gravities that are higher than predicted by even the canonical models, thus arguing against an anomalously bright blue HB + RR




FIG. 1.— The first deep CMD of NGC 6388 clearly revealing the TO point of the cluster (*red \times symbols*) is overplotted on the 47 Tuc CMD (*plus signs*) in the V , $B-I$ plane. To produce this plot, the 47 Tuc CMD was shifted by +3.2 in V and by +0.85 in $B-I$, so as to align their red HB components.

Lyrae component in these clusters. However, a recent re-assessment of the spectroscopic gravities of blue HB stars in NGC 6388 by Moehler & Sweigart (2006) indicates that the actual gravities should, in fact, be lower than the canonical values. It appears that the 1999 values must have been in error by a substantial amount, probably due to unresolved blends in the crowded inner regions of these massive ($M_V \sim -9.5$; Harris 1996) GC's.

In an effort to shed light on this puzzling situation, we have made use of the data obtained for NGC 6388 in the course of our snapshot *HST* program to study stellar variability in the cluster, and also of archival data, to produce its deepest-ever color-magnitude diagram (CMD). In §2 we describe this dataset and reduction procedures. In §3 we compare our CMD with 47 Tucanae's (NGC 104). We close in §4 by discussing the implications of our results for our understanding of the origin of the peculiar HB morphology of NGC 6388.

2. OBSERVATIONAL DATA AND REDUCTION PROCEDURES

The NGC 6388 data used in this paper were obtained under *HST* program SNAP-9821 (PI B. J. Pritzl), which used the Wide-Field Channel of the *Advanced Camera for Surveys* (ACS) to obtain six B (F435W), V (F555W), I (F814W) exposure triptychs on separate dates ranging from 2003 October to 2004 June. In addition, we have retrieved data from the *HST* Archives, as obtained under GO-9835 (PI G. Drukier). These consist of 12 V and 17 I exposures obtained with the High-Resolution Channel of ACS on 2003 October 30.

The results employed here for 47 Tuc were obtained from the ground-based imagery used by Stetson to define his secondary photometric standards. They consist of (125, 136, 94) images in (B , V , I) from 12 distinct observing runs; these photometric indices should be on the system of Landolt (1992) to well under 0.01 mag. Stetson's ground-based data also include (84, 137, 99) images in (B , V , I) for NGC 6388. Although these data are not included in the plots in this paper,




FIG. 2.— Similar to Figure 1, but using the reddening-independent quantity Q instead of V , enforcing a match of the TO colors of the two clusters, and zooming in around the TO/SGB level.

they were used to establish a network of photometric standards in the cluster field which could be used to establish accurate photometric zero-points for the ACS images. Color transformations for the ACS data were based on these local standards and similar ACS-ground-based comparisons for 47 Tuc, NGC 2419, NGC 6341 (M92), and NGC 6752.

The data were reduced in the standard manner, using the DAOPHOT-ALLFRAME software packages, following commonly understood reduction procedures (e.g., Stetson 1987, 1990, 1994). A complete description of our dataset, reduction and calibration procedures will be described in a future paper (Stetson et al. 2006, in preparation).

Before closing, we note that the ACS filter bandpasses are not identical to the standard B , V , I bandpasses of Landolt (1992)—but then again, even among the various ground-based observing runs where the clusters were observed, the filters and detectors are not identical: bandpass mismatch is an unavoidable complication in filter photometry when one does not have a privately owned photometer. Even *with* a private photometer, bandpasses can drift with atmospheric conditions, or as filters, detectors, and mirror coatings age. We have removed the effects of mismatch to second order by including linear and quadratic color terms, both in the transformation equations that relate our ground-based observations to Landolt's photometric system *and* in the equations that relate the ACS magnitudes to our ground-based system. Extensive experience suggests that the residual effects of bandpass mismatch in these filters reach an irreducible minimum scatter of 0.01 – 0.02 mag on a star-by-star basis, and considerably less than this on an ensemble average of many stars.

3. THE FIRST DEEP CMD OF NGC 6388

Our deep V , $B-I$ CMD of NGC 6388 is shown in Figure 1 (*red \times symbols*), overplotted on the 47 Tuc CMD (*plus signs*). To obtain this plot, we have applied shifts of +3.2 in V and +0.85 in $B-I$ to the 47 Tuc data, in order to register its red HB to NGC 6388's. The magnitude and color of the TO in each

cluster were determined by an iterative robust numerical fit of a parabola to the data in a magnitude range ± 0.5 mag of the TO; we obtained $(V, B-I)_{\text{TO}} = (20.93, 2.04)$ for NGC 6388, and $(17.70, 1.27)$ for 47 Tuc. Note that we obtained clean, sharp, well-populated sequences in the CMD by plotting only the stars with the best photometry. Our selection criteria will be described in Stetson et al. (2006).

3.1. The Main Sequence TO and Subgiant Branch (SGB)

The TO points and SGB's of 47 Tuc and NGC 6388 coincide in brightness rather well once their red HB's are registered. This suggests that any age difference between the two clusters is at the level of $\Delta t \lesssim 1$ Gyr, assuming that their red HB's have the same absolute luminosity.

Stetson, Vandenberg, & Bolte (1996) and Vandenberg, Larson, & De Propris (1998) computed isochrones for different values of $[\text{Fe}/\text{H}]$, $[\alpha/\text{Fe}]$, helium abundance Y , the mixing length parameter α_{MLT} , and rotation rates ω . As is already well known, the difference in color between the base of the RGB and the main-sequence TO is a good relative age indicator, due to its small dependence on $[\text{Fe}/\text{H}]$, $[\alpha/\text{Fe}]$, and ω . On the other hand, these calculations also show that this color difference, as well as the detailed shape of the SGB, do present some dependence on Y and α_{MLT} .

To perform a more meaningful differential comparison that minimizes the effects of differential reddening in NGC 6388, we have replaced V with the reddening-independent quantity $Q = V - 1.153(B-I)$, and then registered the TO points of the two clusters in color. Although numerical fitting of parabolas to Q as a function of $B-I$ at the TO's of the two GC's suggests that their bluest colors differ by 0.77 mag, we still performed a sanity check by directly overplotting the two CMDs with a range of horizontal shifts, and concluded that $\Delta(B-I) \sim 0.80 \pm 0.05$ (1σ error bar) appeared best to the eye (see Fig. 2). We should note that the detection limit in these data is near $Q \approx 19.5$, and that past experience (e.g., Stetson & Harris 1988) leads us to expect that the observed colors may be biased too blue by ~ 0.1 mag as this limit is approached. Accordingly, we have been careful to restrict our comparisons to the immediate vicinity of the TO itself, $Q \sim 18.2$, where any systematic bias is expected to be much smaller.

Figure 2 shows that, when the TO points of the clusters are registered, the difference in color $\Delta(B-I)_{\text{TO}}^{\text{RGB}}$ between the base of the RGB and the TO is very similar for both clusters. This quantity is only slightly larger for NGC 6388 than for 47 Tuc, at a level $\simeq +0.010 \pm 0.035$ mag (1σ error bar). If interpreted in terms of an age difference, this translates into $\Delta t \simeq -0.5 \pm 1.6$ Gyr (NGC 6388 being younger). Assuming that the clusters have similar ages, $[\text{Fe}/\text{H}]$, and $[\alpha/\text{Fe}]$, which appears most consistent with the TO/red HB ("vertical method") and the RGB data (see also the next subsection), one also finds $\Delta Y \simeq -0.014 \pm 0.024$ (in the sense that NGC 6388 should have a lower Y), or $\Delta\alpha_{\text{MLT}} \simeq -0.07 \pm 0.12$ (again in the sense that NGC 6388 should have a lower α_{MLT}). Such a difference in Y between the two GC's would be the opposite of that needed to explain the NGC 6388 HB morphology. Stronger constraints on Y and α_{MLT} variations between the two clusters are derived from their RGB properties (see §3.2).

3.2. The Red Giant Branch

The RGB's of 47 Tuc and NGC 6388 have very similar morphologies, according to Figure 1. The larger scatter in

the NGC 6388 CMD than in 47 Tuc's might be due in part to photometric errors and in part to differential reddening, thus making it difficult to determine if the stars scattered toward the red of the main RGB in NGC 6388 represent a minor metal-rich component. A sizeable metal-poor component is clearly not present (see also R02). A small deviation of the bulk of the brighter NGC 6388 RGB stars towards redder colors compared with the 47 Tuc CMD, if real, would suggest that NGC 6388 might be slightly more metal-rich (i.e., by $\lesssim 0.25$ dex) than 47 Tuc ($[\text{Fe}/\text{H}] = -0.76$ dex; Harris 1996). When their red HB's are registered, one also finds that their TO colors become somewhat offset (with NGC 6388's being bluer) and that their RGB's actually cross just above the HB level (Fig. 1). Theoretical isochrones show that these effects are to be expected if NGC 6388 is more metal-rich than 47 Tuc at the level indicated above.

The "bump" in the RGB luminosity function lies at a $V \approx 17.70$ in NGC 6388, and 14.50 in 47 Tuc. Thus, the TO-to-bump V -magnitude differences are 3.26 and 3.20, respectively. If real, this difference would suggest that NGC 6388 is more metal-poor (by $\lesssim 0.1$ dex) than 47 Tuc. On the other hand, the values of $\Delta V_{\text{bump}}^{\text{RHB}}$ measured by Zoccali et al. (1999) differ by only 0.03 mag between 47 Tuc and NGC 6388; this could be produced by a very small metallicity difference, by a $\Delta Y \simeq 0.007$ (NGC 6388 being more helium-rich; see Fig. 5 in R02), and/or by a small difference in age of $\Delta t \simeq 1$ Gyr (NGC 6388 being younger).

Note, finally, that near-infrared photometry (e.g., Frogel et al. 2001; Valenti, Ferraro, & Origlia 2004) shows that NGC 6388 and NGC 6441 have, if anything, slightly bluer RGB's than 47 Tuc, contrary to what would be expected in the R02 scenario. Indeed, Ferraro et al. find a normal value of $\alpha_{\text{MLT}} \approx 2.0$ for both NGC 6441 and 47 Tuc (NGC 6388 is not in their studied sample). The lack of a large difference in α_{MLT} and $[\text{Fe}/\text{H}]$ between the two GC's, while not unexpected, already rules out the R02 scenario, in which a combination of high metallicity and low α_{MLT} is invoked, leading to an underluminous red HB as mentioned in §1.

3.3. The Red Horizontal Branch

An overluminous red HB, as would be implied by a high primordial Y (SC98), is ruled out if, as appears likely, the two clusters have similar ages and metallicities. The HB luminosity function is also known to be affected by the stars' evolutionary parameters; in particular, a higher Y in NGC 6388 should produce more luminosity evolution away from the zero-age HB (e.g., Dorman, Vandenberg, & Laskarides 1989), which is clearly not present in the observed CMD. Likewise, the fact that there is no significant component fainter than the bulk of the NGC 6388 red HB stars suggests that any metal-rich component in this cluster (i.e., with $[\text{Fe}/\text{H}] \gtrsim -0.5$ dex) should be minor.

Figure 3 shows a more detailed comparison of the CMDs of NGC 6388 and 47 Tuc around the red HB region, again using the reddening-independent quantity Q . The 47 Tuc Q values were shifted by $\Delta Q = +2.22$ mag. Adding random scatter at a level of $\sigma(B-I) \simeq 0.03$ mag to the 47 Tuc CMD allows one to reproduce the overall appearance of the RGB and the red HB quite well, thus implying a $\Delta Y \lesssim 0.03$ (in the sense of NGC 6388 having a higher Y) based on the luminosity width of the HB. The main difference between the two HB's is the tendency for the bluer of the NGC 6388 red HB stars to be slightly brighter than the average line that de-




FIG. 3.— Same as in Figure 1, but using Q instead of V and focusing around the red HB region.

finer the 47 Tuc red HB. This, along with the fact that the TO points and SGB's of the two GC's match well in brightness when the two red HB's are registered, strongly suggests that both the RR Lyrae and blue HB components (and, to a lesser extent, also the bluer red HB stars) of NGC 6388 are indeed intrinsically overluminous with respect to the 47 Tuc red HB.

3.4. The Asymptotic Giant Branch “Clump”

Well-populated AGB “clumps” are seen in both the 47 Tuc and NGC 6388 CMD's. Ferraro et al. (1999) have shown that, the bluer the HB morphology of a GC, the less pronounced the resulting AGB clump. Therefore, the majority of the stars in this phase originate from red HB stars. It is interesting to note that the difference in V magnitude between the AGB clump and the red HB is basically indistinguishable between NGC 6388 and 47 Tuc. Unfortunately, this does not provide

us with strong constraints on the difference in Y or metallicity between the two GC's, since the difference in magnitude between the AGB clump and the red HB is not very sensitive to these evolutionary parameters (e.g., Bono et al. 1995).

4. DISCUSSION

In the present *Letter*, we have shown that, apart from the blue HB and RR Lyrae components, the CMD's of NGC 6388 and 47 Tuc are very similar, thus strongly suggesting that the bulk of the stars in these two clusters are very similar. Differences in age, metallicity, $[\alpha/\text{Fe}]$, Y , and α_{MLT} between the two GC's, if present, should be small. Our results suggest that the red HB component of NGC 6388 is neither underluminous (as in the “canonical tilt” scenario of R02) nor overluminous (as in the high- Y scenario of SC98), except for its bluest stars. On the other hand, both the RR Lyrae and blue HB stars in NGC 6388 are significantly overluminous compared to field RR Lyrae stars of similar $[\text{Fe}/\text{H}]$, thus explaining NGC 6388's anomalously long RR Lyrae periods. The lack of a sizeable luminosity difference between the red HB's of NGC 6388 and 47 Tuc indicates that *the 2ndP that leads to the production of an overluminous RR Lyrae + blue HB component in NGC 6388 must be non-canonical in nature*—that is, it must be neither age nor RGB mass loss, or else a sloped HB would not result (SC98). However, since only a relatively small fraction ($\simeq 17\%$; Catelan 2006) of the HB stars are in the RR Lyrae strip or on the blue HB, only a minor fraction of the cluster stars should be affected by this non-canonical 2ndP, which could be either Y or/and the helium-core mass at the He-flash. (A fraction of the red HB stars, especially the bluer 25% or so, could also be affected, though to a much smaller extent.) Additional, detailed studies of the blue HB, RR Lyrae, and main sequence components will be required before we are in a position to conclusively decide what 2ndP(s) is (are) responsible (see Catelan 2006 for a recent review).

We thank the referee, G. Piotto, for a very helpful report. M.C. acknowledges support by Proyecto FONDECYT Regular No. 1030954. B.J.P. would like to thank NASA for the support for the SNAP-9821 project through a grant from the Space Telescope Science Institute. H.A.S. acknowledges support from the CSCE and NSF under AST 02-05813.

REFERENCES

- Bono, G., Castellani, V., Degl'Innocenti, S., & Pulone, L. 1995, *A&A*, 297, 115
- Catelan, M. 2006, in *Resolved Stellar Populations*, ASP Conf. Ser., ed. D. Valls-Gabaud & M. Chávez, in press (astro-ph/0507464)
- Corwin, T. M., Sumner, A. N., Pritzl, B. J., Smith, H. A., Catelan, M., Sweigart, A. V., & Stetson, P. B. 2006, *AJ*, 132, 1014
- Dorman, B., O'Connell, R. W., & Rood, R. T. 1995, *ApJ*, 442, 105
- Dorman, B., Vandenberg, D. A., & Laskarides, P. G. 1989, *ApJ*, 343, 750
- Ferraro, F. R., Messineo, M., Fusi Pecci, F., de Palo, M. A., Straniero, O., Chieffi, A., & Limongi, M. 1999, *AJ*, 118, 1738
- Ferraro, F. R., Valenti, E., Straniero, O., & Origlia, L. 2006, *ApJ*, 642, 225
- Frogel, J. A., Stephens, A., Ramírez, S., & DePoy, D. L. 2001, *AJ*, 122, 1896
- Harris, W. E. 1996, *AJ*, 112, 1487 (Feb. 2003 update)
- Holtzmann, J. A., et al. 1995, *PASP*, 107, 1065
- Landolt, A. U. 1992, *AJ*, 104, 340
- Layden, A. C., Ritter, L. A., Welch, D. L., & Webb, T. M. A. 1999, *AJ*, 117, 1313
- Moehler, S., Sweigart, A. V., & Catelan, M. 1999, *A&A*, 351, 519
- Moehler, S., & Sweigart, A. V. 2006, *A&A*, 455, 943
- Piotto, G., et al. 2002, *A&A*, 391, 945
- Pritzl, B., Smith, H. A., Catelan, M., & Sweigart, A. V. 2000, *ApJ*, 530, L41
- Pritzl, B. J., Smith, H. A., Catelan, M., & Sweigart, A. V. 2001, *AJ*, 122, 2600
- Pritzl, B. J., Smith, H. A., Catelan, M., & Sweigart, A. V. 2002, *AJ*, 124, 949
- Pritzl, B. J., Smith, H. A., Stetson, P. B., Catelan, M., Sweigart, A. V., Layden, A. C., & Rich, R. M. 2003, *AJ*, 126, 1381
- Raimondo, G., Castellani, V., Cassisi, S., Brocato, E., & Piotto, G. 2002, *ApJ*, 569, 975 (R02)
- Ree, C. H., Yoon, S.-J., Rey, S.-C., & Lee, Y.-W. 2002, in *Omega Centauri, A Unique Window into Astrophysics*, ASP Conf. Ser. 265, ed. F. van Leeuwen, J. D. Hughes, & G. Piotto (San Francisco: ASP), 101
- Rich, R. M., Minniti, D., & Liebert, J. 1993, *ApJ*, 406, 489
- Rich, R. M., et al. 1997, *ApJ*, 484, L25
- Silbermann, N. A., Smith, H. A., Bolte, M., & Hazen, M. L. 1994, *AJ*, 107, 1764
- Stetson, P. B. 1987, *PASP*, 99, 191
- Stetson, P. B. 1990, *PASP*, 102, 932
- Stetson, P. B. 1994, *PASP*, 106, 250
- Stetson, P. B., & Harris, W. E. 1988, *AJ*, 96, 909
- Stetson, P. B., Vandenberg, D. A., & Bolte, M. 1996, *PASP*, 108, 560
- Sweigart, A. V., & Catelan, M. 1998, *ApJ*, 501, L63 (SC98)
- Valenti, E., Ferraro, F. R., & Origlia, L. 2004, *MNRAS*, 351, 1204
- Vandenberg, D. A., Larson, A. M., & De Propriis, R. 1998, *PASP*, 110, 98
- Zoccali, M., Cassisi, S., Piotto, G., Bono, G., & Salaris, M. 1999, *ApJ*, 518, L49