Information-theoretic model selection applied to supernovae data

Marek Biesiada

Department of Astrophysics and Cosmology,
University of Silesia
Uniwersytecka 4, 40-007 Katowice, Poland
biesiada@us.edu.pl

Abstract

Current advances in observational cosmology suggest that our Universe is flat and dominated by dark energy. There are several different theoretical ideas invoked to explain the dark energy with relatively little guidance of which one of them might be right. Therefore the emphasis of ongoing and forthcoming research in this field shifts from estimating specific parameters of cosmological model to the model selection.

In this paper we apply information-theoretic model selection approach based on Akaike criterion as an estimator of Kullback-Leibler entropy. Although this approach has already been used by some authors in similar context, this paper provides more systematic introduction to the Akaike criterion. In particular, we present the proper way of ranking the competing models based on Akaike weights (in Bayesian language - posterior probabilities of the models). This important ingredient is missing in alternative studies dealing with cosmological applications of Akaike criterion.

Out of many particular models of dark energy we focus on four: quintessence, quintessence with time varying equation of state, brane-world and generalized Chaplygin gas model and test them on Riess' Gold sample.

As a result we obtain that the best model - in terms of Akaike Criterion - is the quintessence model. The odds suggest that although there exist differences in the support given to specific scenarios by supernova data most of the models considered receive similar support. The only exception is Chaplygin gas which is considerably less supported. One can also notice that models similar in structure i.e. Λ CDM, quintessence and quintessence with variable equation of state are closer to each other in terms of Kullback-Leibler entropy. Models having different structure i.e. Chaplygin gas or brane-world scenario are more distant (in Kullback-Leibler sense) from the best one.

Keywords: classical tests of cosmology, dark energy theory

1 Introduction

The problem of "dark energy" in the Universe is one of the most important issues in modern cosmology. It appeared after the discovery of accelerated expansion of the Universe as inferred from the SNIa Hubble diagram [1]. Since then a lot of specific scenarios have been put forward as an explanation of this puzzling phenomenon. They fall into two broad categories: searching an explanation among hypothetical candidates for dark energy (cosmological constant Λ [1], quintessence - evolving scalar fields [2], Chaplygin gas [3]) or modification of gravity theory (supergravity [4], brane world scenarios [5]).

In the problem of statistical inference from empirical data one very often encounters the problem of selecting the best approximating model [6]. This is exactly the problem one has in the context of dark energy where there exist a variety of theoretical ideas of what could be the cause of accelerating Universe and at the same time there is relatively little theoretical guidance of which specific model (or a class thereof) is preferred. Therefore, it is interesting to ask which cosmological model is the most supported by the data which triggered the problem.

In this paper we approach the above mentioned question from the perspective of information theoretic model selection. One of the approaches available is that of Akaike initiated by him in early seventies [7] and developed during subsequent years into a simple to use diagnostic called Akaike Information Criterion (AIC). In cosmology Akaike criterion has first been used by Liddle [8] and then in papers [9]. Since AIC is relatively unknown among cosmologists while being being popular, if not standard, in other branches of science (e.g. biostatistics [6]) Section 2 contains rudimentary introduction to the ideas underlying AIC closely following [6]. In Section 3 we briefly outline cosmological models which are then compared by means of the Akaike criterion. We illustrate the model selection ideas on Riess' Gold Sample of supernovae [10]. The final section contains results and conclusions.

2 Information theoretical model selection criteria

Akaike information-theoretical model selection criterion is based on Kullback-Leibler information. Kullback-Leibler information between two distributions f(x) and g(x) is defined as

$$I(f,g) = \int f(x) \ln \frac{f(x)}{g(x)} dx \tag{1}$$

The intuitive meaning of I(f,g) (also called K-L divergence) is the information lost when g is used to approximate f. It can be viewed as an extension of Shannon's entropy and sometimes is thus referred to as relative entropy. In cosmology, a very interesting application of this concept has been made by Hosoya et al. [11] who proposed the Kullback Leibler Relative Information Entropy as a measure of the distinguishability of the local inhomogeneous mass density field from its spatial average on arbitrary compact domains.

Let us assume that f(x) denotes the true mechanism behind the data and $g(x|\theta)$ its approximating model (parametrized by θ). The problem is that K-L divergence cannot be assessed without prior knowledge of the true model f(x) as well as parameters θ of the approximating model $g(x|\theta)$. However, given f(x) and $g(x|\theta)$ there exists the "best" value of θ for which Kullback-Leibler divergence is minimized. The observation that

maximum likelihood estimator $\hat{\theta}$ of θ parameter is exactly this K-L "best" one was the crucial ingredient in Akaike's derivation of his criterion.

The core result of Akaike was in showing that an approximately unbiased estimator of K-L divergence is $ln(\mathcal{L}(\hat{\theta}|data)) - K$ where \mathcal{L} is the likelihood function (more precisely its numerical maximum value - taken at $\hat{\theta}$) and K is the number of estimable parameters (θs) in approximating model $g(x|\theta)$. For historical reasons Akaike formulated this result in the following form:

$$AIC = -2ln(\mathcal{L}(\hat{\theta}|data)) + 2K \tag{2}$$

which became known as Akaike information criterion. Heuristically one may think of it as of an estimator of K-L divergence between the model at hand $g(x|\theta)$ and an unknown true model f(x) which generated the data. In the expression for AIC one can recognize two terms: the first measuring goodness of model fit (or more precisely the lack thereof) and the second one (competing) measuring model complexity (number of free parameters).

Defined in this way AIC value has no meaning by itself for a single model (simply because the true model f(x) is unknown). What is useful, instead are the differences $\Delta_i := AIC_i - AIC_{min}$ calculated over the whole set of alternative candidate models i = 1, ..., N where by AIC_{min} we denoted $min\{AIC_i; i = 1, ..., N\}$. Comparing several models, the one which minimizes AIC could be considered the best. Then the relative strength of evidence for each model can be calculated as the likelihood of the model given the data $\mathcal{L}(g_i|data) \propto exp(-\frac{1}{2}\Delta_i)$. Relative likelihoods of the models $\mathcal{L}(g_i|data)$ normalized to unity are called Akaike weights w_i . In Bayesian language Akaike weight corresponds to the posterior probability of a model (under assumption of equal prior probabilities). The (relative) evidence for the models can also be judged by the evidence ratios of model pairs $\frac{w_i}{w_j} = \frac{\mathcal{L}(g_i|data)}{\mathcal{L}(g_j|data)}$. If referred to the best model, evidence ratio gives odds against the given model. One can easily see (with these definitions) that AIC differences of 2,4,8,10 correspond to the odds ratios 2.7, 7.4, 54.6 and 148.4 respectively. This justifies the rules of Akaike model selection that Δ_i in the range 0-2 mean that model i has almost the same support from data as the best one, for the range 2-4 this support is considerably less and with $\Delta_i > 10$ model i is practically irrelevant.

A very similar criterion was derived by Schwarz [12] in a Bayesian context. It is known as the so called Bayesian information criterion (BIC) ([12]):

$$BIC = -2ln(\mathcal{L}(\hat{\theta}|data)) + Kln(n)$$
(3)

where n is sample size and as previously K denotes number of parameters. BIC is not an estimator of K-L divergence – its derivation stems from estimating the marginal likelihood of the data (marginalized over parameters). In cosmological model selection context BIC was used in [9, 13]. Its interpretation, however, should go along similar routes as presented above: values, differences, weights and odds. BIC does not take the full advantage offered by Bayesian techniques. Bayesian model averaging approach, although computationally demanding is by far better. In cosmology it was pursued e.g. by Kunz, Trotta and Parkinson [14] (see also references therein).

It should be noticed that according to some authors [15] in the limit of large data (large n) AIC tends to favor models with more parameters while BIC tends to penalize them.

3 Cosmological models fitted to supernovae data

Our aim is to find out what is the degree of support (in terms of AIC) given by supernovae data to different cosmological scenarios which might (at least phenomenologically) describe presently accelerating Universe.

The sample we use is the so called "Gold" sample of Riess et al. [10] comprising 157 supernovae compiled from a set of previously observed SNIa with reduced systematic errors from differences in calibrations.

To proceeded with fitting the SNIa data we need the magnitude-redshift relation $m(z, \mathcal{M}, \theta_i) = \mathcal{M} + 5 \log_{10} D_L(z, \theta_i)$ where: by θ_i we denoted symbolically cosmological parameters of fitted scenario and $D_L(z, \theta_i) = (H_0/c)d_L(z, \theta_i)$ is the luminosity distance with H_0 factored out and the intercept here is defined as: $\mathcal{M} = M - 5 \log_{10} H_0 + 25$ where M is the absolute magnitude of SNIa. The fitting is performed according to procedure equivalent to marginalization over the intercept (as described in [16]).

In the framework of Friedman-Robertson-Walker cosmology the luminosity distance reads:

$$d_L(z) = (1+z)\frac{c}{H_0} \frac{1}{\sqrt{|\Omega_k|}} \mathcal{F}\left(H_0 \sqrt{|\Omega_k|} \int_0^z \frac{dz'}{H(z')}\right)$$
(4)

where $\Omega_k := -\frac{k}{a_0^2 H_0^2}$ is the curvature term. The $\mathcal{F}(u)$ function is defined as $\mathcal{F}(u) = \sin u$ for k = +1, $\mathcal{F}(u) = u$ for k = 0 and $\mathcal{F}(u) = \sinh u$ for k = -1.

The estimation of cosmological model parameters was performed using the maximum likelihood approach. We assumed that supernovae measurements came with uncorrelated Gaussian errors and in this case the likelihood function \mathcal{L} could be determined from chi-square statistic $\mathcal{L} \propto \exp(-\chi^2/2)$ [1]. The χ^2 function here is defined as:

$$\chi^2 = \sum_i \frac{(m_i^{th} - m_i^{obs})^2}{\sigma_i^2}$$

where the sum is over the SNIa sample and σ_i denote the (full) statistical error of magnitude determination.

The sections below briefly introduce three types of cosmological models which will then be compared by using the Akaike criterion. Formulae therein are given in general form i.e. including the curvature term. Further on we will restrict our attention to flat model k=0 because the flat FRW geometry is strongly supported by cosmic microwave background radiation (CMBR) data [17].

$3.1 \quad \Lambda CDM \mod el$

Friedman - Robertson - Walker model with non-vanishing cosmological constant and pressure-less matter including the dark part of it responsible for flat rotation curves of galaxies (the co called Λ CDM model) is a standard reference point in modern cosmology. Sometimes it is referred to as a concordance model since it fits rather well to independent data (such like CMBR data, LSS considerations, supernovae data). In this case the expansion equation (also called the Friedman equation) reads:

$$H^{2}(z) = H_{0}^{2}(\Omega_{m} (1+z)^{3} + \Omega_{\Lambda} + \Omega_{k}(1+z)^{2})$$
(5)

The cosmological constant suffers from the fine tuning problem (being constant, why does it start dominating at the present epoch?) and from the enormous discrepancy between facts and expectations (assuming that Λ represents quantum-mechanical energy of the vacuum it should be 55 orders of magnitude larger than observed [18]).

3.2 Quintessence model

The most popular explanation of the accelerating Universe is to assume the existence of a negative pressure component called dark energy. One can heuristically assume that this component is described by hydrodynamical energy-momentum tensor with $p = w\rho$ where -1 < w < -1/3 [19]. In such case this component is called "quintessence".

In quintessential cosmology the Friedman equation reads:

$$H^{2}(z) = H_{0}^{2}(\Omega_{m} (1+z)^{3} + \Omega_{Q} (1+z)^{3(1+w)} + \Omega_{k}(1+z)^{2})$$
(6)

where by Ω_m and Ω_Q we have denoted present values of relative contributions of clumped matter and quintessence to the critical density.

If we think that the quintessence has its origins in the evolving scalar field, it would be natural to expect that w coefficient should vary in time, i.e. w = w(z). An arbitrary function w(z) can be Taylor expanded. Then, bearing in mind that both SNIa surveys or strong gravitational lensing systems are able to probe the range of small and moderate redshifts it is sufficient to explore first the linear order of this expansion. Such possibility, i.e. $w(z) = w_0 + w_1 z$ has been considered in the literature (e.g. [20]). The Friedman equation reads now:

$$H^{2}(z) = H_{0}^{2}(\Omega_{m} (1+z)^{3} + \Omega_{Q} (1+z)^{3(1+w_{0}-w_{1})} \exp(3w_{1}z) + \Omega_{k}(1+z)^{2})$$
 (7)

3.3 Generalized Chaplygin gas cosmology

In this class of models matter content of the Universe consists of pressure-less gas with energy density ρ_m representing baryonic plus cold dark matter (CDM) and of the generalized Chaplygin gas with the equation of state $p_{Ch} = -\frac{A}{\rho_{Ch}^{\alpha}}$ with $0 \le \alpha \le 1$, representing dark energy responsible for acceleration of the Universe.

The Friedman equation can be rearranged to the form:

$$H(z)^{2} = H_{0}^{2} \left[\Omega_{m} (1+z)^{3} + \Omega_{Ch} \left(A_{0} + (1-A_{0})(1+z)^{3(1+\alpha)} \right)^{\frac{1}{1+\alpha}} + \Omega_{k} (1+z)^{2} \right]$$
(8)

where the quantities Ω_i , i = m, Ch, k represent fractions of critical density currently contained in energy densities of respective components.

Generalized Chaplygin gas models have been intensively studied in the literature [21] and in particular they have been tested against supernovae data (e.g. [22] and references therein).

3.4 Brane-world cosmological model

According to brane-world scenarios [5], our 4-dimensional Universe is a surface (a brane) embedded into a higher dimensional bulk space-time in which gravity propagates. As

a consequence there exists a certain cross-over scale r_c above which an observer will detect higher dimensional effects. Cosmological models in brane-world scenarios have been widely discussed in the literature [23]. In particular the Friedman's equation takes here the following form:

$$H(z)^{2} = H_{0}^{2} \left[\left(\sqrt{\Omega_{m}(1+z)^{3} + \Omega_{r_{c}}} + \sqrt{\Omega_{r_{c}}} \right)^{2} + \Omega_{k}(1+z)^{2} \right]$$
(9)

where: $\Omega_{r_c} = \frac{1}{4r_c^2H_0^2}$. It has been shown in [23] that flat brane-world Universe with $\Omega_m = 0.3$ and $r_c = 1.4 H_0^{-1}$ is consistent with current SNIa and CMBR data. Note that in flat (i.e. k = 0.) brane-world Universe the following relation is valid: $\Omega_{r_c} = \frac{1}{4}(1 - \Omega_m)^2$.

4 Results and conclusions

Table 1 displays the results of fitting the above mentioned models to the "Gold" sample of SNIa. As already mentioned the flat prior k=0 was assumed. The reason for taking prior assumptions was that although Ω_k could have been included as a free parameter in statistical analysis (e.g.[22]), the information (even more precise than achievable this way) about cosmological parameters like k, H_0 , etc. comes from other types of experiments (such like CMBR, LSS power spectrum, BBN, gravitational lensing etc.). In observational cosmology our goal is in building a consistent picture of the Universe rather than expanding parameter space for statistical analysis. Matter density Ω_m was taken as a free parameter in analysis. Best fit to the supernova data distinguished an unrealistic value of $\Omega_m = 0.49$ in quintessential models, which is also reflected in the best fitted values of w parameters of cosmic equation of state. If one took a prior on matter density $\Omega_m = 0.3$ (as supported by alternative evidence) one would obtain $w = -1.02 \pm 0.11$ for quintessence and $w_0 = -1.40 \pm 0.25 \ w_1 = 1.67 \pm 0.89$ in models with time varying equation of state, which is similar to the values from combined evidence (SNIa, LSS, CMBR, lensing) reported in the literature. However, taking this prior would give an unfair weighting of models considered so we have assumed matter density to be a free parameter. The author thanks the referee for clarifying this point.

A comparison of $\Omega_m = 0.3$ prior fit to Chaplygin gas model, which gives $A_0 = 0.99 \pm 0.03$ $\alpha = 1.0 \pm 0.59$, and respective values from Table 1 are also worth noting. In both cases the "correct" value for Ω_m is singled out, best fits for A_0 are similar but α fits are drastically different. The best fitted Chaplygin gas model with Ω_m prior relaxed is physically equivalent to Λ CDM while taking a rigid prior prefers original Chaplygin gas model. This effect of priors in generalized Chaplygin gas models was also noted and discussed in [22].

From Table 2 one can see that the best model - in terms of Akaike Criterion - is the quintessence model. Therefore this model should be identified as a reference for calculating Akaike differences, weights and odds against alternative models considered. The odds suggest that although there exist differences in the support given to specific scenarios by supernova data almost all models considered receive similar support by the data. The support given to the Chaplygin gas model is considerably less — odds are almost 15 to 1 against it when compared to quintessential model. It is somewhat surprising if one recalls that best fitted Chaplygin model is phenomenologically equivalent to Λ CDM which is the K-L closest one to the quintessential model.

One can also notice that models similar in structure i.e. Λ CDM, quintessence and quintessence with variable equation of state are closer to each other in terms of Kullback-Leibler entropy. In fact, from purely statistical point of view (apart from different physical motivations behind each one of them), they can be considered as a family of nested models. Models having different structure i.e. brane-world scenario or Chaplygin gas are more distant (in K-L sense) from the best one. Consequently one loses more information while fitting these models to supernovae Gold sample. It should be noticed that in other papers [9] referring to Akaike criterion as a tool for cosmological model selection authors ignored the issue of odds against competing model (with respect to the best fitted one) - treating AIC merely as a tool for ranking based just on numerical value of AIC. This paper fills this gap in showing how the model selection procedure should be implemented.

It should be stressed that although it might be tempting to claim that AIC is equivalent to comparing likelihoold functions (which is true for models with the same number of parameters fitted to the data) such altitude is not correct. First of all the likelihood techniques were developed to estimate parameters of a given model best fitted to the data. Consequently, the likelihoods measure goodness of fit not the support for a given model as compared to competing models. It could intuitively be expected that these two distinct concepts are connected, and indeed they are. The proper way to do that is given exactly by the Akaike criterion, which has a sound theoretical background behind. Moreover, as demonstrated in this paper AIC results can not be simply predicted by the number of parameters in the model – the best one was found to be quintessence model (with 2 parameters) closest to it is Λ CDM (1 parameter) and the next one is Var Quintessence (3 parameters). They were found to have a similar support, while the Chaplygin gas model (the same number of parameters as Var Quintessence) is considerably less supported by the data.

Table 3 displays analogous information as Table 2 with respect to the Bayesian Information Criterion (BIC). One can see the differences in both identifying the best fitted model (now it is ΛCDM) and in ranking of remaining ones. The order of ranking reflects dimensionality of the model — 1 parameter models are BIC preferred over 2 or 3 parameter ones (Chaplygin gas now with 157 to 1 odds against is practically ruled out according to BIC).

In Table 4 ranking of the models considered according to different criteria: AIC, BIC and $\chi^2/\text{d.o.f.}$ are presented. Reasons behind the ranking for AIC were already revealed: models similar in structure to the best fitted receive comparable support. Usually used criterion of $\chi^2/\text{d.o.f.}$ gives similar ranking as AIC and BIC criterion apparently penalizes models too much for over fitting (by giving less support to those with larger number of free parameters).

One can hope that the future will shed more light on the nature of dark energy in the Universe. Special surveys (e.g. SNAP) are designed for this purpose. One should realize however, that the emphasis of the ongoing and forthcoming research is shifting from estimating specific parameters of the cosmological model (like the Hubble constant or deceleration parameter or any other physical parameter of the theory) to the model selection. Along with Bayesian techniques (e.g. model averaging) [14] information-theoretic model selection approaches are the most promising for this purpose.

 ${\bf Table~1} \\ {\bf Values~of~best~fitted~parameters~of~four~models~tested}.$

Model	Best fit model parameters (with 1σ ranges)
$\Lambda \mathrm{CDM}$	$\Omega_m = 0.31 \pm 0.04$
Quintessence	$\Omega_m = 0.49 \pm 0.06 \ w = -2.40 \pm 1.12$
Var Quintessence	$\Omega_m = 0.48 \pm 0.14 \ w_0 = -2.48 \pm 1.38 \ w_1 = 1.88 \pm 2.59$
Chaplygin Gas	$\Omega_m = 0.31 \pm 0.04 \ A_0 = 1.00 \pm 0.035 \ \alpha = 0.002 \pm 0.088$
Braneworld	$\Omega_m = 0.21 \pm 0.03$

Table 2 Values of AIC, Akaike differences, Akaike weights w_i (in Bayesian language equivalent to posterior model probabilities) and odds against the model (with respect to the best fitted one).

Model	AIC	Δ_i	w_i	Odds against
$\Lambda \mathrm{CDM}$	179.072	1.368	0.224	1.982
Quintessence	177.704	0.	0.443	1.
Var Quintessence	179.645	1.941	0.168	2.639
Chaplygin Gas	183.072	5.368	0.030	14.644
Braneworld	180.075	2.371	0.135	3.272

Table 3 Analogous values of Bayesian Information Criterion (BIC), BIC differences, BIC weights and BIC odds against the model (with respect to the best fitted one).

Model	BIC	$BIC\Delta_i$	BIC w_i	BIC Odds against
$\Lambda \mathrm{CDM}$	182.128	0.	0.481	1.
Quintessence	183.816	1.69	0.207	2.33
Var Quintessence	188.814	6.68	0.017	28.30
Chaplygin Gas	192.241	10.11	0.003	157.
Braneworld	183.131	1.00	0.292	1.65

Table 4 Ranking of cosmological models fitted to SNIa data according to AIC, BIC and χ^2/dof criteria.

Ranking	AIC	BIC	$\chi^2/\mathrm{d.o.f.}$
1.	Quintessence	$\Lambda { m CDM}$	Quintessence
2.	$\Lambda { m CDM}$	Braneworld	Var Quintessence
3.	Var Quintessence	Quintessence	$\Lambda { m CDM}$
4.	Braneworld	Var Quintessence	Braneworld
5.	Chaplygin Gas	Chaplygin Gas	Chaplygin Gas

References

- S. Perlmutter, G. Aldering, G. Goldhaber, et al., Astrophys. J 517, 565, 1999
 A. Riess, A.V. Filipenko, P. Challis, et al., Astron. J 116, 1009, 1998
- [2] B. Ratra, P.J.E. Peebles, Phys.Rev.D 37, 3406, 1988
 R.R. Caldwell, R. Dave, P.J. Steinhardt, Phys. Rev. Lett. 75, 2077, 1995
 J. Frieman, C. Hill, A. Stebbins, I. Waga, Phys. Rev. Lett. 75, 2077, 1995
 R. Caldwell, R. Dave, and P.J. Steinhardt, Phys. Rev. Lett. 80,1582, 1998
 I. Zlatev, L. Wang and P. J. Steinhardt, Phys. Rev.Lett. 82, 896, 1999
- [3] A. Kamenshchik, V. Moschella, V. Pasquier, Phys.Lett. B 511, 256, 2000;
 J.C. Fabris, S.V.B. Gonçalves, P.E. de Souza, Gen.Rel.Grav. 34, 53, 2002
- [4] Ph. Brax, J. Martin, Phys.Lett.B 468, 40, 1999
- [5] G.Dvali, G.Gabadadze, M.Porrati, Phys. Lett.B 485, 208, 2000
 G.Dvali and G.Gabadadze, Phys.Rev.D 63, 065007, 2001
- [6] K.P. Burnham, D.R. Anderson, "Model Selection and Multimodel Inference", Springer - New York, 2002
- H. Akaike, IEEE Transactions on Automatic Control AC 19, 716-726, 1974
 H. Akaike, Journal of Econometrics 16, 3-14, 1981
- [8] A.R. Liddle, Mon.Not.Roy.Astron.Soc. 351, L49, 2004
- W. Godłowski and M. Szydłowski, Phys.Lett.B 623, 10, 2005
 M. Szydłowski and W. Godłowski, Phys.Lett.B 633, 427, 2006
- [10] A.G. Riess et al. [Supernova Search Team Collaboration], Astrophys. J 607, 665, 2004
- [11] A. Hosoya, T. Buchert, M. Morita, Phys.Rev.Lett. 92, 141302-1-4, 2004
- [12] G. Schwarz, Annals of Statistics 6, 461 464, 1978
- [13] M. Szydłowski, A. Kurek, A. Krawiec, Phys.Lett. B 642, 171–178, 2006
 M. Szydłowski, A. Kurek, "Testing and selection of cosmological models with dark energy" [astro-ph/0603538]
- [14] M. Kunz, R. Trotta, D. Parkinson, Phys.Rev. D 74, 023503, 2006
- [15] D. Parkinson, S. Tsujikawa, B.A. Bassett, L. Amendola, Phys.Rev. D 71, 063524, 2005
- [16] M. Goliath, R. Amanullah, P. Astier, A. Goobar, R. Pain, Astron. Astrophys. 380, 6-18, 2001
 R. Lazkoz, S. Nesseris, L. Perivolaropoulos, J.C.A.P. 0511, 010, 2005
- [17] A. Benoit, et al., Astron. Astrophys. 399, L25-L30, 2003
- [18] S. Weinberg, Rev. Mod. Phys. **61**, 1, 1989

- [19] T. Chiba, N. Sugiyama, T. Nakamura, Mon.Not.Roy.Astron.Soc. 301, 72, 1998
 M.S. Turner, M. White, Phys.Rev.D 56, 4439, 1997
- [20] J. Weller, A. Albrecht, Phys.Rev.Lett. 86, 1939-1942, 2001
 I. Maor, R. Brustein, P.J. Steinhardt, Phys.Rev.Lett. 86, 6-9, 2001
- [21] M. Makler, S.Q. de Oliveira, I. Waga, *Phys.Lett. B* 555, 1, 2003
 P.P. Avelino, L.M.G. Beça, J.P.M. de Carvalho, C.J.A.P. Martins, P. Pinto, *Phys. Rev. D* 67, 023511, 2003
- [22] M. Biesiada, W. Godłowski, M. Szydłowski, Astrophys. J 622, 28–38, 2005
- [23] D. Jain, A. Dev, J.S. Alcaniz, *Phys.Rev.D* 66, 083511, 2002
 J.S. Alcaniz, D. Jain, A. Dev, *Phys.Rev.D* 66, 067301, 2002
 C. Deffayet, S.J. Landau, J. Raux, M. Zaldarriaga, P. Astier, *Phys.Rev.D* 66, 024019, 2002