The Identification of EGRET Sources with Flat-Spectrum Radio Sources

J. R. Mattox^{1,2}, J. Schachter³ L. Molnar⁴, R. C. Hartman⁵, A. R. Patnaik⁶

ABSTRACT

We present a method to assess the reliability of the identification of EGRET sources with extragalactic radio sources. We verify that EGRET is detecting the blazar class of AGN. However, many published identifications are found to be questionable. We provide a table of 42 blazars which we expect to be robust identifications of EGRET sources. This includes one previously unidentified EGRET source, the lensed AGN PKS 1830–210 near the direction of the Galactic center. We provide the best available positions for 16 more radio sources which are also potential identifications for previously unidentified EGRET sources. All high Galactic latitude EGRET sources ($|b| > 3^{\circ}$) which demonstrate significant variability can be identified with flat spectrum radio sources. This suggests that EGRET is not detecting any type of AGN other than blazars. This identification method has been used to establish with 99.998% confidence that the peak γ -ray flux of a blazar is correlated with its average 5 GHz radio flux. An even better correlation is seen between γ -ray flux and the 2.29 GHz flux density measured with VLBI at the base of the radio jet. Also, using high confidence identifications, we find that the radio sources identified with EGRET sources have larger correlated VLBI flux densities than the parent population of flat radio spectrum sources.

Subject headings: Gamma Rays: observations — galaxies: active — quasars: general

1. Introduction

The Energetic Gamma Ray Experiment Telescope (EGRET) aboard the Compton Gamma Ray Observatory is sensitive in the energy range 30 MeV to 30 GeV (Thompson *et al.* 1993). A catalog of EGRET sources based on the first 30 months of exposure is given by Thompson *et al.* (1995, henceforth TH95). It includes 129 sources. Five are pulsars, 40 are listed as "high-confidence identifications" of AGN, and 11 are listed as "lower-confidence identifications" of AGN. A total of 71 are unidentified.

¹Astronomy Department, Boston University, 725 Commonwealth Ave., Boston, MA 02215

²Former affiliations: Astronomy Department, University Of Maryland; Universities Space Research Association; Compton Observatory Science Support Center, NASA/GSFC; Gamma-Ray Astrophysics Branch, NASA/GSFC

³Center for Astrophysics, Cambridge, MA 02138

⁴Dept. Of Phys. & Astron, University of Iowa, Iowa City, IA 52242

⁵Code 662, NASA/GSFC, Greenbelt MD 20771

⁶Max-Planck Institut für Radioastronomie, Bonn, GERMANY

We demonstrate quantitatively in this paper that the claim that EGRET is detecting the γ -ray emission of some members of the blazar class of AGN is valid (see von Montigny *et al.* 1995 for a compendium of EGRET blazar results for the first $\sim 1/2$ of the EGRET mission). By blazars, we mean AGN with strong, compact, flat-spectrum (α be $\gtrsim -0.5$, where $S(\nu) \propto \nu^{\alpha}$) radio emission; and which also show continuum domination of the optical emission, and/or significant optical polarization, and/or significant changes in optical flux on short time scales (designated Optically Violently Variable, or OVV). These observational properties are believed to result from emission by material in a relativistic jet which is directed within $\sim 10^{\circ}$ of the line of sight. The blazar class includes objects classified as BL Lacertae type objects, high polarization quasars (HPQ), and OVV quasars. The apparent γ -ray luminosity is as much as one hundred times larger than that at all other wavelengths for some flaring EGRET blazars. Variability of the γ -ray flux from some blazars on a time-scale as short as 4 hours (Mattox *et al.* 1997) implies that the γ -rays are emitted from a compact region. The γ -ray emission is not understood. It may offer fresh insight into the blazar phenomenon, and useful diagnostics.

A careful analysis of the EGRET source identifications is in order because the source of the γ -ray emission is not well located. EGRET position estimates are imprecise because of the wide point spread function of the EGRET instrument. The half-angle of a cone which contains 68% of the EGRET events from a point source at a specific energy is well fit (Mattox *et al.* 1996a) by

$$\theta_{68} = 5.85^{\circ} [E_{\gamma}/100 \,\mathrm{MeV}]^{-0.534} \tag{1}$$

The localization expected for a source is approximately $\theta = \theta_{PSF}/\sqrt{N}$ where N is the number of γ -rays detected from the source (Thompson 1986). For weak EGRET sources, the 95% confidence position error ellipses are as large as ~5 square degrees.

In the first EGRET catalog (Fichtel et al. 1994), EGRET blazars were classified as "positive detections" or "marginal detections" depending on whether the significance of point-source γ -ray emission exceeded 5σ or 4σ respectively. The confidence of the identification was not explicitly addressed. The second EGRET catalog (TH95) did explicitly address the confidence of identification. TH95 classified an identification as "high-confidence" if the radio source was located within the 95% confidence contour of the EGRET position estimate. Also, TH95 classified some radio sources beyond the 95% confidence contour as "lower-confidence" identifications. The 5 GHz flux density (S_5) of a radio counterpart was generally required by TH95 to be $\gtrsim 1$ Jy; and the radio spectral index to be generally $\alpha \gtrsim -0.5$. No formal consideration was given to the size of the EGRET error region and where the radio source lay within that error region (other than whether the source was located within the 95% confidence contour of the EGRET position estimate region or not). No formal consideration was given to the number density of potentially confusing radio sources (other than a loose requirement that S_5 be $\gtrsim 1$ Jy). No formal consideration was given to the radio spectral index (other than a loose requirement that α be $\gtrsim -0.5$). No consideration was given to the *a priori* probability of detecting a radio source by EGRET — it is less than unity because many blazars which are bright at other wavelengths have been observed by EGRET without being detected. Also, the decision by TH95 on which radio sources beyond the 95% confidence contour to include as "lower-confidence" identifications was subjective.

We have developed an analysis of EGRET radio source identification which quantitatively incorporates all of this information. We avoid a cutoff in the search for an identification at an arbitrary confidence contour of the EGRET position estimate region. Also, we avoid terminating the search for candidates at a specific minimum radio flux density or radio spectral index. Bayes' theorem is used to incorporate the *a priori* probability of detecting a radio source (see Sturrock 1973 and Loredo 1990 for discussions of the use of Bayesian statistics in astronomy). It is reasonable to use a Bayesian analysis here for the following reasons: 1) we can demonstrate with very high confidence (see §3.1) that EGRET is detecting flat-spectrum radio sources without making any *a priori* assumptions; 2) we can derive a reasonable estimate for the *a priori* probability of EGRET detecting a radio source as a function of radio flux (see §3.2).

The method is described in §2. In §3 we demonstrate conclusively that EGRET is detecting blazars and obtain an estimate for the *a priori* probability of EGRET detecting a radio source as a function of radio flux; and then provide a quantitative study of the reliability of the EGRET identifications of TH95 and Thompson *et al.* (1996), and search for additional radio identifications among the previously unidentified EGRET sources. Limitations of this analysis are discussed in §3.6. Implications are discussed in §4. In §3.3.1, we use the high probability identifications of §3.3 to demonstrate that the EGRET blazars have higher VLBI flux densities than the parent population of flat-spectrum radio sources.

2. EGRET Source Identification

We analyze the probability of a correct identification of an EGRET source with a radio source. The calculation considers the size of the EGRET error region and where the radio source lies within that error region. Bayes' theorem is used to include consideration of the number density of potentially confusing radio sources and the *a priori* probability of detecting a specific radio source by EGRET. This method is an extension of that used by de Ruiter, Willis and Arp (1977) to identify optical counterparts of radio sources. This method has also been applied to identification of IRAS sources (Wolstencroft *et al.* 1992) and X-ray sources (Schachter, *et al.* 1997).

2.1 Radio Catalogs

Because blazars are flat-spectrum radio sources, high frequency surveys are appropriate for this study since they efficiently detect flat spectrum sources. We have used the 4.85 GHz Greenbank (GB) survey (Condon, Broderick, & Seielstad 1989) for the northern sky and the 4.85 GHz Parkes–MIT–NRAO (PMN) survey (Griffith & Wright 1993) for the southern sky. Both surveys have a threshold flux density of ~ 30 mJy and give positions with $\sim 20''$ uncertainty. They are both confusion limited at Galactic latitudes less than 3°. Therefore, we do not attempt to identify the 19 unidentified EGRET sources at latitudes less than 3° in this paper. Also, the PMN survey is confusion limited near the Large Magellanic Cloud and Centaurus A. For this reason, EGRET sources 2EG J0532-6914 and 2EG J1324-4317 are also excluded from this analysis.

The GB survey covered the declination range $0^{\circ} < \delta < 75^{\circ}$. We have used the catalog of Becker, White, and Edwards (1991) of 53522 sources derived from this survey. The White and Becker (1992) catalog of sources from the 1.4 GHz GB survey was used to obtain an $\alpha_{1.4-4.85}$ spectral index for the 22384 sources which are found in both catalogs. Because the 1.4 GHz survey preceded the 4.85 GHz survey by ~3 years, variability of blazar type AGN can result in this index differing by as much as ~1 from what would be obtained with simultaneous measurements.

The PMN survey (Griffith & Wright 1993) covered the declination range $-87.5^{\circ} < \delta < 10^{\circ}$. There are small gaps in the survey due to observational problems. The PMN catalog contains 49329 sources with $S_5 > 30$ mJy. Spectral indices are provided for the ~10% of the PMN sources which were detected in the Parkes 2.7 GHz survey (Bolton, Wright, and Savage 1979). Again, variability can cause a substantial error in this spectral index. Only one unidentified source (2EG J1332+8821) falls completely outside of these surveys. The error ellipses of two EGRET sources (J1409–0742 and J1513–0857) fall partially in gaps in the PMN survey, but both have strong identifications from the PKS catalog. We retain the original eight digit position names of both catalogs. The GB source names are prefixed with "B" because the position names are based on B1950 celestial coordinates. The PMN source names are prefixed with "J" corresponding to J2000 celestial coordinates. We use the GB survey in the region where the surveys overlap, $0^{\circ} < \delta < 10^{\circ}$.

2.2 Probability of Correct Identification

For a radio source found at an angle r from the center of an EGRET error ellipse, there are two possibilities: the source is the radio counterpart of the EGRET source; or, the source is a confusing source which is close to the EGRET source by accident. We denote the probabilities of these two hypotheses by p(id|r) and p(c|r) respectively.

If the radio source is merely a confusing source, the Poisson distribution pertains. The probability that one or more radio sources are accidently within the angle r is

$$p(r|c) = 1 - e^{-r^2/r_0^2},$$
(2)

where r_0 is the characteristic angle between confusing sources,

$$r_0 = (\pi \rho(S_5, \alpha))^{-\frac{1}{2}},\tag{3}$$

and $\rho(S_5, \alpha)$ is the number density of radio sources with at least the flux density of the candidate radio source, and a spectrum at least as flat. The differential probability is

$$dp(r|c) = 2\frac{r}{r_0^2} e^{-r^2/r_0^2} dr$$
(4)

For sources without spectral indices, we assume a spectral index of -0.5. This results in a conservative estimate of the significance of the identification for radio sources which are indeed flat spectrum. A refined estimate of the probability of correct identification can be made after further radio observations yield a spectral index. We obtain $\rho(S_5, \alpha)$ from the GB survey for the 16300 square degree region $0^{\circ} < \delta < 75^{\circ}$, and $|b| > 10^{\circ}$. The integral of $\rho(S_5, \alpha)$ over α is observed to have a Euclidean slope between 100 and 1000 mJy in agreement with the result of Kellerman and Wall (1987). We use this number density also for analysis of PMN counterparts.

If the radio source is the counterpart of an EGRET source, it is appropriate to use the EGRET data to estimate the probability distribution, p(r|id), of the angle r. A likelihood analysis of the EGRET data (Mattox et al. 1996a) yields the 95% confidence position contour from the likelihood ratio test using Wilks' theorem. These contours are usually fit accurately with ellipses. The ellipse is centered on the centroid of the 95% confidence location region which is given by TH95 as the position estimate. This position is observed to differ by only a small amount from the position of maximum likely which is formally the correct position estimate. The ellipse is specified by the semi-major axis a, semi-minor axis b, and the position angle of the major axis ϕ . The departure from circularity is often substantial. This results from source confusion (nearby γ -ray sources below the EGRET detection threshold), error in the model of Galactic diffuse γ -ray emission, and statistical fluctuation. The elliptical fits given by TH95 were used in our analysis. The parameters for the elliptical fits which are not given in TH95 (because of an imprecise fit) have also been used in this analysis: 2EG J0426+6618, a = 66', b = 54', $\phi = 27^{\circ}$; 2EG J0511+5523, a = 135', b = 36', $\phi = 24^{\circ}$; 2EG J0545+3943, a = 48', b = 47', $\phi = 94^{\circ}$; These fits are all accurate to within $\sim 30\%$ in radius which is not substantially worse than the potential effect of error in the Galactic diffuse model and undetected nearby γ -ray sources.

For strong sources, we observe that the dependence of the log of the likelihood of the EGRET data on the assumed source position is well represented by a paraboloid. The 95% confidence contour corresponds to a drop in the log of the likelihood of 3.00 from the maximum (Mattox *et al.* 1996a). Therefore, we represent the log likelihood surface corresponding to an elliptical error region as an elliptic paraboloid $\ln L_{max} - \ln L = 3r^2/\Psi^2$, where

$$\Psi = [a^{-2}\cos^2(\theta - \phi) + b^{-2}\sin^2(\theta - \phi)]^{-1/2},$$
(5)

and θ is the position angle. For a circular error region, Ψ is simply the 95% confidence radius. From Wilks' theorem, $2(lnL_{max} - lnL)$ is distributed as χ_2^2 in the null hypothesis (Mattox *et al.* 1996a). Thus, the confidence with which a source may be expected within a region delineated by a $\Delta \equiv \ln L_{max} - \ln L$ decrease in log likelihood is

$$C = 1 - \int_{2\Delta}^{\infty} \chi_2^2(\zeta) d\zeta = 1 - e^{-\Delta}$$
(6)

Therefore, the probability of the EGRET source being located at an angle smaller than r is

$$p(r|id) = 1 - e^{-3r^2/\Psi^2}$$
(7)

and

$$dp(r|id) = 6\frac{r}{\Psi^2} e^{-3r^2/\Psi^2} dr$$
(8)

Bayes' theorem allows the observed angle between the radio source and the EGRET position estimate, r, to be used with an *a priori* probability and the expected distributions of r under the two possible hypotheses (equations 4 and 8) to obtain p(id|r), the *a posteriori* probability that a radio source is the correct identification of an EGRET source.

$$p(id|r) = \frac{p(id)dp(r|id)}{dp(r)} = \frac{p(id)dp(r|id)}{p(id)dp(r|id) + p(c)dp(r|c)}$$
(9)

With the substitution $\eta \equiv p(id)$ for the *a priori* probability, and using the fact that one of the two hypotheses is true, either a source is the identification or it is a confusing source, p(c) = 1 - p(id); we obtain

$$p(id|r) = \frac{\frac{\eta}{1-\eta}LR}{\frac{\eta}{1-\eta}LR+1},\tag{10}$$

where the likelihood ratio, LR, is defined to be

$$LR \equiv \frac{dp(r|id)}{dp(r|c)} = 3\frac{r_0^2}{\Psi^2} e^{-r^2(3/\Psi^2 - 1/r_0^2)}$$
(11)

Figure 1 shows the LR and η dependence of p(id|r) given by equation 10.

We note that LR diverges at large r if $\Psi > \sqrt{3} r_0$. In this case, the differential probability of r under the confusion hypothesis falls off more rapidly with r than it does under the identification hypothesis. This reflects the fact that the size of the EGRET error region, Ψ , is larger than the characteristic angle between confusing sources, r_0 . A meaningful identification is not possible. We note that LR < 1 at r = 0 in this case. Therefore, we require $\Psi < \sqrt{3} r_0$ before we consider a radio source as a candidate identification of an EGRET source. This limits the number density of sources for which an attempted identification is meaningful. The consequence is that a 5 GHz flux density of $\gtrsim 50$ mJy is required for a source to be a meaningful candidate for identification given the error ellipse sizes typical of the unidentified EGRET sources.

3. Results

Table 1 presents the results of the analysis described in §2 for the 102 EGRET sources from TH95 which are not identified as pulsars, and which are not in regions where the radio surveys are source confused $(|b| < 3^{\circ}, \text{ and near the LMC and Cen A})$. All radio sources are given for which an *a posteriori* probability, p(id|r) > 0.001 is found. The radio source with the highest value of p(id|r) is included for all EGRET sources which fall within the radio survey regions. We note that sources with radio flux densities as small as ~50 mJy appear, but with very small probabilities of being EGRET sources. The same analysis is presented in Table 2 for the sources in the Supplement to the Second EGRET Catalog (Thompson *et al.* 1996). The content of Tables 1& 2 is:

Column (1) contains the J2000 position name of the EGRET source from TH95 or Thompson *et al.* (1996).

Column (2) is the Galactic position of the EGRET source (from TH95 or Thompson *et al.* 1996). For the five sources where TH95 accidently gave the radio position (0528+134, Mkn 421, 3C 279, 1406-076, 3C 454.3), the EGRET positions have been substituted.

Column (3) is the variability index which is $-\log_{10}(P)$, where P is the probability that the source is invariant based on a χ^2 test of flux for all available viewing periods (McLaughlin *et al.* 1996). A larger variability index corresponds to a higher probability that the source is variable.

Column (4) is the identification given (for some sources) by TH95. A question mark following the B1950 radio position name means that TH95 indicated that it was a "lower-confidence identification"; otherwise, the identification was indicated by TH95 to be a "high-confidence identification".

Column (5) is the name of the potential radio counterpart. The original B1950 position names are used for Greenbank sources, and the original J2000 position names are used for PMN sources.

Column (6) is a more common name for some potential radio counterpart.

Column (7) is the 4.85 GHz flux density in mJy from the survey catalog.

Column (8) is the radio spectral index, α , from the radio survey flux densities if available, $S(\nu) \propto \nu^{\alpha}$.

Column (9) is r_0 , the characteristic angle (in arcmin) between confusing sources which are at least as bright and flat as this radio source (see equation 3).

Column (10) is $\eta = p(id)$, the *a priori* probability that the radio source is a γ -ray source from equation 13.

Column (11) is the angle, r (in arcmin), between the EGRET position estimate and the radio position.

Column (12) is Ψ , the radius (in arcmin) of the 95% confidence contour in the direction of the radio source from equation 5.

Column (13) is the position confidence contour at the radio position (from equation 6).

Column (14) is the likelihood ratio (equation 11) indicating the strength of the indication for the identification.

Column (15) is the *a posteriori* probability that the identification is correct as determined with equation 10.

3.1 The Statistical Significance of the EGRET Detection of Radio Loud AGN

We can now address an important question: with what confidence can we conclude that EGRET is detecting radio loud AGN? At this point, it is not appropriate to use Bayesian statistics. We have not yet determined the *a priori* probability of a flat spectrum radio source being an EGRET source. As illustrated by Figure 1, the *a posteriori* probabilities depends very strongly on the *a priori* probability. Note that for *a priori* probabilities $\eta = 1$ or $\eta = 0$, equation 10 shows that Bayes' theorem implies that the EGRET data do not influence the expectation about whether or not a radio source emits γ -rays.

Since Bayesian statistics are of no use at this point, we apply a "frequentist" ansatz, a Monte Carlo calculation. We use LR (equation 11) as an empirical indication of the strength of a potential identification. We find that the distribution of LR for the most likely identification of each source in Table 1 differs dramatically from the distribution found by Monte Carlo calculation assuming no EGRET detections of radio loud AGN. Simulated EGRET source positions were chosen randomly with $|b| > 3^{\circ}$ and used with the error ellipse parameters of TH95. The analysis of §2 for ~3000 simulated sources yielded an estimate for the distribution of LR in the null hypothesis. A long tail at large LR was observed. Confusing radio sources with $LR > 10^3$ were found at a frequency of 0.010 ± 0.002 . This corresponds to the occurrence at a non-negligible frequency of spurious coincidences which look fairly solid. We find no strong dependence of LR on the size of the error ellipse, supporting the validity of the method we describe in §2.

The strongest identification (and coincidently the first, Hartman *et al.* 1992) is 3C 279 with $LR = 4.6 \times 10^5$. The large value of LR reflects the fact that 3C 279 is centrally located in a small error ellipse, and that sources as bright and flat as 3C 279 are very rare: $\rho(S_5 \ge 17 \text{Jy}, \alpha \ge 0.3) = 4 \times 10^{-3}$ per square degree. The largest observed LR in the ~3000 Monte Carlo trials was 6×10^3 . Therefore, the 3C 279 detection alone indicates that AGN are seen by EGRET with a confidence well above (1-102/3000)=97%. With the admission of an *a priori* probability as small as $\eta = 10^{-3}$, equation 10 implies an *a posteriori* probability of 0.998 that 3C 279 is the EGRET source counterpart; and with the self-consistent *a priori* probability of $\eta = 0.2$ which we derive in §3.2, the *a posteriori* probability that the identification is correct is 0.999991.

Even stronger evidence that EGRET is detecting radio loud AGN is obtained by considering all sources in Table 1. From the Monte Carlo calculation, LR > 100 occurs with a frequency of 0.05 ± 0.004 in the null hypothesis. In comparison, 32 of the 102 EGRET sources in Table 1 have an identification with LR > 100. The probability to obtain this by chance is

$$\sum_{n=32}^{102} 0.95^{(102-n)} 0.05^n \frac{102!}{(102-n)!n!} = 2 \times 10^{-17}$$
(12)

This is a very conservative estimate of the significance because many of the values of LR are much larger than 100. However, there are "trials" made in the selection of the type of source to try to identify with EGRET sources. The number of such trials cannot exceed the ~100 independent types of astronomical objects. There are also possible "trials" made in the selection of the energy range (e.g. 2022-077) or diffuse model (e.g. 1622-253) which was used to obtain an EGRET position for an identification. This number is ≤ 2 . The choice of the minimum value of LR of 100 for this analysis also corresponds to a number of "trials", but the number is less than ≤ 10 . Also, the use of the radio spectral index in the determination of the number density of radio sources can be regarded as a trials factor of size ~2. (We demonstrate in §4.2 that EGRET is generally not detecting steep-spectrum radio sources, so the use of the radio spectral index in the determination of the number density of radio sources is justified.) The number of potential "trials", the product of these, is far less than the inverse of the result of equation 12. Therefore, we conclude with very high confidence that EGRET is detecting radio loud AGN. We note that this result does not depend on any *a priori* assumption about the emission of γ -rays by AGN.

3.2 The A Priori Probability of EGRET Detecting a Radio Source

Now that we have clearly established that EGRET is detecting some radio-loud AGN, it is sensible to estimate the *a priori* probability, η , for EGRET to detect a particular flat-spectrum radio source. This will allow the use of equation 10 to estimate the *a posteriori* probability of a correct identification of any specific source. We do not attempt here to model the effect on η of the variation across the sky of the depth of the EGRET exposure. We expect that this cannot effectively be done because of the large uncertainty of the gamma-ray luminosity function of AGN (Chiang *et al.* 1995), but that it is not a huge effect. Also, we assume that η is not dependent on the radio spectral index, although we derive η for $\alpha > -0.5$. It is appropriate to thus give sources with $\alpha \leq -0.5$ the "benefit of the doubt" because of the potential error in α obtained from the surveys.

We seek a functional dependence on radio flux for η which is self-consistent as done by de Ruiter *et al.* (1977) in their identification of optical counterparts of radio sources. Specifically, we solve for a value of η such that the integral of p(id|r) given by equation 10 (for radio sources with $\alpha > -0.5$) divided by the number of radio sources considered (with $\alpha > -0.5$) yields the assumed value of η . We analyze η for several S_5 intervals. These intervals are chosen to provide adequate statistics while preserving sensitivity to the S_5 dependence of η .

For the flux density range $S_5 > 5$ Jy, iteration leads to the self-consistent value of $\eta = 0.19 \pm 0.08$ with $\Sigma p(id|r) = 6.0$ for 32 radio sources. For the flux density range $2 < S_5 < 5$ Jy, iteration leads to the self-consistent value of $\eta = 0.13 \pm 0.04$ with $\Sigma p(id|r) = 14.7$ for 109 radio sources. For the flux density range $1 < S_5 < 2$ Jy, iteration leads to the self-consistent value of $\eta = 0.034 \pm 0.011$ with $\Sigma p(id|r) = 9.1$ for 266 radio sources. And, for the flux density range $0.6 < S_5 < 1$ Jy, iteration leads to the self-consistent value of $\eta = 0.0047 \pm 0.0032$ with $\Sigma p(id|r) = 2.2$ for 473 radio sources. The uncertainty of η is simply assumed to be $\eta [\Sigma p(id|r)]^{-1/2}$. This is the approximate expectation for large S_5 because of Poisson fluctuation. For small S_5 , it might be an overestimate, but we expect that it is reasonable given potential systematic error.

We have chosen an analytical representation for η which has appropriate limiting values for both high and low values of S_5 . The high limiting value is chosen to be consistent with $\eta = 0.19 \pm 0.08$ for $S_5 >$ 5 Jy. An additional two parameters have been fit to match the relatively well determined values of η for $1 < S_5 < 2$ Jy and $2 < S_5 < 5$ Jy.

$$\hat{\eta}(S_5) = 0.2(1 - e^{-0.07(S_5/\mathrm{Jy})^{2.3}})$$
(13)

This fit is shown along with the data in Figure 2. It is slightly in excess of the data for $0.6 < S_5 < 1$ Jy, but only by a small amount relative to the uncertainty. It is not inappropriate to thus give small S_5 sources the "benefit of the doubt" because of variability. We have also done this analysis for an alternate choice of S_5 bins: $0.5 < S_5 < 0.8$ Jy, $0.8 < S_5 < 1.5$ Jy, $1.5 < S_5 < 4$ Jy, & $S_5 > 4$ Jy. The values of η thus obtained are consistent with equation 13. Also, the value of η for S_5 bins a factor of 2 narrower is consistent with equation 13. Thus, the derived *a priori* probability is not strongly dependent on binning. In §4.5, we discuss the strong probability that the very low value of η for $0.6 < S_5 < 1$ Jy implies a correlation between γ -ray flux and radio flux.

3.3 Verification of Previous Identifications

With the method of §2 and the prescription for an *a priori* probability of equation 13, we can now look objectively at the radio source identifications of TH95 which are noted in column 4 of Table 1. Most of the "high-confidence identifications" of TH95 are indeed so. Of these 40 identifications; six have $p(id|r) \ge 0.999$, seven have $0.99 \le p(id|r) < 0.999$, nine have $0.90 \le p(id|r) < 0.99$, ten have $0.50 \le p(id|r) < 0.90$, and six have $0.10 \le p(id|r) < 0.50$. Only two have p(id|r) < 0.10: 2356+196 and 1604+159. The 2EG J0000+2041 EGRET 95% confidence position contour contains 3.2 square degrees and radio sources as flat and bright as PKS 2356+196 have a number density of 0.01 per square degree. We find that the fraction of flat-spectrum radio sources with the 5 GHz flux density of PKS 2356+196 (700 mJy) which are detected by EGRET is only about 0.006. Thus we find with 93% confidence that PKS 2356+196 is a confusing source rather than the identification of 2EG J0000+2041. It is certainly possible that 2EG J0000+2041 is PKS 2356+196, but we suggest that it should not be considered to be a "high-confidence identification". Similarly, PKS 1604+159 is also a weak 5 GHz sources in a large EGRET error ellipse.

Four of the ten "lower-confidence identifications" of TH95 are found to be very unlikely with p(id|r) < 0.005. All are modest 5 GHz sources well beyond the 95% contour of large EGRET error ellipses. On the other hand, two of the ten "lower-confidence identifications" of TH95 are found to be quite likely. With $p(id|r) \ge 0.95$, the identifications of 0458-020 and 0234+285 are found to be better than 55% of the "high-confidence identifications". Both are bright, flat-spectrum 5 GHz sources which have a low number density and the fraction of sources this bright at 5 GHz which are detected by EGRET is large.

Although TH95 classify PKS 1406-076 as a "high-confidence identification", their method would also allow it to be classified as a "lower-confidence" identification because PKS 1406-076 lies right on the 95% confidence contour of TH95 (see the ApJ CD ROM). We find that the identification of PKS 1406-076 is correct with 86% confidence. Although PKS 1406-076 is not near the center of the error region, the 95% confidence position contour contains only 0.2 square degrees. The variability of the EGRET flux and the finding of a possible correlation of the variability of optical flux with the γ -ray flux (Wagner *et al.* 1995) indicates strongly that the identification is correct.

Table 2 shows the analysis of §2 for the sources in the Supplement to the Second EGRET Catalog (Thompson *et al.* 1996). The three AGN identifications in the Supplement to the Second EGRET Catalog are found to be compelling.

3.3.1 The VLBI Flux Density of EGRET Blazars

We have examined the 2.29 GHz VLBI flux densities (Preston *et al.* 1985) for the sources in Tables 1&2 which either have $p(id|r) \ge 0.9$, or have $p(id|r) \ge 0.7$ for a EGRET source which is highly variable (PKS



Fig. 1.— The relationship of equation (3) for the probability of a correct identification p(id|r) as a function of the log of LR for the indicated values of η .



Fig. 2.— The measured *a priori* probability $\eta = p(id)$ of a radio source being detected by EGRET as a function of S_5 . The fit of equation 13 is also shown.

1406-076) or for which the confidence of the identification is otherwise enhanced (Mrk 421). Thirty of these 31 sources (all except PKS 1830-210) were measured by Preston *et al.* (1985). The 30 EGRET blazars are shown in comparison to the parent population in Figure 3. The marginal distribution of the VLBI flux density is shown in Figure 4 for the sources in Figure 3 with $\alpha \ge -0.5$. It is immediately apparent that the EGRET blazars have higher VLBI flux densities than the parent population. A Kolmogorov-Smirnov (KS) test shows that these distributions differ with 99.9994% confidence. VLBI upper limits were treated as detections in this KS test. This is conservative because Preston *et al.* (1985) obtained VLBI detections for all 30 of the EGRET blazars they observed.

We note that the amount of VLBI flux density is a much better discriminant of EGRET blazars than the VLBI core visibility. Figure 5 shows the distribution of the visibilities from the catalog of Preston *et al.* (1985) for the EGRET blazars. A KS test indicates with only 81% confidence that the EGRET blazars have higher visibilities than the parent population. The VLBI visibility is influenced by the amount of flux from the radio lobes. If the properties of radio lobes do not reflect whether γ -rays are visible, it is expected that the variance of lobe radio luminosity will cause the observed effect.

This is the first report of a radio property which distinguishes the EGRET blazars from the parent population of flat-spectrum radio sources. It supports arguments that the γ -ray emission is taking place at the base of a relativistic jet. This finding also has a practical application. A large VLBI flux density enhances the confidence of an EGRET blazar identification.

3.4 Identification Analysis of Previously Unidentified EGRET Sources

An obvious hypothesis is that some of the unidentified EGRET sources are also blazar type AGN. Our results are shown in Table 1 for the unidentified EGRET sources in the Second EGRET Catalog (TH95) and in Table 2 for the unidentified EGRET sources in the Supplement to the Second EGRET Catalog (Thompson *et al.* 1996). We find evidence of additional blazar identifications. The strongest new identification is lensed blazar PKS 1830–210 near the direction of the Galactic center with p(id|r) = 0.98. A total of four new potential identifications with p(id|r) > 0.5 appear in Tables 1 & 2. They include the ~2 Jy sources J1650–5044 and J1802–3940 which are within 10° of the Galactic plane and have not yet been sufficiently studied to be certain that they are not Galactic sources. These 4 likely identifications are given in Table 4 along with an additional 11 possible identifications with $p(id|r) \ge 0.05$. Two additional sources are also included in Table 4 which have 0.02 < p(id|r) < 0.05 and an indication of variability with confidence in excess of 99% (McLaughlin *et al.* 1996) — variable EGRET sources are more likely to be blazars.

3.5 Summary of Blazar Identifications

Table 3 gives the 42 EGRET blazar identifications which we expect to be robust. All of these sources have been established to be AGN. Most of these identifications have $p(id|r) \ge 0.9$. Several sources are included with 0.3 < p(id|r) < 0.9 if there is high confidence of variability of the EGRET source (in excess of 99%), or if a large VLBI flux density has been observed. Our minimum requirement is 300 mJy in the 2.29 GHz survey of Preston *et al.* (1985) or in a subsequent VLBI observation (details given in the footnotes of Table 3). From Figure 4, the requirement of a 2.29 GHz VLBI flux density > 300 mJy excludes 54% of potential confusing sources. Table 3 includes 33 of the 40 "high-confidence identifications" of TH95, 2 of 10 "lower-confidence identifications" of TH95, 1 new identification of an unidentified TH95 source,



Fig. 3.— A scatter plot of the 2.29 GHz VLBI flux densities and radio spectral indices from Preston *et al.* (1985) for the 1259 of 1398 sources in their catalog with radio spectral indices. Upper limits for VLBI flux densities are shown for 430 of these. Those which are robustly identified with EGRET sources (p(id|r) > 0.9, or p(id|r) > 0.7 for PKS 1406–076 and Mrk 421) are circled. The EGRET blazar shown with $\alpha = +1.3$ is 3C 454.3 — the index of Preston *et al.* (1985) probably differs from the GB survey value $\alpha = +0.1$ because of variability.



Fig. 4.— The marginal distribution of the VLBI flux densities of Figure 3 for $\alpha \ge -0.5$. The 882 sources in the catalog of Preston *et al.* (1985) with $\alpha \ge -0.5$ are shown with the histogram using thin lines. Upper limits are treated as detections (at the value of the upper limit). The peak at ~120 mJy results from this treatment of upper limits. The 30 EGRET blazars circled in Figure 3 are shown with the histogram using bold lines.

3 identifications from Thompson *et al.* 1996 (one of these, NRAO 190, was first identified by McGlynn *et al.* 1996), and 3 identifications from subsequent EGRET exposures. Table 4 comprises the potential identifications for which we expect further investigation will be most useful. Four of these potential identifications are very compelling with p(id|r) > 0.5. Some sources warrant specific comments.

B0115+0242: EGRET source 2EG J0119+0312 is identified with 6% confidence with radio source B0115+0242. Because there is a 92% confidence indication of variability for 2EG J0119+0312, this identification is actually more likely than 6%. The optical magnitude is 17.5. The red shift is 0.672. It shows OVV behavior. This source is also known as PKS 0115+02 and 3C 37. The $\alpha_{1.4-4.85}$ radio spectral index is -0.7. It would be worthwhile to conduct more radio and optical flux monitoring observations of this source to obtain a radio spectral index from simultaneous observations, to characterize variability, and to determine the amount of mm radio flux density. Also, it would be worthwhile to observe the parsec scale radio structure of this source, and determine a spectral index of possible compact radio emission.

0219+4248 (3C 66A): We find p(id|r) of 4.0% for the identification of EGRET source J0220+4228 with 3C 66A assuming a radio spectral index of -0.5. However, a radio index of ~ 0 . is indicated by 22 and 37 GHz monitoring (Teräsranta private communication). Using this radio spectral index, p(id|r) is found to increase from 4.0% to 6.2%. We note that 3C 66A is a BL Lac type blazar which shows optical variability and that the VLBI flux density at 2.29 GHz from Preston *et al.* (1985) is 0.27 ± 0.05 Jy. Thus, this identification is credible. However, Verbund *et al.* (1996) identify this source with binary ms pulsar J0218+4232 after finding $\sim 2\sigma$ evidence of modulation of the EGRET flux after epoch folding.

0336–019: Source 0336–019 (CTA 26) was detected by EGRET during the interval 05/09/95 - 05/23/95 (Mattox *et al.* 1995). The EGRET 95% confidence contour is well fit by an ellipse centered on l=188.25, b= -42.34, and with semi-major axis a=44', semi-minor axis b=37', and $\phi=27^{\circ}$. A complete description of multiwavelength observation is in preparation.

0446+112: This identification by TH95 is plausible, but narrowly fails to meet our criteria with p(id|r) = 0.31 and $S_{VLBI} = 200 \pm 20$ mJy. It is possible that this 2.29 GHz VLBI flux density measurement (Preston *et al.* 1985) is under representative because of variability.

1229–021: This identification by TH95 is plausible, but fails to meet our criteria with p(id|r) = 0.32 and $S_{VLBI} = 90 \pm 30$ mJy. It is possible that this 2.29 GHz VLBI flux density measurement (Preston *et al.* 1985) is under representative because of variability.

B1324+2226: EGRET source 2EG J1324+2210 is identified with 63% confidence with this GB source. The $\alpha_{1.4-4.85}$ radio spectral index indicated by the (non-simultaneous) GB survey flux densities is +1, very strongly inverted. Recent simultaneous cm radio measurements indicate a radio spectral index of ~0. (M. Aller, private communication). A VLBI position was published by Johnston *et al.* (1995) but a VLBI flux was not given.

J1650-5044: Our analysis shows a strong probability (92%) that this PMN source is EGRET source 2EG J1648-5042. However, observations at other wavelengths are required to ascertain that this radio source is an AGN rather than a Galactic radio source. This source was previously detected as radio source MRC 1646-506. Otherwise, it does not appear in the literature. An observation with the Australian Telescope Compact Array would be useful (it is too far south for the VLA).

J1802–3940: 2EG J1800-4005 is identified with PMN J1802–3940 with 81% confidence. Observations at other wavelengths are required to ascertain that this radio source is an AGN rather than a Galactic radio source.

1830–210: 2EG J1834–2138 is identified with PKS 1830–210 with 98% confidence. No VLBI observation has been reported for this source. This 8 Jy source has not been identified optically (it is 6° from the Galactic disk near the Galactic center). A VLA radio image shows what appears to be a steep-spectrum Einstein Ring and two parity-reversed images of a flat-spectrum core-jet structure separated by 1" (Jauncey *et al.* 1991). This structure is interpreted as a gravitationally lensed AGN jet. Thus, it could be an EGRET blazar. Wiklind and Combes (1996) established through mm observations of hydrocarbon absorption lines that the redshift of the lensing galaxy is z=0.89. If a detailed lens model can be developed, the autocorrelation function of future high-temporal density γ -ray flux measurements may reveal information about the location of the γ -ray emitting region. The expected time delay between the two components is ~10⁷ seconds (Jauncey *et al.* 1991).

1908–201: PKS 1908–201 is identified with 2EG J1911–1945 with 92% confidence. This 2 Jy source has not yet been identified optically (it is 21° from the Galactic center). The 8.4 GHz flux density of PKS 1908–201 is 2.3 Jy (Wright *et al.* 1991). Using the consequent index $\alpha = 0$ (instead of assuming $\alpha = -0.5$ without a 2.7 GHz detection) increases the probability of the identification to ~95%. The correlated VLBI 2.29 GHz flux density is 0.46 ± 0.05 Jy (Preston *et al.* 1985). It would be worthwhile to observe the the parsec scale radio structure of this source. An optical (or IR) identification and spectrum is also of interest. Preston *et al.* (1985) give a position which is accurate to 1": B1950, 19 8 12.6 -20 11 57. or J2000, 19 11 09.8 -20 06 57.

1933–400: This identification by TH95 is plausible, but narrowly fails to meet our criteria with p(id|r) = 0.34 and $S_{VLBI} = 280 \pm 30$ mJy. It is possible that this 2.29 GHz VLBI flux density measurement (Preston *et al.* 1985) is under representative because of variability.

2022–077: This identification by TH95 is plausible, but fails to meet our criteria with p(id|r) = 0.61 and $S_{VLBI} = 130 \pm 20$ mJy. It is possible that this 2.29 GHz VLBI flux density measurement (Preston *et al.* 1985) is under representative because of variability.

2155–304: An EGRET detection of this X-ray selected BL Lac type object was reported by Vestrand, Stacy and Sreekumar (1995). Since these authors do not provide a specification of the 95% confidence source location region, a likelihood analysis (Mattox *et al.* 1996a) of the VP 404 data for E> 100 MeV was done. The 95% confidence contour is well fit by an ellipse centered on l=17.26, b= -52.28, and with semi-major axis a=45', semi-minor axis b=40', and $\phi=79^{\circ}$. The analysis of §2 indicates that B2155–304 ($S_5=407$ mJy, $\alpha = 0.3$) can be identified with the EGRET source with p(id|r)=12% confidence. A test for variability of the EGRET source using the average flux of phase 1&2 of the mission, VP 209, and VP 404 results in $\chi^2=7.3$ for 2 DOF. Thus, a constant γ -ray flux can only be rejected at the 97.5% confidence level. Preston *et al.* (1985) measured a 2.29 GHz VLBI flux density of 144±8 mJy. The identification is very plausible, but fails to meet our criteria for inclusion in Table 3.

2200+420: A 4.4 σ EGRET detection of 2200+420 (BL Lacertae) is reported by Catanese *et al.* (1997). The 95% confidence contour for the event selection E> 100 MeV is well fit by an ellipse centered on l=92.59, b= -10.44, and with semi-major axis a=89', semi-minor axis b=58', and $\phi=35^{\circ}$. The analysis of §2 indicates that B2200+420 ($S_5=3.6$ Jy, $\alpha = -0.2$) can be identified with the

EGRET source with p(id|r)=99.3% confidence. A high confidence identification is obtained in spite of the large error region because the 5 GHz flux density is large.

3.6 Limitations of this Analysis

Although it is a significant improvement over previous EGRET identification work, we note that our technique is subject to a number of potential sources of systematic error. The error ellipses of TH95 indicate statistical error only. Substantial systematic error is also expected, but was not given because of the difficulty of quantifying it. As discussed by TH95 and Mattox *et al.* (1996a), error in the model of the Galactic diffuse γ -ray emission and undetected nearby γ -ray point sources can have a substantial effect on EGRET position estimates. Because the present investigation is limited to $|b| > 3^\circ$, the effect of error in the Galactic diffuse model of Hunter *et al.* (1996) is not severe for the EGRET sources we study in this paper.

The radio flux densities of blazars vary by factors as large as ~10. Therefore, it is problematic to use the radio survey flux for the identification analysis because the flux at the survey epoch could differ significantly from the average flux. Also, the radio spectral index we use was obtained from non-simultaneous surveys at 4.85 GHz and 1.4 GHz. It is subject to error because of flux variability. The error that this introduces can be mitigated by follow-up radio observations of sources which are potential EGRET source identifications. For this reason, potential identifications with values of p(id|r) as small as ~0.05 should not be dismissed until follow-up observations have been made.

The necessary simplification of assuming that the *a priori* probability, η , is independent of EGRET exposure and radio spectral index also contributes some error to our estimates of the probabilities of correct identifications, but we expect that they are not severely compromise. An additional minor error in our identification analysis may result from some deviation of the dependence of the log of the likelihood of the EGRET data on the source position from the assumed paraboloid. Also, a minor error in our identification analysis is introduced by the discernible but small differences between the elliptical fits and the actual position confidence contours of TH95 which are available on the ApJ CD ROM corresponding to TH95.

For these reasons, the probabilities of correct identification derived here are indicative rather than definitive. We expect that for identifications with an *a posteriori* probability p(id|r) > 0.9, 1-p(id|r) could be in error by as much as a factor of 5. Thus, p(id|r) = 0.9 implies 0.5 < p(id|r) < 0.98. Similarly we expect that for identifications with an *a posteriori* probability p(id|r) < 0.1, p(id|r) could also be in error by as much as a factor of 5. Thus, p(id|r) = 0.9 implies 0.02 < p(id|r) < 0.1, p(id|r) could also be in error by as much as a factor of 5. Thus, p(id|r) = 0.1 implies 0.02 < p(id|r) < 0.5. Although the large uncertainty of p(id|r) is not ideal, it is not so large as to render this analysis useless. We expect that our method offers an order of magnitude improvement in precision over that of TH95.

For latitudes less than ~15°, a substantial fraction of the radio sources will be Galactic radio sources. We search for counterparts in this range because correct blazar identifications could result nonetheless. In fact, one planetary nebulae and one H_{II} region appear in Table 1 (and are noted in the footnotes), but are not thought to be γ -ray source counterparts. We find that no excess of compelling identifications results at low latitudes. Therefore, undiscerned Galactic radio sources are apparently not a major problem for Galactic latitudes $3^{\circ} < |b| < 15^{\circ}$.

We do not quantitatively incorporate in our identification analysis the variability of EGRET sources, nor the amount of correlated VLBI flux density — although these certainly pertain. They are considered in the selection of the high probability EGRET identifications of Table 3 and the "most interesting" potential identifications of Table 4.

Also, there is potentially more information for some sources which would be difficult to quantitatively incorporate in our identification analysis. We note that the analysis described in §2 results in a formal probability of 70% that MRK 421 is correctly identified as an EGRET source. This results from the low radio flux of MRK 421, and the large EGRET position uncertainty for this weak source. Additional information drastically improves the confidence of this identification. MRK 421 has distinguishing characteristics which effectively reduce the number density of such sources. In addition to a substantial correlated VLBI flux density and domination of the optical emission by continuum synchrotron radiation, MRK 421 shows a large and highly variable X-ray flux. It has been detected at TeV energies through the observation of Čerenkov light produced in the atmosphere by γ -ray induced electromagnetic cascades (Buckley *et al.* 1996). The TeV flux is highly variable and it has been observed to vary in correlation with the X-ray flux (Buckley *et al.* 1996). The TeV identification is very secure because the position of the TeV emission has been determined to $\sim 4'$ and is consistent with the MRK 421 position (Buckley *et al.* 1996).

The identifications of Table 1 indicate that the OVV property is common to many of the sources detected by EGRET which have been frequently observed with optical telescopes. Thus, the OVV property may serve to distinguish sources and improve identification confidence. This may also be true for optical polarization and BL Lac classification. However, careful work remains to be done to ascertain that these characteristics distinguish EGRET sources. Wagner (in preparation) finds that 80% of all sources which have a flat radio spectrum up to 90 GHz are OVV irregardless of whether or not they are detected by EGRET. The fraction of EGRET sources which are believed to be OVV is not significantly larger than this; however not all have been yet been extensively monitored in the optical band. Also, the fraction of blazars detected by EGRET which are radio-selected BL Lac type objects is not significantly different from the fraction of flat-spectrum radio sources which are BL Lacs. Also, optical polarization may be generic for flat-spectrum radio sources. Fugmann (1988) has studied optical polarization observations of the Kühr *et al.* (1981) sample. He finds that the probability of measuring significant optical polarization for a flat-spectrum radio sources would have the property of high optical polarization in at least one observation if sufficient observations were made.

4. Discussion

We find that a substantial number of the identifications of TH95 are far from certain. This has implications for most studies of the "EGRET blazars". It impacts the study of the γ -ray luminosity function and evolution (Chiang *et al.* 1995). It should be considered in studying the correlation of γ -ray and radio luminosity (Padovani *et al.* 1993; Stecker, Salamon, and Malkan 1993; Dondi and Ghisellini 1995; Mücke *et al.* 1996), and in the use of the EGRET detections in calculations of the production of isotropic diffuse high-energy γ -ray background by AGN (Padovani *et al.* 1993; Stecker *et al.* 1993; Salamon and Stecker 1994; Chiang *et al.* 1995; Stecker and Salamon 1996; Kazanas and Perlman 1996). Careful consideration of the confidence of the identification is also in order when seeking distinguishing properties of the "EGRET blazars". Unless explicit consideration is given to potential misidentification, we suggest that these analyses be restricted to the 42 EGRET blazars of Table 3 for which p(id|r) exceeds 90%, or a high confidence identification is otherwise indicated. We estimate that ~1 of these identifications is specious. Additional EGRET exposure and additional analysis and observation at other wavelengths may eventually contribute ~10 more sources to this population.

4.1 On the Detection of Blazars by EGRET

We have demonstrated in §3.1 with confidence far beyond question that EGRET is detecting radio loud AGN. Six of the identifications in Table 1 have a probability of being correct of at least 99.9% (0420–014, 0528+134, 3C 273, 3C 279, 1633+382, and 3C 454.3) All of these have flat radio spectra ($\alpha \geq -0.5$). All have substantial correlated VLBI flux density. All 6 show significant changes in optical flux on short time-scales. Of these 6, the 4 which have been extensively observed show significant optical polarization. Also, the 5 which have been observed sufficiently with VLBI to discern it show superluminal motion of the parsec scale structure of radio jets. Therefore, it is apparent that EGRET is detecting the blazar class of AGN.

The other identifications of Table 3 all have radio spectra with $\alpha \geq -0.5$, except for 0521–365 for which $\alpha = -0.7$ from the PMN catalog. This source has not been extensively observed. The report of an EGRET detection lead to VLBI observations which reveal that the spectrum of the core is $\alpha_{4.9-8.4} = +0.3$ (Tingay *et al.* 1996). Thus, if this source is truly steep-spectrum in single dish observations, it is due to the flux of the radio lobes. Therefore, it appears that all high confidence EGRET identifications are flat-spectrum radio sources. Fugmann (1988) concludes from his study of the optical polarization that at least 2/3 of all flat-spectrum radio sources are HPQ sources and therefore blazars. Therefore, it is plausible that all extragalactic EGRET source are blazars.

It is interesting that EGRET is detecting only ~10% of the blazars. It is possible that this results from time variability of the gamma-ray emission. However, a "duty cycle" of ~30% is observed for the strong EGRET sources — too large to explain the fact that only 10% of the blazars are detected, unless there is strong variability on time scales longer than ~1 year. Another possibility is that the gamma-ray luminous blazars differ intrinsically from the blazar parent population. E.g., the gamma-ray emission may be more narrowly beamed than the longer wavelength emissions so that the γ -ray blazar sub-population has a smaller angle between the jet and the line of sight than the blazar parent-population (Salamon & Stecker 1994, Dermer 1995).

4.2 Non-Detection of Steep Radio Spectrum and Radio Quiet AGN by EGRET

In Figure 6 we present a scatter plot of S_5 and α for all radio sources considered. The identifications of Table 1 & 2 with $p(id|r) \ge 0.02$ are surrounded by circles. The size of the circle increases with the confidence of the identification. It appears that the distribution of observed spectral indices for the identified sources is consistent with that of flat-spectrum sources ($\alpha > -0.5$) given the errors in the spectral indices due to the non-simultaneity of the radio surveys. Thus, it is plausible that EGRET is detecting only flat-spectrum radio sources.

It is a possible concern that the non-detection of steep-spectrum sources results from our use of the reduced number density of flat-spectrum radio sources in the calculation of the likelihood ratio. Therefore, we have repeated the analysis described in §2 with a number density of radio sources $\rho(S_5)$ which depends only on flux density. This results in values of LR smaller by a factor of ~2. However, significant detections persist and the list of strong identifications is nearly unchanged. The sum for the sources in Table 1 is then $\Sigma p(id|r) = 31.7$ (down from 36.6 with the radio index dependent number density described in section 2). In the null hypothesis, the Monte Carlo calculation with random EGRET positions described in section 3.1 yields $\Sigma p(id|r) = 4 \pm 2$. Thus, the result that EGRET is detecting radio loud AGN does not depend upon the assumption that the flat spectrum sources are more likely to be detected. It follows that the steep-spectrum sources are generally not being detected. Therefore, the use of the spectral index in the



Fig. 5.— The distribution of the visibilities (VLBI flux density/total flux density). The 451 sources in the catalog of Preston *et al.* (1985) with $\alpha \ge -0.5$ and a visibility are shown with the histogram using thin lines. Upper limits are treated as detections at the upper limit of the visibility. The 19 EGRET blazars with visibilities in the catalog of Preston *et al.* (1985) are shown with the histogram using bold lines.



Fig. 6.— The distribution of radio flux density (S_5) and radio spectral index (α) for all sources in the PMN and GB surveys with $S_5 > 100$ mJy. We plot an index of $\alpha = -0.5$ if an index is not available from the survey catalogs (as assumed for the analysis of §2). The consequent enhanced source density at $\alpha = -0.5$ is apparent. The identifications of Table 1 and Table 2 with $p(id|r) \ge 0.02$ are surrounded by circles (using α from Table 1 or 2). The size of the circle increases with p(id|r). The five different sizes correspond to the following ranges: $0.02 \le p(id|r) < 0.1$, $0.1 \le p(id|r) < 0.5$, $0.5 \le p(id|r) < 0.9$, $0.9 \le p(id|r) < 0.99$, $0.99 \le p(id|r) < 1.00$.

calculation of confusing source number density is justified.

We also suggest that EGRET is not detecting radio-quiet AGN. If an unidentified EGRET source is actually a blazar, that blazar is very likely to be among the possible identifications given here. If the EGRET flux density is variable, the likelihood of a blazar identification is enhanced. We note that every variable EGRET source in Table 1 (with variability index ≥ 2 corresponding to confidence of variability $\geq 99\%$), has a plausible radio-loud identification. If we assume that the γ -ray emission of all AGNs would vary, we can conclude that EGRET is not detecting any radio-quiet AGN. We note that there are several unidentified EGRET sources at $|b| < 3^{\circ}$ which are excluded from our analysis which are probably Galactic (e.g. Tavani *et al.* 1997).

4.3 Suggested Observations at Other Wavelengths of Potential Gamma-Ray Selected Blazars

Although we have found only one new EGRET identification in this analysis (PKS 1830–210) which qualifies for inclusion in the high confidence identifications of Table 3, it is apparent that a substantial number of the previously unidentified EGRET sources are also blazars. Further study at other wavelengths (primarily radio and optical) is required to ascertain whether or not some of these candidates have blazar properties. Most of these potential EGRET sources have not been studied extensively. Upon further investigation, they may be found to be "gamma-ray selected blazars". In Table 4 we list the 17 "most interesting" new potential identifications. This table also gives the best positions available for these sources.

Further radio observation can establish variability and a radio spectrum from simultaneous observations. Radio interferometric images (VLBI, or VLA/AT) can be obtained to establish that the radio source is an AGN. A large correlated VLBI flux density is indicative of EGRET sources. Measurements of flux densities in the mm radio band may help to distinguish blazars. Measurements of optical variability and polarization can be used to establish membership in the OVV or HPQ blazar class. Spectra to establish redshifts, and equivalent widths of emission lines are of interest, as are X-ray and infrared observations. Simultaneous multiwavelength observation will be of interest once identifications are firmly established. In principle, the identification could be secured by observation of gamma-ray flux variation which correlates with variation at a longer wavelength. However, the EGRET sensitivity is generally not sufficient for this unless the γ -ray flux is exceptionally large.

In many cases, an optical counterpart is obvious from an examination of the POSS (or ESO) plates for the GB or PMN error region. In other cases, an optical counterpart is not clearly apparent in this $\sim 20''$ region because of source confusion. Radio observations to obtain an interferometric position with the VLA or the Australian Telescope Compact Array (AT) are then appropriate. Positions accurate to $\sim 0.01''$ are given in Table 4 for sources of likely interest which were observed by Patnaik *et al.* (1992 & 1996) with the VLA for potential use as MERLIN phase calibrators.

4.4 Pulsar Candidates

The non-variable EGRET sources for which a radio-loud identification is not apparent are more likely than other EGRET sources to be pulsars. Tables 1&2 may thus be used to to select optimal pulsar candidates. Romani and Yadigaroglu (1995) predict that 70% of gamma-ray luminous pulsars are in fact radio-quiet. Direct detection of periodicity for more than ~ 2 through a FFT of the EGRET data is unlikely (Mattox *et al.* 1996b). More work is required to determine if the predicted spatial distribution of pulsars (Yadigaroglu and Romani 1996) matches that of the pulsar candidates of Tables 1&2.

We note that both EGRET sources in Table 1 which coincide with SNRs (2EG J0008+7307 and 2EG

J0618+2234) have no flat-spectrum radio identifications with $p(id|r) > 2 \times 10^{-6}$. This is consistent with the claim (Sturner and Dermer 1994, Esposito *et al.* 1996, Sturner, Mattox, and Dermer 1996) of a correlation between EGRET sources and SNRs. This correlation could be either the result of enhanced interaction of cosmic rays with the interstellar medium near SNR, or pulsars associated with the SNRs or with young stellar associations co-located with a SNR.

4.5 The Correlation of Gamma-ray Flux with Radio Flux

With the limited EGRET data, the dependence of the *a priori* probability, η , on S_5 cannot be understood in detail. A simple hypothesis is that ~20% of all flat spectrum radio sources emit γ -rays, and that the observed decrease in η with radio flux is the result of detecting a fraction of γ -ray blazars which decreases with radio flux because γ -ray flux is correlated with radio flux and EGRET has a relatively high detection threshold. The initial reports of a correlation between γ -ray luminosity and radio luminosity (Padovani *et al.* 1993; Stecker *et al.* 1993; Salamon and Stecker 1994; Stecker and Salamon 1996) are flawed in that they fail to consider the effect of a common redshift on the apparent correlation. An analysis which overcomes this defect through partial correlation analysis (Dondi and Ghisellini 1995) indicated a correlation with 99.5% confidence. However, a partial correlation analysis which also considers upper limits (Mücke *et al.* 1996) does not find significant correlation. We present a simple argument below which establishes with 99.998% confidence that the maximum observed γ -ray flux of EGRET blazars correlates with the average 5 GHz radio flux.

We simply note that all of the EGRET blazars which are γ -ray bright are also bright radio sources. The ten EGRET blazars with a peak γ -ray flux above 10^{-6} cm⁻²s⁻¹ at E>100 MeV (0208-512, CTA 26, 0528+134, 1156+295, 3C 279, 1406-076, 1622-297, 1633+382, 1730-130, and 3C 454.3) all have an average radio flux density S_5 which exceeds 1.0 Jy. From the GB and PMN catalogs, we estimate that 58%of the $S_5 > 1.0$ Jy radio sources have flat spectra ($\alpha \geq -0.5$), and that there are 183 of them away from the Galactic plane ($|b| > 3^{\circ}$). The fraction detected as bright γ -ray blazars is 0.05±0.02. Using Poisson statistics, a 99.998% confidence lower limit on the fraction of these which are bright γ -ray sources is 0.0098. Examining the GB and PMN catalogs over narrow intervals ($250 < S_5 < 350$ mJy, $350 < S_5 < 500$ mJy, and $500 < S_5 < 1000$ mJy) and correcting for the variable fraction of sources which have measured spectral indices, and the variable fraction which are flat spectrum; we estimate that there are 1130 flat spectrum radio sources with 250 mJy $< S_5 < 1.0$ Jy away from the Galactic plane ($|b| > 3^\circ$). If the γ -ray flux of EGRET sources were not correlated with radio flux, we would expect $5\pm 2\%$ or 62 ± 20 of these to be bright γ -ray sources. Using the 99.998% confidence lower limit on the fraction detected, at least 11.1 of these radio sources should be bright γ -ray sources. None are observed. The Poisson probability of observing 0 when 11.1 are expected is 1.5×10^{-5} . We thus reject the hypothesis that the γ -ray flux of EGRET sources is not correlated with radio flux with 99.998% confidence. It is possible that the correlation is non-linear as suggested by Mücke et al. (1996).

This analysis assumes that the distribution of α is the same for the sources where it was not measured as it is for sources where it was measured. This may not be true in detail, but it is expected to be a good approximation for sources with $S_5 > 250$ mJy. We have effectively analyzed the correlation between the maximum observed γ -ray flux of EGRET blazars and the average 5 GHz radio flux because that is essentially what the current data offers. Most EGRET blazars are only detected by EGRET during a flaring state. And the GB and PMN surveys with a 5 GHz observation at one epoch offer approximately an average flux.

If bright EGRET blazars with 250 mJy $< S_5 < 1.0$ Jy existed, the method of §2 could identify them

because the γ -ray error regions would still be small. We analyzed the 10 bright EGRET sources listed above with the counterpart radio flux artificially altered to $S_5=250$ mJy. All identifications remain above the threshold for inclusion in Table 1 (p(id|r) > 0.001), and 9 of 10 above the threshold for variable EGRET sources in Table 4 (p(id|r) > 0.02). Two sources (3C 279 and 0528+134) would still be above the threshold for variable EGRET sources in Table 3 (p(id|r) > 0.3). This a mute point however. There are no bright EGRET sources (peak γ -ray flux above 10^{-6} cm⁻²s⁻¹, E>100 MeV) at high Galactic latitude ($|b| > 3^{\circ}$) which do not coincide with either a pulsar or a radio source with $S_5 > 1.0$ Jy. In fact, this argument can be extended to the myriad radio sources with $S_5 < 250$ mJy. The prevalent interpretation of the isotropic high-energy γ -ray diffuse radiation is that it is due to the γ -ray emission of these sources (Padovani *et al.* 1993; Stecker *et al.* 1993; Salamon and Stecker 1994; Chiang *et al.* 1995; Stecker and Salamon 1996; Kazanas and Perlman 1996). If this interpretation is correct, there must be a correlation between γ -ray and radio flux or else the weakest radio sources would also be detected as bright γ -ray sources.

We note that a Spearman rank-order correlation test of the sources in Table 3 for the 5 GHz radio flux of Table 3 and the peak γ -ray flux (without doing a K-correction) suggests a correlation (with 67% confidence). The same test shows a correlation with 99.2% confidence for the peak γ -ray flux and correlated VLBI flux of Table 3. This is consistent with the finding of §3.3.1 that the EGRET blazars are distinguished by a large VLBI flux density; and adds more weight to arguments that the γ -ray emission is taking place at the base of a relativistic jet. A tighter correlation might be obtained using simultaneous γ -ray and VLBI fluxes (perhaps allowing for a delay in VLBI flux because of synchrotron self-absorption of radio flux at the base of the jet). We note that the same analysis with the non-simultaneous 8.4 GHz VLBI flux densities of the 28 sources in Table 3 which were measured by Morabito *et al.* (1986) shows much less correlation (the Spearman rank-order correlation test indicates a 34% probability of a correlation). To some extent this is caused by the reduced number of sources included in the correlation analysis. Perhaps it also reflects higher radio variability at higher frequency.

5. Summary

We present a means to quantitatively assess the probability that an identification of an EGRET source with a radio source is correct. We also demonstrate conclusively that EGRET is detecting γ -ray emission from the blazar class of AGN. Our analysis suggests that EGRET is not detecting radio quiet and steep radio spectrum AGN. This identification method is used to ascertain with 99.998% confidence that the γ -ray and radio flux of EGRET blazars are correlated. Previous EGRET identifications are re-examined. Table 3 gives the 42 EGRET blazars for which the confidence of a correct identification is high. It is found that the peak γ -ray flux and the VLBI radio flux of these sources is correlated with 99% confidence. One high confidence AGN identification (PKS 1830-210) is found among the unidentified EGRET sources of TH95. Sixteen additional potential identifications are presented in Table 4 with the best available position for study at other wavelengths.

Acknowledgments

This research has made extensive use of the NASA/IPAC Extragalactic Database (NED) which is operated by the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration. J. Mattox acknowledges support from NASA Grants NAG5-2833 and NAG5-3384.

REFERENCES

- Becker, R.H., White, R.L., and Edwards, A.L., 1991, Ap.J.S., 75,1
- Bolton, J.G., Wright, A.E., and Savage, A., 1979, AuJPA, 46
- Bloom, S.D., et al., 1997, ApJ S, submitted
- Brinkman, W., et al., AA,281, 355
- Buckley, J., et al., 1996, ApJL, in preparation
- Catanese, M., et al., 1997, ApJ, in press
- Chiang, J., et al., 1995, ApJ, 452, 156
- Condon, Broderick, and Seielstad, 1989, A.J., 97, 1064
- Dermer, C.D., 1995, ApJL, 446, L63
- Dingus, B.L., et al., 1996, ApJ, 467,589
- Dondi, L., Ghisellini, G., 1995, MNRAS, 273, 583
- Esposito, J.A., et al., 1996, ApJ, 461, 820
- Fichtel, C.E., et al., 1994, ApJ S, 94, 551
- Fich, M., 1993, ApJS, 86, 475
- Fugmann, W., 1988, A&A, 205,86
- Griffith, M.R., and Wright, A.E., 1993, A.J., 105, 1666
- Gregory, P.C., Condon, J.J., 1991, ApJ Sup., 75, 1011
- Halpern, J, 1996, private communication.
- Hartman, R.C., et al., 1992, ApJ L, 385, L1
- Hewitt, A., & Burbidge, G., 1987, ApJ S, 63, 1
- Hunter, S.D., Digel, S.W., de Gues, E.J., Kanbach, G., 1994, ApJ 436, 216
- Hunter, S.D., et al., 1996, in preparation
- Jauncey, D. et al., 1991, Nature, 352, 132
- Johnston, K.J. et al., 1995, AJ, 110, 880
- Kazanas, D. and Perlman, E., 1996, in preparation
- Kellerman, K.I., & Wall, J.V., 1987, in Observational Cosmology, IAU Symp. #124, eds. A. Hewitt, G. Burbidge, & L. Z. Fang (Dordrecht: Reidel), p. 545
- Kuehr, H., Witzel, A., Pauliny-Toth, I.I.K., and Nauber, A., 1981, Astr. Ap. Suppl., 45, 367
- Loredo, T.J., 1990, in Maximum-Entropy and Bayesian Methods, ed. P. Fougere, Kluwer, Dordrecht, p. 81
- Lundgren, S.C., et al., 1995, IAUC #6258
- Mattox, J.R., et al., 1995, IAU circ. #6161
- Mattox, J.R., et al., 1996a, ApJ, 461, 396
- Mattox, J.R., et al., 1996b, AA S, 120, xxx
- Mattox, J.R., et al., 1997, ApJ, 476, xxx
- McGlynn, T. A., et al., 1996, ApJ, submitted

- McLaughlin, M. A., Mattox, J. R., Cordes, J. M., and Thompson, D.J., 1996, ApJ, in press
- Mücke, A., et al., 1996, in press
- von Montigny, C., et al., 1995, ApJ, 440, 525
- Morabito, D.D., et al., 1986, Ast. J., 91, 1038
- Mukherjee, R., et al., 1995, ApJ, 445, 189
- Nolan, P., et al., 1996, ApJ, 459, 100
- Patnaik, 1996, private communication
- Patnaik, A.R., Browne, I.W.A., Wilkinsen, P.N., Wrobel, J.M., 1992, MNRAS, 254, 655
- Preston, R.A., et al., 1985, AJ 90,1599
- de Ruiter, H.R., Willis, A.G., and Arp, H.C., 1977, AA Sup., 28, 211
- Padovani, P., Ghisellini, G., Fabian, A.C., and Celotti, A., 1993, MNRAS, 260, L21
- Romani, R.W., and Yadigaroglu, I.A., 1995, ApJ 438, 314
- Salamon, M.H. and Stecker, F.W., 1994, ApJ L., 430, L21
- Schachter, J., et al., 1997, in preparation
- Stickel, M., and Kuehr, H., 1994, AAS, 105, 211
- Stecker, F.W., Salamon, M.H., and Malkan, M.A., 1993, ApJ L., 410, L71
- Stecker, F.W. and Salamon, M.H., 1996, ApJ, 464, 600
- Sturner, S.J., Dermer, C.D., 1995, AA L, 293, 17
- Sturner, S.J., Mattox, J.R., Dermer, C.D., 1996, AA S, 120, xxx
- Sturrock, P.A., 1973, Ap.J., 182, 569
- Tavani, M., 1997, ApJ, submitted
- Thompson, D.J., 1986, NIM, A251, 390
- Thompson, D.J., et al., 1993, Ap J S, 86, 629
- Thompson, D. J., et al., 1995, Ap J S, 101, 259 (TH95)
- Thompson, D. J., et al., 1996, Ap J S, 107, xxx
- Tingay, S.J., 1996, Ap.J., 464, 170
- Verbund, F., et al., 1996, AAL, in press
- Vestrand, W.T., Stacy, J.G., Sreekumar, P., 1995, ApJL, 454, L93
- Wagner, S., et al., 1995, ApJ L, 454, L97
- Wiklind, T., and Combes, F., 1996, Nature, 379, 11
- White, R. L. and Becker, R. H., 1992, Ap.J.Supp., 79, 331
- Wolstencroft, R.D., MacGilivray, H.T., Lonsdale, C.J., Conrow, T., Yentis, D.J., Wallin, J.F., & Gau, G. 1992, in Digitized Optical Sky Surveys, eds. H. T. MacGilivray and E. B. Thomson (Dordrecht: Kluwer), p. 471.
- Wright, A.E., et al., 1991, MNRAS, 251,330
- Wright, A.E., Griffith, M.R., Hunt A., Troup E., Burke, B.F., and Ekers, R.D., 1996, in preparation

This preprint was prepared with the AAS ${\tt IATEX}$ macros v4.0.