FAUST observations of UV sources toward the Virgo cluster

Noah Brosch¹, Liliana Formiggini^{2 3} and Elchanan Almoznino⁴

The Wise Observatory and the School of Physics and Astronomy

Raymond and Beverly Sackler Faculty of Exact Sciences

Tel Aviv University, Tel Aviv 69978, Israel

and

Timothy Sasseen ⁵, Michael Lampton ⁶ and Stuart Bowyer ⁷ Space Sciences Laboratory and Center for EUV Astrophysics University of California Berkeley CA 94720, U.S.A.

Received ____

accepted _

ABSTRACT

We analyze three UV images covering a ~ 100 square degree field toward the Virgo cluster, obtained by the FAUST space experiment. We detect 191 sources to a signal-to-noise ratio of 4.4 and identify 94% of them. Most sources have optical counterparts in existing catalogs and about half are identified as galaxies. Some sources with no listed counterpart were observed at the Wise Observatory. We present the results of low resolution visible spectrophotometry and discuss the foreground 101 stellar sources and the 76 detected galaxies, both in the cluster and in the fore- or background. We derive conclusions on star-formation properties of galaxies and on the total UV flux from discrete and diffuse sources in the cluster. We test for the presence of intra-cluster dust, determine the clustering properties of UV emitting galaxies, and derive the UV luminosity function of Virgo galaxies.

Key words: Ultraviolet: Stars, Galaxies – Galaxies: Clusters: Individual: Virgo – ISM: General

1. Background

The ultraviolet (UV) segment of the spectrum is an excellent tool to identify and quantize active star formation (SF) regions (Donas *et al.* 1987). It is also a sensitive probe of evolved hot stellar populations (*e.g.*, Burstein *et al.* 1988). Therefore, UV observations may serve to understand processes of SF and stellar evolution. These processes are amplified and enhanced in a cluster environment, due to interactions among galaxies, and between galaxies and the intracluster medium. In the case of the Virgo cluster (VC) the observations are optimized by the proximity of the region, despite the VC not being a fully relaxed cluster. Throughout this work we **assume** a uniform distance of 18 Mpc to all the galaxies in the VC. A smaller cluster distance, ~ 14 Mpc, is suggested by the [OIII] distances to planetary nebulae (Jacoby *et al.* 1990), but Visvanathan and Griersmith (1979) give 17.9 Mpc, in agreement with Sandage and Tammann (1974). We note that the depth of the cluster (~4 Mpc) is significant in comparison to the distance to the cluster.

The VC was studied intensively by many groups; we shall not give here a thorough review of all publications but mention in particular the extensive compilation of Binggeli, Sandage and Tammann (1985, BST). Their VC catalog relied mainly on the morphological appearance of a galaxy on large plate-scale photographs, to determine whether an object is member of the VC or lies in its fore- or background. In addition, the many papers by Hoffman and collaborators (Hoffman *et al.* 1989a, 1989b; Hoffman 1989; Hoffman *et al.* 1995) used HI observations from the Arecibo Observatory to determine the large-scale-structure in the direction of the VC. The combination of HI and optical redshifts for many of the galaxies indicates the existence of a void in the distribution of galaxies located behind the VC and in front of the Great Wall at $5500 \le cz \le 13500 \text{ km s}^{-1}$.

The complex structure of the VC region was analyzed by Hoffman *et al.* (1989a) in the context of the blue compact dwarf galaxies (BCDs) in the cluster. Their Fig. 4 emphasizes the positions of the various groups and clouds of galaxies in this region. They also established upper limits to the number of HI clouds devoid of optical counterparts in the cluster, and considered other observational properties of these galaxies, such as their FIR emission from IRAS and optical properties. Hoffman *et al.* (1995) studied the background galaxies beyond the VC, in the context of their clustering properties.

A reanalysis of the velocity distribution of VC galaxies (Binggeli, Popescu and Tammann 1993) showed a slight difference in the cluster redshift (1050±35 km s⁻¹) from the previously accepted value (1094 km s⁻¹; Binggeli, Tammann and Sandage 1987). In addition to the void behind Virgo mentioned by Hoffman *et al.* (1995), they found a significant lack of galaxies behind Virgo, from 3000 to 3500 km s⁻¹. We consider here all galaxies in the direction of the VC and with $v_{\odot} < 3000$ km s⁻¹ as cluster members, disregarding their membership in the Virgo core or the various sub-clusters.

X-ray observations with the ROSAT PSPC (Trinchieri *et al.* 1994) showed that at least one Virgo cluster galaxy (NGC 4636) may be interacting with VC gas to produce significant X-ray flux at \sim 3 Mpc from M87. Other such galaxies may be M86 (Forman *et al.* 1979) and possibly NGC 4472 (Trinchieri *et al.* 1986). In addition, we mention the discovery of very soft X-ray emission from M87 (Lieu *et al.* 1996) by the EUVE satellite, with 31 counts per 1000 sec in the 100Å band. The X-ray and far-infrared emission from M87, together with subtle optical asymmetries seen mainly in the blue, induced White *et al.* (1991) to propose that this galaxy is stripped of its interstellar material by some hot gas, present at the core of the VC.

High resolution HST observations of Ly α absorptions in the UV spectrum of 3C273 at the redshift of the VC (Weymann *et al.* 1995 and references therein) indicate that the absorptions are probably produced by low-column-density clouds with low metallicity. One need they clearly identify is to evaluate the metagalactic background ionizing radiation field, away from the vicinity of normal galaxies; we attempt this by using new UV observations from approximately the entire Virgo cluster, obtained by the FAUST payload.

1.1. Previous UV observations

A few galaxies in the Virgo cluster were observed in the vacuum UV by the ANS satellite (Wesselius *et al.* 1982). More were observed by the rocket payloads of Smith and Cornett (1982) and Kodaira *et al.* (1990), and recently by the FAUST imager (see section 2, below) on board the ATLAS–1 Shuttle experiment (Bowyer *et al.* 1993). Additional measurements from a balloon–mounted telescope were reported by Donas *et al.* (1987).

Smith and Cornett (1982) observed the cluster with a wide field camera (11°.4 FOV) with a CsTe intensifier coupled with photographic film behind a quartz window, which define a passband with FWHM \simeq 1150Å and constant–energy mean wavelength of 2421Å. Their absolute calibration relies of ρ Vir, calibrated during the TD–1 mission against Vega, and UV magnitudes or upper limits are given for 201 galaxies.

Donas *et al.* (1987) used the SCAP 2000 balloon-borne telescope to image various regions of the sky in a band ~ 125 Å wide centered near 2000Å. The SCAP telescope was a 17 cm diameter reflector with a 6° field-of-view, imaging on film behind a UV-to-visible image converter/amplifier. Among the galaxies listed by Donas *et al.* 12 are members of the Virgo cluster.

The GUV experiment (Onaka *et al.* 1989) consisted of two 17 cm diameter imagers with a 4° field of view and was launched in 1987 on board a sounding rocket. The GUV sensitivity response had a FWHM $\simeq 200$ Å and peaked near 1450Å. The results relevant to Virgo cluster galaxies were presented by Kodaira *et al.* (1990) and a discussion of the diffuse UV light (mainly galactic) in the direction of the Virgo cluster was given by Onaka

and Kodaira (1991). The galaxies' paper gives UV fluxes for 42 galaxies observed by GUV and upper limits for another 47 galaxies. Note that the reported final angular resolution of the GUV experiment was $16'.0 \times 8'.2$ (Kodaira *et al.* 1990) and the equivalent exposure time on the Virgo cluster was 176 sec.

2. FAUST observations of the Virgo cluster

FAUST (FAr Ultraviolet Space Telescope) is a wide-field (~ 8° field of view) telescope with a bandpass between 1400 and 1800Å and an angular resolution of 3.5 arcminutes. It utilizes a microchannel plate detector with wedge and strip anode, which records the position of each detected photon. Descriptions of FAUST and its operation aboard ATLAS-1 were given by Lampton *et al.* (1993) and Chakrabarti *et al.* (1993). Details of the image construction and subsequent reductions are given in Bowyer *et al.* (1993). A catalog of 4660 UV sources detected by FAUST was produced by Bowyer *et al.* (1995).

The Virgo cluster region was imaged on three different FAUST frames, with typical exposures ranging from 630 sec (Northernmost frame) to 1250 sec (Southernmost frame). The images are separated by ~ 4° in declination, with the middle image centered on M87 and each pixel is ~1.11 arcmin. The point source fluxes can be converted to physical units with a calibration derived in flight from observations of stars that were observed by IUE (Bowyer *et al.* 1993). The total sky coverage of the three Virgo cluster frames, defined as the region with non-zero exposure and allowing for image overlap, is ~ 100 square degrees. The entire region is at high galactic latitude (b \geq 65°) and is shown in Figure 1. A general discussion of UV properties of some galaxies observed by FAUST, in which 29 Virgo cluster objects are included, was given by Deharveng *et al.* (1994).

An impartial source detector algorithm, described in Brosch *et al.* (1995), operated on the three FAUST images of the Virgo cluster and produced a list of 191 UV sources, some appearing on more than one image, with their FAUST UV flux and equatorial coordinates (Table 1). Objects for which the coordinates were calculated from an astrometric solution based on bright, well-identified stellar images, have their coordinates in Table 1 marked with an asterisk. Note that, because of the character of the FAUST images, different regions of the images are exposed for different times. Therefore, the detection of sources here is not flux-limited, but rather signal-to-noise limited. The flux was measured with a simulated round aperture of diameter 13.2 arcmin. This is large enough to include all the flux from any galaxy in the VC, thus we did not distinguish between point and extended sources. A few very bright UV sources (stars brighter than ~ 6.5) were measured with a larger simulated aperture. The sky contribution was derived from a square box with 44 arcmin sides, except for the few bright sources, where the sides were 77 arcmin.

Table 1 lists also the error in the flux; this was calculated by combining in quadrature the instrumental and systematic errors. The first was obtained from the "photon statistics" distribution of the events from which the sky brightness was calculated. The systematic errors, originating from the laboratory calibration and the intrinsic errors in the IUE scale, are estimated by Bowyer *et al.* (1993) to be 15.8% of the measured flux, *i.e.*, an additional error of 0.16 mag to the monochromatic magnitude used here. The full analysis of the UV source list, the identification of UV sources with catalogued objects, and follow-up optical observations of non-catalogued sources, are the subject of this paper.

2.1. Internal consistency of the FAUST measurements and source identification

The three FAUST images have considerable overlap. This allows the comparison of magnitudes for a significant number of sources, which appear on two or three images. The comparison of different magnitudes for the same objects is shown in Figure 2. Objects measured three times, which appear in the region of overlap of all images, are represented by three points connected with straight lines, each corresponding to one pair of measurements. Fig. 2 shows 133 pairs of values, after deleting the six brightest and the three faintest pairs of points, for which observational biases could modify the correlation, we find that the correlation coefficient between pairs of measurements of the same object is 0.94. The slope of the relation is consistent with unity within 2σ (0.93±0.03). The mean error in the comparison among different observations of the same object is 0.28 mag. We conclude that the FAUST measurements in the Virgo cluster area are internally consistent over at least a six-magnitude range.

We compared our detections with those listed by Bowyer *et al.* (1995) in the FAUST catalog. We find that the absolute majority of our sources have been listed in the catalog, but 8% (16) were detected here while having no catalog counterpart. As mentioned above, we used here a different automatic detection criterion than used by Bowyer *et al.* In most cases, these sources are faint, or appear in a single FAUST image, or both; they may be artifacts but we accept them here as genuine as there is no clear reason to reject them. Table 1 lists in its last column the FAUST catalog number(s) of sources corresponding to our detections. Again, in most cases the proposed identifications coincide.

We found that a number of sources listed in the FAUST catalog of Bowyer *et al.* (1995) could be "merged" as a single source. These presumably are the result of the automated processing of FAUST catalog sources and in all cases are sources which appear in more than a single FAUST image. An extreme example is V66=N4438, which appears on three FAUST images and was listed in Bowyer *et al.* as three separate sources.

The identification of the detected sources is based mainly on positional coincidence with catalogued objects. We used various catalogs of stars, such as the Smithsonian Astrophysical Observatory's (SAO 1966), the Hipparcos Input Catalog (Turon *et al.* 1993), the TD-1 catalog (Thompson *et al.* 1978)), *etc.* We identified galaxies using the Third Revised Catalog of Galaxies (de Vaucouleurs *et al.* 1991; RC3) and the Principal Galaxy Catalog (PGC; Paturel *et al.* 1992), mainly through the LEDA data bank. In addition to positional coincidence (3' error box), we applied some astrophysical constraints. For example, if near the position of the UV source the catalogs listed a bright late-type star, but slightly further away we found a UV-bright object such as a sub-dwarf, we accepted the latter as the proper counterpart.

In cases when no suitable catalogued counterpart could be found within 3' of the FAUST position, we inspected the Palomar Sky Survey (PSS) prints at the Wise Observatory. We searched in the immediate neighborhood of a UV source location for (a) relatively bright objects, with (b) bluish colors, as could be determined from a comparison of the E and O plates. These candidate counterparts were observed at the Wise Observatory (see section 3, below).

2.2. Comparison of FAUST and GUV measurements

Among the various UV observations toward the Virgo cluster, those by GUV (Kodaira *et al.* 1990) have a very similar bandpass and may be used to check external consistency, along with those objects observed by IUE. The GUV experiment operated during two observation periods. The "acquisition" (A) phase and the "pointing" (P) phase. During the A phase, with the payload scanning and being spin-stabilized, 39 sources and one upper limit were detected and identified with known stars (Table 1, Onaka *et al.* 1989). The FAUST experiment detected 8 of these stars; the remaining GUV sources lie outside the field observed by FAUST. In the pointed phase the telescopes were kept within one degree of their intended pointings, with a drift rate of 0'.5 s⁻¹. The final resolution of the GUV P phase images, after stacking 70 images with 2.52 sec exposure each, was $\Delta \alpha = 16'.0$ and

 $\Delta \delta = 8^{\circ}.2.$

Among the sources detected during the pointing phase and listed in Table 2 of Onaka *et al.* (1989), five were detected by FAUST and identified with stars. An improved reduction of the P phase data led to the identification of some 42 galaxies and of further anonymous sources (Table 1 and Table 4 of Kodaira *et al.* 1990). FAUST detected 23 of the galaxies of Kodaira *et al.* (1990) and four of the anonymous sources have been identified with known stars. Thus the total number of detected sources in common by the two experiments is 39 and one upper limit.

Nine galaxies detected by GUV were missed by FAUST: IC3255, N4360, IC3371, N4458, N4459, N4473, IC 3425, IC3457, N4503, IC3483, IC3501. IC3255 is listed as FAUST source 3110 by Bowyer *et al.* (1995) with a S/N<4.4. For this reason it had not been selected by the algorithm which produced the present list. The remaining sources do not appear in the Bowyer *et al.* (1995) catalogue. As for the anonymous objects of Table 4 of Kodaira *et al.* (1991), five (A1221+09.7, A1221+12.7, A1224+14.2, A1231+10.7, A1235+10.2) were missed here as well as in the FAUST catalog (Bowyer *et al.* 1995). The P10 source (Table 2 of Onaka *et al.* 1989) is listed as catalog source 3195, but is missed in the present list, due to low S/N.

Below we comment on several sources, for which the better resolution of FAUST allows an improved identification. The two GUV sources identified as N4387 and N4402, and the upper limit source identified as N4406, overlap. FAUST detected only one source, V51, better identified as N4406. The GUV source identified as N4435 is interacting with N4438, which is tidally disturbed and is FAUST V66. IC3416 is 5' from V74, which is identified here as a F0 star. N4497 is not a good identification. The FAUST source V90 is identified with IC3446, which is nearer the UV location than N4497. N4564 was identified as the GUV source, but it overlaps with N4567 and N4568. N4567 is FAUST V126. IC3583 overlaps with N4569=V129. The GUV source A1224+08.8 lies in the field of V56 and V62. See the comment in section 3.1. The source A1225+11.5 (P2) is 4' south of V64 and can be identified with HD108452 (A0). Note that Fig. 5 of Onaka *et al.* (1989) is reversed and North is at the bottom. A1225+11.8 (P3) is close to V65. A1227+08.6 (P7) corresponds to V82. The anonymous object suggested by Onaka *et al.* (1989) has been observed to be a late-B star [S122710+083901]. A1230+13.2 is V102, observed to be a late-B star [S123030+131240]. A1232+09.7 (P17) corresponds to V122. The object near the UV position has been observed to be a late-B star [S123252+094327].

Our analysis shows that, unfortunately, GUV cannot serve as an external confirmatory source for FAUST. This is a result of the FAUST exposures being deeper and having better resolution, as well as being calibrated against IUE measurements. Despite those aspects, it is reassuring to see that (a) most "real" sources of GUV were found by us, and (b) spurious GUV sources, with low S/N, were not detected by two different source-finding algorithms in the FAUST images; that of Bowyer *et al.* (1995) and the one used by us.

3. Optical observations

One hundred and forty six sources were identified with stars or galaxies, as explained above. The remaining 45 FAUST detections were inspected on both PSS copies at the Wise Observatory, to identify possible counterparts below the magnitude limit of the catalogs we consulted. Spectroscopic observations of these candidates from the Wise Observatory resulted in 31 sources identified as stars, which we classified as hot stars. In 7 FAUST fields we did not observe all possible stellar candidates (V1, V65, V92, V99, V112, V119, and V180). Two other FAUST sources were found to have no reliable optical counterpart on pure astrophysical reasons: these are V101 and V154, which are discussed in sub-section 3.1 below. The remaining 5 sources (V56, V62, V104, V116, and V125) are discussed below is section 3.1. All these 14 sources are labelled "NO ID" in Table 1.

We used the FOSC (Faint Object Spectrometer Camera) in spectroscopic mode for these observations. The instrument consists of a focal reducer camera with a collimated beam section and long-slit focal plane entrance apertures. The FOSC operates at the f/7 focal position of the 1 meter telescope of the Wise Observatory. It is possible to insert grisms in the collimated beam section to disperse the incoming light. We used grisms with 300 gr/mm or 600 gr/mm, and a wedged window as beam steerer to center the desired spectral region onto a TI CCD chip. The CCD has a 1024x1000 pixel format, but for spectral observations only the relevant part of the chip, containing the spectrum of the object and that of the nearby sky, was read and recorded. The FOSC spectra in the first configuration cover the region from ~4000Å to ~8000Å with ~4Å per pixel. Subsequently, the TI CCD was changed to a Tektronix 1024×1024 pixel back-illuminated CCD with a Metachrome II coating to enhance its blue-UV response. The pixels of this chip are larger than those of the TI. In this configuration the spectra cover the range from ~3750Å to ~7100Å with ~3.5Å/pixel.

Each object was observed in a slitless configuration (thus the seeing disk defined the entrance aperture). We observed spectrophotometric standard stars and obtained calibration exposures of a He–Ar arc with a 2" slit together with the candidate sources. Flat fields were obtained of the twilight sky and of an internal incandescent filament lamp. The spectra were extracted from the CCD images after debiasing and were flat fielded and divided pixel-by-pixel by a similarly processed spectrum of the standard star. Figures 3a to 3d show representative spectra obtained for some survey sources. In the spectrum of V24 it is possible to see a flux depression near 4650Å. This is an artefact caused by the standard star used to reduce the observations of that particular night. The flux depression appears in all spectra obtained during that specific night. Most spectra show only Balmer absorption features. The lines are pronounced, but not extremely wide, indicating that the objects are probably not DA white dwarfs. No HeII features are observed. We compared the FOSC spectra with "template" spectra from Jacoby *et al.* (1984) and from Silva and Cornell (1992). Most appear to be early–type stars, from mid–B to late–A or F. The resolution is not sufficient to distinguish between main sequence and giant luminosity classes, but supergiants appear to be ruled out in most cases, because supergiant B stars have almost no H α absorption while late A supergiants show reasonably deep HeI λ 5876Å absorption. The WiseObs spectral types of all the candidate optical counterparts are listed in Table 2, along with those typing data obtained from the literature.

The distribution of all FAUST UV sources in the Virgo fields, detected here and identified as described above, is summarized in Table 3. We bin the sources by spectral type and UV magnitude (if they are stars), or just by magnitude if they are extragalactic. We also log the 14 non-identified sources.

3.1. Individual objects

Here we discuss selected individual objects that are worth some extra entry that could not be fitted in the tables.

V44=N4374=M84 is an elliptical galaxy with a skewed, extended dark lane in its central region. The dark lane is associated with a rotating gas disk with a LINER spectrum. The extinction by dust in the dark lane was studied by Goudfrooij *et al.* (1994), who concluded that the dust was similar with that in the Galaxy. The total amount of dust was estimated at ~3 10⁴ M_☉. The galaxy also contains significant amounts of hot gas, being an EINSTEIN X-ray source (Forman *et al.* 1985; Matilsky

et al. 1985). Despite its dark lane, its outer isophotes are remarkably uniform (Malin 1994).

- V46=N4383=U7505=Mrk 769 is a bright Seyfert galaxy with a starlike, blue nucleus (Barbieri and Benvenuti 1974). A comparison of the [FAUST-B] and [UV-B], using the Smith and Cornett (1982) magnitude, indicates a very blue spectral energy distribution (SED).
- In the field of V56 and V62 there are some possible candidate counterparts of UV sources. The faint galaxy between the two sources has been observed spectroscopically. We found a single emission line at λ5960Å. One of the two stars is SAO119413 (K0) and the other, S122425+085025, has been observed spectroscopically once from the WiseObs and was identified as a late A star. Kodaira *et al.* (1990) found a UV source which they did not identify (A1224+08.8) about 3' South of V56; it is possible that this GUV source and V56 are the same, and coincide with the late-A star.
- V65: The two m≈13 mag candidates were not observed. There is one dwarf galaxy (VCC 1020; dE) which is nearby. If it is adopted as a possible conterpart, its [FAUST-B] color (-5.67) would be anomalous for a dwarf elliptical.
- V74: The observed candidate S122644+105941 is a V≃12.5 late F star. In the field there is a faint blue star near the UV source position, and two other candidates 6 arcmin farther away. The faint V>15 blue candidate, S122629+105705, has not been observed.
- V79: The star S122700+063417 has been observed and it is a late F star. The suggested identification is near the UV position, the V=8.7 mag G0 star SAO119429.
- V92: The S122903+080353 object near the UV position has been observed spectroscopically and identified as a ~10 mag late G star. There are three other

candidates in the field, which were not observed. One of these, fainter than V=15, appears extremely blue when compared on the PSS E and O copies.

- V99: On the PSS there is a faint blue and slightly compact galaxy about 1'.4 E of the UV position, which was not observed.
- V101: The faint object S123017+094747, which is near the UV position, has been observed and identified as a late G star, fainter than V=15 mag.
- V104: The S123031+051936 object near position has been observed to be a V≃13.5 mag F/G star. There are other fainter candidates in the field, which were not observed.
- V106 was identified as the QSO Q1230+0947, a V=16.2, z=0.420 AGN found in the APM objective-prism survey (Foltz et al. 1987).
- V116: The V=14 star S123153+154409 has been observed and it is an early G star. There are some other faint candidates in the field.
- V119: A m>15 blue star is located ~1' away from the UV position, and was not observed.
- V121: A very weak spectrum was obtained from the m>15 mag candidate in 5400 sec, which could not be used for classification. A dwarf elliptical galaxy (VCC 1600) is nearby and if selected as counterpart would have a very anomalous color [FAUST-B]=-5.68 for a dwarf elliptical, leaving the association of V121 with VCC 1600 uncertain.
- V125: The S123404+075544 candidate observed by us is a V≃13 mag late-G star. This is the brightest candidate object among the star-like images on the PSS near the location of the UV source and it has a faint bluish companion to the West-South West.

- V129=NGC 4569=M90=UGC 7786 is an Sab galaxy. It was observed during a snapshot FOC-HST program (Maoz et al. 1995) to have a nuclear UV source at 2300Å whose size is FWHM≤0".22=2.2h⁻¹ pc (h is the Hubble constant in units of 100 km s⁻¹). Spectroscopy of the source indicates it to be a transition between an HII nucleus and a LINER. There is residual UV emission from an extended source 0".65 South of the bright nuclear region. The flux density measured for the nuclear source is 10⁻¹⁴ erg/sec/cm²/Å, or a monochromatic magnitude at 2270Å of 13.82. The difference from the integrated UV flux measured by FAUST is due to emission at much larger nucleocentric distances than sampled by the FOC.
- V138=NGC4579=M58=UGC 7796 is an Sab galaxy with a LINER nuclear spectrum and evidence of weak broad H α and X-ray emission (Fillipenko and Sargent 1985, Halpern and Steiner 1983). It was observed in the same FOC-HST snapshot survey to have two near-nuclear sources, a bright nucleus and a nearby slightly extended source (Maoz *et al.* 1995). The FOC nuclear source flux density corresponds to a monochromatic magnitude at 2270Å of 16.22 mag. The slightly extended source is two magnitudes fainter. However, the FAUST measurement indicates that most UV emission does not originate from the nuclear region.
- V144=VCC 1791=UGC 7822=IC3617=MCG 01-32-127=DDO 140 has strong Hα emission and equivalent width, indicating vigorous star formation activity (Almoznino 1995). We have mapped it with the VLA and found an HI companion in contact with the main galaxy (Brosch *et al.* 1996).
- V152: A very weak spectrum was obtained in a one hour exposure for the observed candidate (S123754+101551), a V>15 bluish object which coincides with the location of the UV source. The color [FAUST-V] is very blue for the assigned spectral type; this may indicate some source peculiarity.

- V154: The S123837+071338 candidate observed is a V≃13 mag K star. No other candidates are present in the field. The UV flux appears to be fairly well established, however note that the source appears on a single FAUST image. In view of the faintness in the visible, *i.e.*, the very blue [FAUST–V] color of this K-star, we adopt a "no identification" label for the source.
- V169: A 5400 sec spectrum showed Balmer lines and a steep continuum, as expected from a B type star. The object is very faint (V>15 mag) and, if selected as counterpart, would imply [FAUST-B]<-1.75.
- V171=N4649+N4647=Arp 116 is a contact pair of galaxies consisting of the $S0_1(2)$ N4649=M60 and the Sc N4647. The S0 galaxy is included in the list of objects for which Storchi-Bergmann, Kinney and Chalis (1994) produced the UV to near-IR spectral distributions. This is based on a combination of IUE and ground-based spectrophotometry using an entrance aperture shaped and oriented to match that of IUE's. Using their flux density at 1583Å (the spectral region closest to FAUST's) we obtain a monochromatic magnitude of 15.03, three magnitudes fainter than measured by FAUST but more than one magnitude brighter than the FOC HST snapshot measurement (Maoz, private communication). Part of the difference could be attributed to the shape of the spectrum in the UV, but note that the spectral slope as shown by Storchi-Bergmann et al. stays reasonably flat shortwards of 3000Å. This implies that the UV emission is this LINER is extended, with a small fraction emitted in the nuclear region sampled by the FOC or IUE, and the entire emission detected by FAUST. Our FAUST total magnitude is consistent with measurements, albeit at different UV bands, from SCAP ($m_{UV}=12.0$) and Smith and Cornett (1982) $(m_{UV}=11.87).$

- V172=NGC4651 is an Sc galaxy that, at v_☉=800 km s⁻¹, is probably in the VC. Deep plates (*e.g.*, Sandage *et al.* 1985) show the presence of a jet-like feature. Malin (1994) found a faint outer shell at the end of this structure. Giuricin *et al.* (1990) include V172 in their LINER list, while Kenney and Young (1988) note that it has a high SFR.
- V180: A 3600 sec spectrum of a faint bluish candidate identified on the Palomar prints shows only a weak continuum with no emission lines. No spectral classification is possible from this spectrum.

4. Discussion

4.1. Stars in the direction of the Virgo cluster

Table 3 gives the distribution of stars according to spectral type and magnitude. In comparison with our previous FAUST study of the North Galactic Pole region (NGP: Brosch *et al.* 1995), one item is immediately apparent: no white dwarfs were positively identified in the VC fields. The NGP FAUST field covered 69 square degrees; we detected there 81 objects, thus 1.17 per square degree (\Box°). The source density in the ~ 100 \Box° VC fields is higher, 1.91 per \Box° . However, the VC images are toward a cluster of galaxies, whereas the NGP image is of the general field. Excluding galaxies from both fields, and concentrating only on identified objects, we find a UV star density of 1.03 stars per \Box° at the NGP *vs.* 0.96 stars per \Box° toward the VC.

The density of detected and identified FAUST stellar UV sources in the direction of the VC does not differ from that to the NGP. Despite this, no white dwarfs (WDs) are identified in the VC UV images. The space distribution of (DA) WDs was studied by Boyle (1989) and was included in the model of the UV sky (Brosch 1991). We modeled the stellar distribution for the VC central location using the FAUST spectral bandpass and show a comparison of the detected vs. the predicted stellar density in Figure 4. The model follows the actual stellar distribution down to $m_{UV} \simeq 11$, where incompleteness sets in, but does not reproduce it with great fidelity (half the data points up to the completeness limit are more than 1σ from the model). It is possible that this is a "small-numbers" effect because of the size of the sample, since according to our model, there should be at least three WDs in a $100\Box^{\circ}$ wide field up to a UV magnitude limit of 14. About 140 stars of all types, to this magnitude limit, should have been detected. Considering the completion limit of FAUST, the model and the total number of detected stars agree. The discrepancy with the expected number of WDs remains unsolved. One possibility is that the missing WDs are among the NO ID sources, another is that some were mis-classified by us as A or B stars, where in reality they were DAs (*e.g.*, V152, V169 or V186). Another is that WDs are observationally discriminated against, when searched for in the vicinity of a cluster of galaxies. A similar "avoidance" of Galactic stars by QSOs (in objective-prism surveys) was demonstrated by Gould, Bahcall and Maoz (1993).

V24=S121757+101353 has been shown by our spectroscopic observations to be an early-G V=13.2 object. Its spectrum is shown in Fig. 3a. Note that the 4650Å absorption-like feature is an artefact (see section 3). The [FAUST-V] color, -5.72, is very blue for such a spectral type, where more yellowish colors are expected. The source was detected on two separate FAUST images and its count statistics is reasonable. We have therefore reasonable confidence of its reality and of the accuracy of the UV magnitude. The visual magnitude is from the HST Guide Star Catalog; its accuracy is of order 0.1–0.2 mag (Ratnatunga 1990). Such a blue UV–V color index is expected only from a "mixed" source, where a late-type optically-bright star is paired with a hot companion, which outshines the late-type star in the UV. This star is, therefore, a candidate for such a mixed binary object. Note that our spectrum does not show a trace of such a hot companion; spectra reaching further into the blue-violet are required to detect this.

V143=SAO100195 is an A0m star, for which we found [FAUST-V]=4.42. Am stars are those "A or early F stars for which no unique spectral type can be assigned" (Jaschek and Jaschek 1987). The color we measured is much redder than expected for "regular" A0 stars, which have colors near 0.0; in absence of interstellar reddening, which is not expected at the galactic latitude of this star, the possibilities could be either circumstellar reddening or some photospheric activity that reddens the UV-optical colors. However, Table 2 shows that three other Am stars (V3, V120 and V178) also have positive and fairly large [FAUST-V] color indices. It is possible that this is caused by additional metallic lines present in the UV spectrum of Am stars; this has been noted by Jascheck and Jaschek (1987) for the optical spectra and is probably true also in the UV.

4.2. Virgo galaxies: UV and other observational properties

In Table 4 we present the UV properties of the galaxies detected by FAUST with morphological types from the BST catalog. We include UV measurements from the GUV experiment (Kodaira *et al.* 1990), from the SCAP balloon telescope (Donas *et al.* 1987), and from Goddard rocket-borne telescope (Smith and Cornett 1982, labelled S+C). For consistency, we transformed all GUV measurements into monochromatic magnitudes as explained above. Note that each experiment measured a different spectral segment; GUV measured effectively at 1560Å with a bandwith of FWHM=230Å, SCAP measured at \sim 2000Å and the bandpass was \sim 200Å, and the Goddard imager had a 1150Å wide bandpass centered near 2421Å.

We mentioned above that, in a number of cases, the UV emission detected by FAUST is extended, and from a comparison with HST FOC observations only a small fraction of the UV is centrally produced. This was shown to be the case for a number of elliptical galaxies in the VC by Kodaira *et al.* (1990), who compared the GUV flux density with that measured by the large IUE aperture. They found that, for five galaxies, the ratio of central to total UV flux ranged from 14% to more than 63%. This is confirmed here by a comparison of FOC (Maoz *et al.* 1995) and other UV observations for three of the galaxies, as mentioned above. Observations by UIT during the ASTRO I and ASTRO II missions (e.g., Neff *et al.* 1994, Cornett *et al.* 1994) detected complex morphologies in the far-UV emission of some galaxies. In the case of M31 and M33, far-UV observations found also that their central regions were bluer than their outer parts (O'Connell 1992).

The galactic latitude of all sources discussed here is higher than 65°. Despite this, the UV measurements could be affected by foreground galactic extinction. Burstein and Heiles (1982) showed that small but significant reddening may exist over the Virgo cluster. We consulted the reddening catalog of Burstein and Heiles (1984) and found that our objects have small reddening values [E(B-V)<0.04]. This corresponds to $A_{1650\mathring{A}} \leq 0.33$ mag, using the wavelength dependence of the extinction from Savage and Mathis (1979), *i.e.*, $\frac{A(1650)}{E(B-V)} = 8.0$. We corrected the FAUST magnitudes for extinction, using the 4×E(B-V) values listed for the galaxies by Burstein and Heiles, or as appropriate for the nearmost galaxy listed by them in cases where no specific entry appeared in the list of Burstein and Heiles. We corrected by $A_{UV}=2\times$ the listed value if positive, or left the UV magnitude unchanged if Burstein and Heiles listed a zero or negative value. The UV magnitudes, still called "FAUST magnitudes" and corrected for extinction, are listed in Table 5, where galaxies are discussed separately from Galactic objects. The table also presents the total magnitude of the galaxy (usually in the B band and from the BST or the RC3 catalogue), corrected for extinction, and adds the T-type of each and the HI flux integral FI(HI), from Huchtmeier and Richter (1986), Haynes and Giovanelli (1986), Hoffman et al. (1989a and 1989b), or Hoffman (private communication). In the case whenever a number of FI(HI) values are given we adopted their straight average. The correlation between FI(HI) and

the FAUST UV magnitude is shown in Figure 5. We find a correlation coefficient of -0.74 when regressing log FI(HI) against the FAUST magnitude, and including only spiral and Irregular galaxies (T>0). This has been known from other UV–HI studies (*e.g.*, Buat *et al.* 1989) and has been linked to a correlation between total gas content and the star formation rate (see below).

In Table 5 we include also the far infrared (FIR) measurements of the same galaxies by the IRAS telescope. The data are mainly from the IRAS Faint Source Catalog. We estimate the total FIR flux as in Lonsdale *et al.* (1985):

$$F_{FIR} = 1.26 \left(2.58S_{60} + S_{100} \right) 10^{-14} W m^{-2} \tag{1}$$

using the 60 and 100 μ m flux densities, which are also given in Table 5. The correlation between the FAUST and the FIR emissions is shown in Figure 6. We plot in Figure 7 the distance-independent "colors" $\frac{FIR}{HI}$ vs. [FAUST–B]. Excluding three background galaxies at high cz (V12, V63, and V118), which may be exceptional, we find a correlation coefficient of 0.62.

The color [FAUST-B] can be used, as done by Deharveng *et al.* (1994), to compare the VC galaxies with the stellar population models of galaxies which are used for deriving the K-correction in cosmological models. Deharveng *et al.* compared the colors with the spectral energy distributions of Coleman, Wu and Weedman (1980). Our study, like that by Deharveng *et al.*, and previously by Faber (1982) and by Burstein *et al.* (1988), shows that elliptical galaxies exhibit a large dispersion in their (UV-B) color index relative to other, late-type galaxies. Thus, it is hard to establish a universal K-correction for their UV segment of the spectrum, which could be useful when studying in the optical elliptical galaxies at $z\simeq 2$. Hence, the elliptical galaxies do not form a uniformly bright population of objects at high redshifts, making their use in optical surveys or studies of high redshifts objects problematic, due to Malmquist and other selection effects.

4.3. Total UV flux from cluster galaxies

In the background section we explained that the discovery of nearby Lyman α absorptions in the spectra of 3C273 prompted the evaluation of the metagalactic ionizing background. The presence of such absorbers in the direction of the VC and at the redshift of the cluster (Weymann *et al.* 1995, Shull *et al.* 1996) prompts us to re-evaluate the ionizing flux, based on the measured UV emission from the cluster galaxies.

Here we restrict the discussion to the total UV flux detected from galaxies in the direction of the VC by different UV experiments. We estimate the total number of available ionizing photons, based on published UV data of Virgo cluster galaxies. As the bandpasses of observations were always *longward* of the Lyman limit, we estimate the Lyman continuum (Lyc) photon flux by scaling the measured fluxes to the wavelength of the Lyman break using the stellar population models of evolving galaxies of Bruzual and Charlot (1993).

Each publication with UV observations of the VC provides a list of galaxies towards the Virgo cluster and either a UV "monochromatic" magnitude or a flux density in a specific bandpass. In order to estimate the amount of diffuse UV contribution when the locations of the galaxies are not known in three-dimensional space, we *assume* that the UV contribution from all galaxies comes from a single point located at the cluster center. We calculate the UV contribution from each galaxy from the given flux, we scale it for the distance of each galaxy to the center of the cluster (to account statistically for nearer and farther galaxies than the center of Virgo), and add it to the total flux densities (TFD). The reference position for the center of the Virgo cluster was given by van den Bergh (1977): $\alpha = 12^h 25^m .4$, $\delta = +12^\circ 40^\circ$. Objects for which only an upper limit is given are included as if their emission would equal this upper limit (except for FAUST). This yields an upper limit to the total galactic UV emission of the Virgo cluster.

The TFD reached for the different instruments/wavebands are remarkably similar; for

the detected FAUST galaxies the TFD is $4.82 \ 10^{-12} \text{ erg sec}^{-1} \text{ cm}^{-2} \text{ Å}^{-1}$, whereas for the objects listed by Smith and Cornett (1982) the TFD is $\sim 2 \ 10^{-12} \text{ erg sec}^{-1} \text{ cm}^{-2} \text{ Å}^{-1}$. The Virgo galaxies measured by SCAP and listed in Donas *et al.* (1987) total to a TFD of 1.1 $10^{-12} \text{ erg sec}^{-1} \text{ cm}^{-2} \text{ Å}^{-1}$, but only a few brighter objects are included in their list. The TFD for the GUV galaxies is $\sim 10^{-12} \text{ erg sec}^{-1} \text{ cm}^{-2} \text{ Å}^{-1}$, close to the SCAP 2000 total.

The TFD measured by FAUST is $\sim 2.5 \times$ larger than that estimated by Weymann *et al.* (1995) for the 1550Å flux of VC galaxies. Weymann *et al.* calculated the flux from the B-band luminosity of early type galaxies, compounded with a UV–V transformation derived from Burstein *et al.* (1988). The difference from the FAUST measurement is the result of the neglect of spiral and other late types by Weynmann *et al.* ; these galaxies produce most of the UV flux of the VC in the FAUST band.

We adopt the highest value of TFD from FAUST, and scale it to the Lyman limit, using the maximal ratio of 1650Å flux to Lyc flux just short of the Lyman limit allowed by the galaxy models of Bruzual and Charlot (1993), which is ~2. From this, we conclude that the TFD at the Lyman limit which is produced by galaxies detected at Earth can be at most ~8 10^{-12} erg sec⁻¹ cm⁻² Å⁻¹, the equivalent of a single m_{UV}=7.2 mag star.

4.4. Dust in the Virgo cluster ?

Girardi *et al.* (1992) found, by analyzing loose groups, an anticorrelation between the E(B-V) color excess of galaxies and the velocity difference with respect to the brightest group member. They interpreted this as evidence for the presence of dust within the intra-group space, which reddens the light from galaxies in the anti-Earth location. Similar claims, of significant amounts of dust within Abell clusters, were made by Karachentsev and Lipovetski (1969), Bogart and Wagoner (1973), Boyle, Fong and Shanks (1988), and Romani and Maoz (1992). In the context of the VC, a search for inter-cluster dust can

be linked to the possible presence of "almost invisible" galaxies, which give rise to the absorption lines at the redshift of the VC in spectra of background QSOs.

We attempted to test this by first checking five background galaxies, with $v_{\odot} \geq 3000$ km s⁻¹, against cluster and foreground galaxies with $v_{\odot} < 3000$ km s⁻¹, hereafter called "comparison galaxies" (CGs). The assumption here is that background galaxies, seen through the cluster, should have redder [FAUST–B] colors that the CGs of similar morphological types (within 2 T subclasses), because of dust reddening. We found that there is no consistent evidence for the background galaxies being significantly redder than the cluster and foreground galaxies.

We then tested the CGs against themselves, by splitting the sample into a "near" bin if $v_{\odot} > 1100 \text{ km s}^{-1}$ (redshift of the Virgo core), but $v_{\odot} < 3000 \text{ km s}^{-1}$, and a "far" bin, with galaxies which have $v_{\odot} \leq 1100 \text{ km s}^{-1}$. The assumption here is that galaxies which belong to the cluster have their cosmological recession velocity modulated by the gravitational attraction of the cluster (see, *e.g.*, Binggeli *et al.* 1993). Therefore, galaxies that are between us and the VC core will be accelerated by the cluster attraction and will have, on average, velocities higher than that of the cluster core. Galaxies in the VC but beyond the core will be slowed by the cluster attraction and may even have negative heliocentric velocities. Although the gravitational influence of the cluster is certainly much more complex, this crude separation may show some significant result. The separation, and subsequent comparison of cluster members, compensates for such effects as "anaemic" spirals, stripped galaxies, and such, which should apply equally to both the "far" and the "near" sub-samples. In addition, Malmquist biases in the selection of background objects should be minimized.

We tested whether the "near" bin galaxies are bluer in [FAUST–B] than the "far" bin galaxies, but did not find significant differences. The conclusion is that the total amount of

dust in the VC is rather small, probably negligible. A similar result for a sample of clusters was obtained recently by Maoz (1995), from a study of the (V–I) color distribution of QSOs beyond the clusters.

4.5. Large-scale distribution and UV luminosity function of UV-emitting galaxies

In the context of the three-dimensional distribution of galaxies in the direction of the VC it is interesting to test the distribution of UV-emitting objects. One finding, emphasized here as well as by Deharveng *et al.* (1994), is that UV-emitting galaxies are mainly of late-type. These galaxies show usually less clustering than early-type objects, such as elliptical, lenticular and Sa galaxies. It is therefore to be expected that UV-bright objects will tend to line at the edges of the main clusterings, and in the voids.

In order to check the three-dimensional distribution of UV galaxies we plot in Figure 8 a diagram of all detected objects. The vertical axis is the heliocentric velocity as listed by BST and the horizontal axis is the right ascension of each galaxy. It is clear that most of the galaxies detected by FAUST are within the redshift range of the VC and only a few are in its background.

In addition, we tested the distribution of the cluster galaxies in Virgocentric projected distance (degrees from the adopted cluster center position), by binning the galaxies in one-degree wide rings, despite the fact that our observations cover an ~elliptical shaped sky area. Considering that galaxies are actually distributed in a three-dimensional volume, the distribution indicates a high over-density of UV emitting galaxies in the inner four degrees of the VC.

The optical luminosity function (LF) of galaxies in the VC has been studied by Sandage,

Binggeli and Tammann (1985). The LF for bright galaxies is well represented by a function bounded at both bright and faint ends. The optical LF for field galaxies was studied by Marzke *et al.* (1994) and was fitted by single Schechter functions (Schechter 1976) over the entire magnitude range. In the Coma cluster, Biviano *et al.* (1996) found that the best fit is by a Schechter function combined with an Erlang function.

We plot in Figure 9 the distribution of UV magnitudes for cluster member galaxies, *i.e.*,those with $v_{\odot} < 3000 \text{ km s}^{-1}$, excluding V121 and V155 (uncertain identifications), after binning it into half magnitude-wide bins. In view of the small number of galaxies with UV magnitudes, we include all our objects, instead of splitting them into morphological subclasses.

We note that, for the VC's bright galaxies, Sandage, Binggeli and Tammann found that a Gaussian best represents the luminosity function. Only when including the lower luminosity objects, the dwarfs, were they justified in fitting a Schechter function. The Gaussians fitted by Sandage *et al.* peak at B*=13.3 and 13.0 for spirals and E+S0 galaxies, respectively. The dwarf galaxies were fitted by Sandage *et al.* as Schechter functions, with α =-1.35 and B*=14.3 (for dEs). Studies of the LF of field galaxies (*e.g.*,Efstathiou *et al.* 1988, Loveday *et al.* 1992) found less steep α values (-1.07 and -1.11, respectively).

We fitted a Gaussian distribution to the histogram of FAUST galaxies of the VC, and allowed the peak, the average, and the width of the Gaussian to change. A best fit was estimated by minimizing χ^2 , where the average UV*=12.37 and the full width at half-maximum of the Gaussian was 2.02 mag.

5. Conclusions

We presented a comprehensive study of three UV FAUST images in the direction of the Virgo cluster of galaxies. We detected 191 UV sources and identified them, through comparison with catalogued optical sources, with (mostly) early-type stars or galaxies. We showed that, within the completeness limit of FAUST observations, the UV stellar distribution follows that predicted by our UV sky model. Am stars were shown to be UV-faint, in comparison to normal stars of the same spectral sub-class.

For galaxies, we confirmed that significant UV flux is emitted outside the central regions. We confirmed earlier results that elliptical galaxies have widely varying UV brightness, making their use in studies of high redshift galaxy samples problematic. Star formation properties were demonstrated through UV–IR and UV–HI correlations. These confirm essentially the accepted pattern of star formation. We estimated the total ionizing flux by galaxies in the VC and showed that it is consistent with earlier estimates. We could not put significant new limits on dust in the VC but our results are consistent with a "no dust" case. The UV galaxies were found to concentrate in the central region of Virgo. The UV luminosity function was fitted with a Gaussian. A better determination of the UV luminosity function in the Virgo cluster requires deeper UV observations.

Acknowledgements

The UV astronomy effort at Tel Aviv is supported by special grants to develop a space UV astronomy experiment (TAUVEX) from the Ministry of Science and the Arts, and from the Austrian Friends of Tel Aviv University. Observations at the Wise Observatory are supported by a Center for Excellence Grant from the Israel Academy of Sciences. This paper made use of the LEDA galaxy data bank. Many optical observations for this project were mostly performed by Mr. E. Goldberg and the reductions by Ms. Ana Heller. Some of the identification, using computerized data banks, were done by Ms. Susanna Steindling. NB is grateful to J. Donas for providing a list of galaxies observed by SCAP, to G. Lyle Hoffman for an updated list of galaxies in the VC region with HI measurements, and to B. Bilenko for running the UV sky model for the VC.

References

- Almoznino, E. 1995 PhD thesis, Tel Aviv University.
- Barbieri, C. and Benvenuti, P. 1974 Astron. Astrophys. Suppl. 13, 269.
- Binggeli, B., Popescu, C.C. and Tammann, G.A. 1993 Astron. Astrophys. Suppl. 98, 275.
- Binggeli, B., Sandage, A. and Tamman, G.A. 1985 Astron. J. **90**, 1681 (BST).
- Binggeli, B., Tammann, G.A. and Sandage, A. 1987 Astron. J. 94, 251.
- Biviano, A. Durret, F., Gerbal, D., Le Fevre, O., Lobo, C., Mazure, A. and Slezak, E. 1996 Astron. Astrophys. **311**, 95.
- Bogart, R.S. and Wagoner, R.V. 1973 Astrophys. J. 181, 609.
- Boyle, B. 1989 Mon. Not. R. astr. Soc. **240**, 533.
- Boyle, B.J., Fong, R. and Shanks, T. 1988 Mon. Not. R. astr. Soc. 231, 897.
- Bowyer, S., Sasseen, T.P., Lampton, M. and Wu, X. 1993 Astrophys. J. 415, 875.
- Bowyer, S., Sasseen, P.T., Xiauyi, W. and Lampton, M. 1995 Astrophys. J. Suppl. 96, 461.
- Brosch, N. 1991 Mon. Not. R. astr. Soc. 250, 780.

- Brosch, N., Almoznino, E., Leibowitz, E.M., Netzer, H., Sasseen, T., Bowyer, S., Lampton, M. and Wu, X. 1995 Astrophys. J., 450, 137.
- Brosch, N., Almoznino, E. and Hoffman, G.L 1996, submitted.
- Bruzual, G.A. and Charlot, S. 1993 Astrophys. J. 405, 538.
- Buat, V., Deharveng, J.M. and Donas, J. 1989 Astron. Astrophys. 223, 42.
- Burstein, D., Bertola, F., Buson, L.M., Faber, S.M., and Lauer, T.R. 1988 Astrophys. J. **328**, 440.
- Burstein, D. and Heiles, C. 1982 Astron. J. 87, 1165.
- Chakrabarti, S., Sasseen, T.P., Lampton, M. and Bowyer, S. 1993, Geophys. Res. Lett. **20**, 535.
- Coleman, G.D., Wu, C.C. and Weedman, D.W. 1980 Astrophys. J. Suppl. 43, 393.
- Cornett, R.H., O'Connell, R.W., Greason, M.R., Offenberg, J.D., Angione, R.J., Bohlin, R.C., Cheng, K.P., Roberts, M.S., Smith, A.M., Smith, E.P., Talbert, F.D. and Stecher, T.P 1994 Astrophys. J. 426, 553.
- de Vaucouleurs, G., de Vaucouleurs, A., Corwin, H.G., Buta, R.J., Paturel, G. and Fouqué,P. 1991 Third Reference catalog of Bright Galaxies, New York: Springer Verlag (RC3).
- Deharveng, J.M., Sasseen, T.P., Buat, V., Bowyer, S., Lampton, M. and Wu, X. 1994, Astron. Astrophys. 289, 715.
- Donas, J., Deharveng, J.M., Milliard, B., Laget, M. and Huguenin, D. 1987 Astron. Astrophys. 180, 12.
- Efstathiou, G., Ellis, R.S., and Peterson, B.A. 1988 Mon. Not. R. astr. Soc. 232, 431.

- Elmegreen, B.G., Elmegreen, D.M. and Montenegro, M. 1992 Astrophys. J. Suppl. 79, 37.
 Faber, S. 1982 Highlights Astr. 6, 165.
- Fillipenko, A.V. and Sargent, W.L.W. 1985 Astrophys. J. Suppl. 57, 503.
- Foltz, C.B., Chaffee, Jr. F.H., Hewett, P.C., Macalpine, G.M., Turnshek, D.A., Weymann, R.J. and Anderson, S.F. 1987 Astron. J. 94, 1423.
- Forman, W., Schwarz, J., Jones, C., Liller, W. and Fabian, A.C. 1979 Astrophys. J. Lett. 234, L27.
- Forman, W., Jones, C. and Tucker, W. 1985 Astrophys. J. 293, 102.
- Girardi, M., Mezzetti, M., Giuricin, G. and Mardirossian, F. 1992 Astrophys. J. 394, 442.
- Giuricin, G., Bertotti, G., Mardirossian, F. and Mezzetti, M. 1990 Mon. Not. R. astr. Soc. 247, 444.
- Goudfrooij, P., de Jong, T., Hansen, L. and Nøgaard-Neilsen, H.U. 1994, Mon. Not. R. astr. Soc., **271**, 833.
- Gould, A., Bahcall, J.N. and Maoz, D. 1993 Astrophys. J. Suppl. 88, 53.
- Halpern, J.P. and Steiner, J.E. 1983 Astrophys. J. 269, 37.
- Haynes, M.P. and Giovanelli, R. 1986 Astrophys. J. 306, 466.
- Hoffman, G.L. 1989 in Large Scale Structure and Motions in the Universe M. Mezzetti *et al.* (eds.), Dordrecht: Kluwer, p. 365.
- Hoffman, G.L., Helou, G., Salpeter, E.E. and Lewis, B.M. 1989a, Astrophys. J. 339, 812.
- Hoffman, G.L., Lewis, B.M., Helou, G., Salpeter, E.E. and Williams, H.L. 1989b Astrophys. J. Suppl. 69, 65.

- Hoffman, G.L., Lewis, B.M. and Salpeter, E.E. 1995 Astrophys. J. 441, 28.
- Huchtmeier, W.K. and Richter, O.-G. 1986 Astron. Astrophys. Suppl. 64, 111.
- Jacoby, G.H., Hunter, D.A. and Christian, C.A. 1984 Astrophys. J. Suppl. 56, 257.
- Jacoby, G.H., Ciardullo, R and Ford, H. 1990 Astrophys. J. 356, 332.
- Jaschek, C. and Jaschek, M. 1987 The Classification of Stars, Cambridge University Press, p.220.
- Karachentsev, I.D. and Lipovetski, V.A. 1969 Sov. Astron. 12, 909.
- Kenney, J.D. and Young, J.S. 1988 Astrophys. J. 326, 588.
- Kodaira, K., Watanabe, T., Onaka, T. and Tanaka, W. 1990 Astrophys. J. 363, 422.
- Lampton, M., Sasseen, T.P., Wu, X. and Bowyer, S. 1993, Geophys.Res.Lett. 20, 539.
- Lieu, R., Mittaz, J.P.D., Bowyer, S., Lockman, F.J., Hwang, C.-Y., Schmitt, J.H.M.M. 1996 Astrophys. J. **458**, 5.
- Lonsdale, C.J., Helou, G., Good, J.C. and Rice, W. 1985 Catalog of galaxies and quasars observed in the IRAS survey, Jet Propulsion Laboratory, Pasadena, CA.
- Loveday, J., Peterson, B.A., Efstathiou, G., Maddox, S.J., and Sutherland, W.J. 1992 Astrophys. J. 390, 338.
- Malin, D. 1994 in Astronomy from Wide Field Imaging (H.T. MacGillivray, E.B. Thomson, B. Lasker, I.N. Reid, D.F. Malin, R.M. West and H. Lorenz, eds.), Dordrecht: Kluwer Academic Press, p. 567.
- Maoz, D. 1995 ApJ, 455, L115.

- Maoz, D., Fillipenko, A.V., Ho, L.C., Rix, H.-W., Bahcall, J.N., Schneider, D.P. and Duccio Machetto, F. 1995, Astrophys. J. 440, 91.
- Marzke, R.O., Geller, M.J., Huchra, J.P. and Corwin, H.G. Jr. 1994 Astron. J. 108, 437.
- Matilsky, T., Jones, C. and Forman, W. 1985 Astrophys. J. 291, 621.
- Neff, S.G., Fanelli, M.N., Roberts, L.J., O'Connell, R.W., Bohlin, R., Roberts, M.S., Smith, A.M. and Stecher, T.P 1994 Astrophys. J. 430, 545.
- O'Connell, R.W. 1992 in **The stellar populations of galaxies** IAU Symp. no. 149 (B. Barbuy and R. Renzini, eds.), Dordrecht: Kluwer, p.233.
- Onaka, T. and Kodaira, K. 1991 Astrophys. J. 379, 532.
- Onaka, T., Tanaka, W., Watanabe, T., Watanabe, J., Yamaguchi, A., Nakagiri, M., Kodaira, K., Nakano, M., Sasaki, M and Tsujimura, T. 1989 Astrophys. J. 342, 238.
- Paturel, G., Vauglin, I., Garnier, R., Marthinet, M.C., Petit, C., Di Nella, H., Bottinelli, L., Gouguenheim, L. and Durand, N. 1992 *LEDA: The Lyon-Meudon Extragalactic Database*, CD-ROM Observatorie de Lyon.
- Ratnatunga, K.U. 1990 Astron. J. 100, 280.
- Romani, R.W. and Maoz, D. 1992 Astroph. J. 386, 36.
- Sandage, A., Binggeli, B., and Tammann, G. 1985 Astron. J. 90, 1759.
- Sandage, A. and Tammann, G.A. 1974 Astrophys. J. 194, 559.
- Savage, B.D. and Mathis, J.S. 1979 Ann. Rev. Astron. Astrophys. 17, 73.
- SAO 1966 Smithsonian Astrophysical Observatory Star Catalog Washington: Smithsonian Institution (SAO).

- Schechter, P. 1976 Astrophys. J. 203, 297.
- Shull, J.M., Stocke, J.T. and Penton, S. 1996 Astron. J. 111, 72.
- Silva, D.R. and Cornell, M.E. 1992, Astrophys. J. Suppl. 81, 865.
- Smith, A.M. and Cornett, R.H. 1982 Astrophys. J. 261, 1.
- Storchi-Bergmann, T., Kinney, A. and Chalis, P. 1994, Astrophys. J. Suppl. 98, 103.
- Thompson, G.I., Nandy, K., Jamar, C. Monfils, A., Houziaux, L., Carnochan, D.J. and Wilson, R. 1978 Catalog of Stellar Ultraviolet Fluxes, The Science Research Council (TD-1).
- Trinchieri, G., Fabbiano, G. and Canizares, C.R.C. 1986 Astrophys. J. 310, 637.
- Trinchieri, G., Kim, D.-W., Fabbiano, G. and Canizares, C.R.C. 1994 Astrophys. J. 428, 555.
- Turon, C. et al. 1993 Bull. Inf. CDS 43 (Version 2 of the Hipparcos Input Catalog).
- van den Bergh, S. 1977 Vistas in Astronomy 21, 71.
- Visvanathan, N. and Griersmith, D. 1979 Astrophys.J. 230, 1.
- Weymann, R., Rauch, M., Williams, R., Morris, S. and Heap, S. 1995 Astrophys. J. **438**, 650..
- Wesselius, P.R., van Duinen, R.J., deJonse, A.R.W., Aalders, J.W.G., Luinse, W. and Wildeman, K.J. 1982 Astron. Astrophys. Suppl. 49, 427.

Figure captions

- Figure 1: Combined image of the three FAUST fields toward the Virgo cluster. The UV sky brightness was different at the time the images were obtained, thus the change in background appearance. M87 is near the center of the combined image.
- Figure 2: Internal consistency check: UV magnitudes for FAUST sources measured more than once. Straight lines connect measurements of the same source observed three times.
- Figure 3a to 3d: Examples of spectra for typical Virgo cluster objects. Note that the broad "absorption" near ~4650Å for V24 is an artefact originating from the standard star observation of the specific night.
- Figure 4: UV sky model from Brosch (1991) calculated for the FAUST spectral response (solid line), vs. the actual measured stellar densities (squares).
- Figure 5: Comparison of the HI flux density and the UV magnitude of galaxies. Early-type galaxies are shown as triangles, intermediate types as squares, and late types as diamonds. Open symbols indicate upper limits.
- Figure 6: Comparison of the far infrared flux density (FIR) and the UV magnitude of galaxies. Symbols are as in Fig. 5.
- Figure 7: The change of the FIR/HI ratio as a function of (FAUST-B) for galaxies.
- Figure 8: The large-scale distribution of galaxies measured by FAUST and with redshift available from the literature is shown in a diagram of right ascension *vs.* heliocentric velocity as a depth indicator. The cluster center is marked by an asterisc at the velocity of M87.

• Figure 9: UV luminosity function of Virgo galaxies (cluster and field galaxies are included). The dashed line indicates the Gaussian fit to the observational data.

Id. no.	α	δ	m_{UV}	N_{obs}	Proposed id.	FAUST no.
1	*12 09 35.2	10 23 07	12.79 ± 0.43	1	No ID	-
2	*12 10 11.7	$11 \ 08 \ 26$	11.61 ± 0.19	1	N4178	3038
3	*12 10 50.7	$10 \ 31 \ 46$	$8.40 {\pm} 0.16$	1	SAO099997	3043
4	12 11 17.0	$09\ 15\ 02$	$8.83 {\pm} 0.016$	1	SAO119277	3046
5	*12 12 28.4	10 15 39	$12.69 {\pm} 0.25$	1	SAO100006	3049
6	12 13 16.0	$09\ 18\ 45$	$11.06 {\pm} 0.17$	1	SAO119294	3056
7	12 13 52.8	$07 \ 37 \ 58$	$10.70 {\pm} 0.17$	1	SAO119300	3059
8	$12 \ 13 \ 53.3$	$15 \ 06 \ 35$	$6.10 {\pm} 0.16$	1	SAO100012	3058
9	*12 14 24.8	08 16 41	$9.84{\pm}0.16$	1	S121426+081714	3061
10	*12 14 56.3	$06 \ 53 \ 18$	$11.65 {\pm} 0.23$	1	SAO119308	3065
11	*12 15 01.6	$15 \ 49 \ 03$	$10.42 {\pm} 0.18$	1	S121449+155138	3066
12	*12 15 05.7	08 44 11	$13.03 {\pm} 0.32$	1	IC3111	3068
13	*12 15 10.8	$09 \ 54 \ 10$	$12.67 {\pm} 0.24$	1	S121508 + 095341	3069
14	$12 \ 15 \ 17.0$	$10 \ 23 \ 51$	$7.62 {\pm} 0.16$	2	SAO100024	3072
15	*12 15 25.6	$07 \ 27 \ 27$	$12.83 {\pm} 0.35$	1	N4246	3074
16	$12 \ 15 \ 40.0$	10 08 46	$9.86 {\pm} 0.77$	2	SAO100029	3078
17	*12 16 12.3	$14 \ 40 \ 53$	$10.08 {\pm} 0.17$	2	N4254	3079
18	*12 16 17.1	10 23 11	$13.40 {\pm} 0.26$	1	SAO100034	3080
19	*12 16 28.5	$13 \ 12 \ 47$	$11.94{\pm}0.23$	2	HD107116	3082
20	*12 16 32.6	09 08 01	$13.42 {\pm} 0.31$	1	IC0776	3083
21	12 16 52.0	07 16 10	$9.43 {\pm} 0.16$	1	SAO119328	3085
22	*12 17 15.5	$08 \ 57 \ 13$	$12.94{\pm}0.26$	1	HD107227	-
23	*12 17 35.5	$07 \ 57 \ 52$	$12.90 {\pm} 0.31$	1	N4276	3089

Table 1. Detected FAUST sources in the Virgo images

Table 1—Continued

Id. no.	α	δ	δ m _{UV}		Proposed id.	FAUST no.
24	*12 17 51.2	$10 \ 14 \ 50$	8.92 ± 0.16	2	S121757+101353	3090/1
25	*12 18 36.4	$09 \ 28 \ 08$	$13.63 {\pm} 0.41$	1	S121836 + 092743	3093
26	*12 18 48.5	11 48 15	$11.57 {\pm} 0.17$	2	N4294	3094/6
27	*12 18 57.8	$10 \ 41 \ 17$	12.41 ± 0.72	2	S121860+104153	3098
28	*12 19 01.1	$11 \ 47 \ 58$	$11.58 {\pm} 0.17$	2	N4299	3099
29	*12 19 01.8	14 53 27	12.21 ± 0.31	2	N4298	3097
30	12 19 30.0	$17 \ 01 \ 20$	$6.81 {\pm} 0.16$	1	SAO100062	3101
31	*12 20 23.6	$16 \ 04 \ 55$	$10.07 {\pm} 0.17$	2	N4321	3104/5
32	*12 20 32.8	10 55 55	$13.73 {\pm} 0.97$	3.73 ± 0.97 2 N4325		3106
33	*12 20 43.4	$13 \ 56 \ 08$	11.12 ± 0.20	11.12±0.20 3 SAO100071		3108
34	*12 20 47.3	11 38 44	13.28 ± 0.35	13.28±0.35 1 N4330		3109
35	*12 21 15.8	$12 \ 48 \ 05$	$12.46 {\pm} 0.24$	2	IC3258	3111
36	12 21 18.0	$08 \ 37 \ 17$	$11.38 {\pm} 0.16$	1	SAO119371	3113
37	*12 21 28.7	$06\ 27\ 45$	$12.71 {\pm} 0.26$	1	S122129+062746	-
38	*12 21 34.1	12 31 10	$12.77 {\pm} 0.38$	2	N4351	3114
39	*12 21 36.4	$06 \ 52 \ 47$	$12.60 {\pm} 0.25$	1	IC3268	3115
40	*12 21 52.6	$06 \ 30 \ 43$	$12.67 {\pm} 0.25$	1	S122154 + 063148	3116
41	*12 21 58.0	$07 \ 35 \ 46$	$13.98 {\pm} 0.61$	1	N4365	3117
42	*12 22 11.8	$08 \ 48 \ 15$	$14.10 {\pm} 0.60$	1	VCC 740	3118
43	*12 22 27.0	$11 \ 55 \ 45$	$13.84 {\pm} 0.44$	1	N4371	-
44	*12 22 39.9	13 12 19	$13.56 {\pm} 0.63$	1	N4374 ?	3119
45	*12 22 50.9	$10 \ 19 \ 07$	$12.81 {\pm} 0.38$	2	N4380	3122
46	*12 22 54.8	$16 \ 45 \ 08$	$11.75 {\pm} 0.21$	1	N4383	3125

Table 1—Continued

=

Id. no.	α	δ	m_{UV}	N _{obs} Proposed id.		FAUST no.
47	*12 23 11.5	$07 \ 29 \ 20$	$12.93 {\pm} 0.24$	1	IC3322A	3127
48	*12 23 19.4	$10 \ 45 \ 46$	$12.73 {\pm} 0.40$	2	N4390	3129/31
49	*12 23 20.7	12 59 07	$12.35 {\pm} 0.27$	2	N4388	3128
50	*12 23 31.2	$15 \ 57 \ 42$	12.42 ± 0.45	2	N4396	3132
51	*12 23 49.3	$13 \ 15 \ 47$	$13.16 {\pm} 0.79$	3	N4406	3133
52	*12 24 05.3	$12 \ 57 \ 21$	$12.07 {\pm} 0.31$	3	N4413 or HD108285 ?	3134
53	*12 24 09.6	$09\ 07\ 19$	11.32 ± 0.23	1	N4411A	3136
54	*12 24 12.7	09 10 28	$11.85 {\pm} 0.17$	1	N4411B	3137
55	*12 24 19.3	$08 \ 12 \ 43$	$12.41 {\pm} 0.20$.41±0.20 1 N4416		3139
56	*12 24 35.9	08 52 31	$13.66 {\pm} 0.25$	1 No ID		3140
57	*12 24 41.8	$07 \ 33 \ 17$	$12.27 {\pm} 0.17$	0.17 1 VCC 975 ?		3142
58	*12 24 42.4	$09 \ 42 \ 49$	$14.16 {\pm} 0.59$	1	N4424	3144
59	*12 24 44.3	$16 \ 10 \ 34$	$12.77 {\pm} 0.64$	2	IC3365	3143
60	12 24 49.0	$07 \ 25 \ 30$	$12.14 {\pm} 0.18$	1	SAO119404	3145
61	12 24 52.0	07 39 31	$9.55 {\pm} 0.16$	1	SAO119405	3147
62	*12 24 57.0	$08 \ 54 \ 44$	$13.50 {\pm} 0.26$	1	No ID	3151
63	*12 24 57.5	06 32 02	$12.02 {\pm} 0.21$	1	N4432	3148
64	*12 25 02.8	$11 \ 35 \ 35$	$11.11 {\pm} 0.20$	3	HD108452	3150
65	*12 25 15.2	$11 \ 55 \ 25$	11.23 ± 0.24	3	No ID	3154
66	*12 25 17.3	13 20 26	$12.38 {\pm} 0.39$	3	N4438	3152/3/5
67	$12 \ 25 \ 27.0$	$09 \ 37 \ 44$	$12.21 {\pm} 0.49$	2	SAO119416	3156/7
68	*12 25 41.2	$15 \ 52 \ 29$	$13.01 {\pm} 0.72$	2	SAO100112	3158
69	*12 25 51.4	09 01 34	$12.69 {\pm} 0.23$	1	VCC 1091 ?	3159

Table 1—Continued

Id. no.	α	$lpha$ δ m_{UV}		N_{obs}	Proposed id.	FAUST no.
70	*12 25 52.4	$12 \ 13 \ 04$	15.46::	3	S122551+121247	-
71	*12 26 12.7	09 33 03	$13.77 {\pm} 0.46$	1	N4451	3162
72	*12 26 16.6	$07 \ 26 \ 10$	$9.12 {\pm} 0.16$	1	S122614 + 072530	3164
73	*12 26 30.9	$05 \ 57 \ 10$	$13.29 {\pm} 0.76$	1	SAO119423	3165
74	*12 26 32.0	10 58 46	$13.03 {\pm} 0.52$	2	S122644 + 105941	3166
75	*12 26 33.8	$06\ 15\ 13$	$11.89 {\pm} 0.21$	1	SAO119424	3167
76	*12 26 45.3	11 58 58	$13.85 {\pm} 0.98$	2	PG 1226+119	3168
77	12 26 49.0	$17 \ 35 \ 56$	$7.35 {\pm} 0.16$	1	SAO100122	3169
78	*12 26 59.9	$07 \ 03 \ 11$	13.11±0.31 1 IC3414		3174	
79	*12 27 01.3	$06 \ 31 \ 44$	12.64 ± 0.26 1 SAO119429		3173	
80	$12\ 27\ 00.0$	11 46 06	11.97±0.33 3 SAO100126		3172	
81	*12 27 08.5	08 07 08	12.13 ± 0.19	1	N4470	3177
82	*12 27 12.4	$08 \ 37 \ 48$	11.12 ± 0.24	2	S122710+083901	3179/80
83	*12 27 18.0	08 17 31	$12.17 {\pm} 0.19$	1	N4472	3181
84	*12 27 24.4	$13 \ 06 \ 02$	$13.05 {\pm} 0.67$	3	S122720+130503	3182
85	*12 28 04.3	10 01 23	$12.80 {\pm} 0.36$	2	SAO 119436	3188
86	*12 28 09.3	08 49 02	$12.50 {\pm} 0.20$	1	SAO119438	3189
87	*12 28 11.6	09 02 23	$12.34{\pm}0.19$	1	S122810+090112	3190
88	*12 28 18.4	12 41 15	12.02 ± 0.30	3	N4486	3191
89	*12 28 45.9	$05\ 21\ 17$	$9.76 {\pm} 0.16$	1	SAO119454	3197
90	*12 28 57.5	11 48 40	13.45 ± 1.28	3	IC3446	3200
91	*12 29 06.1	17 08 33	$12.09 {\pm} 0.26$	1	N4489	3207
92	*12 29 07.1	$08 \ 03 \ 37$	$13.90 {\pm} 0.38$	1	No ID	3204

Table 1—Continued

Id. no.	α	$lpha$ δ ${ m m}_{UV}$		N _{obs}	Proposed id.	FAUST no.
93	12 29 07.0	12 24 13	11.02 ± 0.19	3	SAO100140	3206
94	*12 29 20.5	$15 \ 24 \ 18$	$13.13 {\pm} 0.65$	2	IC0797	3214
95	*12 29 27.4	$14 \ 42 \ 34$	$11.10 {\pm} 0.18$	2	N4501	3211
96	*12 29 35.8	$12 \ 34 \ 10$	13.07 ± 0.44	2	S122936 + 123224	3216
97	*12 29 38.7	08 02 36	12.92 ± 0.25	1	S122935+080140	-
98	*12 29 47.4	06 00 28	12.35 ± 0.28	1	SAO119459	3219
99	*12 29 55.8	09 41 09	$13.10 {\pm} 0.80$	2	No ID	3221
100	*12 30 12.9	$14 \ 20 \ 47$	12.07 ± 0.38	3	IC3476	3222
101	*12 30 16.9	09 50 33	$13.89 {\pm} 0.39$	1	No ID	3224
102	*12 30 29.4	13 13 11	$13.13 {\pm} 0.73$	3	S123030+131240	3228
103	*12 30 30.4	08 12 41	12.49 ± 0.21	1	S123028+081156	3226
104	*12 30 34.3	$05 \ 18 \ 07$	$9.43 {\pm} 0.16$	1	No ID	-
105	*12 30 51.7	$06 \ 41 \ 56$	$8.74 {\pm} 0.16$	1	PG 1230+067	3231
106	*12 30 54.0	09 49 10	$13.92{\pm}1.20$	2	Q $1230 + 0947$	3233
107	*12 30 59.5	08 55 05	11.13 ± 0.23	2	N4519/4519a	3235/6
108	*12 31 05.8	09 11 50	$13.81 {\pm} 0.52$	1	SAO119467	3237
109	*12 31 07.9	09 26 30	12.26 ± 0.45	2	N4522	3238
110	*12 31 13.7	$15 \ 25 \ 36$	$11.87 {\pm} 0.30$	2	N4523	3240
111	*12 31 21.8	08 18 11	$14.17 {\pm} 0.63$	1	VCC 1525	-
112	* 12 31 37.0	$17 \ 13 \ 01$	$11.53 {\pm} 0.20$	1	No ID	3248
113	*12 31 46.9	$06 \ 44 \ 51$	$10.95 {\pm} 0.16$	1	N4532	3249
114	*12 31 49.4	08 19 13	$10.18 {\pm} 0.16$	1	N4535/4535a	-
115	12 31 48.0	11 11 46	$10.91 {\pm} 0.18$	3	SAO100149	3250

Table 1—Continued

Id. no.	lpha	δ	m_{UV}	N_{obs}	Proposed id.	FAUST no.
116	*12 32 00.4	$15 \ 45 \ 23$	12.81 ± 0.52	2	No ID	-
117	$12 \ 32 \ 20.0$	$11 \ 06 \ 43$	$12.74{\pm}0.18$	3	SAO100156	3258
118	*12 32 26.7	15 50 25	$13.16 {\pm} 0.36$	1	IC3528	3256
119	*12 32 34.0	$13 \ 36 \ 32$	$12.65 {\pm} 0.48$	3	No ID	3260
120	12 32 37.0	$18 \ 39 \ 07$	$8.92 {\pm} 0.16$	1	SAO100159	3260
121	*12 32 41.3	10 48 22	$13.32 {\pm} 0.52$	2	VCC 1600 ?	3263
122	*12 32 53.3	$09 \ 44 \ 57$	$10.49 {\pm} 0.16$	2	S123252+094327	3268
123	*12 32 55.9	14 46 40	$11.90 {\pm} 0.27$	2	N4548	3267
124	*12 33 15.4	$12 \ 51 \ 41$	$13.10 {\pm} 0.68$	3	N4552	-
125	*12 33 59.5	$07 \ 54 \ 46$	$13.41 {\pm} 0.37$	1	No ID	3278
126	*12 34 03.7	11 33 19	$11.57 {\pm} 0.26$	3	N4567 or N4568	3280
127	*12 34 05.0	06 53 07	$12.63 {\pm} 0.25$	1	IC3576	3279
128	12 34 13.0	$15 \ 32 \ 21$	12.11 ± 0.33	2	SAO100173	3287
129	*12 34 16.4	$13\ 28\ 44$	$11.38 {\pm} 0.21$	3	N4569	3281
130	12 34 17.0	08 12 30	$12.37 {\pm} 0.19$	1	SAO119494	3284
131	12 34 31.0	14 31 38	$11.86 {\pm} 0.25$	2	N4571 or SA0100177	3286
132	*12 34 33.8	$07 \ 11 \ 44$	$12.99 {\pm} 0.31$	1	IC3591	3288
133	12 34 35.0	09 04 19	$9.28 {\pm} 0.16$	2	SAO119497	3290
134	*12 35 09.4	$10 \ 04 \ 44$	$12.25 {\pm} 0.30$	2	S123507+100344	3299
135	*12 35 10.8	15 50 16	$10.73 {\pm} 0.18$	2	S123515 + 155104	3297
136	*12 35 11.7	08 50 27	$13.10 {\pm} 0.27$	1	VCC 1725	3298
137	*12 35 14.2	$07 \ 22 \ 44$	13.62 ± 0.51	1	U7795	3300
138	*12 35 14.9	12 07 11	12.03 ± 0.32	3	N4579	3302

Table 1—Continued

Id. no.	α	δ	m_{UV}	N_{obs}	Proposed id.	FAUST no.
120	*10.26.00.0	10 20 22	14.22	0	C102550 + 102120	
139	12 30 00.0	10 32 33	14.33::	2	5123558+105139	-
140	*12 36 24.6	16 41 21	11.47 ± 0.26	2	SAO100194	3311
141	*12 36 29.1	$14 \ 03 \ 36$	$11.16 {\pm} 0.19$	2	SAO100192	3319
142	12 36 32.0	$17 \ 42 \ 42$	$11.62 {\pm} 0.23$	1	SAO100193	3323
143	12 36 39.0	$14 \ 36 \ 06$	$11.72 {\pm} 0.23$	2	SAO100195	3322
144	*12 36 53.3	08 14 14	13.17 ± 0.32	1	IC3617	3331
145	*12 37 09.5	$13 \ 14 \ 25$	$10.42 {\pm} 0.17$	3	PG 1237+132	3336
146	*12 37 19.7	$15 \ 33 \ 59$	12.57 ± 0.42	2	N4595	3339
147	*12 37 34.8	$10\ 26\ 31$	12.50±1.04 2 N4596		3342	
148	*12 37 35.1	$11 \ 49 \ 24$	11.80±0.28 3 PG 1237+		PG 1237+118	3347
149	*12 37 41.6	$15 \ 05 \ 17$	12.72 ± 0.47	2	S123734 + 150560	3343
151	*12 37 49.7	09 24 00	$13.68 {\pm} 0.47$	1	SAO119524	3352
152	*12 37 55.5	$10\ 15\ 57$	$11.76 {\pm} 0.26$	2	S123754 + 101551	3355
150	$12 \ 37 \ 59.0$	$16 \ 33 \ 12$	$12.50 {\pm} 0.19$	2	SAO100201	3357
153	12 38 22.0	09 06 13	$11.50 {\pm} 0.17$	1	SAO119530	3360
154	*12 38 34.8	$07 \ 15 \ 16$	$10.49 {\pm} 0.16$	1	No ID	3365
155	*12 38 39.5	08 28 07	$13.04 {\pm} 0.32$	1	KARA 68181 ?	3367
156	*12 38 52.3	09 58 23	$13.08 {\pm} 0.36$	1	S123853+095836	-
157	*12 38 52.3	$10 \ 53 \ 13$	$12.41 {\pm} 0.21$	1	S123847+105318	3371
158	12 39 03.0	$10 \ 42 \ 02$	$7.63 {\pm} 0.16$	2	SAO100207	3375
159	12 39 17.0	10 09 36	$9.76 {\pm} 0.16$	2	SAO100210	3380
160	12 39 21.0	10 30 39	$5.54 {\pm} 0.16$	2	SAO100211	3382
161	12 39 25.0	$07 \ 04 \ 51$	$6.17 {\pm} 0.16$	1	SAO119538	3381

Table 1—Continued

Id. no.	α	δ	m_{UV}	N _{obs}	Proposed id.	FAUST no.
162	*12 39 27.0	$17 \ 47 \ 19$	$6.95 {\pm} 0.16$	1	PG 1239+178	3385
163	*12 39 31.0	08 55 18	14.10 ± 0.84	1	S123934 + 085455	_
164	*12 39 37.5	$16 \ 37 \ 19$	12.10 ± 0.36	2	TD1 26442 ?	3389
165	12 39 39.0	11 09 26	11.08 ± 0.19	2	SAO100213	3388
166	*12 39 42.3	08 18 15	$13.68 {\pm} 0.55$	1	SAO119543	3387
167	*12 40 02.2	$12 \ 00 \ 58$	11.14 ± 0.23	3	PG 1240+120	3394
168	*12 40 13.6	$14 \ 35 \ 59$	$12.54 {\pm} 0.44$	2	N4634	3393
169	*12 40 23.6	$13 \ 53 \ 25$	$13.25 {\pm} 0.72$	13.25 ± 0.72 2 S124024+135325		3399
170	*12 40 24.7	$13 \ 32 \ 07$	11.71 ± 0.24	11.71 ± 0.24 2 N4639		3400
171	*12 41 07.6	$11 \ 50 \ 42$	11.40 ± 0.20 2 N4649 or N4647		3409	
172	*12 41 16.6	$16 \ 40 \ 09$	$10.97 {\pm} 0.18$	10.97 ± 0.18 1 N4651		3414
173	12 41 20.8	$15 \ 23 \ 09$	$11.51 {\pm} 0.25$	2	SAO100228	3415
174	*12 41 28.9	$13\ 24\ 24$	$10.86 {\pm} 0.18$	2	N4654	3417
175	12 41 30.0	11 11 42	$11.05 {\pm} 0.19$	2	SAO100229	3419
176	*12 41 57.0	$10 \ 37 \ 21$	$11.93 {\pm} 0.30$	2	S124155 + 103724	3426
177	*12 43 01.6	14 39 18	$5.57 {\pm} 0.16$	2	SAO100235+100236	3441
178	*12 43 01.9	$07 \ 56 \ 12$	$8.65 {\pm} 0.16$	1	SAO119574=FM Vir	3442
179	*12 43 070	$13 \ 37 \ 43$	12.72 ± 0.55	2	IC3742	-
180	*12 43 10.0	$10 \ 35 \ 35$	$10.74 {\pm} 0.19$	1	No ID	-
181	*12 43 50.4	09 32 03	$10.66 {\pm} 0.17$	1	SAO119582	3454
182	12 44 08.8	$16 \ 51 \ 01$	$10.45 {\pm} 0.17$	1	SAO100249/252/257	3456
183	*12 44 31.8	11 19 14	$10.05 {\pm} 0.17$	2	PG 1244+113	3460
184	12 44 42.6	$12 \ 13 \ 52$	$6.74{\pm}0.16$	1	SAO100260	3461

Id. no.	$lpha$ δ		m_{UV} N_{obs}		Proposed id.	FAUST no.	
185	*12 44 42.3	$13 \ 53 \ 47$	$11.54 {\pm} 0.36$	2	SAO100258	3463	
186	*12 45 11.8	$13 \ 35 \ 55$	$11.98 {\pm} 0.22$	1	S124507+133509	3469	

Table 1—Continued

Id. no.	α	δ	m_{UV}	N _{obs}	Proposed id.	FAUST no.
107	*10 AF 01 C	14 00 10	11 50 10 91	0	N4690	2472
187	12 45 21.0	14 02 13	11.39 ± 0.31	2	N4089	3473
188	*12 45 41.0	$09\ 21\ 21$	$11.30 {\pm} 0.21$	1	SAO119602	3480
189	$12 \ 45 \ 44.1$	13 49 33	$6.23 {\pm} 0.16$	2	SAO100269	3483
190	*12 46 10.8	$12 \ 26 \ 28$	$9.10 {\pm} 0.16$	1	S124615+122650	3491
191	12 46 23.8	$14 \ 23 \ 43$	$7.20{\pm}1.54$	2	SAO100283	3497

Table 1—Continued

Note. — 1. The adopted UV magnitude is the average of the different FAUST measurements. The error is the root-mean-square of the the individual errors. 2. The : indicates a very large measurement error. 3. N_{obs} is the number of times a source was observed, *i.e.*, the number of FAUST images in which it appears. 4. The question mark ? at N44, N57, and N69 indicates that the proposed identification is a bit off the FAUST source. 5. N52: The UV source may be a blend of the emission from the galaxy and the HD108285 (A2) star. 6. N155: The three stars near the FAUST source have been observed spectroscopically and identified with late G or K star. The proposed identification is the faint KARA 68181 galaxy marked on the overlays of the PSS. Note that there is a dE galaxy (VCC 1866) a bit far from the FAUST source. 7. N164: This source is near the TD-1 26442 from the Extended TD-1 catalog (preprint). On the PSS there is a faint V> 15 blue looking candidate 123945+163743 not observed.

Source no	ID	Sp. Type	V	R V	II B	FAUST	Other IDa
Source no.	ID	Sp. Type	v	D - V	U = D	PAUSI	Other IDs
3	SAO099997	A2m	5.8	0.26	.09	$8.40 {\pm} 0.16$	HIC 59608
4	SAO119277	A2	7.6	0.16		$8.83 {\pm} 0.16$	HIC 59636
5	SAO100006	F5	8.7	0.48		$12.69 {\pm} 0.25$	HIC 59737
6	SAO119294	A2	8.85	0.28		$11.06 {\pm} 0.17$	HIC 59804
7	SAO119300	A3	8.40	0.32		$10.70 {\pm} 0.17$	HIC 59852
8	SAO100012	A3	5.10	0.06		$6.10 {\pm} 0.17$	HIC 59819
9	S121426+081714	B(e)	11.9			$9.84{\pm}0.16$	
10	SAO119308	F2	7.9	0.40		$11.65 {\pm} 0.23$	HIC 59916
11	S121449+155138	B(l)	11.7			$10.42 {\pm} 0.18$	
13	S121508+095341	A(l)	14.4			$12.67 {\pm} 0.24$	
14	SAO100024	B9	9.81	-0.12		$7.62 {\pm} 0.16$	HIC 59955
16	SAO100029	A3	8.89	0.17		$9.86{\pm}0.77$	HIC 59992
18	SAO100034	F0	9.3			$13.40 {\pm} 0.26$	HD107089
19	HD107116	A3	10.38	0.23		$11.94{\pm}0.23$	HIC 60065
21	SAO119328	A0	8.60	0.09		$9.43 {\pm} 0.16$	HIC 60095
22	HD107227	A3	10.8			$12.94{\pm}0.26$	
24	S121757+101353	G(e)	13.2			$8.92 {\pm} 0.16$	
25	S121836+092743	A(e)	12.6			$13.63 {\pm} 0.41$	
27	S121860+104153	B(l)	12.9			$12.41 {\pm} 0.72$	
30	SAO100062	A2	6.66	0.05	.05	$6.81 {\pm} 0.16$	HIC 60313
33	SAO100071	F5	8.55	0.33		11.12 ± 0.20	HIC 60419
36	SAO119371	F0	8.83	0.32		$11.38 {\pm} 0.16$	HIC 60473
37	S122129+062746	A(e)	12.5			$12.71 {\pm} 0.26$	
40	S122154 + 063148	B(l)	12.2			$12.67 {\pm} 0.25$	
60	SAO119404	F2	8.43	0.44		12.14 ± 0.18	HIC 60766

Table 2. Stars towards the Virgo cluster

Table 2—Continued

Source no.	ID	Sp. Type	V	B-V	U-B	FAUST	Other IDs
61	SAO119405	A0	8.35	0.43		$9.55{\pm}0.16$	HIC 60772
64	HD108452	A0	10.0			$11.11 {\pm} 0.20$	
67	SAO119416	F5	9.4			$12.21 {\pm} 0.49$	HD108535
68	SAO100112	F8	8.6	0.42		$13.01{\pm}0.72$	HIC 60836
70	S122551+121247	A(l)	> 15			15.46::	
72	S122614+072530	A(l)	10.8			$9.12{\pm}0.16$	
73	SAO119423	A2	8.7			$13.29 {\pm} 0.74$	HD108663
74	S122644 + 105941	F(l)	12.6			$13.03 {\pm} 0.52$	
75	SAO119424	F2	8.1	0.44		$11.89 {\pm} 0.21$	HIC 60915
76	PG 1226+119	HBB	15.37 (B)			$13.85 {\pm} 0.98$	
77	SAO100122	A0	7.7	0.11	01	$7.35 {\pm} 0.16$	HIC 60933
79	SAO119429	$\mathrm{G0}$	8.7			$12.64{\pm}0.26$	HD108738
80	SAO100126	F0	8.53	0.40		$11.97 {\pm} 0.33$	HIC 60945
82	S122710+083901	B(l)	11.5			$11.12 {\pm} 0.24$	
84	S122720+130503	A(l)	10.5			$13.05 {\pm} 0.67$	
85	SAO119436	F2	7.5	0.48		$12.80{\pm}0.36$	HIC 61035
86	SAO119438	A5	8.8			$12.50 {\pm} 0.20$	HD108876
87	S122810+090112	A(l)	11.2			$12.34{\pm}0.19$	
89	SAO119454	F2	7.8	0.36		$9.76 {\pm} 0.16$	HIC 61104
93	SAO100140	$\mathrm{F0}$	8.1	0.34		$11.02{\pm}0.19$	HIC 61135
96	S122936+123224	A(e)	10.9			$13.07 {\pm} 0.43$	
97	S122935+080140	A(e)	11.			$12.92{\pm}0.25$	
98	SAO119459	$\mathrm{F0}$	8.9			$12.35 {\pm} 0.28$	HD109129
102	S123030+131240	B(l)	13.2			$13.13 {\pm} 0.73$	
103	S123028+081156	B(l)	10.5			$12.49 {\pm} 0.21$	

Table 2—Continued

=

C	ID	C T	X 7				
Source no.	ID	Sp. Type	v	D - V	U - D	FAUST	Other IDs
105	PG 1230+067	sdO-B	13.4 (B)			$8.74 {\pm} 0.16$	
108	SAO119467	F8	8.9	0.56		$13.81 {\pm} 0.52$	HIC 61297
115	SAO100149	A5	7.6	0.40		$10.91{\pm}0.18$	HIC 61353
117	SAO100156	F0	7.9	0.30		$12.74{\pm}0.18$	HIC 61397
120	SAO100159	A7m	6.6	0.27		$8.92{\pm}0.16$	HIC 61415
122	S123252+094327	B(l)	11.4			$10.49 {\pm} 0.16$	
128	SAO100173		8.7			$12.11 {\pm} 0.33$	
130	SAO119494	F8	8.9	0.56		$12.37 {\pm} 0.19$	HIC 61561
133	SAO119497	A2	6.6	0.25		$9.28{\pm}0.16$	HIC 61579
134	S123507 + 100344	B(l)	10.6			$12.25 {\pm} 0.30$	
135	S123515 + 155104	B(l)	10.0			$10.73 {\pm} 0.18$	
139	S123558 + 103139	A(e)	11.2			14.33::	
140	SAO100194	F3	7.6	0.45		$11.47 {\pm} 0.26$	HIC 61727
141	SAO100192	F8	8.9			$11.16{\pm}0.19$	HD109997
142	SAO100193	F0	9.45	0.31		$11.62 {\pm} 0.23$	HIC 61718
143	SAO100195	A0m	7.3	0.99		$11.72 {\pm} 0.23$	HIC 61728
145	PG 1237+132	sdB	13.96 (B)			$10.42 {\pm} 0.17$	TON 97
148	PG 1237+118	sdB-O	15.7	13	-1.21	$11.80{\pm}0.28$	TON 98
149	S123734 + 150560	A(e)	13.5			$12.72 {\pm} 0.47$	
150	SAO100201	F5	7.9	0.39		$12.50 {\pm} 0.19$	HIC 61844
151	SAO119524	F8	8.9			$13.68 {\pm} 0.47$	HD110197
152	S123754 + 101551	A(e)	> 15			$11.76 {\pm} 0.26$	
153	SAO119530	F2	7.2	0.43		$11.50{\pm}0.17$	HIC 61889
156	S123853+095836	A(e)	12.7			$13.08 {\pm} 0.36$	
157	S123847+105318	A(l)	12.2			$12.41 {\pm} 0.21$	

Table 2—Continued

Source no.	ID	Sp. Type	V	B-V	U - B	FAUST	Other IDs
159	SAO100207	Δ.7	6 9	0.10		7.62 ± 0.16	HIC 61027
150	SAO100207		0.2	0.19		7.03 ± 0.10	
159	SAO100210	FU	7.0	0.32		9.70±0.10	HIC 61950
160	SAO100211	A0	4.9	0.08		5.54 ± 0.16	HIC 61960
161	SAO119538	A2	5.6	0.00	-0.01	6.17 ± 0.16	HIC 61968
162	PG 1239+178	sdO	10.95	32	-1.22	$6.95 {\pm} 0.16$	BD+18°2647
163	S123934 + 085455	A(l)	12.2			14.10 ± 0.84	
164	TD1 26442	?				$12.10 {\pm} 0.36$	
165	SAO100213	A0	8.8			$11.08 {\pm} 0.19$	HD110466
166	SAO119543	F8	7.9	0.50		$13.68 {\pm} 0.55$	HIC 61990
167	PG 1240+120	sdO-B	15.3 (B)			$11.14 {\pm} 0.23$	Ton 109
169	S124024 + 135325	В	> 15			$13.25 {\pm} 0.72$	
173	SAO100228	F2	8.8	0.27		$11.51 {\pm} 0.25$	HIC 62116
175	SAO100229	F2	8.6	0.32		$11.05 {\pm} 0.19$	HIC 62133
176	S124155+103724	B(l)	12.0			$11.93 {\pm} 0.30$	
177	SAO100235+100236	A0				$5.57 {\pm} 0.16$	
178	SAO119574	A8m	5.2	0.32		$8.65 {\pm} 0.16$	FM Vir
181	SAO119582	F0	7.7			$10.66 {\pm} 0.17$	HD111043
182	SAO100249/252/257	F-K	5			$10.45 {\pm} 0.17$	HD111067
183	PG 1244+113	sd B	13.28	23	-1.19	$10.05 {\pm} 0.17$	
184	SAO100260	A3	6.1	0.12		$6.74{\pm}0.16$	HIC 62394
185	SAO100258	F5	8.2			11.54 ± 0.36	HD111155
186	S124507 + 133509	B(e)	12.1			11.98 ± 0.22	
188	SAO119602	E(0)	7.0	0.42		11.00 ± 0.22 11.30 ± 0.21	HIC 62480
100	SAO100260	1 U A 1	1.3 6 F	0.42		6.92 ± 0.16	HIC 69479
189	SAU100209		6.0	0.02		0.23 ± 0.10	піс 02478
190	S124615 + 122650	B(I)	10.4			9.10 ± 0.16	

Source no.	ID	Sp. Type	V	B - V	U - B	FAUST	Other IDs
191	SAO100283	A1	5.7	0.02		$7.20{\pm}1.54$	HIC 62541

Table 2—Continued

m_{UV}	B0 - B9	A0-A9	F0-F9	G0-G9	HBB, sdO/B	AGN/Galaxies	No ID	No Sp
5-6	0	2	0	0	0	0	0	0
6-7	0	5	0	0	1	0	0	0
7–8	1	3	0	0	0	0	0	0
8-9	0	4	0	1	1	0	0	0
9–10	2	5	2	0	0	0	1	0
10-11	3	2	2	0	2	6	2	0
11 - 12	3	6	14	0	2	16	2	0
12–13	4	8	8	1	0	31	3	2
13–14	2	5	6	0	1	20	6	0
14 - 15	0	2	0	0	0	3	0	0
>15	0	1	0	0	0	0	0	0
Total	15	43	32	2	7	76	14	2

Table 3. Distribution of UV sources

Note. — Sources marked "No Sp." are tentatively identified with objects on the PSS, which have not been observed spectroscopically.

Source no.	Name	Type	FAUST	4xE(B-V)	GUV	SCAP	S+C
2	N4178	$\mathrm{SBc}(\mathrm{s})\mathrm{II}$	00	$11.61 {\pm} 0.19$			12.34
12	IC3111	Sb(s)II	05	$13.03 {\pm} 0.32$			
15	N4246	Sc(s)I-II	06	$12.83 {\pm} 0.35$			
17	N4254	Sc(s)I.3	.14	$10.08 {\pm} 0.17$		10.1	10.74
20	IC0776	$\mathrm{SBcd}(\mathbf{s})\mathrm{III}$	03	$13.42 {\pm} 0.31$			
23	N4276	Sc(s)II	02	$12.90 {\pm} 0.31$			15.82
26	N4294	SBc(s)II-III	.02	$11.57 {\pm} 0.17$			13.07
28	N4299	Scd(s)III	.02	$11.58 {\pm} 0.17$			12.92
29	N4298	Sc(s)III	.10	$12.21 {\pm} 0.31$			12.94
31	N4321	Sc(s)I	.04	$10.07 {\pm} 0.17$		10.1	10.54
32	N4325	E4	01	$13.73 {\pm} 0.97$			>16.30
34	N4330	Sd(on edge)	.03	$13.28 {\pm} 0.35$			14.39
35	IC3258	ScIII-IV	.07	$12.46 {\pm} 0.24$	$14.69 {\pm} 0.75$		13.37
38	N4351	Sc(s)II.3	.01	$12.77 {\pm} 0.38$	$13.16 {\pm} 0.20$		13.97
39	IC3268	Sc(s)III-IV or Sm	02	$12.60 {\pm} 0.25$			
41	N4365	E3	03	$13.98 {\pm} 0.61$			13.64
42	VCC 740	SBmIII	03	$14.10 {\pm} 0.60$			
43	N4371	$\mathrm{SB0}_2(\mathrm{r})(3)$.02	$13.84{\pm}0.44$	>14.58		15.39
44	N4374 ?	E1	.13	$13.56 {\pm} 0.63$	$14.38 {\pm} 0.6$		12.59
45	N4380	Sab(s)	01	$12.81 {\pm} 0.38$	$13.17 {\pm} 0.22$		13.97
46	N4383	Amorphous	.02	$11.75 {\pm} 0.21$		12.1	12.59
47	IC3322A	Sc(on edge)	02	$12.93 {\pm} 0.24$			14.79
48	N4390	$\mathrm{Sbc}(\mathrm{s})\mathrm{II}$.00	$12.73 {\pm} 0.40$	$13.70 {\pm} 0.35$		13.62
49	N4388	Sab	.11	$12.35 {\pm} 0.27$	$12.83 {\pm} 0.18$		13.42
50	N4396	Sc(s)II	.04	$12.42 {\pm} 0.45$		12.6	13.04

Table 4. Galaxies towards the Virgo cluster-UV data

Table 4—Continued

Source no.	Name	Type	FAUST	4 x E(B-V)	GUV	SCAP	S+C
51	N4406	$S0_1(3),E3$.11	$13.16 {\pm} 0.79$	>14.82		13.22
52	N4413 ?	SBbc(rs)II-III	.11	$12.07 {\pm} 0.31$	$13.11 {\pm} 0.20$		14.32
53	N4411A	SBc(s)II	03	$11.32{\pm}0.23$			15.39
54	N4411B	Sc(s)II	02	$11.85 {\pm} 0.17$	$9.90{\pm}0.13$		13.49
55	N4416	SBc(s)II.2	02	$12.41 {\pm} 0.20$	14.08 ± 0.39		>16.30
57	VCC 975 ?	Scd(s)II	02	$12.27 {\pm} 0.17$			
58	N4424	Sa pec	03	$14.16 {\pm} 0.59$	$13.61 {\pm} 0.31$		14.14
59	IC3365	Scd(s)III	.03	$12.77 {\pm} 0.64$			
63	N4432	Sc(s)I-II	.12	$12.02 {\pm} 0.21$			
66	N4438	Sb(tides)	.10	$12.38 {\pm} 0.39$	$13.83 {\pm} 0.31$		13.49
69	VCC 1091 ?	Sbc(s)I.8	03	$12.69 {\pm} 0.23$			
71	N4451	Sc(s)III	03	$13.77 {\pm} 0.46$	>14.82		14.64
78	IC3414	Sc(s)II	03	$13.11 {\pm} 0.31$			
81	N4470	ScIII pec	02	$12.13 {\pm} 0.19$	$12.60 {\pm} 0.13$		13.84
83	N4472	$E2/S0_{1}(2)$	02	$12.17 {\pm} 0.19$	$13.01 {\pm} 0.20$		12.47
88	N4486	E0	.09	$12.02 {\pm} 0.30$	12.24 ± 0.12		12.68
90	IC3446	SmIII/BCD	.03	13.45 ± 1.28			
91	N4489	$S0_1(1)$.03	$12.09 {\pm} 0.26$			
94	IC0797	SBc(s)II-III	.09	$13.13 {\pm} 0.65$			
95	N4501	$\mathrm{Sbc}(\mathrm{s})\mathrm{II}$.09	$11.10{\pm}0.18$	$12.87 {\pm} 0.22$		11.29
100	IC3476	Sc(s)II.2	.09	$12.07 {\pm} 0.38$	$12.87 {\pm} 0.30$		13.64
106	Q 1230+0947		03	$13.92{\pm}1.29$			
107	N4519/4519a	SBc(rs)II.2, dSO?	02	$11.13 {\pm} 0.16$	$11.59{\pm}0.07$		13.04
109	N4522	$\rm Sc/Sb$	04	$12.26 {\pm} 0.45$	$13.52 {\pm} 0.24$		14.29
110	N4523	SBd(s)III	.09	$11.87 {\pm} 0.30$			13.07

Table 4—Continued

_

=

	Source no.	Name	Type	FAUST	4xE(B-V)	GUV	SCAP	S+C
_	111	VCC 1525	Sbc(r) II	- 01	14.17 ± 0.63			
	111	N4532	Speciel II	01	14.17 ± 0.03 10.05 ± 0.16			
	115	N4535 / 4535a	$S = S B_{\alpha}(a) I 2$	04	10.95 ± 0.10 10.18 \pm 0.16	10 00±0 04		11 20
	114	114000/4000a	S, SDC(S)I.3	01	10.10 ± 0.10	10.90 ± 0.04		11.39
	110	103028 VCC 1600 2		.04	13.10 ± 0.30			
	121	VCC 1600 ?	dE	01	13.32 ± 0.52			
	123	N4548	SBb(rs)I-II	.07	11.90 ± 0.27			12.77
	124	N4552	$S0_1(0)$.14	13.10 ± 0.68	13.30 ± 0.30		13.42
	126	N4567/N4568	Sc(s)II-III+Sc(s)III	.01	11.57 ± 0.26	12.15 ± 0.10		12.22
	127	IC3576	SBdIV	03	$12.63 {\pm} 0.25$			
	129	N4569	Sab(s)I-II	.08	$11.38 {\pm} 0.21$	$12.46 {\pm} 0.13$		11.39
	131	N4571	Sc(s)II-III	.07	$11.86 {\pm} 0.25$			
	132	IC3591	IB	03	$12.99 {\pm} 0.31$			
	136	VCC 1725	SmIII/BCD	02	$13.10 {\pm} 0.27$			
	137	U7795	Sdm	05	$13.62 {\pm} 0.51$			
	138	N4579	Sab(s)II	.14	$12.03 {\pm} 0.32$	$12.84{\pm}0.18$		12.37
	144	IC3617	SmIII/BCD	01	$13.17 {\pm} 0.32$			14.54
	146	N4595	Sc(s)II.8	.02	$12.57 {\pm} 0.42$			14.39
	147	N4596	SBa	03	12.50 ± 1.04			14.64
	155	KARA 68181		.02	$13.04{\pm}0.32$			
	168	N4634	Sc (on edge)	03	1254 ± 044		135	
	170	N/630	SBb(r)II	.00	11.71 ± 0.24		19.0	19 79
	170	N4640/N4647	SDO(1)	.00	11.71 ± 0.24 11.40 ± 0.20		12.4	11.07
	170	114045/114047 N4651	$SU_1(2) + SU_1(15)$.04	11.40 ± 0.20		12.0	11.07
	1(2	N4051		.03	10.97±0.18			44 OF
	174	N4654	SBc(rs)II	.05	10.86 ± 0.18		11.2	11.67
	178	IC3742	$\mathrm{SBc}(\mathrm{s})\mathrm{II}$.06	$12.72 {\pm} 0.55$			

Source no.	Name	Туре	FAUST	4xE(B-V)	GUV	SCAP	S+C
187	N4689	Sc(s)II.3	.04	$11.59 {\pm} 0.31$		12.2	

Table 4—Continued

Ν	Name	T-Type	FAUST	FI(HI)	Ir [60]	Ir [100]	V_r	Bc_{tot}
2	N4178	5	$11.61 {\pm} 0.19$	56.2	2.11	8.08	381	11.89
12	IC3111	3	$13.03 {\pm} 0.32$.24540	.67	0.72	14755	14.78
15	N4246	5	12.83 ± 0.35	21.5			3724	13.38
17	N4254	5	$9.80 {\pm} 0.17$	105.6	25.3	76.01	2470	10.30
20	IC0776	6	13.42 ± 0.31	10.2			2464	14.30
23	N4276	5	$12.90 {\pm} 0.31$	4.66	.48	1.64	2628	13.25
26	N4294	5	11.53 ± 0.17		.28	9.40	359	12.51
28	N4299	6	$11.54{\pm}0.17$	13.5	2.64	8.09	221	12.86
29	N4298	5	12.01 ± 0.31	12.9	5.09	21.27	1133	11.94
31	N4321	5	$9.99 {\pm} 0.17$	50.0	18.95	56.19	1560	10.01
32	N4325	-5	$13.73 {\pm} 0.97$.75	1.77	7786	14.30
34	N4330	7	13.22 ± 0.35		.63	2.78	1552	13.06
35	IC3258	5	12.32 ± 0.24	3.69	.49	.97	-437	13.57
38	N4351	5	$12.75 {\pm} 0.38$	4.17	.71	2.01	2320	13.02
39	IC3268	$5,\!9$	$12.60 {\pm} 0.25$	8.54	.69	1.65	697	14.22
41	N4365	-5	$13.98 {\pm} 0.61$	$\leq \! 2.65$			1240	10.52
42	VCC 740	9	$14.10 {\pm} 0.60$				874	15.70
43	N4371	-1	$13.80 {\pm} 0.44$	≤ 3.02			941	11.77
44	N4374 ?	-5	$13.30 {\pm} 0.63$	≤ 2.19	.50	.98	1033	9.96
45	N4380	2	$12.81 {\pm} 0.38$	3.02	.63	3.12	963	12.66
46	N4383		$11.71 {\pm} 0.21$	44.2	7.95	12.32	1694	12.65
47	IC3322A	5	$12.93 {\pm} 0.24$	22.7	1.71	5.39	1001	13.55
48	N4390	4	12.73 ± 0.40	7.07	.75	1.88	1118	13.30

Table 5. Galaxies towards the Virgo cluster-Multispectral data

Table 5—Continued

Ν	Name	T-Type	FAUST	FI(HI)	Ir [60]	Ir [100]	v_r	Bc_{tot}
10	NUCCO	2		0.40	10.04	10.1		44.05
49	N4388	2	12.13 ± 0.27	6.46	10.24	18.1	2535	11.65
50	N4396	5	12.34 ± 0.45		1.15	3.89	-155	13.02
51	N4406	-1, -5	$12.94{\pm}0.79$	≤ 2.51			-221	9.72
52	N4413 ?	4	$11.85 {\pm} 0.31$	3.39	1.04	3.19	96	12.14
53	N4411A	5	$11.32 {\pm} 0.23$	7.50			1280	13.41
54	N4411B	5	$11.85 {\pm} 0.17$	21.88	.40	1.78	1269	12.91
55	N4416	5	12.41 ± 0.20	4.79	.93	2.70	1380	13.14
57	VCC 975 ?	6	12.27 ± 0.17	17.7			932	13.58
58	N4424	1	$14.16 {\pm} 0.59$	2.82	3.31	5.92	432	12.34
59	IC3365	6	$12.71 {\pm} 0.64$	3.52			2336	14.33
63	N4432	5	$11.78 {\pm} 0.21$	5.1	.25	1.27	6403	14.70
66	N4438	3	12.18 ± 0.39	7.76	3.76	11.27	86	10.92
69	VCC 1091 ?	4	12.69 ± 0.23	16.13	.38	.82	1117	14.6
71	N4451	5	$13.77 {\pm} 0.46$	2.68	1.68	4.56	860	13.29
78	IC3414	5	13.11 ± 0.31	3.63	.23	.73	525	13.94
81	N4470	$5,\!13$	12.13 ± 0.19	8.32	1.82	4.46	2360	13.02
83	N4472	-5, -1	12.17 ± 0.19	≤ 0.72			997	9.37
88	N4486	-5	$11.84{\pm}0.30$.39	1.02	1292	9.50
90	IC3446	9	13.39 ± 1.28				1251	14.87
91	N4489	-1	12.03 ± 0.26		1.24	4.03	1507	12.81
94	IC0797	5	12.95 ± 0.65	3.94	.74	2.18	2100	13.46
95	N4501	4	$10.92 {\pm} 0.18$	33.88			2321	10.27
100	IC3476	5	$11.89 {\pm} 0.38$	4.57	1.35	3.24	-169	13.10

Table 5—Continued

Ν	Name	T-Type	FAUST	FI(HI)	Ir [60]	Ir [100]	v_r	Bc_{tot}
100	0.1000.004		12.02 1.20					
106	Q 1230+0947		13.92 ± 1.30					
107	N4519/4519a	5,0	11.13 ± 0.23	51.29	3.88	6.65	1229	12.34
109	N4522	$5,\!3$	12.26 ± 0.45	7.08	1.30	4.20	2316	12.99
110	N4523	7	$11.69 {\pm} 0.30$	25.9	.38	1.30	262	14.33
111	VCC 1525	4	$14.17 {\pm} 0.63$.36	.83	11382	14.81
113	N4532	9	$10.95 {\pm} 0.16$	42.3	8.95	16.21	2010	12.30
114	N4535/4535a	3?,5	$10.18 {\pm} 0.16$	89.13	6.88	23.61	1962	10.50
118	IC3528	4	13.08 ± 0.36	1.4	1.53	5.52	13830	15.20
121	VCC 1600 ?	-5	13.32 ± 0.52					19.00
123	N4548	3	$11.76 {\pm} 0.27$	9.25	1.47	9.44	550	10.89
124	N4552	-1	12.82 ± 0.68	≤ 2.51			322	10.59
126	N4567 or N4568	5	$11.55 {\pm} 0.26$	21.38	16.91	47.88	2186	12.05 +
								11.67
127	IC3576	7	12.63 ± 0.25	14.5			1076	13.70
129	N4569	2	11.22 ± 0.21	12.3	7.56	23.66	-223	10.18
131	N4571	5	$11.72 {\pm} 0.25$	12.8	1.10	5.78	343	11.75
132	IC3591	10	$12.99 {\pm} 0.31$				1635	14.40
136	VCC 1725	9	$13.10 {\pm} 0.27$				1065	14.51
137	U7795	8	$13.62 {\pm} 0.51$	4.82			61	14.54
138	N4579	2	$11.75 {\pm} 0.32$	8.91			1520	10.34
144	IC3617	9	13.17 ± 0.32	6.3	.27	.64	2079	14.37
146	N4595	5	12.53 ± 0.42		.91	2.81	630	12.89
147	N4596	1	$12.50{\pm}1.04$	≤ 3.71	.49	1.28	1870	11.35

Table 5—Continued

								_
Ν	Name	T-Type	FAUST	FI(HI)	Ir [60]	Ir [100]	v_r	Bc_{tot}
155	KARA 68181		$13.00 {\pm} 0.32$					
168	N4634	5	12.48 ± 0.44		4.04	11.42	144	13.13
170	N4639	3	$11.61 {\pm} 0.24$	15.8	1.41	4.63	990	12.19
171	N4649 or N4647	-1, 5	11.32 ± 0.20	$\leq 6.88 \text{ or } 2.1$	5.33	15.42	1095	9.77 +
								11.90
172	N4651	3?	$10.17 {\pm} 0.18$	59.9	5.45	15.57	800	11.36
174	N4654	5	$10.76 {\pm} 0.18$	68.3	13.08	34.91	1044	11.05
179	IC3742	5	$12.60 {\pm} 0.55$	4.24	.25	.94	872	14.00
187	N4689	5	$11.51 {\pm} 0.31$	6.0	2.55	10.99	1616	11.56

Note. — FI(HI) is the HI flux integral, in units of Jy km $\rm s^{-1}.$

















