# THE 1996 SOFT STATE TRANSITIONS OF CYGNUS X-1

S. N. Zhang<sup>1,2</sup>, W. Cui<sup>3</sup>, B. A. Harmon<sup>1</sup>, W. S. Paciesas<sup>1,4</sup>, R.E. Remillard<sup>3</sup> and J. van Paradijs<sup>4,5</sup>

<sup>1</sup>ES-84, NASA/Marshall Space Flight Center, Huntsville, AL 35812, USA <sup>2</sup>Universities Space Research Association

<sup>3</sup>Center for Space Research, Massachusetts Institute of Technology <sup>4</sup>University of Alabama in Huntsville

<sup>5</sup>University of Amsterdam and Center for High-Energy Astrophysics

Accepted for ApJL publication on 1/4/97.

Received \_\_\_\_\_\_\_\_\_\_\_\_\_\_\_\_\_; accepted

# ABSTRACT

We report continuous monitoring of Cygnus  $X-1$  in the  $1.3 - 200 \text{ keV}$  band using ASM/RXTE and BATSE/CGRO for about 200 days from 1996 February 21 to 1996 early September. During this period Cygnus X-1 experienced a hard-to-soft and then a soft-to-hard state transition. The low-energy X-ray (1.3-12 keV) and high-energy X-ray (20-200 keV) fluxes are strongly anti-correlated during this period. During the state transitions flux variations of about a factor of 5 and 15 were seen in the 1.3-3.0 keV and 100-200 keV bands, respectively, while the average 4.8-12 keV flux remains almost unchanged. The net effect of this pivoting is that the total 1.3-200 keV luminosity remained unchanged to within ∼15%. The bolometric luminosity in the soft state may be as high as 50-70% above the hard state luminosity, after color corrections for the luminosity below 1.3 keV. The blackbody component flux and temperature increase in the soft state is probably caused by a combination of the optically thick disk mass accretion rate increase and a decrease of the inner disk radius.

Subject headings: stars: individual (Cygnus X-1, Cyg X-1) — X-rays: stars — black hole physics — accretion disks

## 1. Introduction

Cygnus X-1 is one of the brightest high-energy sources in the sky, with an average 1-200 keV energy flux of  $\sim 3 \times 10^{-8}$  erg/cm<sup>2</sup>/sec. Most of the time its X-ray spectrum is very hard, with a high-energy (> 20 keV) X-ray flux around or above that of the Crab Nebula while its low-energy (< 10 keV) X-ray flux is around 0.5 Crab. Occasionally, the spectrum of Cyg X-1 becomes much softer; then its low-energy X-ray flux increases to  $1 - 2$  Crab while the hard X-ray flux decreases to 0.5 Crab or less. On the basis of the radial-velocity curve of its O9.7 Iab companion star, Webster and Murdin (1971) and Bolton (1971) concluded that the compact star in Cyg X-1 may be a black hole (BH); their results have been confirmed by later detailed work of Gies and Bolton (1986) who concluded that the mass of the compact object in Cyg X-1 is greater than  $7 M_{\odot}$ , but more probably 16  $M_{\odot}$ , far exceeding the theoretical (and observational) upper mass limit of 3.2 M<sub> $\odot$ </sub> for a neutron star. It is thus the first stellar mass BH candidate (BHC) (cf. Liang and Nolan 1984 and Lewin and Tanaka 1995, for reviews and references therein).

Cyg X-1 is often considered as the canonical BHC; many of its characteristics, such as its high X-ray luminosity above 100 keV, the ultra-soft component in its X-ray spectrum, the hard/soft X-ray flux anti-correlation, and the rapid X-ray flux variability, are shared by other systems, which on the basis of a dynamical mass estimate may contain a BH. It is, however, currently not understood why accreting BHs show these characteristics.

In spite of extensive studies over the last three decades, the mass accretion conditions near the central compact object in Cyg X-1 and other BH systems, and the high-energy radiation mechanism are still poorly understood. This remains an outstanding issue in high-energy astrophysics. State transitions

fully covered over a large range in X-ray photon energy may provide us with important clues towards a better understanding of these BH X-ray binaries. During a transition, a rich collection of information may be obtained, such as flux variations at many time scales in all energy bands, energy spectral evolution, correlations between different energy bands, etc; such data should provide useful tests of various theories and models.

Previously, several transitions between the hard state (HS) and soft state (SS) of Cyg X-1 have been observed. In 1971, a soft-to hard (S-to-H) state transition was observed in the 2-20 keV band (Tananbaum *et al.* 1972). In 1975, a complete transition was observed with Ariel V between 3 and 6 keV (Holt et al. 1976), and by Vela between 3 and 12 keV (Priedhorsky et al. 1983). These observations were limited to rather low X-ray energies. The only broad-band observations of a transition were obtained in 1980, with Hakucho between 1 and 12 keV (Ogawara *et al.* 1981) and with HEAO-3 between 48 and 183 keV (Ling *et al.* 1983). However, the SS onset was observed only with HEAO-3 as a rapid decrease in the 48-183 keV flux, and then later the SS was observed with Hakucho only when the 1-12 keV flux was already a factor of 2-3 higher than its HS level before the transition. Therefore, no simultaneous low- and high-energy X-ray observations during S-to-H or hard-to-soft (H-to-S) transitions of Cyg X-1 have been made so far.

In this paper we report results obtained from the near continuous monitoring of Cyg X-1 during a complete state transition from the HS to the SS (Cui 1996; Cui, Focke and Swank 1996; Zhang *et al.* 1996a), and then back to the HS (Zhang et al. 1996b, 1996c), observed simultaneously with ASM/RXTE (1.3-12 keV) and BATSE/CGRO (20-200 keV) for about 200 days from February to September 1996. Our results are complemented by detailed pointing observations with ASCA (0.5-10 keV) (Dotani *et al.* 1996), and with the PCA and HEXTE (2-250 keV) aboard RXTE (Cui, Focke zhang Swank 1996; Belloni *et al.* 1996; Cui *et al.* 1997), and with SAX  $(0.1-300 \text{ keV})$  (Piro *et al.* 1996) and with  $OSSE/CGRO$  (50-600 keV) (Phlips *et al.* 1997). We will focus on variations in the total luminosity during this period and on the correlations between the low-energy and high-energy X-ray fluxes.

# 2. Observational Data

The Rossi X-ray Timing Explorer (RXTE) was launched on 1995 December 30. The All-Sky Monitor (ASM) began normal operation on 1996 February 21. More than 70 sources have been routinely monitored in three energy bands (1.3-3.0, 3.0-4.8 and 4.8-12 keV). A detailed description of the performance of the ASM, as well as its calibration and data reduction procedures, has been given by Levine et al. (1996).

The BATSE experiment is one of the four instruments aboard the Compton Gamma Ray Observatory, and has operated continuously since launch in April 1991. BATSE can monitor the entire hard X-ray (HXR) sky with almost uniform sensitivity for the detection of gamma-ray bursts, solar flares, pulsars and other persistent and transient HXR sources (Fishman et al. 1989). The persistent and transient HXR source monitoring is achieved by using the Earth occultation technique (Harmon et al. 1992) and Earth occultation transform imaging technique (Zhang et al. 1993). Cyg X-1 has been one of the brightest sources in the BATSE database, in which it is detected from 20 to above 300 keV (see Crary et al. 1996, and Paciesas et al. 1996, for recent papers describing results of analyses of the BATSE data on Cyg X-1).

## 3. Light curves and total luminosity variations

The daily averaged ASM light curves in the three energy bands are plotted in figure 1 (upper panel), in units of the total Crab Nebula counting rates detected by ASM in the same energy bins. (The three-channel data on each day would overlap each other for an energy spectrum with the same shape as that of the Crab Nebula (power law with a photon index of  $\sim$  –2). For a steeper spectrum the lower energy channel data will lie above the higher energy data and vise versa.) In the middle panel of the figure we plot the three energy band (20-50, 50-100 and 100-200 keV) BATSE light curves. There are a total of 8-9 energy channels covering the 20-200 keV energy range. For each energy band, we use the detector response matrices and integrate over the relevant energy channels to produce the photon fluxes, by assuming a power law shape of a photon index of –2.0. The fluxes are then divided by the fluxes of the Crab Nebula detected with BATSE in the same energy range. By converting the fluxes to 'Crab' units, we avoid possible absolute detector calibration problems between ASM and BATSE.

Two spectral state transitions occurred near TJD 10220 and TJD 10307, characterised by the rapid increase and decrease of the ASM low-energy flux. We call the state "hard state" before TJD 10220 and after TJD 10307, since the overall spectral shape is harder than that of the Crab Nebula. Similarly we call the state between the transitions the "soft state". The HS spectrum is characterised by a hard power law between 1.3-100 keV with a photon index of around –1.8. The 20-50 keV flux is above the power law by 20-30%. Above 100 keV, a spectral cutoff is observed. In the SS, the overall 3-200 keV spectrum is dominated by a power law with a photon index of about  $-2.5$ . The first ASM energy band is contaminated significantly by a soft excess. A flux excess

between 20-50 keV above the power law, similar to the HS spectrum, makes the overall 20-200 keV spectrum fit reasonably well with a cutoff power law model. The details of the spectral evolution will be presented elsewhere (Zhang *et al.*) 1996d)

In the bottom panel of the figure we plot the total luminosity observed with ASM and BATSE between 1.3-200 keV. For the ASM bands, the luminosity is calculated in each band separately by assuming the Crab Nebula spectrum shape. A hydrogen column density  $N_H \sim 5.5 \times 10^{21}$  atoms/cm<sup>2</sup> (Ebisawa *et* al. 1996) is used for correcting the absorption in the 1.3-3.0 band. This correction is not sensitive to the exact value of  $N_{\rm H}$  we used. The luminosity would be underestimated in the 1.3-3.0 keV only by  $\sim$ 5% if the true  $N_{\rm H}$  is  $7.0\times10^{21}$  atoms/cm<sup>2</sup>. The luminosity in the BATSE energy range (20-200 keV) is calculated by assuming that the energy spectrum is a power law with an exponential cutoff (e.g., Sunyaev and Truemper 1979; Sunyaev and Titarchuk 1980) (the power law index and the cutoff energy on each day were separately determined from spectral fits to the daily averaged count rates in 8 or 9 spectral energy bands). The gap between the ASM and the BATSE energy bands is filled by interpolating the 4.8-12 keV and the 20-30 keV fluxes and assuming again a power law energy spectrum. Overall, we estimate that the deviations of the calculated luminosity should be less than 10% of the true luminosity in the 1.3-200 keV energy band. A distance of 2.5 kpc is assumed for Cyg X-1 in these calculations.

There is a general anti-correlation between the ASM fluxes and the BATSE fluxes. The fractional r.m.s. variations, calculated from the daily averaged fluxes, in the six energy bands are  $62\%$  (1.3-3.0 keV),  $40\%$  (3.0-4.8 keV),  $17\%$ (4.8-12 keV), 43% (20-50 keV), 54% (50-100 keV) and 62% (100-200 keV). As a comparison, the 1.3-200 keV luminosity r.m.s. fractional variation is about 15%. This behaviour indicates a spectral pivoting at around 10 keV.

To obtain the bolometric luminosity, we need to correct for the flux emitted beyond both the low-energy (1.3 keV) and high-energy (200 keV) limits of our observations. The high-energy correction is not important due to the rather steep spectral cutoff around 100 keV in the HS and the rather steep power law (photon index around  $-2.5$ ) in the SS. The total luminosity above 200 keV is comparable in both the HS and SS and amounts to less than 10% of the bolometric luminosity.

The low-energy cutoff correction is necessary because a significant portion of the ultra-soft component is not detected with ASM. This component can be fitted with a blackbody (BB). For the HS, values of  $kT_{\text{bb}} \sim 0.12$ -0.16 keV were obtained from one ROSAT (0.1-2.0 keV) observation on 1991 April 18-20 (Balucinska-Church *et al.* 1995) and 11 ASCA  $(0.5\n-10 \text{ keV})$  observations between 1993 October and 1994 December (Ebisawa et al. 1996). Values of  $kT_{\rm bb} \sim 0.34$  keV were obtained in the SS, obtained from one ASCA observation on 1996 May 30 during the SS (Dotani et al. 1996), and from 11 RXTE PCA and HEXTE observations throughout the whole SS (Cui *et al.* 1996b). The RXTE data indicate that the BB temperature is slightly higher in some RXTE observations (Cui et al. 1996b). We thus take the 0.34 keV as the lower limit of the BB temperaturein the SS. It is not possible to estimate the BB component in the HS from the ASM data since the power law component dominates the ASM HS detector counting rates. Previous ROSAT and ASCA observations of Cyg X-1 in the HS revealed a BB component luminosity level of about  $5\times10^{36}$ erg/s. We added this value to the HS 1.3-200 keV luminosity as shown in the bottom panel of Fig. 1. The soft excess accounts for about 50-70% of the

1.3-3.0 keV ASM counts in the SS. Taking the BB temperature  $kT \geq 0.34$  keV, we estimate that the BB luminosity below 1.3 keV is  $\leq 1.2\times10^{37}$  erg/s (the total SS BB luminosity is  $\leq 2.2\times10^{37}$  erg/s). The value of  $1.7\times10^{37}$  erg/s is added to the SS 1.3-200 keV luminosity as depicted in the bottom panel of Fig. 1. In summary, SS bolometric luminosity is between the thin and thick curves.

The bolometric luminosity displays an increase and decrease by  $\leq 50-70\%$ during the H-to-S and S-to-H state transitions. In the middle of the SS, the luminosity is however, only  $≤10-20%$  higher than the HS luminosity, because of the gradual decrease of the HXR flux after the initial SS transition and the almost symmetric recovery after reaching the minimum HXR flux between TJD 10260 and 10275, while the soft excess in the 1.3-3.0 keV band, averaged over several days, stayed at about 1.5 Crab during the entire SS. Therefore the bolometric luminosity variations throughout the state transitions are less than 50-70%.

#### 4. Discussion

Compared to the high-energy observations of the previous Cyg X-1 SS transitions, the data presented here are the first set covering the whole transition episode continuously over a broad energy band. The observed 1.3-15 keV (Ariel V) spectral evolution (spectral pivoting and power law index changes) (Chiappetti *et al.* 1981) during the 1975 S-to-H state transition is quite similar to that presented here. The initial HXR flux and spectral steepening observed with HEAO 3 between  $48-183$  keV (Ling *et al.* 1983) and the soft X-ray flux level between 1-12 keV observed later with Hakucho (Ogawara et al. 1981) are similar to that during the SS presented here. Both previous SS lasted between 60-100 days, again similar to the 1996 SS duration. So this 1996 SS transition is qualitatively similar to the previous ones. It is thus reasonable to assume that the same physical mechanism is responsible for all of them.

GX 339-4 is the only other BHC observed to have recurrent state transitions. From a comparison with its 1981 H-to-S state transition (Maejima *et al.* 1984), we find that the HS spectra of this source and Cyg X-1 are quite similar. The ratio between the SS and HS total luminosity observed from GX 339-4 (e.g., Ricketts 1983) is also similar to that of Cyg X-1 presented here. During a SS observation of GX 339-4 in 1983 (Makishima et al. 1986), the HXR flux increased significantly while the soft X-ray flux remained nearly unchanged, also similar to the HXR flux increase in the second half of the SS of Cyg X-1. The photon index of the power law changed from  $-0.9$  to  $-2.1$  following the HXR flux increase of GX 339-4, different from the near constant value of  $-2.5$  during much of the SS in Cyg X-1. The power law tail during a 'very high' state of GX 339-4 (Miyamoto *et al.* 1991), when the overall flux was about a factor of 2-3 higher than during the previous SS observations, is however, very similar to that of Cyg X-1 in the SS. Qualitatively similar S-to-H state transitions have also been observed from the low-mass X-ray binary BH system GS 1124–683 (Ebisawa *et al.* 1994) and the neutron star system 4U 1608–52 (Mitsuda *et al.* 1989). Therefore similar physical mechanisms might operate in all of them.

The 'hard' and 'soft' states we refer to in this paper are usually called the 'low' state, and the 'high' (and 'very high') states. This is due to the fact that the early observations of them were made in the low-energy X-ray band (<20 keV; see e.g., Tananbaum et al. 1972), so the terms 'low' and 'high' refer to the low and high values of the low-energy X-ray fluxes, respectively. It is generally believed that the low-energy X-ray flux tracks the total mass accretion rate of the system; therefore, these 'low' and 'high' states have been considered to

correspond to low and high values of  $\dot{M}$ , respectively. This idea gained strong support in the unified scheme of source states of both neutron star and BH X-ray binaries proposed by Van der Klis (1995).

This picture may be incomplete in view of the lack of a strong variation of the total luminosity during the state transitions which we observed in 1996. The blackbody component detected in both the hard and soft states is generally thought to originate from an optically thick and geometrically thin accretion disk near the BH (see, e.g., Mitsuda et al. 1984). According to current models the hard power law spectral component is likely produced in a very hot and optically thin region, through Comptonization of low-energy X-ray photons; the detailed nature of the optically thin region and the mechanism that makes it very hot distinguish these different models. The near constant luminosity during the transitions indicates that they are driven by a redistribution of the gravitational energy release between the optically thick and the optically thin regions. Therefore a H-to-S state transition, and its reverse, probably reflects a change in the relative importance of the energy release in the optically thin and thick regions of the accretion disk near the BH and this may not require a substantial change in the total accretion rate.

Spectra emitted by accretion disks around black holes are well described by the 'multi-temperature disk blackbody model' of Mitsuda et al. (1984), which has as fit parameters the inner disk radius,  $R_{\rm in}$ , and the temperature,  $T_{\rm in}$ , at that radius. It is usually assumed that the inner disk radius equals three times the Schwarzschild radius, and this provides acceptable mass estimates for the black hole (see Tanaka and Lewin 1995, for a review).

The blackbody fits made to the high-energy tail of this multi-temperature disk BB model model provide good fits for a blackbody temperature,  $T_{\rm bb}$ , which turns out to be equal to the temperature in the disk at a radius of about 7 Schwarzschild radii (Ross et al. 1992); correspondingly, one has  $T_{\text{bb}} \simeq 0.7T_{\text{in}}$ . Independent of the ratio of inner disk radius to the Schwarzschild radius, one has  $T_{\text{bb}} \propto T_{\text{in}}$ . Since the multi-temperature disk model the bolometric luminosity,  $(L_{\text{bol,disk}})$  follows the proportionality relation  $L_{\text{bol,disk}} \propto R_{\text{in}}^2 T_{\text{in}}^4$ , one has  $(L_h/L_s) = (R_h/R_s)^2 (T_h/T_s)^4$ , by applying the proportionality relation in both the HS and SS, where the subscripts 'h' and 's' denote the hard and soft states respectively.

From the observations we infer that  $L_s/L_h \leq 6$ , and  $T_s/T_h \geq 2.1$  to 2.8. Consequently, we obtain  $R_h/R_s \ge 1.8$  to 3.2. This result suggests that *during the* H-to-S transition the inner radius of the geometrically thin and optically thick disk changed from  $\gtrsim$  170 km to ~ 70 km (the latter value has been obtained by Dotani et al. 1997, from ASCA observations during the soft state). Applying the relationship between the mass accretion rate, inner disk radius and the inner disk temperature in the multi-temperature disk model we have  $\dot{M}_{\rm s}/\dot{M}_{\rm h} \leqslant 2.0$  to 3.4.

This change in inner disk radius accompanying the spectral state changes suggests a picture in which during the hard state advection of internal disk energy into the black hole, as proposed by Narayan (1996), dominates within a radial distance of  $\gtrsim$  170 km from the hole. In the soft state the flow in the inner disk may still be advection dominated, with only a moderate fraction of the mass flow passing through an optically thick inner disk.

Variations of the inner disk radius have also been discussed by Ebisawa et al. (1996), in the context of the mass accretion and high-energy radiation model of Chakrabarti and Titarchuk (1995), and have recently been proposed by Belloni et al. (1996) as an explanation for the rapid variability of the black-hole candidate GRS 1915+105 (albeit at much higher mass accretion rates).

A potential problem for any model that purports to explain the spectral transitions in terms of an instability in the inner disk region is that the transitions seem to be accompanied by the gradual changes in the slope and the flux of the HXR power law component, which occur over a very long time scale (∼ weeks). In fact, the 1996 S-to-H transition was predicted from the HXR flux increase (Zhang et al. 1996c). It is possible that Cyg X-1 underwent a very slow change in the mass accretion rate, which initially only affected the properties of the hard X-ray emission region. The sudden soft  $(1.3-3.0 \text{ keV})$ X-ray flux increase and decrease over a much shorter time scale, during the state transitions, may then represent the crossing of a threshold, at which the radiative efficiency exceeds a value required for the formation of an optically thick disk down to the innermost stable orbit.

In view of the observational limitations the above remarks are necessarily somewhat speculative. It would appear that a better understanding of the cause of the spectral state changes in Cyg X-1 requires that further monitoring of the long-term behaviour of Cyg X-1 include low-energy ( $\leq 1$  keV) coverage of the source as well.

We thank the ASM/RXTE and BATSE/CGRO teams for providing excellent technical supports. SNZ thanks W. Chen, K. Ebisawa, L. Titarchuk, S. Chakrabarti, R. Narayan and C. Robinson for many stimulating discussions. JvP acknowledges support from NASA through grant NAG5-3003. We also appreciate very much the constructive comments and suggestions from the anonymous referee.

## REFERENCES

- Balucinska-Church, M., Belloni, T., Church, M.J., Hasinger, G. 1995, A&A  $302(1)$ , L5-8
- Belloni, T. et al. 1996, ApJL 472, L107-110
- Bolton, C.T. 1972, Nature 235, 271
- Chakrabarti, S.K. & Titarchuk, L. 1995 ApJ 455(2), 623-39
- Chiappetti, L. et al. 1981, MNRAS 197, 139
- Crary, D.J. et al. 1996, ApJ 462, L71
- Cui, W. 1996, IAU Circ. 6404
- Cui, W., Focke, W., & Swank, J. 1996, IAU Circ. 6439
- Cui, W., et al. 1997, ApJL 474(1), 57
- Dotani, T., Negoro, H., Mitsuda, K., Inoue, H., and Nagase, F. 1996, IAU Circ. 6415
- Dotani, T., *et al.* 1997, submitted to Nature
- Ebisawa, K., Ueda, Y., Inoue, H., Tanaka, Y., & White, N. E. 1996, ApJ, 459
- Ebisawa, K., et al. 1994, PASJ 46, 375-395
- Fishman, G.J. et al. 1989, in Gamma Ray Observatory Science Workshop (eds Shrader, C.R., Gehrels, N. & Dennis, B.) 2-39–2-50 (CP-3137, NASA, Greenbelt, Maryland, USA, 1989)
- Gies, D.R. & Bolton, C.T. 1986, ApJ 304, 371
- Harmon, B.A. et al. 1992, AIP Conf. Proc. 280 314
- Levine, A., et al. 1996, ApJ 469, L33-36
- Liang, E. P., & Nolan, P. L. 1984, Space Sci. Rev., 38, 353
- Ling, J. C., Mahoney, W. A., Wheaton, W. A., & Jacobson, A. S. 1983, ApJ, 275, 307
- Maejima, Y. et al. 1984, ApJ 285, 712
- Makishima, K. et al. 1986, ApJ 308, 635-643
- Miyamoto, S., & Kitamoto, S. 1989, Nature 342, 773-4
- Miyamoto, S., et al. 1991, ApJ 383, 784-807
- Mitsuda, K. et al. 1984, PASJ 36, 741
- Mitsuda, K. et al. 1989, PASJ 41, 97-111
- Narayan, R. 1996, ApJ 462(1), 136-41
- Ogawara, Y., et al. 1982, Nature 295, 675
- Paciesas, W.S. et al. 1996, 31st COSPAR Scientific Assembly, Birmingham, UK, July 1996
- Phlips, B. et al. 1997, in prepsration
- Piro, L. et al. 1996, IAU Circ. 6431
- Ricketts, M. 1983, A&A 118, L3
- Ross, R.R., Fabian, A.C. & Mineshige, S., 1992, MNRAS 258, 189
- Sunyaev, R. & Truemper, J. 1979, Nature279, 506
- Sunyaev, R. & Titarchuk, L. 1980, A&A 86, 121
- Tanaka, Y., & Lewin, W. H. G. 1995, in "X-ray Binaries", eds. W. H. G. Lewin,
- J. van Paradijs, & E. P. J. van den Heuvel (Cambridge U. Press, Cambridge) p.
- 126

van der Klis, M. 1995, in "X-ray Binaries", eds. W. H. G. Lewin, J. van

Paradijs, & E. P. J. van den Heuvel (Cambridge U. Press, Cambridge) p. 126

Webster, B.L. & Murdin, P. 1972, Nature 235, 37

Zhang, S.N., Fishman, G.J., Harmon, B.A. & Paciesas, W.S. 1993, Nature 366, 245-247

Zhang, S. N., Harmon, B. A., Paciesas, W. S., & Fishman, G. J., 1996a, IAU Circ. 6405

Zhang, S. N., Harmon, B. A., Paciesas, W. S., 1996b, IAU Circ. 6447

Zhang, S. N., et al. 1996c, IAU Circ. 6462

Zhang, S. N., et al. 1996d, to be submitted to ApJ

This manuscript was prepared with the AAS IATEX macros  $\rm v4.0.$ 

Fig. 1.— Cyg X-1 light curves and the  $1.3\text{-}200$  keV luminosity variations during its 1996 SS transition.



Fig. 1.— Cyg X-1 light curves and luminosity variations during its 1996 SS transition.