

ASCA AND ROSAT X-RAY SPECTRA OF IC 5063 AND NGC 3998

C. VIGNALI¹, A. COMASTRI², M. CAPPI³, G.G.C. PALUMBO^{1,3}

¹*Dipartimento di Astronomia, Università di Bologna, via Zamboni 33, I-40126 Bologna, Italy*

²*Osservatorio Astronomico, via Zamboni 33, I-40126 Bologna, Italy*

³*ITeSRE/CNR, via Gobetti 101, I-40129 Bologna, Italy*

ABSTRACT. The results of the 0.1-10 keV spectral analysis of IC 5063 (Narrow Line Radio Galaxy) and NGC 3998 (LINER) are presented. In both cases there is evidence of a FeK α emission line, but its width is poorly constrained. The highly absorbed spectrum of IC 5063 is well described by a double power law model. The X-ray luminosity of a few $\times 10^{43}$ ergs s⁻¹ reveals a Sey 1 nucleus seen nearly edge-on, as confirmed by optical and IR studies. NGC 3998 is well fitted by a power law model ($\Gamma \sim 1.9$) without reflection and weak absorption. The weak X-ray emission detected by ASCA and ROSAT suggests a low-luminosity AGN or an active nucleus in a low-accretion period.

1. IC 5063

IC 5063 is classified as a Narrow Line Radio Galaxy ($z=0.011$) (Caldweel & Phillips 1981). IC 5063 was observed with the gas imaging spectrometer (GIS) and solid state spectrometer (SIS) on the ASCA satellite (Tanaka et al. 1994) in April 1994 and with the ROSAT PSPC (Pfeffermann et al. 1987) from November 1991 to April 1992. After applying standard criteria to the data, about 19 Ks of useful data for ROSAT and $\sim 21(24)$ Ks for each SIS(GIS) detector were collected.

The combined analysis of ASCA and ROSAT data reveals a soft X-ray excess, a hard continuum and a reflection component. A double power law model plus a reflection component (fixed to cover 2π steradians at the source) provides a good fit to the ROSAT and ASCA data, but the spectral slope of the soft power law ($\Gamma_S=2.16_{-0.31}^{+0.29}$, errors at 90%, $\Delta\chi^2=2.71$), steeper than the hard ($\Gamma_H=1.70_{-0.19}^{+0.18}$), indicates a more complex soft X-ray emission. A $L_{\text{soft}}/L_{\text{hard}} \sim 1\%$ ratio is obtained by a partial covering model. A bremsstrahlung model for the soft emission is equally good (see table 1). The broad H α line in polarized flux (FWZI ~ 6000 -9000 Km s⁻¹, Axon et al. 1994) and the anisotropy of the radiation field (“X” geometry), with an opening angle of $\sim 50^\circ$ (Colina et al. 1991), suggest that the observed soft X-ray component may be emission scattered along the line of sight from ionized gas above an absorbing torus. The column density value, $N_H=2.15\pm 0.18 \times 10^{23}$ cm⁻², and the dust lane across the galaxy, likely responsible for the IR emission (Axon et al. 1982), agree with a nearly edge-on configuration. The 2-10 keV luminosity $L \sim 2 \times 10^{43}$ ergs s⁻¹ indicates a Sey1-like active nucleus for IC 5063. Strong evidence for the FeK α line is given only by the SIS1 detector. The line param-

ters are given in table 2.

Tab. 1 - Spectral parameters

source	model	Γ_S/kT	Γ_H	N_H	χ^2/dof
IC5063	po+po	$2.16^{+0.29}_{-0.31}$	$1.70^{+0.18}_{-0.19}$	$2.15 \pm 0.18 \times 10^{23}$	260/263
.....	brem+po	$1.39^{+1.02}_{-0.42}$	$1.66^{+0.20}_{-0.18}$	$2.07^{+0.20}_{-0.19} \times 10^{23}$	262/263
NGC3998	po	1.88 ± 0.02	$\sim N_{H_{gal}}$	1036/688

2. NGC 3998

NGC 3998 is a LINER galaxy ($z=0.0035$) with strong UV emission (Reichert et al. 1992), but weak X-ray luminosity ($L_X \sim 3.5 \times 10^{41}$ ergs s^{-1} in the 2-10 keV energy range). NGC 3998 was observed twice by ROSAT (May 91-May 92), for a total of ~ 65 Ks of observation, and once by ASCA (May 94), for which only GIS data were available (40 Ks/GIS). The joint ASCA and ROSAT analysis gives a good fit with a power law model ($\Gamma = 1.88 \pm 0.02$) and a weak excess of absorption above the galactic value. An upper limit of 0.88 is obtained for the reflection component. There is also evidence of an iron line, whose parameters are given in table 2. No flux or spectral variability have been detected by ASCA and ROSAT. The low X-ray flux may be due to the fact that NGC 3998 is a low-luminosity AGN, but the high UV state suggests the presence of a massive black hole ($M \sim 10^8 M_\odot$, Fabbiano et al. 1994), therefore in a steady-quiet state, or a more elusive advection-dominated AGN (Fabian & Rees 1995).

Tab. 2 - FeK α parameters

source	$E_{K\alpha}$	$\sigma_{K\alpha}$	EW
IC 5063	$6.35^{+0.09}_{-0.11}$	<0.31	168^{+138}_{-105}
NGC 3998	$6.44^{+0.16}_{-0.30}$	<0.41	200^{+161}_{-139}

References

- Axon, D.J., et. al.: 1994, *Astrophys. Space Sci.* **216**, 379
Axon, D.J., Bailey, J., Hough, J.H.: 1982, *Nature* **299**, 234
Caldwell, H., Phillips, M.M.: 1981, *Astrophys. J.* **244**, 447
Colina, L., Sparks, W.B., Macchetto, F.: 1991, *Astrophys. J.* **370**, 102
Fabbiano, G., Fasnacht, C., Trinchieri, G.: 1994, *Astrophys. J.* **434**, 67
Fabian, A.C., Rees, M.J.: 1995, *Mon. Not. R. Astr. Soc.* **277**, L55
Pfeffermann, E., et al.: 1987, *Proc. SPIE* **733**, 519
Reichert, G.A., et al.: 1992, *Astrophys. J.* **387**, 536
Tanaka, Y., Inoue, H., Holt, S.S.: 1994, *Publ. Astron. Soc. Japan* **46**, L37