EVOLUTION OF X-RAY CLUSTERS OF GALAXIES AND SHOCK HEATING OF INTRACLUSTER MEDIUM

MOTOKAZU TAKIZAWA AND SHIN MINESHIGE

Department of Astronomy, Faculty of Science, Kyoto University, Sakyo-ku, Kyoto 606-01, JAPAN

_:

Received _____

accepted _____

ABSTRACT

Evolutions of X-ray clusters of galaxies are studied by N-body (shell model) + mesh code (TVD) simulations on the assumption of spherical symmetry. We consider a density perturbation of $10^{15} M_{\odot}$ composed of dark matter and gas in cold dark matter dominated universe with the cosmological density parameter, $\Omega_0 = 1$ or 0.2. A shock front appears during its initial collapse, moving outward as ambient gas accretes towards cluster center. The shock front separates the inner X-ray emitting, hot region, where gas is almost in hydrostatic equilibrium but with small radial infall ($\sim 100 \text{km s}^{-1}$) being left, from the outer cool region, where gas falls almost freely and emits no X-rays. Gas inside the shock is strongly compressed and heated by shock so that X-ray luminosity rapidly rises in the early stage (until temperature reaches about virial). In the late stage, however, the X-ray luminosity rises only gradually due partly to the expansion of the inner high temperature region and partly to the increase of X-ray emissivity of gas as the results of continuous adiabatic compression inside the shock. We also find for clusters in lower density universe that the density distribution is generally less concentrated and, hence, X-ray luminosity more slowly rises than in higher density universe.

The shock front structure, which was not clearly resolved in the previous SPH simulations, is clearly captured by the present simulations. Our results confirm that shock heating plays an important role in the heating process of intracluster medium. In addition, we find a sound wave propagating outward, thereby producing spatial modulations with amplitudes of ~ 10 % in the radial temperature and density profiles and time variations in the strength of the shock. Such modulations, if observed, could be used as a probe to investigate

the structure of clusters.

Subject headings: galaxies: clustering — galaxies: intergalactic medium — galaxies: X-ray — hydrodynamics

1. INTRODUCTION

Clusters of galaxies (CGs) are luminous X-ray sources (Sarazin 1988). X-rays are thought to be emitted through thermal bremsstrahlung from very hot, optically thin plasma gas in intergalactic space, which is called intracluster medium (ICM). Recent X-ray and optical observations have revealed dynamical aspects of CGs (Fabricant, Kent & Kurtz 1989; Kneib et al. 1995). Time evolution of the cluster X-ray luminosity function has been confirmed; each CG became brighter and the number density of CGs with a fixed X-ray luminosity increased through mergers towards zero redshift (Edge et al. 1990; Gioia et al.1990; Henry et al. 1992; Bower et al. 1993; Castander et al. 1994). To understand the physical structure of CGs, therefore, it is essential to consider non-equilibrium processes involved with the dynamical evolution of CGs.

Recent extensive X-ray observations have established that the temperature profiles of ICM are nearly flat in many clusters (Fabian 1994; Ohashi et al 1996). Although there is no widely accepted explanation for this, it is, at least, reasonable to expect that shock heating through gravitational collapse plays an important role. It then follows that the isothermality of gas may reflect the shape and time evolution of the gravitational potential well. What then governs the total dynamics of CGs? It is galaxies and dark matter (DM), both of which can be regarded as a collisionless self-gravitating many-body system. Dynamics of ICM, collisional fluid on a CG scale, simply follows the dynamical evolution of the gravitational potential exerted mainly by DM and galaxies. Thus, the thermal history of ICM is rather sensitive to the physical processes involving violent time variation of gravitational potential field (such as formation, merger, etc), Violent relaxation (Lynden-Bell 1967) is thought to play a crucial role there, although its physical significance is still poorly understood (Funato, Makino & Ebisuzaki 1992a,b; Takizawa & Inagaki 1996). CGs were believed to be formed from overdense perturbations in the universe. They grew through gravitational instability and collapsed at $z \sim 1$. CGs have been growing by accreting ambient matter, which is still an ongoing process at the present. Gunn & Gott (1972) quantitatively discussed the growth of a spherical symmetric perturbation consisting of only collisionless particle in an expanding universe. Bertschinger (1985) found the self-similar solution describing evolution of spherical density fluctuation consisting of DM and gas in the Einstein-de Sitter universe (where the cosmological density parameter is $\Omega_0 = 1$ and the cosmological constant is $\lambda_0 = 0$).

There have been plenty of numerical studies performed so far regarding the formation and evolution of CGs by using N-body and hydrodynamic codes. Perrenod (1978) was the first to calculate the evolution of a spherical symmetric cluster with standard mesh hydrodynamic code. Three-dimensional calculations have been carried out recently, which mostly use smooth particle hydrodynamics (SPH) codes (Evrard 1990; Thomas & Couchman 1992; Bryan et al. 1994; Metzler & Evrard 1994; Navarro, Frenk & White 1995, hereafter NFW). According to Evrard (1990), a shock front moved outward when gas around the cluster accreted onto the cluster center and a relatively flat temperature profile was realized within the shock front. This result was significant in the sense that the above expectation was confirmed qualitatively. However, quantitative estimation may be problematic because of limited spatial resolutions and poor reliability of SPH code in calculations of shocks. It might be kept in mind that SPH codes are easy to work with and could give reasonable accuracy, however, they are not better for problems with discontinuities (such as strong shock and contact discontinuity) than mesh codes (Monaghan 1992). On a larger scale, numerical simulations of cosmological structure formation by using N-body and hydrodynamic mesh code have been carried out mainly to investigate the statistical properties of CGs (Cen & Ostriker 1994; Kang et al. 1994; Anninos & Norman

1996).

In this paper, we focus on the dynamical aspects of ICM, such as time-dependent properties of shock waves and the effects of shock heating on the evolution of CGs. For this purpose, we perform numerical simulations of spherical CGs with N-body (shell model) + mesh code (TVD). Note that the TVD code is one of the most useful tools to deal with spatial discontinuities. Since we assume spherical symmetry, the problem can be reduced to one-dimensional. Better spatial resolutions can be achieved, therefore. Using these codes, we calculate dynamical evolutions of a density perturbation of $10^{15} M_{\odot}$ consisting of DM and gas and collapsing at $z \sim 1$ in universe with $\Omega_0 = 1$ or 0.2. It is also interesting to investigate how different cosmological models affect evolution of CGs.

The rest of this paper is organized as follows. In section 2 we describe our numerical methods and the adopted initial conditions. In section 3 we present our results, discussing physical processes underlying the structural evolution of calculated CGs. In section 4 we summarize our results and discuss their implications.

2. THE SIMULATIONS

In the present study we regard CG consisting of two components: collisionless particles corresponding to galaxies and DM, and gas corresponding to ICM. When calculating gravity both components are considered, although the former component dominates over the latter. We also assume spherical symmetry in all the calculations.

2.1. Basic Equations for Collisionless Particles

For calculations of collisionless particles we adopt a shell model (Hénon 1964). The distribution function of spherical symmetric stellar systems can be expressed as f(r, u, v, t), where r is radial distance, u is radial velocity, v is tangential velocity, and t is time, respectively. So the state of the system is represented by N points in the (r, u, v) space (with N being the number of shells) and a trajectory of the *i*-th point at (r_i, u_i, v_i) is calculated according to the equations of motion;

$$\frac{dr_i}{dt} = u_i, \tag{1}$$

$$\frac{du_i}{dt} = \frac{A_i^2}{r_i^3} - \frac{GM_i}{r_i^2},\tag{2}$$

for $i = 1 \sim N$, where G is the gravitational constant, $A_i \equiv r_i v_i$ (= constant in time) is angular momentum of the *i*-th shell, and M_i is total mass (also including mass of gas) interior to r_i , respectively. Since it is convenient to carry out numerical calculations using comoving coordinates for our purpose, we transform (r_i, u_i, v_i) to (R_i, U_i, v_i) as follows;

$$r_i = a(t)R_i, (3)$$

$$u_i = \dot{a}R_i + U_i, \tag{4}$$

where a(t) is the dimensionless scale factor of the universe and \dot{a} represents the derivative of a(t) with respect to time. Equations (1) and (2) are then transformed into

$$\frac{dR_i}{dt} = \frac{U_i}{a},\tag{5}$$

$$\frac{dU_i}{dt} = \frac{A_i^2}{(aR_i)^3} - \frac{GM_i}{(aR_i)^2} - \frac{\dot{a}}{a}U_i - \ddot{a}R_i,$$
(6)

respectively, by using equations (3) and (4). Equations (5) and (6) are integrated by using leap-frog method. As to the inner boundary condition we set a reflecting wall at $r_{\rm rw} = 0.02/(1+z)$ and impose that when a shell reaches the wall, $r_i < r_{\rm rw}$, that shell is elastically reflected. Only shells with rather small angular momentum is influenced by the wall.

2.2. Basic Equations for Gas

For gas components basic equations in the comoving frame are as follows,

$$\frac{\partial \rho}{\partial t} + \frac{1}{a} \frac{\partial}{\partial R} (\rho v_{\text{gas}}) = -3 \frac{\dot{a}}{a} \rho - \frac{2}{aR} \rho v_{\text{gas}}, \tag{7}$$

$$\frac{\partial}{\partial t}(\rho v_{\rm gas}) + \frac{1}{a}\frac{\partial}{\partial R}(\rho v_{\rm gas}^2 + P) = -4\frac{\dot{a}}{a}\rho v_{\rm gas} - 2\frac{\rho v_{\rm gas}^2}{aR} + \rho g_R, \tag{8}$$

$$\frac{\partial}{\partial t}(\rho E) + \frac{1}{a}\frac{\partial}{\partial R}(\rho H v_{\text{gas}}) = -\frac{\dot{a}}{a}\left(\frac{5}{2}\rho v_{\text{gas}}^2 - \frac{3\gamma}{\gamma - 1}P\right) - 2\frac{\rho H v_{\text{gas}}}{aR} + \rho g_R v_{\text{gas}}, \quad (9)$$

where ρ is gas density, v_{gas} is radial velocity of gas, P is gas pressure, and γ is the adiabatic exponent, respectively. The total energy of gas per unit mass, E, and the enthalpy per unit mass, H, are given by

$$E = \frac{P}{\rho(\gamma - 1)} + \frac{v_{\text{gas}}^2}{2},$$
 (10)

$$H = \frac{\gamma}{\gamma - 1} \frac{P}{\rho} + \frac{v_{\text{gas}}^2}{2}, \qquad (11)$$

respectively, and g_R is defined by

$$g_R \equiv -\frac{GM_R}{(aR)^2} - \ddot{a}R,\tag{12}$$

where, M_R is the total mass (including gas and collisionless particles) inside R.

We neglect viscosity and angular momentum of gas. We also assume that gas is ideal gas with $\gamma = 5/3$. A second order up-wind TVD code (minmod limiter) is used for our simulations (Hirsch 1990). Number of mesh point is 500 and one mesh spacing corresponds to $\Delta r = 0.02/(1+z)$ Mpc with z being a redshift [1 + z = a(0)/a(t)]. We set boundary condition as follows. The inner edge is assumed to be a perfectly reflecting point;

$$\rho_{-1} = \rho_0 = \rho_1, \tag{13}$$

$$P_{-1} = P_0 = P_1, (14)$$

$$v_{\rm gas,-1} = v_{\rm gas,0} = 0,\tag{15}$$

where ρ_i , P_i and $v_{\text{gas},i}$ is gas density, gas pressure and radial velocity of gas at the *i*-th mesh point, and the first mesh point corresponds to the inner boundary. The 0-th and -1-st points are necessary for calculations (to derive spatial derivatives of physical quantities). The outer edge is assumed to be a perfectly transmitting surface;

$$q_{n+2} = 3q_n - 2q_{n-1}, (16)$$

$$q_{n+1} = 2q_n - q_{n-1}, (17)$$

where q_i is any physical quantity of gas at the *i*-th mesh point and the *n*-th corresponds to the outer boundary. Again, the (n+1)-th and (n+2)-th points are necessary for calculation purpose.

2.3. Models and Initial Conditions

In this paper, calculations are carried out from $z_{ini} = 10$ to the present time $(z_0 = 0)$. We adopt cosmological models with no cosmological constant, $\lambda_0 = 0$. When $\Omega_0 = 1$, therefore, we find

$$a(t) = \left(\frac{t}{\frac{2}{3}H_0^{-1}}\right)^{\frac{2}{3}},\tag{18}$$

while when $0 < \Omega_0 < 1$ we have

$$a(t) = \frac{\Omega_0}{2(1-\Omega_0)} (\cosh \psi - 1),$$
 (19)

$$H_0 t = \frac{\Omega_0}{2(1 - \Omega_0)^{3/2}} (\sinh \psi - \psi), \qquad (20)$$

where H_0 is the Hubble constant and is set to be $H_0 = 100 \text{ (km s}^{-1} \text{ Mpc}^{-1}\text{)}$ in transformations of length and time.

We make initial density profiles in the same manner as Peebles (1982). At first we prepare N concentric shells with uniform density being equal to the mean density of the universe at $z_{ini} = 10$. Then a density fluctuation is introduced by perturbing the radius and velocity of each shell as

$$r_i = r_i^{(0)} \left[1 + \delta(r_i^{(0)}) \right], \qquad (21)$$

where $r_i^{(0)}$ is the unperturbed coordinate and $\delta(r)$ represents the perturbation as a function of r (specified below). The velocity perturbation is written, using Zel'dovich approximation, by

$$u_i = H(z_{\rm ini})r_i^{(0)} \left[1 + 2\delta(r_i^{(0)})\right], \qquad (22)$$

where $H(z_{\rm ini})$ is the Hubble constant at the initial epoch, and we used $\delta \propto t^{2/3}$; the relation which holds exactly in the Einstein-de Sitter universe. The functional form of the perturbation is assumed as

$$\delta(r) = \begin{cases} -\frac{\delta_0}{3} \cos^2(\frac{\pi}{2} \frac{r}{r_0}) & (0 \le r \le r_0) \\ 0 & (r_0 < r) \end{cases},$$
(23)

where r_0 and δ_0 are, respectively, the initial size of the fluctuation on the comoving scale and the parameter for displacement (see Nakamura 1996). We take $(\delta_0, r_0) = (0.4, 9.0$ Mpc) for Model A (with $\Omega_0 = 1.0$) and $(\delta_0, r_0) = (0.5, 15.4 \text{ Mpc})$ for Model B (with $\Omega_0 = 0.2$), respectively. In both models the fluctuation contains a mass of about $10^{15} M_{\odot}$ and correspond to typical density peaks with ~ 1.5σ in the CDM power spectrum with $\sigma_8 = 0.96$. Here, σ is the rms density fluctuations on a cluster scale and σ_8 is the rms of mass fluctuation on scale 8 Mpc; $\sigma_8 = 0.96$ was obtained from the observation of nearby galaxy distribution (see Suto 1993). The profile of the initial density perturbation and the ratio of $(\delta M/M)/\sigma_{8,\text{CDM}}$ are illustrated in figure 1 for Model A (by the solid line) and B (by the dotted line), respectively. Here, $(\delta M/M)$ is the mass fluctuation averaged over a scale r, whereas $\sigma_{8,\text{CDM}}$ is the rms of mass fluctuation averaged over a scale r obtained from CDM power spectrum with $\sigma_8 = 0.96$.

Fig. 1.—

The initial conditions of gas are set as follows. At first density of gas was everywhere set to be 1/10 of the mean density of the universe at z = 10 and temperature of gas $(T_{\text{gas},i})$ was everywhere taken to be 10⁷ K except in Model LT, where we assumed $T_{\text{gas},i} = 10^6$ K initially. Then adiabatic fluctuation (cf. Eq. 16 and 17) was imposed so that gas density became always 1/10 of that of DM. Note that gas temperature distribution becomes nonuniform, accordingly. We calculated 3 models in total. Model parameters are listed in table 1.

3. RESULTS

3.1. Overview of Evolution

At first we overview the evolution of Model A cluster. The evolution of density and radial velocity profiles of DM are displayed in figure 2. The different types of lines correspond to different redshifts (and different times): z = 2.5 (t = 1 Gyr) by the long dash line; z = 1.2 (t = 2 Gyr) by the short dash line; z = 0.4 (t = 4 Gyr) by the dotted line; and z = 0 (t = 6.5 Gyr) by the solid line, respectively. Density profile basically obeys a power law, $\rho_{\rm DM} \propto r^{-2.6}$, and evolves in a self-similar fashion after $z \simeq 1$. The evolutions in the radial distributions of radial velocity, density, temperature, pressure, and entropy (each of the gas component) are summarized in figure 3. The different types of lines correspond to same redshifts as in figure 2.

Fig. 3.—

There are fundamental features commonly seen in the course of evolution of all the models, which can be summarized as follows.

- 1. Before the initiation of DM collapse (at $z \ge 2.5$; $t \le 1$ Gyr) gas continues to expand, following the cosmological expansion. When DM begins to collapse, a shock wave emerges in the central part and moves outwards, accreting ambient gas towards the center.
- 2. The shock front separates the inner hot region from the outer cool region. In the inner region, gas is almost in hydrostatic equilibrium, although bulk velocity of radial infall∼ 100 km s⁻¹ still remains. The temperature profile is relatively flat and gas is hot enough to emit X-ray. In the outer region, in contrast, gas falls almost freely and is too cool to emit X-ray.
- 3. Density profile of gas evolves self-similarly as $n_{\text{gas}} \propto r^{-2.4}$ inside the shock front except near the center, where density profile is rather flat. Even after the passage of the shock, density, temperature, and, therefore, pressure, gradually increase with time. This indicates that the inner region is not perfectly in hydrostatic equilibrium (will be discussed in §3.3).
- 4. Entropy profile shows an overall increase outwards, suggesting larger entropy production taking place as the shock moves outward. There are wavy features seen,

especially, in the distributions of temperature and entropy. These seem to be related to sound wave propagation (will be discussed in $\S3.4$).

Let us next examine each item in more details and discuss similarities and differences between different models.

3.2. Density Profiles

Density profiles of DM at z = 0 can well be fitted with the β -model;

$$\rho_{\rm DM}(r) = \rho_{\rm DM,0} \left[1 + \left(\frac{r}{r_{\rm DM}}\right)^2 \right]^{-3\beta_{\rm DM}/2}.$$
(24)

Here, $r_{\rm DM}$ and $\beta_{\rm DM}$ are fitting parameters. We fit the results of simulations inside r_{100} , the radius where the mean interior density is 100 times of critical density at z = 0. The results of the fitting are summarized in table 2. Note that these results may depend on initial density profile.

In the same way density profiles of gas at z = 0 inside the shock fronts can be fitted with the β -model,

$$n_{\rm gas}(r) = n_{\rm gas,0} \left[1 + \left(\frac{r}{r_{\rm gas}}\right)^2 \right]^{-3\beta_{\rm gas}/2},\tag{25}$$

where $r_{\rm gas}$ and $\beta_{\rm gas}$ are fitting parameters. The results are listed in table 3. In all models we find $\beta_{\rm DM} \approx \beta_{\rm gas} \sim 0.9$ and $r_{\rm DM} \approx r_{\rm gas}$.

We also depict nondimensional density profiles at z = 0 in figure 4 for Model A (by the solid line) and Model B (by the dotted line), respectively, where density is scaled with ρ_{c0} , critical density of the universe at z = 0, and radius is scaled with r_{100} . Nondimensional DM density profiles look similar among two models, whereas gas component expands slightly in Model B, compared with that in Model A.

Fig. 4.—

Let us finally compare our results with the previous results (NFW). As for the density profiles of the clusters in Einstein-de Sitter Universe (Models A and LT), the core radii are smaller than those of NFW (who obtained $r_{\rm DM} \sim r_{\rm gas} \sim 0.2$), while $\beta_{\rm DM}$ and $\beta_{\rm gas}$ are similar. On the other hand, central DM density, $\rho_{\rm DM,0}$, is similar to that of NFW, whereas central gas density, $\rho_{\rm gas,0}$, of model A is lower than that of NFW (our model LT is consistent with their model; $n_{\rm gas,0} \sim 5 \times 10^{-2}$). This can be explained in terms of different initial gas temperatures (see section 3.7).

3.3. Temperature Profiles

Temperature and entropy profiles are shown in figure 5 in the upper and lower panels, respectively. The different types of lines correspond to different models: Model A by the solid line; Model B by the dotted line; and Model LT by the short dash line, respectively. There are common characters as follows. Temperatures gradually fall outwards until the shock front. There are small temperature fluctuations seen. If we evaluate the propagation speed of fluctuation pattern, it is ~ 700 km/s, of the order of the sound velocity inside the shock. In addition, entropy pattern does not change much during the course of wave propagation, indicating that structural variation is adiabatic. We may thus conclude that the temperature fluctuation arises due to sound wave propagation.

Inside the shock fronts temperature is nearly virial, high enough to emit X-ray. Entropy monotonically rises from the center to the shock front. That is, near the center gas is heated up mainly through adiabatic compression, thus possessing relatively lower entropy. Gas in the outer region, on the other hand, will eventually be heated through shocks, which effectively transform kinetic energy of accreting gas into thermal energy at their surface. This explains relatively large entropy just inside the shock. Note that regions with entropy decreasing outward are convectively unstable. Convective motions, if occur, will smear out such a feature.

Fig. 5.—

To understand why temperature, density and entropy steadily increase with time even after the passage of the shock front and how sound waves arise and propagate outward, we check the balance between pressure gradient and gravity in the inner parts. Figure 6 shows the ratio of the absolute value of pressure gradient force, |dP/dr|, and gravitational force inside the shock surface, $F_{\rm g} \equiv GM_r \rho_{\rm gas}/r^2$, from t = 5.5 Gyr to 6.5 Gyr. If the system is perfectly in hydrostatic equilibrium, $|dP/dr|/F_{\rm g}$ should be unity.

This figure shows a clear tendency that gravitational force overcomes pressure gradient force inside the shock front, thus inducing radial gas inflow from outside. Therefore, gas is adiabatically compressed by the infalling material from outside. Importantly, the ratio changes with time. Since DM density profile hardly changes with time near the core, so does the gravity force; the ratio changes are purely due to time and spatial variations in pressure profiles. When ambient gas suddenly falls towards the center, pressure at the core abruptly increases. This gives rise to an outwardly propagating sound wave, since the central point is a reflecting boundary due to spherial symmetry (cf. §2.2). Figure 7 shows radial velocity profiles of gas inside the shock from t = 5.5 Gyr to 6.5 Gyr. We confirm that radial infall systematically remains even after the passage of shock. In addition we confirm fluctuation pattern propagating outwards with a speed roughly equal to the sound velocity.

Fig. 6.—

Fig. 7.—

In our simulations thermal conduction is neglected. It is possible, however, that thermal conduction, if efficient, will erase temperature fluctuations on a small scale. It is thus worthwhile evaluating conduction timescale in the simulated clusters. Conduction time scale is generally expressed as (cf. Sarazin 1988)

$$t_{\rm cond} \approx \frac{n_{\rm e} l_{\rm T}^2 k_{\rm B}}{\kappa},$$
 (26)

where $n_{\rm e}$ is electron number density, $l_{\rm T}$ is scale length of temperature gradient, $k_{\rm B}$ is the Boltzmann constant, and thermal conductivity for hydrogen plasma is (Spitzer 1962)

$$\kappa \approx 4.6 \times 10^{13} \left(\frac{T_{\rm e}}{10^8 {\rm K}}\right)^{5/2} \left(\frac{\ln \Lambda}{40}\right)^{-1} ({\rm erg s}^{-1} {\rm cm}^{-1} {\rm K}^{-1}),$$
(27)

where $\ln \Lambda$, Coulomb logarithm, is

$$\ln \Lambda = 37.8 + \ln \left[\left(\frac{T_{\rm e}}{10^8 \rm K} \right) \left(\frac{n_{\rm e}}{10^{-3} \rm cm^{-3}} \right)^{-1/2} \right].$$
(28)

From equations (26) and (27), we derive

$$t_{\rm cond} \approx 3 \times 10^7 {\rm yr} \left(\frac{n_{\rm e}}{10^{-3} {\rm cm}^{-3}}\right) \left(\frac{T_{\rm e}}{10^8 {\rm K}}\right)^{-5/2} \left(\frac{l_{\rm T}}{0.1 {\rm Mpc}}\right)^2 \left(\frac{\ln \Lambda}{40}\right).$$
 (29)

For Model A, for example, we find $t_{\rm cond} \sim 1.5 \times 10^8$ yr at $r \approx 0.1$ Mpc (where $n_{\rm e} \sim 10^{-3} {\rm cm}^{-3}$, $T_{\rm e} \sim 5 \times 10^7$ K, and $l_{\rm T} \sim 0.1$ Mpc) and $t_{\rm cond} \sim 1.5 \times 10^7$ yr at $r \approx 1$ Mpc (where $n_{\rm e} \sim 10^{-5} {\rm cm}^{-3}$, $T_{\rm e} \sim 2 \times 10^7$ K, and $l_{\rm T} \sim 0.1$ Mpc). At both radii, hence, conduction seems to be efficient. The same is true for Model B. However, we should note that the usage of the classical conductivity (Spitzer 1962) is in question for CGs. In fact, we cannot explain the existence of cooling flows as long as we employ the classical one (Binney & Cowie 1981). Instead, it is suggested that tangled magnetic field (Rosner & Tucker 1989) or plasma instabilities (Pistinner & Shaviv 1996) are likely to suppress heat conduction significantly in CGs. Temperature fluctuations can then survive.

3.4. Evolution of Shock Surface

As we have seen in figure 7, sound wave propagates outward, modulating temperature profile. This also affects shock front propagation, since sound velocity inside the shock is three times greater than the front velocity in strong shock limits. We illustrate in figure 8 the radius of a shock surface (r_{shock}) as a function of the look-back time, time measured from the present time; i.e., $t_{\text{lb}} = 0$ corresponds to the present time (z = 0). The different types of lines correspond to same models as in figure 5. Shock surface moves with nearly a constant velocity (~ 200km s⁻¹). The radius of shock surface at z = 0, r_a , and mean propagation velocity, v_a , are listed in table 4. Again, there are wavy features seen in this figure.

Fig. 8.—

To understand the physics causing modulating features, we plot time variation of shock strength and shock radius ($r_{\rm shock}$) for Model A in figure 9. To evaluate shock strength we use $\Delta v = v_{\rm in} - v_{\rm out}$ with $v_{\rm in}$ and $v_{\rm out}$ being radial gas velocities inside and outside the shock surface, respectively, in the upper panel, and $T_7\Delta S$ where T_7 is the pre-shock gas temperature in the unit of 10⁷K and $\Delta S = S_{\rm in} - S_{\rm out}$ with $S_{\rm in}$ and $S_{\rm out}$ being specific entropies inside and outside the shock surface, respectively, in the lower panel. Since this quantity is proportional to heat produced through shock heating, it is a good representation of the shock strength. Both panels show time modulation in shock strengths. This in turn creates spatial modulation in entropy profile, since shock radius moves outward with time; namely, at the radius over which the shock passed with its maximum (minimum) strength in time, radial entropy profile exhibits a rapid (or slow) rise outward. The time modulation in shock strengths is likely to be caused by shock radius oscillation, and this oscillation is, as we discussed above, caused by the sound wave propagation. Note that while the wavy pattern in temperature profile at a fixed radius varies with time because of sound wave propagation, that in entropy profile does not, since entropy profile is unaffected by sound waves.

Fig. 9.—

3.5. Evolution of X-Ray Luminosity

Time evolutions of X-ray luminosity, $L(t_{\rm lb})$, and the normalized luminosity, $L_{\rm n}(t_{\rm lb}) \equiv L(t_{\rm lb})/L(t_{\rm lb} = 0)$, are plotted for each model in figure 10 (a) and (b), respectively. When calculating luminosity, we assume thermal bremsstrahlung of optically thin plasma (Rybicki & Lightman 1979),

$$\varepsilon^{\rm ff} \equiv \frac{dW}{dV \, dt} = 1.4 \times 10^{-27} T^{1/2} n^2 \bar{g}_B \,\,({\rm erg \ s^{-1} cm^{-3}}),\tag{30}$$

where T is gas temperature, n is number density, \bar{g}_B is a frequency average of velocityaveraged Gaunt factor. In this paper we set $\bar{g}_B = 1.2$. Emission from cool gas with temperature $T < 10^7$ K was neglected because we are interested in time variation of X-ray luminosity.

Fig. 10.—

Figure 11 plot evolution of L and L_n against z instead of time.

Fig. 11.—

In all models the luminosity rapidly rises just before the appearance of a shock wave, and then rises gradually afterwards. This behavior is due to the two distinct phases of evolution of gas.

- The first phase (before the emergence of shock wave): Through accretion of ambient gas towards the center gas in the central region is compressed adiabatically so that temperature and density there rapidly rise until temperature reaches about virial temperature. The rapid increase in L is thus due to a rapid rise in temperature of the central region.
- The second phase (after the appearance of shock wave): As the shock wave propagates outwards, high temperature region expands, thus increasing L. At the same time, gas inside the shock wave continues to be compressed adiabatically, thereby its emissivity being increased gradually. Luminosity thus increases faster than $[r_{\text{shock}}(t)]^3 \propto t^{2.5}$.

3.6. Dependence on the Cosmological Density Parameter

In this subsection we compare the results of Model A ($\Omega_0 = 1$) and Model B ($\Omega_0 = 0.2$) to discuss the Ω_0 dependence of the cluster evolution under the condition that the perturbation amplitudes at z = 10 and total masses contained in the perturbations are similar.

From table 2, we see that central DM density, $\rho_{\text{DM},0}$, is proportional to Ω_0 . The DM distribution is less concentrated in Model B than in Model A, because the ambient gas density is lower in a lower density universe so that a larger volume is needed to contain the same amount of mass (~ $10^{15}M_{\odot}$) initially. The absolute values of DM density are thus different among these models, but the shape of the DM density profiles are similar. This explains why nondimensional DM density profiles look similar among both models (figure 4a). On the other hand, central density of gas, $n_{\text{gas},0}$, depends also on its initial entropy and is, hence, not strictly proportional to Ω_0 (see table 3). This is responsible for the differences

in nondimensional gas density profiles among two models (figure 4b).

In both models X-ray luminosity rises, however, the increase in L(t) is slower in Model B than in Model A. This tendency is more clearly seen when L(t) is plotted against times, rather than against redshifts (see Fig. 10b, 11b). In the latter figure, the difference can be recognized at $z \sim 0.5$, but not at z > 1. This is because smaller amount of DM and gas surrounding a central condensation in a lower Ω_0 universe.

3.7. Dependence on the Initial Temperature of Gas

We calculated a model with lower initial temperature (LT) to see how initial temperature affects the later evolution. The temperature and entropy profiles at z = 0 of Models A and LT are depicted in figure 5. There is no significant difference among these models, especially in the structure inside the shock front. The thermal property of gas outside the shock front has little influence on the thermal property of gas inside because of enormous entropy production at the shock surface.

On the other hand, the central densities are different among these models (figure 12). According to $n_{\text{gas},0}$ estimated from the fitting data of table 3 and figure 12, central density of Model LT is about three times greater than that of Model A. In the density profiles at r > 0.3 Mpc, however, any difference can hardly be seen.

Fig. 12.—

In the central region, gas is adiabatically compressed until the temperature reaches about the virial temperature. So lower initial gas temperature results in higher gas density, if the virial temperatures are the same in both models. However, the difference in central gas densities cannot be perfectly explained by this picture. If there is no entropy production (i.e., gas is perfectly adiabatically compressed), central density of Model LT should be about thirty times greater than that of Model A. This means, shock heating took away the information about the initial gas temperature also in the central region.

Because of higher central gas density Model LT cluster is more luminous in X-ray than Model A cluster (figure 10a, 11b). Substantial difference, however, cannot be seen in evolutionary behavior of X-ray luminosity of both model. (figure 10b, 11b).

4. SUMMARY AND DISCUSSION

We carry out the numerical simulations of spherical clusters of galaxies with shell model for DM and second order up-wind TVD scheme for ICM to examine structural evolution of ICM. Shock front moves outwards as gas accretes towards cluster center, yielding a relatively flat temperature profile inside the shock front. Density and pressure profiles evolves in a self-similar fashion.

X-ray luminosity increases with time in two steps. At the initial collapse of DM gas in the central part is at first adiabatically compressed through accretion of ambient gas towards the center. Eventually, a shock wave appears near the center and X-ray luminosity rapidly rises until temperature increases and reaches about virial temperature via shock heating. In the late stage, in contrast, the luminosity rises only gradually, since the inner region already emits strong X-rays. The gradual brightening is due partly to the expansion of the high temperature region and partly to increasing X-ray emissivity of gas as the results of continuous adiabatic compression the gas inside the shock.

If we compare two clusters with the same density fluctuation amplitudes at z = 10 and with the same total masses but in different mean-density unvierses, the DM distribution is less concentrated in clusters in lower density universe. Hence, X-ray luminosity of clusters rises more slowly than in higher density universe.

Initial gas temperature has some influence on the central gas density. Higher initial temperature results in lower central density. This corresponds to the case that reheating of the ICM by, e.g., proto-galaxies is substantial. Density distribution in the outer region and temperature profile inside the shock front are, however, hardly influenced by changes in the initial temperature. Thus, the inclusion of reheating process modifies the scaling law between X-ray luminosity and temperature in the favorite way to reproduce the observed relation (NFW). Note that the epoch of the shock emergence depends on the initial specific entropy at the core; in the presence of reheating process (so that the initial gas temperature is $\sim 10^7$ K), specific entropy at the core is already high enough, and so the appearance of shock may be delayed.

In table 5 we compare the time dependent properties of Model A cluster and those of the self-similar solution by Bertschinger (1985). Both behavior looks very similar. Note that the self-similarity can be seen in all the calculated models. Although Einstein-de Sitter universe is assumed in Bertschinger (1985), we find that the self-similarity can be found for cluster evolution in low density universe ($\Omega_0 = 0.2, \lambda_0 = 0$).

Our results regarding the present profiles of density, temperature, and so on roughly coincide with those of the previous SPH simulations (Evrard 1990; NFW). However, the structure of the shock front, which was not well resolved in the previous SPH simulations, is now clearly captured in the present mesh-code simulations; our results show that hot gas of X-ray CG is separated with a definite boundary and that shock heating plays an important role for the heating process of ICM. We confirmed the persistence of radial infall of gas after the passage of the shock front. This results in adiabatic compression of the inner parts, inducing gradual temperature increase. We also found time variations in the strengths of the shock due to a sound wave propagation over the entire cluster, which modulates the radial distributions of temperature and density with relative amplitudes of about 10 %. Note that the property of the sound wave may depend on our assumption of spherical symmetry. It is open to question how such sound wave behaves in a realistic three-dimensional situation; i.e, when mass accretion takes place in a nonaxisymmetric way (e.g., by merger of multiple density condensations). If the temperature modulation will be observed with future X-ray mission (such as ASTRO-E), this can be used as a good probe to investigate the structure, especially, the mass distribution of CGs, since sound wave properties sensitively depend on the shape and the depth of the gravitational potential well, and thus on the DM mass distribution.

We would like to thank F.E.Nakamura and M.Hattori for helpful discussions and comments. This work is in part supported by Research Fellowships of the Japan Society for the Promotion of Science for Young Scientists (M.T.), and by the Grants-in Aid of the Ministry of Education, Science, Sports, and Culture of Japan, 06233101, 08640329 (S.M.).

Model	δ_0	$r_0 \; ({\rm Mpc})$	$T_{\rm gas,i}$ (K)	Ω_0
А	0.4	9.0	10^{7}	1.0
В	0.5	15.4	10^{7}	0.2
LT	0.4	9.0	10^{6}	1.0

Table 1: Parameters of each model.

Model	$\rho_{\rm DM,0} \ (10^{-26} \ {\rm g \ cm^{-3}})$	$r_{\rm DM}~({\rm Mpc})$	$\beta_{\rm DM}$	r_{100}
А	78.5	0.067	0.87	1.35
В	14.9	0.13	0.93	1.04
LT	83.3	0.070	0.87	1.34

Table 2: Density profiles of DM at z = 0.

Model	$n_{\rm gas,0}~(10^{-3}{\rm cm}^{-3})$	$r_{\rm gas} \ ({\rm Mpc})$	$\beta_{\rm gas}$
А	9.9	0.082	0.80
В	3.9	0.13	0.93
LT	30	0.065	0.87

Table 3: Density profiles of gas at z = 0.

Model	$r_a \ ({ m Mpc})$	$v_a \ (\mathrm{km} \ \mathrm{s}^{-1})$
А	1.2	210
В	1.4	230
LT	1.2	230

Table 4: The arrival radius of shock surface at $z = 0, r_a$, and mean propagating velocity, v_a .

	$ ho_{\rm DM}(r)$	$\rho_{\rm gas}(r)$	P(r)	$r_{\rm shock}(t)$
А	$r^{-2.6}$	$r^{-2.4}$	$r^{-3.0}$	$t^{0.83}$
Bertschinger(1985)	$r^{-2.25}$	$r^{-2.4}$	$r^{-2.9}$	$t^{0.89}$

Table 5: The comparison between the behavior of A1 and the self-similar solution of Bertschinger (1985)

REFERENCES

- Anninos, P., & Norman, M. L. 1996, ApJ, 459. 12
- Bertschinger, E. 1985, ApJS, 58, 39
- Binney, J., & Cowie, L. 1981, ApJ, 247, 464
- Bower, R. G., Böhringer, H., Briel, U. G., Ellis, R. S., Castander, F. J., & Couch, W. J. 1994, MNRAS, 268, 345
- Bryan, G. L., Cen, R., Norman, M. L., Ostriker, J. P., & Stone, J. M. 1994, ApJ, 428, 405
- Castander, F. J., Ellis, R. S., Frenk, C. S., Dressler, A., & Gunn, J. E. 1994, ApJ, 424, L79
- Cen, R. & Ostriker, J. P. 1994, ApJ, 429, 4
- Edge, A. C., Stewart, G. C., Fabian, A. C., & Arnaud, K. A. 1990, MNRAS, 245, 559
- Evrard, A. E. 1990, ApJ, 363, 349
- Fabian, A. C. 1994, ARA&A, 32, 277
- Fabricant, D. G., Kent, S. M. & Kurtz, M. J. 1989, ApJ, 336, 77
- Gioia, I. M., Henry, J. P., Maccacaro, T., Morris, S. L., Stock, J.T., & Wolter, A. 1990, ApJ, 356, L35
- Funato, Y., Makino, J. & Ebisuzaki, T. 1992a, PASJ, 44, 291
- Funato, Y., Makino, J. & Ebisuzaki, T. 1992b, PASJ, 44, 613
- Gunn, J. E., Gott, J. R. 1973, ApJ, 176, 1
- Henry, J. P., Gioia, I. M., Maccacaro, T., Morris, S. L., Stocke, J. T., & Wolter, A. 1992, ApJ, 386, 408
- Hénon, M. 1964, Ann. Astrophys., 27, 83
- Kang, H., Cen, R., Ostriker, J. P., & Ryu, D. 1994, ApJ, 428, 1

- Kneib, J. D. P., Mellier, Y., Pello, R., Miralda-Escude, J., Le Borgne, J. F., Boehringer, H., & Picat, J. P.
- Lynden-Bell, D. 1967, MNRAS, 136, 101
- Metzler, C. A., & Evrard, A. E. 1994, ApJ, 437, 564
- Monaghan, J. J. 1992, ARA&A, 30, 543
- Nakamura, S. 1996 Doctor Thesis of Tokyo Institute of Technology
- Navarro, J. F., Frenk, C. S., & White, S. D. M. 1995, MNRAS, 275, 720
- Hirsch, C. 1990, Numerical Computation of Internal and External Flows Vol.2:Computational Methods for Inviscid and Viscous Flows (John Wiley & Sons)
- Ohashi et al 1996, X-ray Imaging and Spectroscopy of Cosmic Hot Plasmas (ASCA 3rd Anniversary)
- Pistinner, S., & Shaviv, G. 1996, ApJ, 459, 147
- Peebles, P. J. E. 1982, ApJ, 257, 438
- Perrenod, S. C. 1978, ApJ, 226, 566
- Rosner, R., & Tucker, W. H. 1989, ApJ, 338, 761
- Rybicki, G. B., & Lightman. A. P. 1979, Radiative Process in Astrophysics (John Wiley & Sons)
- Sarazin, C. L. 1988, X-ray Emission from Clusters of Galaxies (Cambridge University Press)
- Spitzer, L., Jr. 1962, Physics of Fully Ionized Gases (New York:Wiley)
- Suto, Y. 1993, Prog. Theor. Phys., 90, 1173
- Takizawa, M., & Inagaki, S. 1996, submitted to PASJ

Thomas, P. A., & Couchman, H. M. P. 1992, MNRAS, 257, 11

This manuscript was prepared with the AAS ${\rm \sc Lat}_{\rm E\!X}$ macros v4.0.

Figure Captions

- figure 1 (a) Radial profiles of initial density perturbations and (b) $(\delta M/M)/\sigma_{8,\text{CDM}}$, where $(\delta M/M)$ is the mass fluctuation averaged over a scale r and $\sigma_{8,\text{CDM}}$ is the rms of mass fluctuation of scale r obtained from CDM power spectrum with σ_8 being a normalization. The solid line correspond to Model A and the dotted line corresponds model B.
- figure 2 The evolution of (a) density profile and (b) radial velocity profile of DM of Model A cluster. The different lines correspond to different redshifts (and different times): z = 2.5 (t = 1 Gyr) by the long dash line; z = 1.2 (t = 2 Gyr) by the short dash line; z = 0.4 (t = 4 Gyr) by the dotted line; and z = 0 (t = 6.5 Gyr) by the solid line, respectively.
- figure 3 Evolution of Model A cluster. Radial profiles of representative physical quantities of gas are depicted: (a) radial velocity, (b) density, (c) temperature, (d) pressure, and (e) entropy. The different types of lines correspond to same redshifts as in figure 2
- figure 4 Nondimensional density profiles at z = 0 for Models A (by the solid line) and for B (by the dotted line), respectively. Here, density is scaled with ρ_{c0} , critical density of the universe at z = 0, and radius is scaled with r_{100} .
- figure 5 (a) Temperature profiles and (b) entropy profiles of each model at z = 0. Entropy is normalized to zero at the outer boundary. The different types of lines correspond to different models: Model A by the solid line; Model B by the dotted line; and Model LT by the short dash line, respectively.
- figure 6 The ratio of pressure gradient force and gravitational force inside the shock surface of Model A cluster from t = 5.5 Gyr to 6.5 Gyr. If the system is perfectly in

hydrostatic equilibrium, the ratio, $|dP/dr|/F_{\rm g}$, should be unity.

- figure 7 Radial velocity of gas inside shock front from t = 5.5 Gyr to 6.5 Gyr of model A. Sound wave propagates outwardly. Radial infall systematically remains even after the passage of the shock.
- figure 8 Time evolution of the radius of a shock surface, r_{shock} . The abscissa is the look-back time, time measured from the present time; $t_{\text{lb}} = 0$ at z = 0. Shock surface moves with a nearly constant velocity (~ 200 km s⁻¹). The different types of lines correspond to same models as in figure 5.
- figure 9 Evolution of shock strength and the shock radius of Model A. (a) Time evolution of $r_{\rm shock}$ by the dotted line and that of Δv , jump in the radial velocity of gas over the shock surface. (b) That of $T_7\Delta S$, where T_7 is the pre-shock gas temperature in the unit of 10⁷K and ΔS is the jump in the entropy over the shock surface. This quantity is proportional to heat produced through shock heating.
- figure 10 (a) Time evolution of X-ray luminosity, $L(t_{\rm lb})$, and (b) that of normalized luminosity, $L_{\rm n}(t_{\rm lb}) \equiv L(t_{\rm lb})/L(t_{\rm lb} = 0)$. In all models L rapidly rises just before the generation of a shock wave, and then rises gradually afterwards. The different types of lines correspond to same models as in figure 5.
- **figure 11** Evolution of L and L_n plotted against z.
- figure 12 Gas density profiles for Model A (solid line) and for Model LT (dotted line). Central density is about three times greater in Model LT cluster than in Model A cluster. In the region of r > 0.3Mpc, however, this difference can hardly be seen.

























