# Understanding Gravitational Clustering with Non-Linear Perturbation Theory

Román Scoccimarro

CITA, McLennan Physical Labs, 60 St George Street, Toronto ON M5S 3H8

I discuss new results concerning the evolution of the bispectrum due to gravitational instability from gaussian initial conditions using one-loop perturbation theory (PT). Particular attention is paid to the transition from weakly non-linear scales to the non-linear regime at small scales. Comparison with numerical simulations is made to assess the regime of validity of the perturbative approach.

### 1 Introduction

This work is based on results to be reported in [1]. Here, I briefly present results on the one-loop corrections to the bispectrum and compare them to numerical simulations for CDM initial spectra.<sup>*a*</sup> In particular, we work in terms of the hierarchical amplitude Q defined from the bispectrum,  $B(\mathbf{k}_1, \mathbf{k}_2)$ , and power spectrum, P(k), as follows (superscripts denote, tree-level, one-loop PT, and so on):

$$Q \equiv \frac{B(\mathbf{k}_1, \mathbf{k}_2)}{\Sigma(\mathbf{k}_1, \mathbf{k}_2)} = \frac{B^{(0)}(\mathbf{k}_1, \mathbf{k}_2) + B^{(1)}(\mathbf{k}_1, \mathbf{k}_2) + \dots}{\Sigma^{(0)}(\mathbf{k}_1, \mathbf{k}_2) + \Sigma^{(1)}(\mathbf{k}_1, \mathbf{k}_2) + \dots} = Q^{(0)} + Q^{(1)} + \dots, \quad (1)$$

with:

$$\langle \delta(\mathbf{k})\delta(\mathbf{k}')\rangle = \delta_D(\mathbf{k} + \mathbf{k}') P(k),$$
 (2)

$$\langle \delta(\mathbf{k}_1)\delta(\mathbf{k}_2)\delta(\mathbf{k}_3) \rangle = \delta_D(\mathbf{k}_1 + \mathbf{k}_2 + \mathbf{k}_3) \ B(\mathbf{k}_1, \mathbf{k}_2).$$
(3)

$$\Sigma(\mathbf{k}_1, \mathbf{k}_2) \equiv P(k_1) \ P(k_2) + P(k_1) \ P(k_3) + P(k_2) \ P(k_3)$$
(4)

where we have used that the bispectrum is defined for closed triangle configurations,  $\sum_{i=1}^{3} \mathbf{k}_{i} = 0$ . The perturbative quantities in Eq. (1) can be calculated from the standard machinery of PT.<sup>1,2</sup> In the following, we consider Q for configurations where  $k_1/k_2 = 2$ , as a function of  $\theta$ , the angle between  $\hat{\mathbf{k}}_1$  and  $\hat{\mathbf{k}}_2$ .

## 2 Results

Figure 1 shows Q in numerical simulations compared to tree-level PT and oneloop PT. Error bars in these plots are estimated from the number of independent Fourier modes contributing to each configuration assuming gaussianity.<sup>1</sup>

<sup>&</sup>lt;sup>*a*</sup> with  $\Omega = 1$ ,  $\Gamma = 0.25$ . The simulation data is publically available through the Hydra Consortium Web page (http://coho.astro.uwo.ca/pub/consort.html).



Figure 1: The hierarchical amplitude Q for triangle configurations with  $k_1/k_2 = 2$  as a function of the angle  $\theta$  between  $\hat{\mathbf{k}}_1$  and  $\hat{\mathbf{k}}_2$  in the Hydra CDM numerical simulations (symbols), tree-level PT (dotted lines) and one-loop PT (solid lines).

The degree of non-linearity in each case can be inferred from the dimensionless power spectrum,  $\Delta(k) \equiv 4\pi k^3 P(k)$ . In the top panels ( $\sigma_8 = 0.21$ ), we note a clear deviation of the N-body results from the tree-level PT prediction for Q, and a good agreement with the one-loop correction. At this stage of nonlinear evolution, the dynamics is dominated by large-scale power and therefore an enhancement of Q at collinear configurations ( $\theta = 0, \pi$ ) develops.<sup>2</sup> In the bottom panels ( $\sigma_8 = 0.33$ ), where already  $\Delta(k_1) > 1$ , we use the ratio of oneloop quantities in Eq. (1) (denoted as "one-loop (s)" in Fig. 1) for the one-loop prediction.<sup>2</sup> We see very good agreement for configurations close to collinear, and a progressively flattening of  $Q(\theta)$  as we look at smaller scales. The flattening is due to configurations close to equilateral becoming more probable due to random motions at small scales.<sup>1</sup> At even more non-linear scales, Q becomes configuration independent, in rough agreement with the hierarchical ansatz for the three-point function.<sup>1</sup>

### Acknowledgments

This work is based on a project in collaboration with S. Colombi, J.N. Fry, J.A. Frieman, E. Hivon, & A. Mellot. I would like in addition to thank F. Bernardeau, E. Gaztañaga, B. Jain, R. Juszkiewicz, C. Murali for conversations, and especially Hugh Couchman for numerous helpful discussions. The CDM simulations analyzed in this work were obtained from the data bank of cosmological N-body simulations provided by the Hydra consortium (http://coho.astro.uwo.ca/pub/data.html) and produced using the Hydra N-body code.<sup>3</sup>

### References

- 1. R. Scoccimarro, S. Colombi, J.N. Fry, J.A. Frieman, E. Hivon, and A. Melott, in preparation (1997).
- 2. R. Scoccimarro, submitted to Astrophys. J., astro-ph/9612207, (1996).
- H. Couchmann, P.A. Thomas, and F. Pearce, Astrophys. J. 452, 797 (1995).