

Spectroscopy of Close Companions to QSOs and the Ages of Interaction-Induced Starbursts¹

Gabriela Canalizo² and Alan Stockton²

Institute for Astronomy, University of Hawaii, 2680 Woodlawn Drive, Honolulu, HI 96822

ABSTRACT

We present low-resolution absorption-line spectra of three candidate close ($< 3''$) companions to the low redshift QSOs 3CR 323.1, PG 1700+518 and PKS 2135–147. The spectra were obtained with LRIS on the Keck telescopes and with the Faint Object Spectrograph on the University of Hawaii 2.2 m telescope. For 3CR 323.1 and PG 1700+518, we measure relative velocities that are consistent with an association between the QSOs and their companion galaxies. The spectral features of the companion galaxy to 3CR 323.1 indicate a stellar population of intermediate age (~ 2.3 Gyr). In contrast, the spectrum of the companion object to PG 1700+518 shows strong Balmer absorption lines from a relatively young stellar population, along with the Mg *Ib* absorption feature and the 4000 Å break from an older population. By modeling the two stellar components of this spectrum, it is possible to estimate the time that has elapsed since the end of the most recent major starburst event: we obtain ~ 0.1 Gyr. This event may have coincided with an interaction that triggered the QSO activity. Finally, our spectroscopy shows conclusively that the supposed companion to PKS 2135–147 is actually a projected Galactic G star.

Subject headings: galaxies: evolution—galaxies: interactions—quasars: individual (3C 323.1, PG 1700+518, PKS 2135–147)

¹Based on observations made with the NASA/ESA Hubble Space Telescope, obtained from the data archive at the Space Telescope Science Institute, which is operated by the Association of Universities for Research in Astronomy, Inc., under NASA contract NAS 5-26555.

²Visiting Astronomer, W.M. Keck Observatory, jointly operated by the California Institute of Technology and the University of California.

1. Introduction

There is considerable circumstantial evidence that strong interactions can provide the means to feed material into a galaxy nucleus to fuel activity in luminous QSOs (see, e.g., Stockton 1990; Hutchings & Neff 1992). Such evidence includes tidal tails and bridge-like asymmetries in QSO host galaxies, extended emission around QSOs, and close companion galaxies. Stockton (1982) proposed that some QSO companions might be tidally-stripped cores from galaxies that had interacted with the QSO host galaxies.

Interest in QSO companions has been reawakened recently by the HST results from a nearby luminous QSO sample. Bahcall *et al.* (1997) found at least one such companion lying within 25 kpc for 13 of the 20 QSOs in their sample. This result underlines the importance of companion galaxies as potential triggerers of the QSO activity.

However, imaging observations alone cannot distinguish true companions from projections of other objects, so it is important to make spectroscopic observations as well. With precise redshift measurements we can calculate the relative velocities between the QSO and its companion(s) and determine whether the latter are likely to be in gravitationally bound orbits. The relatively slow passages one obtains in bound systems appear to be important in bringing fresh material deeply into the center of galaxies.

Stockton (1982) measured redshifts from spectra of close companions to Markarian 205, 3CR 323.1, and PKS 2135–147 and found all to have redshifts very close to those of the respective QSOs. He argued that all three were probably tidally-limited companion galaxies. However, the redshifts of the latter two objects were determined from emission lines alone. Later, Stockton & MacKenty (1987) found that the emission lines seen in their spectra come from the general extended emission around the QSO and not specifically from the supposed companion, so the redshifts for these two objects are no longer valid.

The main objective of this study was therefore to obtain better quality spectra and measure accurate redshifts for QSO companions from stellar absorption lines. We remeasured the redshifts of the objects near 3CR 323.1 and PKS 2135–147. In addition, we measured for the first time the redshift of an object recently found near another QSO, PG 1700+518.

2. Observations and Data Reduction

3CR 323.1 was observed on 1991 July 19 using an image slicer in the Faint Object Spectrograph with a Tektronix 1024×1024 CCD on the University of Hawaii 2.2 m telescope

on Mauna Kea. The spectra were centered at 5500 \AA and had a resolution of $2.17 \text{ \AA pixel}^{-1}$. The three image slicer slits were oriented perpendicularly to the line joining the companion to the QSO, with the companion falling on the central slit. The other two slits were used to monitor the QSO light.

PG 1700+518 and PKS 2135–147 were observed on 1996 February 15 and 1996 October 13, respectively, with the Low-Resolution Imaging Spectrometer (LRIS) on the Keck I (February) and the Keck II (October) telescopes. We used a $600 \text{ groove mm}^{-1}$ grating blazed at 5000 \AA to obtain a dispersion of $1.24 \text{ \AA pixel}^{-1}$ and a useful wavelength range of $4400\text{--}6900 \text{ \AA}$ for PG 1700+518 and $4200\text{--}6700 \text{ \AA}$ for PKS 2135–147. The slit ($1''$ wide, projecting to ~ 5 pixels on the Tektronix 2048×2048 CCD) was oriented in both cases along the line joining the companion and the QSO. The total exposure times were 60 minutes for 3CR 323.1 and 40 minutes each for PKS 2135–147 and PG 1700+518.

The spectra were reduced with IRAF, using standard reduction procedures. Scattered QSO light sometimes contaminated the companion spectra, and it had to be subtracted before any companion spectral features were visible. For 3CR 323.1, no subtraction was necessary. For PG 1700+518 and PKS 2135–147, the contamination amounted to 56% and 38%, respectively, at rest-frame 4500 \AA . The scattered light was removed by reversing the QSO continuum profile about its center and subtracting it from the companion spectrum.

3. Results and Discussion of Individual Objects

The fields of the three QSOs are shown in Figs. 1 and 2 (Plates 00 and 00), on which objects of interest are labelled and the slit positions indicated. Table 1 summarizes the observed properties of the three objects. Columns 2 and 3 give the redshift of the QSO and the companion respectively, both of them obtained from our data. In columns 4 and 5 we list the projected separations between the QSOs and their close companions in arcseconds and kiloparsecs, respectively. In column 6 we list the velocity of the companion in the rest frame of the QSO.

3.1. 3CR 323.1

Figure 3 shows the spectrum of the companion (*a* in the upper panel of Fig. 1) and the QSO in their rest frame. While [O II] and [O III] emission lines from the extended ionized region contaminate the spectrum of the companion, absorption features are clearly present, including the Ca II H and K lines, the G-band, and the CN feature near 3830 \AA .

The redshift obtained from the absorption features is $z = 0.2664$, which gives a velocity of $470 \pm 230 \text{ km s}^{-1}$ in the QSO reference frame defined by the narrow emission lines. Given the uncertainty in both the velocity measurements and in the true systemic velocity of the QSO host galaxy, there is a good possibility that the object is bound to the QSO, especially if the QSO host has a fairly high mass.

Superposed on the spectrum of the companion in Fig. 3 is a Bruzual & Charlot (1993) model for a 2.3 Gyr-old instantaneous burst stellar population with solar metallicity. The discrepancy in the fit is mainly due to the lower resolution in the model, particularly evident around the Ca II features. The intermediate-age population we find dominating the optical spectrum could be consistent with an aged starburst from the time of a previous interaction between the companion and the QSO host galaxy, but better spectra over a wider wavelength range and consideration of more complex models would be necessary to make such a scenario compelling.

3.2. The demise of the companion to PKS 2135–147

The object $2''$ southeast of PKS 2135–147 (a in the lower panel of Fig. 1) was first noticed by Kristian (1973). Stockton (1982) found that this object had a stellar or nearly stellar profile. He identified it as a compact companion associated with the QSO by measuring its redshift from emission lines. As noted previously, this redshift is invalid since the emission comes from an extended region surrounding the QSO, not from the companion (Stockton & MacKenty 1987). Later, Hickson & Hutchings (1987) found weak evidence for the Mg Ib absorption feature at the redshift of the QSO in the spectrum of the companion, and Hutchings & Neff (1992) thought that the object appeared extended along the radial direction towards the QSO.

On WFPC2 images of PKS 2135–147 from the HST archive, the object has a completely stellar profile (see Fig. 1). This lack of extension made us suspect that the object was actually a close projection of a Galactic star. The high signal-to-noise LRIS spectrum in the top panel of Fig. 4 confirms this suspicion. Superposed on the spectrum of the object is the spectrum of a G 5 star, which matches closely the flux distribution and absorption features of the supposed companion.

In Fig. 4 we also show the spectrum of galaxy b , $5''$ to the southeast of the QSO, and having a radial velocity with respect to the QSO of $-150 \pm 80 \text{ km s}^{-1}$. This galaxy and galaxy d were among those previously identified as associated galaxies (Stockton 1978). A much fainter galaxy on our slit $15''$ to the northwest of the QSO (c in Figs. 1 and 4) has a

relative velocity of $-200 \pm 40 \text{ km s}^{-1}$. Finally, we observe a strong emission region (marked by arrows and labelled “em” in Fig. 1) beyond companion galaxy b , $\sim 12''$ southeast of the QSO; its velocity with respect to the QSO is $-64 \pm 20 \text{ km s}^{-1}$.

3.3. PG 1700+518

PG 1700+518 is one of the brightest low-redshift QSOs, one of the few low-redshift broad-absorption-line (BAL) QSOs, and a luminous IR object. A peculiar characteristic of this object is an arc-like structure $\sim 2''$ north of the QSO (Hutchings *et al.* 1992; Stickel *et al.* 1995; Stockton *et al.* 1997; see Fig. 2). It is not clear from the imaging data whether this object is a distorted companion or a tidal feature from the host galaxy, though we will generally refer to it as a “companion” for convenience. Both we and Stickel *et al.* (1995) find evidence for a fairly symmetric main component to the host galaxy, aligned approximately E–W (see Fig. 2).

Figure 5 shows the spectrum of the companion as well as that of the QSO. We measure a redshift $z = 0.2929$ for the companion and $z = 0.2923$ for the QSO, giving a relative velocity of $140 \pm 150 \text{ km s}^{-1}$ in the reference frame of the QSO. The strong [O II] and [O III] lines in the spectrum come from the extended ionized gas around the QSO and not specifically from the companion. The apparent absorption features marked with an asterisk are a result of an over-subtraction of the QSO Fe II emission. The Fe II lines seem to be spatially variable in PG 1700+518 (Hickson & Hutchings 1987), leading to an uncertainty in their subtraction when we correct for the QSO scattered light contribution.

The companion shows a classic “E+A” spectrum. The presence of strong Balmer absorption lines is indicative of a relatively young stellar population, whereas the Mg I b feature and the 4000 Å break suggest an older population. We used the Bruzual & Charlot (1993) isochrone synthesis models, with solar metallicity and a Scalo (1986) IMF, to model the two suggested components of the companion spectrum. The ages of the two isochrone stellar populations, determined from a χ^2 fit to the data, are 0.09 (+0.04, –0.03) Gyrs and 12.25 Gyrs, respectively. The younger population contributes 68% of the total light at 4500 Å (rest frame) and comprises $\sim 6\%$ of the total luminous mass. We caution that we are making a very simple approximation in assuming instantaneous bursts and using only two discrete stellar populations. Accordingly, our best-fit model comprising these two populations cannot be claimed to be unique, but, as shown by the lighter curve superposed on our observed spectrum in Fig. 5, it does fit the data reasonably well.

E+A galaxies are characterized by strong Balmer absorption, fairly blue continuum,

and weak [O II] emission. At least some E+A galaxies seem to result from mergers (Lavery et al. 1992, Oegerle et al. 1991), and Liu & Kennicutt (1995) find a large fraction of E+A-like galaxies in their sample of nearby merging galaxies. Together with the unusual morphology of the companion, this spectrum gives us fairly strong evidence for an interaction that triggered both the QSO activity and an extensive starburst.

4. Age-Dating Starbursts associated with QSOs

While determining spectral ages for stellar populations produced in a starburst may be the most direct approach to placing QSO hosts in an age sequence, correlations among other observable parameters may ultimately give insights into the nature of their evolution.

The spectrum of PG 1700+518 (Fig. 5) shows unusually strong Fe II emission lines for a QSO. Although the Fe II emission in AGNs has been extensively studied (Joly 1991 and references therein), its precise nature is still poorly understood. However, from optical spectral characteristics and relative positions on the far-IR $\alpha[60, 25]$, $\alpha[100, 60]$ diagram, L ipari (1994) suggests an evolutionary sequence connecting the strength of the Fe II emission with the time elapsed since a major starburst. This scenario fits well with the proposal of Sanders et al. 1988, who have suggested an evolutionary connection between ultraluminous IRAS galaxies (virtually all of which are interacting or merging systems) and the classical QSO population.

Extreme Fe II objects like Mrk 231 (Hutchings & Neff 1987; L ipari et al. 1994), close to the blackbody region in the FIR color–color plot, are believed to be in the last stages of merger and in the midst of a major starburst. Their narrow-line regions (NLR) are weak or nonexistent because the heavy obscuration of the QSO nucleus by dust strongly attenuates its UV continuum. Objects like IRAS 07598+6508 have slightly less extreme Fe II emission and a position in the FIR color–color diagram intermediate between the black-body and the power-law regions; they are presumed to be right at the end of the starburst stage. Finally, strong (but not extreme) Fe II emitters like PG 1700+518, which are still closer to the power law region of the two-color plot, but which also still have very weak NLR, are in the early post-starburst stage. The theoretical underpinning of this model is the supposition that the strong Fe II emission arises principally in a superwind generated by the starburst (Terlevich et al. 1992). This assumption is supported by our confirmation of the spatial variability of the Fe II emission in PG 1700+518 (Hickson & Hutchings 1987).

Such an evolutionary sequence is at least qualitatively consistent with another correlation pointed out by Stockton & Ridgway (1991), who noticed that the three

low-redshift QSOs that fall near the region of the FIR two-color plot occupied by luminous FIR galaxies (Mrk 231, Mrk 1014, and 3CR 48) are also the three QSOs that show the most unambiguous evidence for tidal tails.

Whether the arc-like object associated with PG 1700+518 is actually a tidal tail associated with either the host galaxy or a companion is still uncertain, but the age we estimate since the end of the most recent starburst in this feature is consistent with the evolutionary scenarios mentioned above. If a close interaction induced a starburst in the host galaxy as well as in the companion 9×10^7 years ago, one might well expect the system now to be a post-starburst object with strong Fe II emission.

Thus for at least some classes of young QSOs, we have a number of potential indicators for defining an evolutionary sequence connecting ultraluminous IR galaxies with classical QSO population. The strength of the Fe II emission may plausibly be related to the numbers of Type II supernovae produced in a starburst (Terlevich et al. 1992). The position in the FIR two-color plot indicates the relative balance of thermal emission from warm dust associated with a starburst and nonthermal emission from an AGN nucleus. The morphology of the QSO host galaxy and the distribution and morphologies of very close companions may give clues to the interaction history of the immediate environment of the QSO nucleus. Finally, the age of the youngest host galaxy or companion galaxy stellar population may well pinpoint the time since the event that triggered the QSO activity. Working out the relationships among these various parameters will be a difficult task, but one that will almost certainly clarify the relationships between IR galaxies and QSOs.

We thank Susan Ridgway for helping with some of the observations. This research was partially supported by NSF under grants AST92-21909 and AST95-29078.

REFERENCES

- Bahcall, J. N., Kirharos, S., Saxe, D.H., & Schneider, D.P. 1997, ApJ, 479, in press
Bruzual, G., & Charlot, S. 1993, ApJ, 405, 538
Hickson, P., & Hutchings, J. B. 1987, ApJ, 312, 518
Hutchings, J. B., & Neff, S. G. 1992, AJ, 104, 1
Hutchings, J. B., Neff, S. G., & Gower, A. C. 1992, PASP, 104, 62
Joly, M. 1991 A&A, 242, 49
Kristian, J. 1973 ApJ, 179, L61

- Lavery, R. J., Pierce, M. J., & McClure, R. D. 1992, *AJ*, 104, 2067
- Lípari, S. 1994 *ApJ*, 436, 102
- Lípari, S., Colina L., & Macchetto F., 1994 *ApJ*, 427, 174
- Liu, C. T., & Kennicutt, R. C. 1995 *ApJ*, 450, 547
- Oegerle, W. R., Hill, J., & Hoessel, J. 1991, *ApJ*, 381, L9
- Pettini, M., & Boksenberg, A. 1985, *ApJ*, 294, L73
- Sanders, D. B., Soifer, B. T., Elias, J. H., Madore, B. F., Matthews, K., Neugebauer, G., & Scoville, N. Z. 1988, *ApJ*, 325, 74
- Scalo, J. M. 1986, *Fund. Cosmic Phys.*, 11, 1
- Stickel, M., Fried, J. W., McLeod, K. K., & Rieke, G. H. 1995 *AJ*, 109, 1979
- Stockton, A. 1978 *ApJ*, 223, 747
- Stockton, A. 1982 *ApJ*, 257, 33
- Stockton, A. 1990, in *Dynamics and Interactions of Galaxies*, ed. R. Wielen, (Springer-Verlag: Berlin), 440
- Stockton, A., & MacKenty, J. W. 1987, *ApJ*, 316, 584
- Stockton, A., & Ridgway, S. E. 1991, *AJ*, 102, 488
- Stockton, A., Ridgway, S. E., & Kellogg, M., in preparation
- Terlevich, R., Tenorio-Tagle, G., Franco, J., & Melnick, J 1992, *MNRAS*, 255, 713
- Turnshek, D. A., Foltz, C. B., Weymann, R. J., Lupie, O. L., McMahon, R. C., & Peterson, B. M. 1985 *ApJ*, 294, L1

Table 1. Observed Properties of Close Companions

Name	z_{QSO}	z_{comp}	$\Delta r(\prime\prime)$	$\Delta r(\text{kpc})^a$	$\Delta v(\text{km s}^{-1})$
3CR 323.1	0.2644	0.2664	2.7	9.2	470
PKS 2135–147	0.2005	0.0000	2.0
PG 1700+518	0.2923	0.2929	2.0	7.2	140

^a $H_0 = 75 \text{ km s}^{-1} \text{ Mpc}^{-1}$ and $q_0 = 0.5$ assumed.

Fig. 1.— Fields of 3CR 323.1 (*top panel*) and PKS 2135–147 (*bottom panel*), both from HST archival data. Insets show the main regions of interest at lower contrast and $2.5\times$ larger scale. The parallel lines indicate the orientations of the slits for spectroscopy. In both panels, the object labelled “*a*” is the candidate companion galaxy. In the PKS 2135–147 field, galaxies *b*, *c*, and *d* have been spectroscopically confirmed to have essentially the same redshift as the QSO. The arrows labelled “em” indicate the position of a strong emission-line component seen in our spectrum. North is up and East to the left in both panels.

Fig. 2.— The field of PG 1700+518. The main upper panel shows an image obtained with the University of Hawaii 2.2 m telescope (Stockton, Ridgway, & Kellogg 1997) in an essentially line-free continuum band centered at 7248 \AA and having a FWHM of 1260 \AA (5610 \AA and 975 \AA , respectively, in the rest frame). The QSO has been subtracted, using a PSF derived from a nearby star, and the white cross indicates the QSO position. Extensions of the parallel lines indicate the region covered by the slit. The upper-left inset shows the image before PSF subtraction, and the lower-right inset shows a lower-contrast version of the subtracted image at twice the scale of the main panel. The white oversubtracted regions are due to a slight mismatch of the profiles of the star and the QSO; they have been suppressed in the main panel for clarity. The lower panel shows an *H*-band image before (*left*) and after (*right*) PSF subtraction. The small inset shows a star with the stretch adjusted so that the peak is the same as that of the QSO host galaxy in the subtracted image.

Fig. 3.— 3CR 323.1 spectra in rest frame. (*top*) Companion (heavier line) and a 2.3 Gyr-old isochrone synthesis model. (lighter line) (*bottom*) QSO. In this and the following figures, the spectra of the companions have been smoothed with Gaussian filters having σ equal to the projected slit widths. The QSO spectra have not been smoothed.

Fig. 4.— Spectra of objects associated with PKS 2135-147. Panels are identified by object labels on Fig. 1. (a) Comparison of PKS 2135–147 “companion” *a* (heavier line) and a G 5 star (lighter line). The spectrum of the G star has been normalized to the flux of the companion spectrum. (b) Galaxy 5'' southeast of the QSO, at the same redshift as the QSO, transformed to the rest frame. (c) Faint galaxy 15'' northwest of the QSO, also at the same redshift. (Q) QSO spectrum.

Fig. 5.— PG 1700+518 spectra in rest frame. (*top*) Companion (heavier line) and isochrone synthesis model (lighter line). (*bottom*) QSO spectrum.

This figure "gcf1.gif" is available in "gif" format from:

<http://arxiv.org/ps/astro-ph/9702204v1>

This figure "gcfg2.gif" is available in "gif" format from:

<http://arxiv.org/ps/astro-ph/9702204v1>




Fig. 3.— 3CR 323.1 spectra in rest frame. (*top*) Companion (heavier line) and a 2.3 Gyr-old isochrone synthesis model. (lighter line) (*bottom*) QSO. In this and the following figures, the spectra of the companions have been smoothed with Gaussian filters having σ equal to the projected slit widths. The QSO spectra have not been smoothed.




Fig. 4.— Spectra of objects associated with PKS 2135-147. Panels are identified by object labels on Fig. 1. (a) Comparison of PKS 2135-147 “companion” *a* (heavier line) and a G5 star (lighter line). The spectrum of the G star has been normalized to the flux of the companion spectrum. (b) Galaxy 5'' southeast of the QSO, at the same redshift as the QSO, transformed to the rest frame. (c) Faint galaxy 15'' northwest of the QSO, also at the same redshift. (Q) QSO spectrum.




Fig. 5.— PG 1700+518 spectra in rest frame. (*top*) Companion (heavier line) and isochrone synthesis model (lighter line). (*bottom*) QSO spectrum.