Fundamental stellar parameters of ζ Pup and γ^2 Vel from HIPPARCOS data

Daniel Schaerer^{1,2,3}

Werner Schmutz⁴

Michel Grenon²

ABSTRACT

We report parallax measurements by the HIPPARCOS satellite of ζ Puppis and γ^2 Velorum. The distance of ζ Pup is $d = 429^{+120}_{-77}$ pc, in agreement with the commonly adopted value to Vela OB2. However, a significantly smaller distance is found for the γ^2 Vel system: $d = 258^{+41}_{-31}$ pc.

The total mass of γ^2 Vel derived from its parallax, the angular size of the semimajor axis as measured with intensity interferometry, and the period is $M(WR+O) = 29.5 \pm 15.9 M_{\odot}$. This result favors the orbital solution of Pike et al. (1983) over that of Moffat et al. (1986). The stellar parameters for the O star companion derived from line blanketed non-LTE atmosphere models are: $T_{\rm eff} = 34000 \pm 1500$ K, $\log L/L_{\odot} = 5.3 \pm 0.15$ from which an evolutionary mass of $M = 29 \pm 4 M_{\odot}$ and an age of $4.0^{+0.8}_{-0.5}$ Myr is obtained from single star evolutionary models. With non-LTE model calculations including He and C we derive a luminosity $\log L/L_{\odot} \sim 4.7 \pm 0.2$ for the WR star. The mass-luminosity relation of hydrogen-free WR stars implies a mass of $M_{\rm WR} \sim 5 \pm 1.5 M_{\odot}$.

From our data we favor an age of ~ 10 Myr for the bulk of the Vela OB2 stars. Evolutionary scenarios for ζ Pup and γ^2 Vel are discussed in the light of our results.

Subject headings: Stars: distances – Stars: early-type – Stars: fundamental parameters – Stars: mass loss – Stars: Wolf-Rayet

Submitted to ApJ Letter

²Observatoire de Genève, CH-1290 Sauverny, Switzerland

³Observatoire Midi-Pyrénées, 14, Av. E. Belin, F-31400 Toulouse, France

⁴Institut für Astronomie, ETH-Zentrum. CH-8092 Zürich, Switzerland

1. Introduction

Accurate stellar parameters of massive stars are of fundamental importance to guide our understanding of stellar evolution. The closest single O star is ζ Pup (HD 66811, O4I(n)f), which is used as a prototype for radiation-driven wind models (e.g. Pauldrach et al. 1994, Schaerer & Schmutz 1994 for recent models). The closest O and WR stars are found in the system γ^2 Vel (HD 68273). Its binary nature hampers a spectroscopic analysis of its components but it has the advantage to allow direct mass determinations.

With the HIPPARCOS satellite it has been possible to obtain annual parallaxes for these two objects. In the present publication we discuss their fundamental stellar parameters based on the HIPPARCOS distances. Establishing a reliable distance – and hence luminosity – for ζ Pup is of particular interest for quantitative comparisons of hydrodynamic wind models with observations and the understanding of its observed He enrichment (cf. Bohannan et al. 1990). With an accurate distance to the γ^2 Vel system we are in a position to test the mass-luminosity relation for the WR star.

Stellar parameters for ζ Pup and γ^2 Vel based on the HIPPARCOS parallaxes are revised and/or determined from non-LTE atmosphere models in §2. In §3 we discuss evolutionary scenarios for these objects in the light of previous observations and new HIPPARCOS parallaxes of few presumed members of the Vela OB2 and Vela R2 associations.

2. Revised stellar parameters

2.1. ζ Pup

The HIPPARCOS parallax of ζ Pup is 2.33 \pm 0.51 mas corresponding to a distance of $d = 429^{+120}_{-77}$ pc. This value is consistent with the canonically adopted value of d = 450 pc based on its presumed association with Vela OB2 (cf.

Brandt et al. 1971), which may, however, be questionable (see §4). Distance dependent stellar parameters derived so far for ζ Pup are thus compatible with the HIPPARCOS measurement. We therefore only provide an update of these values adopting the new distance measurement and its error.

Rescaled to the new distance the radio massloss rate from Lamers & Leitherer (1993) is $\log \dot{M} = -5.57 \pm 0.15$, where we adopt the same uncertainty as these authors. As mentioned earlier the basic parameters of O stars with important winds are still somewhat uncertain (for ζ Pup cf. Bohannan et al. 1990, Gabler et al. 1989, Schaerer & Schmutz 1994, Puls et al. 1996). Adopting the model results ($T_{\rm eff}=42000$ K, BC) and observational parameter (V, A_V) from Bohannan et al. (1990) the new distance yields the following values (the uncertainties reflect only the distance): $\log L/L_{\odot} = 5.87^{+0.25}_{-0.13}$ and $R/R_{\odot} = 16.3^{+5.4}_{-2.3}$. This value is compatible with the corrected angular diameter ($\alpha =$ $4''.0\,10^{-4}$) from Kudritzki et al. (1983) which gives $R/R_{\odot} = 18.4^{+5.2}_{-3.3}$. The Lyman continuum flux of ζ Pup derived from recent models is $\log Q_0 = 49.7$ photon s^{-1} (Bohannan et al. 1990; Schaerer & Schmutz 1994).

2.2. γ^2 Vel

Interestingly, γ^2 Vel shows a significantly larger parallax than expected: $\pi = 3.88 \pm 0.53$ mas, corresponding to $d = 258^{+41}_{-31}$ pc. The HIPPARCOS parallax could be influenced by the orbital motion. By adopting the orbital orientation from (1986), the inclination from St.-Moffat et al. Louis et al. (1987), the WR/O star luminosity ratio from Conti & Smith (1972), and the angular size of the semi-major axis from Hanbury Brown et al. (1970) we find that the center of light is displaced by 0.3 to 1.2 mas from its average location. An interpretation of the orbital motion as a standard deviation yields 0.8 mas. For smaller magnitude differences (cf. below) or the orbital parameters of Pike et al. (1983) this error

becomes smaller. The contribution of the orbital motion to the error of the HIPPARCOS parallax is $0.8/\sqrt{N} = 0.09$ mas, where N=78 is the number of measurements used minus the number of fitted parameters. The measured parallax of the spectroscopic binary system γ^2 Vel provides therefore an accurate distance indicator. In the following we shall adopt the internal HIPPAR-COS error as the uncertainty on the parallax.

Combining the HIPPARCOS parallax with the observed angular size of the semi-major axis from the interferometric measurements of Hanbury Brown et al. (1970) and the known orbital period (cf. Niemela & Sahade 1980, Pike et al. 1983) allows to derive the total mass of the system: $M(WR + O) = 29.5 \pm 15.9 M_{\odot}$ (see Table 3 for parameters).

In the recent literature, there are two different amplitudes of the orbital motion. Pike et al. (1983) derived a mass ratio of $q = 0.36 \pm 0.09$, whereas Moffat et al. (1986) find a significantly larger mass ratio of $q = 0.54 \pm 0.03$, due to a larger radial velocity amplitude for the O star. Adopting an inclination of $i = 70^{\circ}$ later confirmed by St.-Louis et al. (1987), Pike et al. obtain $M(WR) = 7.7 \pm 2.5 M_{\odot}$ and M(O) = $22 \pm 5M_{\odot}$, while the Moffat et al. data yields $M(WR) = 19^{+7}_{-2}M_{\odot}$ and $M(O) = 35^{+13}_{-3}M_{\odot}$ (see Smith & Maeder 1989). The total mass derived above using the HIPPARCOS parallax favors the solution of Pike et al. (1983). On the other hand for the orbital parameters of Moffat et al. (1986) the observed angular size of the semi-major axis would imply a distance of $d = 308 \pm 37$ pc, which is within 1 σ of the HIPPARCOS distance. We conclude that the orbital parameters from both Pike et al. (1983) and Moffat et al. (1986) are compatible with the HIPPARCOS parallax and the interferometric determination of the semimajor axis from Hanbury Brown et al. (1970). The discrepancy between the two orbital solutions, however, implies a significant uncertainty for the masses of the γ^2 Vel components.

The HIPPARCOS distance measurement places

Table 1: Derived absolute magnitudes for the γ^2 Vel system for different assumptions on the WR-O magnitude difference (CS: Conti & Smith 1972, BC93: Brownsberger & Conti 1993). $M_v(\text{system}) = -5.49, M_V(\text{system}) = -5.39$, where v stands for the Smith and V the Johnson system

ID	$\Delta M_{\rm V}$	$M_{\rm v}({ m WR})$	$M_{\rm V}({\rm O})$
CS72	1.4	-3.83	-5.13
BC93	1.9	-3.42	-5.22
1	1.36	-3.86	-5.12
2	1.75	-3.54	-5.20

 γ^2 Vel considerably closer than the previously adopted distance ($d \sim 450 \text{ pc}$) based on its assumed association with Vela OB2 (e.g. Brandt et al. 1971, Sahu 1992). The apparent magnitude and extinction from the catalogue of van der Hucht et al. (1988), the flux zero-point correction of Schmutz & Vacca (1991) and the revised distance yields $M_v(\text{system}) = -5.49$. Absolute magnitudes obtained for the WR and O star components, assuming different luminosity ratios, are given in Table 1. The values of $\Delta M_V = 1.4$ mag and 1.9 mag result from the comparison of equivalent widths of γ^2 Vel with those from the WC8 star WR135 (Conti & Smith 1972; Brownsberger & Conti 1993). The absolute brightness of the WC8 star is considerably smaller than the average value for WC8 stars given by van der Hucht et al. (1988) ($< M_v(WC8) > = -4.8$).

The bolometric correction of the WC8 star can be obtained from evolutionary or atmosphere models. Adopting the mass-luminosity relation of Schaerer & Maeder (1992) or Smith et al. (1994) one has $\log L/L_{\odot} = 5.06$ (5.69) for the low (high) mass and hence a bolometric correction of BC=-4.2 to -3.7 (-5.8 to -5.3). Smith et al. derived a value of -4.5 for this subtype. From non-LTE atmosphere models (cf. below) we obtain BC=-3.5 to -3.4.

The brightness of the O star is typical for a 1996). Conti & Smith giant (cf. Vacca et al. (1972) concluded that the O star is a supergiant, based on the ratio 4089 Si IV/4143 He I. However, they note that the measurements have been difficult due to blending effects. The fast rotation of the O star makes it indeed difficult to locate a weak HeI line in a spectrum dominated by WR features and to isolate Si IV $\lambda 4089$ from a nearby nitrogen line. Therefore, the spectroscopic luminosity classification is not severely in conflict with the absolute magnitude of the O star. Comparing two spectra from Kaufer et al. (1997)observed at the phases $\phi = 0.24$ and 0.72 we find that HeII $\lambda 4686$ is in absorption. Its equivalent width corrected for the emission of the WR star is compatible with a classification as a giant (Mathys 1988), especially when considering the large uncertainty of measuring a broad absorption superimposed on a strong emission line. A statistical argument against a luminosity class I can be made based on the observed rotation of $v_{\rm rot} \sin i = 220 \text{ km s}^{-1}$ (Baade et al. 1990). For a supergiant such a velocity is extremely rare whereas for giants such high values are more ordinary (cf. Howarth et al. 1997). We determine a 4471 He I/4541 He II ratio of 1.6 ± 0.2 . According to Mathys (1988) this classifies the O star as O8 or O8.5. Conti & Smith (1972) have determined O9 spectral type. We agree in the equivalent width for the 4471 line, but in our spectra, the HeII λ 4541 absorption is much stronger than what is cited by Conti & Smith (1972). We therefore favor a re-classification of γ^2 Vel as WC8+O8III.

The radio mass loss rate of WR11 which was recently re-derived by Leitherer et al. (1997) has to be reduced by a factor of 2.3, which yields $\log \dot{M} = -4.55 \pm 0.16 \,\mathrm{M_{\odot}yr^{-1}}$. This value is more typical of \dot{M} of WC9 stars which seem to show lower mass loss rates than WR stars of other types. It may, however, be underestimated if the ionization in the outer parts of the winds is lower than assumed (Leitherer et al.). Interestingly \dot{M} is now in very good agreement with results from fits of ASCA X-ray observations with hydrodynamic wind-collision models (Stevens et al. 1996) allowing a distance independent determination of $\dot{M} \sim 3.10^{-5} \,\mathrm{M_{\odot}yr^{-1}}$. Agreement between both methods weakens the requirement of important structure ("clumping") in the wind (cf. Stevens et al. 1996), which, if present, leads to an overestimate of the radio mass loss rate.

Depending on the adopted WR mass, the mass dependent mass loss rate formula of Langer (1989) for WC stars yields values between log $\dot{M} = -4.78$ and $-3.80 \text{ M}_{\odot}\text{yr}^{-1}$. With the recently proposed relation of Schmutz (1997) the values are log $\dot{M} = -5.32$ and $-4.54 \text{ M}_{\odot}\text{yr}^{-1}$. A better constraint on M is clearly necessary for an accurate test of such a relation.

A 1.8 MeV γ -ray line emission of 2.3 to $5 \, 10^5$ cm⁻² s⁻¹ has been detected in the direction of the Vela SNR (Diehl et al. 1995). From the ²⁶Al yields of Meynet et al. (1997) the contribution of γ^2 Vel to the γ -ray flux is estimated between 1.5 and 2.6 10^{-5} with our new distance measurement. This values depends, however, quite strongly on the evolutionary scenario of the WR star, which is still uncertain (cf. §4).

3. Non-LTE model analysis for γ^2 Vel stars

From a line blanketed non-LTE analysis of the photospheric classification lines He I λ 4471 and He II λ 4541 of the O star we derive $T_{\rm eff}(O) =$ 33500 ± 1500 K and log $L(O)/L_{\odot} = 5.3 \pm 0.15$ for $M_V(O) = -5.13$. Its Lyman continuum flux is log(Q_0) = 48.7 photon s^{-1} . From these values we derive $M(O) = 29 \pm 4 \ M_{\odot}$ and an age of ~ 4±0.5 Myr from single star evolutionary models. (Meynet et al. 1994). With the age of the O star and the luminosity of the WC companion (cf. below) single star evolutionary models yield an estimate of $M_{\rm ini} \sim 57 \pm 15 M_{\odot}$ for the initial mass of the WC8 star.

A coarse analysis of the WC stellar parame-

Table 2: Stellar parameters for WR11 derivedfrom non-LTE atmosphere models

ID	T_{\star} [kK]	R_{\star} $[R_{\odot}]$	$\frac{\log \dot{M}}{[\rm M_{\odot}yr^{-1}]}$	$\frac{\log L}{[L_{\odot}]}$	BC [mag]	$\log Q_0 \\ \mathrm{s}^{-1}$
1	51.	3.3	-4.2	4.8	-3.5	48.5
2	61.	1.9	-4.3	4.7	-3.4	48.4

ters can be obtained from a model grid following the general procedure of Schmutz et al. (1989). The non-LTE models used for this purpose include He and C with an abundance of C/He=0.25 by number. The adopted terminal velocity is $v_{\infty} = 1450 \text{ km s}^{-1}$ (Eenens & Williams 1994). We use the observed equivalent widths of He I λ 5876 and He II λ 5412 (EW(5876)=9.5 ± 2.5 Å and EW(5412)= 4 ± 1 Å) and correct these values for the contamination by the light of the companion. The analysis then yields the temperature T_{\star} , radius R_{\star} , and \dot{M} . From the uncertainties of the EWs and the fit procedure we estimate an internal error of $\sim \pm 0.2$ dex on the luminosity L. In Table 2 we list the parameters derived for the different assumptions for the magnitude difference between the components (cf. Table 1). We find that the γ^2 Vel system provides approximately a factor of six less ionizing photons than ζ Pup.

We have used a simpler model for carbon than the more elaborate calculations by Koesterke & Hamann (1995). Test calculations with the parameters for the WC8 star WR135 derived by Koesterke & Hamann (1995) showed that our computations agree very accurately for the helium lines but we found less good agreement for the carbon lines. Therefore, we have used only the helium lines for the spectroscopic diagnostic. We have also performed a coarse analysis using the grid of Koesterke & Hamann (1995) for $\beta_C = 0.2$ (C/He= 0.083 by number). Although we get systematically lower temperatures we find that L is essentially identical. We therefore conclude that the luminosity obtained from

Table 3: Summary of parameters for γ^2 Vel. Column 3 gives alternate values for certain parameters

Quantity	value	alt. value	Ref.
π [mas]	3.88 ± 0.53		SSG
$d = 1/\pi$ [pc]	258^{+41}_{-31}		SSG
$a \; [mas]$	4.3 ± 0.5		HB70
P [days]	78.5002		NS80
	± 0.0001		
$M_{\rm tot}$	29.5 ± 15.9		SSG
$K(O) [\rm km s^{-1}]$	40.9 ± 6.5	70 ± 2	P83, M86
q	0.36 ± 0.09	0.54 ± 0.03	P83, M86
$i [^{\circ}]$	70 ± 10		ST87
e	0.38 ± 0.03	0.40	P83, M86
$M(WR) [M_{\odot}]$	7.7 ± 2.5	19^{+7}_{-2}	P83, SM89
$M(O) [M_{\odot}]$	22 ± 5	35^{+13}_{-3}	P83, SM89
$\log L(WR)_{M-L} [L_{\odot}]$	5.06	5.69	SSG
$v_{\infty}(WR) \ [km \ s^{-1}]$	1450		E94
$\log \dot{M}(WR) [M_{\odot} yr^{-1}]$	-4.55 ± 0.16		SSG
$\log L(WR)_{NLTE}[L_{\odot}]$	$\sim 4.7 - 4.8$		SSG
$T_{\rm eff}(O)$ [K]	34000 ± 1500		SSG
$\log L(O)_{\rm NLTE} [L_{\odot}]$	5.3 ± 0.15		SSG
$M(O)_{\text{evol}} [M_{\odot}]$	29 ± 4		SSG
age(O) [Myr]	$4.0^{+0.8}_{-0.5}$		SSG

HB70: Hanbury Brown et al. (1970)
NS80: Niemela & Sahade (1980)
M86: Moffat et al. (1986)
P83: Pike et al. (1983)
ST87: St.-Louis et al. (1987)
SM89: Smith & Maeder (1989)
E94: Eenens et al. (1994)
SSG: this paper

our model is consistent with the WC models from Koesterke & Hamann. The parameters T_{\star} and R_{\star} depend quite strongly on assumptions about the velocity law and are therefore less reliable (e.g. Schmutz 1996, 1997).

For the luminosity derived from the atmosphere models the M-L relation gives $M(WR) \sim 5 \pm 1.5 \ M_{\odot}$, which is compatible with the low mass determination for WR11. On the other hand, if the mass is of the order of 19 M_{\odot} (as mostly adopted in the literature) we are faced with a serious discrepancy (up to an order of magnitude !) between L derived from atmosphere and evolutionary models. The work of Howarth & Schmutz (1992) and Schmutz (1996, 1997) previously revealed differences for some WN stars. According to Schmutz (1996, 1997) this is most likely due to systematic errors in the non-LTE model atmosphere, and it may lead to underestimates of L by typically 50 to 300 %. We conclude that for the low mass determination of WR11 the theoretical M-L relation for hydrogen-free WR stars agrees within the errors with L derived from atmosphere models.

4. Discussion

The distance and the origin of the Gum nebula, the spatial distribution of important objects in this region (Gum nebula, Vela R2 and Vela OB associations, ζ Pup , γ^2 Vel , the Vela SNR etc.) and their possible relation are still relatively poorly known and controversial (cf. e.g. Brandt 1971, Bruhweiler et al. 1983, Franco et al. 1990, Sahu 1992, Sahu & Blaauw 1993, Oberlack et al. 1994, Fitzpatrick & Spitzer 1994, Aschenbach et al. 1995). Combining recent data (including additional HIPPARCOS parallaxes and proper motions) with the most elaborate study of this region done by Sahu (1992) may lead to a refined picture. While such a task is clearly beyond the scope of the present paper we shall in the following briefly discuss a few questions related to ζ Pup and γ^2 Vel.

Based on its proper motion Sahu (1992) suggested that the runaway star ζ Pup originates from the more distant young Vela R2 association. Although the HIPPARCOS distance to ζ Pup is lower than the ~ 730 pc suggested by Sahu, her scenario remains valid provided the distance to Vela R2 is less than the adopted 800 pc⁵, which is compatible with the distance estimate of Liseau et al. (1992) and the HIPPARCOS parallax of two presumed members (HD 76534: 2.43 ± 1.30, HD 76838: 3.39 ± 0.90 mas). Evolutionary scenarios for ζ Pup have been presented by van Rensbergen et al. (1996). It would be highly desirable to include the effects of the rapid rotation of ζ Pup in future calculations.

We obtained additional HIPPARCOS parallaxes for 5 presumed Vela OB2 members (HD 63922: 1.66 \pm 0.53, HD 64740: 4.53 \pm 0.52, HD 68657: 4.48 ± 0.49 , HD 70930: 2.16 ± 0.57 , HD 72485: 3.10 ± 0.52 , all values in mas) for which Sahu (1992) derived proper motions and statistical parallaxes. With the exception of HD 68657 (and γ^2 Vel) her values are within 1 σ of the HIPPARCOS measurements. The unexpectedly large annual parallax of γ^2 Vel lies thus well within the range covered by other stars which, as shown by Sahu (1992), are likely members of the association Vela OB2. On the basis of our limited data (no new proper motions) there is no evidence against the membership of γ^2 Vel to this group of objects. Given the very small number of objects and the apparently large distance spread determined from the annual parallaxes, this association seems, however, to be quite loosely defined. Despite these uncertainties an age of ~ 20 to 30 Myr (Eggen 1980, Sahu 1992) is usually estimated. For the objects mentioned above we obtain individual ages between $\sim 6-30$ Myr from Geneva photometry and the HIPPARCOS data. For the majority we favor ~ 10 Myr, which is also supported by recent kinematic observations of the Vela Shell (Churchwell et al. 1996).

Single star evolution models fail to explain the formation of the γ^2 Vel system for ages $\gtrsim 6$ Myr (cf. Schild & Maeder 1984). A young age is, however, compatible with our estimate for the O star. An older age of ~ 10 Myr as typical for other Vela OB2 stars would exclude a large total mass of the system (~ 50-60 M_{\odot}). On the other hand $M_{\rm tot} \sim 30 M_{\odot}$ might be compatible with 10 Myr if WR11 evolved from a intermediate-mass progenitor through mass transfer in the binary system. However, the relatively large eccentricity of γ^2 Vel may speak against such a scenario.

 $^{^5 \}mathrm{Adopting}$ our one sigma error on d of ζ Pup the distance of Vela R2 must be $\lesssim 580$ pc.

Regrettably, the present data does not allow to draw more definite conclusions about the formation of the γ^2 Vel system.

We thank Drs. A. Kaufer, J. Schweickhardt, O. Stahl, and B. Wolf for the spectra of γ^2 Vel obtained with HEROS at the ESO 0.5m telescope, and Dr. V. Niemela for discussions on spectral classification. DS wishes to thank Georges Meynet, Jean-Claude Mermilliod, Claus Leitherer and Jürgen Knödlseder for useful discussions. Travel support from the Geneva Observatory and an invitation from the Université Paul-Sabatier in Toulouse, where part of this work was carried out, are kindly acknowledged. DS receives a fellowship from the Swiss National Foundation of Scientific Research and partial support from the Directors Discretionary Research Fund of the STScI.

REFERENCES

- Aschenbach B., Egger R., Trümper J., 1995, Nature 373, 587
- Baade D., Schmutz W., van Kerkwijk M., 1990, A&A 240, 105
- Bohannan B., Voels S.A., Hummer D.G., Abbott D.C., 1990, ApJ 365, 729
- Brandt J.C., Stecher T.P., Crawford D.L., Maran S.P., 1971, ApJ 163, L99
- Bruhweiler F.C., Kafatos M., Brandt J.C., 1983, Comments Astroph., 10, 1
- Brownsberger K.R., Conti P.S., 1993, in: Massive Stars: Their Lives in the Interstellar Medium, ASP Conf. Series, Vol. 35, p. 275
- Churchwell E., Winnberg A., Cardell J., Cooper G., Suntzeff N.B., 1996, ApJ 469, 209
- Conti P., Smith L.F., 1972, ApJ 172, 623
- Diehl R., et al., 1995, A&A 298, L25
- Eenens P.R.J., Williams P.M., 1994, MNRAS 269, 1082

Eggen O.J., 1980, ApJ 238, 627

- Fitzpatrick E.L., Spitzer L. Jr., 1994, ApJ 427, 232
- Franco G.A.P., 1990, A&A 227, 499
- Gabler R., Gabler A., Kudritzki R.P., Puls J., Pauldrach A., 1989, A&A 226, 162
- Hanbury Brown R., Davis J., Herbinson-Evans D., Allen L.R., 1970, MNRAS 148, 103
- Howarth I.D., Schmutz W., 1992, A&A 261, 503
- Howarth I.D., Siebert K.W., Hussain G.A.J., Prinja R.K., 1997, MNRAS 284, 265
- van der Hucht K., Hidayat B., Admiranto A.G., Supelli K.R., Doom C., 1988, A&A 199, 217
- Kaufer A., Schweickhardt J., Stahl O., Wolf B., 1997, personal communication
- Koesterke L., Hamann W.-R., 1995, A&A 299, 503
- Kudritzki R.P., Simon K.P., Hamann W.-R., 1983, A&A 118, 245
- Lamers H.J.G.L.M., Leitherer C., 1993, ApJ 412, 771

Langer N., 1989, A&A 220, 135

- Leitherer C., Chapman J.M., Koribalski B., 1997, ApJ , in press
- Liseau R., Lorenzetti D., Nisini B., Spinoglio L., Monetti A., 1992, A&A 265, 577

Mathys G., 1988, A&AS 76, 427

- Meynet G., Arnould M., Prantzos N., Paulus G., 1997, A&A in press
- Meynet G., Maeder A., Schaller G., Schaerer D., Charbonnel C., 1994, A&AS 103, 97
- Moffat A.F.M., Vogt N., Paquin G., Lamontagne R., Barrera L.H., 1986, AJ 91, 1386
- Niemela V.S., Sahade J., 1980, ApJ 238, 244
- Oberlack U., Diehl R., Montmerle T., Prantzos N., von Ballmoos P., 1994, ApJS 92, 433
- Pauldrach A.W.A, Kudritzki R.P., Puls J., Butler K., Hunsinger J., 1994, A&A 283, 525

- Pike C.D., Stickland D.J., Willis A., 1983, The Observatory 103, 154
- Puls J., Kudritzki R.P., Herrero A., Pauldrach A.W.A., Haser S.M., Lennon D.J., Gabler R., Voels S.A., Vilchez J.M., Wachter S., Feldmeier A., 1995, A&A 305, 171
- van Rensbergen W., Vanbeveren D., de Loore C., 1996, A&A 305, 825
- Sahu M., 1992, PhD thesis, University of Groningen, The Netherlands
- Sahu M., Blaauw A., 1993, in: Massive Stars: Their Lives in the Interstellar Medium, ASP Conf. Series, Vol. 35, p. 278
- Schaerer D., Maeder A., 1992, A&A 263, 129
- Schaerer D., Schmutz W., 1994, A&A 288, 231

Schild H., Maeder A., 1984, A&A 136, 237

- Schmutz W., 1996, in: Wolf-Rayet Stars in the Framework of Stellar Evolution, 33rd Liège International Astrophys. Coll., Eds. J.M. Vreux et al., p. 553
- Schmutz W., 1997, A&A, in press
- Schmutz W., Hamann W.-R., Wessolowski U., 1989, A&A 210, 236
- Schmutz W., Vacca W.D.W., 1991, A&AS 89, 259
- Smith L.F., Maeder A., 1989, A&A 211, 71
- Smith L.F., Meynet G., Mermilliod J.-C., 1994, A&A 287, 835
- Stevens I.R., Corcoran M.F., Willis A.J., Skinner S.L., Pollock A.M.T., Nagase F., Koyama K., 1996, MNRAS 283, 589
- St.-Louis N., Drissen L., Moffat A.F.J., Bastien P., Tapia S., 1987, ApJ 322, 870
- Vacca W.D.W., Garmany C.D., Shull J.M., 1996, ApJ 460, 914

This 2-column preprint was prepared with the AAS ${\rm L\!AT}_{\rm E\!X}$ macros v4.0.