Ly α Clouds Associated with Pop III Objects in Cold Dark Matter Models

Benedetta Ciardi¹, Andrea Ferrara²

¹Dipartimento di Astronomia, Università di Firenze, 50125 Firenze, Italy E-mail: ciardi@arcetri.astro.it ²Osservatorio Astrofisico di Arcetri 50125 Firenze, Italy E-mail: ferrara@arcetri.astro.it

Received _

accepted.

ABSTRACT

We present a semi-analytical model to test the hypothesis that Ly α QSO absorption lines originate in gaseous halos produced by multiple supernova explosions occurring in Pop III objects in a CDM cosmological scenario hoping to assess the validity of CDM models and/or constrain their parameters. The preliminary results indicate that the range of N_{HI} , redshift distribution and metallicity of clouds are well reproduced by CDM if they are associated with galaxy halos or groups. Firm conclusions on clouds with $N_{HI} \leq 10^{14}$ cm⁻² need a more refined study.

Subject headings: Cosmology: theory – dark matter –quasars: absorption lines

1. Introduction

One of the most challenging current attempts in cosmology is to relate the origin and evolution of large-scale structure to the observed properties of QSO absorption-line systems. Several groups have now performed sophisticated hydro simulations of various hierarchical models (CDM, CDM+ Λ , HCDM) to clarify the above issues. In spite of the great advances provided by numerical approaches, their results are still not conclusive (and often hard even to compare) since the physics included is necessarily limited by computational economy. In particular, the study of effects as feedback and radiative transfer, whose importance, particularly for small objects – such as, presumably, the very first generation of galaxies (Pop III) – can hardly be overlooked, is still in its youth (Gnedin & Ostriker 1997). These facts motivate the effort to develop versatile semi-analytical models, that could co-operatively complement numerical studies. Here we present one such whose aim is to test the hypothesis that Ly α QSO absorption lines originate in gaseous halos produced by multiple supernova explosions occurring in Pop III objects in a CDM cosmological scenario.

2. Dark halos in CDM models

We assume a CDM power spectrum of fluctuations as in Efstathiou et al. (1992),

$$|\delta_k|^2 = \frac{Ak^n}{\left\{1 + \left[Bk + (Ck)^{3/2} + (Dk)^2\right]^\nu\right\}^{2/\nu}},\tag{1}$$

where $\nu=1.13$; $B=6.4 \ h^{-2}$ Mpc; $C=3.0 \ h^{-2}$ Mpc; $D=1.7 \ h^{-2}$ Mpc; n=1. The previous spectrum describes a "standard" CDM model with $\Omega_0 = 1$ and h = 0.5, our adopted values; we also take $\Omega_b = 0.19$ (from Coma cluster data, White & Frenk 1991). The power spectrum normalization has been determined from *COBE*, yielding $\sigma_8 = 1.07$ (Efstathiou *et al.* 1992). To obtain the number density of dark matter (DM) halos, n, as a function of redshift, z, and of their circular velocity, $v_c = [GM(r)/r]^{1/2}$, we use the Press – Schechter formalism (Press & Schechter 1974, PS) following White & Frenk (1991):

$$n(v_c, z) = -\frac{3\delta_c(1+z)}{(2\pi r_0^2)^{3/2} \Delta(r_0)} \frac{d\ln\Delta(r_0)}{d\ln r_0} \exp\left[-\frac{\delta_c^2(1+z)^2}{2\Delta^2(r_0)}\right];$$
(2)

 $\delta_c = 1.68$ is a critical value of the overdensity and $\Delta^2(r_0)$ is the variance of fluctuations on scale r_0 :

$$\Delta^2(r_0) = \frac{1}{2\pi^2} \int_0^\infty dk \ k^2 |\delta_k|^2 W^2(kr_0).$$
(3)

We used three different window functions $W(kr_0)$: (i) gaussian, (ii) sharp k-space, (iii) top-hat. The results are very weakly dependent on the specific choice of $W(kr_0)$; we adopt a top-hat window function since it gives the best agreement with numerical simulations for $\delta_c = 1.68$ (Lacey & Cole 1994).

3. Evolution of the baryonic component

As DM halos are formed, baryons settle dissipatively into their potential well, being heated by virialization shocks. Both baryonic evolution and galaxy formation can be influenced by another important heating/ionization source, the ultraviolet background radiation field (UVB); its potential importance has been recently discussed by Efstathiou (1992) (see also Thoul & Weinberg 1996), who emphasized its role in delaying or even suppressing the formation of very low v_c objects.

3.1. Reionization epoch and UVB

In the absence of ionizing radiation, baryons can collapse freely in the gravitational potential of the DM, allowing for the formation of the first proto-galaxies. These objects, which we will refer to collectively as Pop III, should have masses $M_P \gtrsim 10^6 M_{\odot}$ in order to be able to cool in a Hubble time (Tegmark *et al.* 1997). According to PS, the formation peak for these objects is at $z_f \sim 30$. Assuming an efficiency of UV photon production per collapsed proton ~ 190 (Tegmark *et al.* 1994), and an average life time of massive stars ~ 10^7 yr, the ionizing photon rate per galaxy is ~ 7×10^{50} s⁻¹. These photons will produce expanding HII regions in the surrounding IGM, eventually overlapping. To determine the epoch of complete reionization we calculate the IGM porosity parameter, Q, defined by $dQ(z) = |dn(M_P, z)/dz|r_i(z)^3 dz$, with $r_i \sim (1 + z)^{-3/2}$ Mpc being the radius of the HII region. Complete overlap (*i.e.* Q = 0.16, Smith 1976) occurs at $z_{ion} \sim 7$; this value depends on M_P ($z_{ion} \sim 4.5[8]$ for $M_P = 5 \times 10^6 [5 \, 10^5] M_{\odot}$). For $z < z_{ion}$ the UVB is approximated as a power law spectrum with index α and intensity at the Lyman limit (Haardt & Madau 1996, Ferrara & Giallongo 1996 [FG]):

$$J_{-21}(z) = 0.53 \begin{cases} 0 & z > 7, \\ \exp[-0.69(z-3)] & 3 < z \le 7, \\ (1+z)^{1/4} & 2 \le z \le 3, \\ (1+z)^3 & z < 2. \end{cases}$$
(4)

where $J_{-21} = J_0/10^{-21} \text{ergs s}^{-1} \text{ cm}^{-2} \text{ Hz}^{-1} \text{sr}^{-1}$.

3.2. Galaxy formation

As the UVB heats the IGM, it can prevent the gas from collapsing and forming proto-galaxies if the IGM temperature, T, is higher than the virial temperature, $T_c \propto v_c^2$, of the corresponding DM halo. We have derived the temperature and ionization evolution of the IGM (assumed of primordial chemical composition) from the standard time-dependent energy and ionization equations. The cooling function is described in FG; we have studied two cases corresponding to values of $\alpha = 1.5(5)$ mimicking a hard, AGN-dominated (soft, galaxy-dominated) spectrum. We have also explored different initial values $T_i = T(z = 7)$ finding that this parameter has essentially no influence on the results, as already stressed by FG: we take therefore $T_i = 10^3$ K. The condition $T < T_c$ defines a minimum circular velocity, $v_{c,min}(z)$, (or equivalently a minimum mass) for collapse, shown in Fig. 1. Galaxies with *total* mass below $1 - 4 \times 10^7 M_{\odot}$ are prevented from collapsing after the onset of the UVB; of course $v_{c,min} = v_{c,min}(M_P) \sim 1.3(M_P/10^6 M_{\odot})^{1/3}$ km s⁻¹ for z > 7.

3.3. Blow away and secondary halos

As the gas collapses towards the center of the proto-galaxy, the density enhancement is very likely to trigger star formation (SF). Star forming regions will inject energy into the galactic ISM, largely in the form of supernova explosions: for a Miller-Scalo IMF, one supernova is expected for each 56 $M_{\odot} = \nu^{-1}$ of stars formed. Type II SNe tend to occur in associations, generating very energetic multiple explosions whose observational signature is an expanding supershell. If the galaxy mass is low, its entire gas content could be swept by the shell, and driven into the surrounding IGM, in a process called *blow-away*. The expanding shell will be eventually confined by the IGM pressure, provided it is high enough. The blow-away condition essentially requires that $v_s > \mathcal{A}(\Omega_0, \Omega_b, g)v_e$, where g is a geometrical factor (details in Ferrara & Tolstoy 1997). The previous condition determines the maximum circular velocity for which the blow-away can occur, $v_{c,max} = 16(1+z)^{1/2}$ km s⁻¹ for our parameters, also shown in Fig. 1; more massive objects will escape disruption. Hence, only objects with $v_{c,min} \leq v_c \leq v_{c,max}$ will be able to generate a secondary halo, via a shell *re*-expansion, as explained below. Since the gas is lost, further SF is inhibited, and the remnant is a faint agglomerate of low mass stars produced in the first (and last) SF episode.

Next we derive the evolution of the (spherical) supershell radius, R_s , in the thin-shell approximation (Kompaneets 1957). The unperturbed external medium has density $\rho_e(r) \propto r^{-2}$ for $r \leq r_{200}$, corresponding to the galaxy ISM, and $\rho_e(r) = \rho(z)$ for $r > r_{200}$, the IGM density. The shell is driven by the pressure, P_b , of the hot gas in the bubble cooling at a rate $\Lambda(T_b)$, and subject to the gravitational pull of the DM halo (we neglect self-gravity of baryons). The SN mechanical luminosity is $L_{SN} \propto \gamma = \nu \dot{M}_* \sim (\nu M_{gas}/\tau t_{ff}) = 1.7 \times 10^{39} (v_c/30 \text{ km s}^{-1})^3 \text{ ergs s}^{-1}$, where γ is the SN rate, $\tau = 0.6\%$ is the SF efficiency and t_{ff} is the free-fall time of the gas. The evolution of R_s is governed by energy and momentum conservation equations (subscripts *b* and *s* refer to bubble and shell, respectively):

$$\frac{dE_b}{dt} = L_{SN} - 3V_b P_b \frac{\dot{R}_s}{R_s} - n_b^2 V_b \Lambda(T_b),\tag{5}$$

$$\frac{d}{dt}\left(V_b\rho_e \dot{R}_s\right) = \frac{3V_b}{R_s}(P_b - P_e) - \frac{GM(R_s)}{R_s^2}V_b\rho_e,\tag{6}$$

where $V_b = (4\pi/3)R_s^3$. Fig. 2 shows the behavior of R_s as a function of z for different values of v_c and corresponding DM halo formation redshift z_f ; the evolution is followed up to the stagnation point, *i.e.* $\dot{R}_s \simeq$ sound speed in the IGM. Secondary halos associated with DM halos formed at z > 7 are larger, since their expansion is unimpeded until the reionization epoch due to the low IGM pressure; the final size increases with v_c , since $L_{SN} \propto v_c^3$. The final value of R_s depends on P_e : sizes of the order of tens of kpc are obtained setting $P_e \sim 10^3 \times$ the IGM pressure, or $P_e/k_B \sim 7$ K cm⁻³ at z = 0. This value is typical of the environments of spiral galaxies and groups: for the Galactic halo at r = 30 kpc, $P_e/k_B \sim 12$ K cm⁻³ (Wolfire *et al.* 1995). Lower values of P_e result in larger sizes and lower densities.

Once the stalling point is reached, as the shell pressure is higher than P_b , the inner boundary of the shell re-expands back into the cavity. The expansion ends when pressure equilibrium across the contact discontinuity has been achieved: from this condition we derive the shell thickness, S_s . We find that the ratio $S_s/R_s > 1/2$ in all cases and it often exceeds 0.8. In addition, the re-expansion factor can be enhanced by a magnetic field (Slavin & Cox 1992) and by accounting for cooling of hot bubble gas. For this reason, we assume that the volume inside R_s is uniformly filled with blown-away baryons with density $n_H = 3\Omega_b M/4\pi (1 - \Omega_b)R_s^3$.

3.4. Ly α clouds

We are now able to test the main hypothesis of this work, *i.e.* that the secondary baryonic halos so found can be responsible for Ly α absorption lines. As a preliminary test (a complete study is devoted to a forthcoming paper) we have derived the neutral hydrogen column density of the clouds and their redshift distribution. Assuming a value for the secondary halo gas (or the Ly α clouds, in our view) temperature of $\sim 4 \times 10^4$ K, consistent with a purely thermal Doppler parameter $b \sim 25$ km s⁻¹ (Lu *et al.* 1996), we have determined the HI density, n_{HI} , from ionization equilibrium, Integrating along a line of sight at impact parameter p, we find hydrogen column densities $10^{12} < N_{HI} < 5 \times 10^{18} \text{cm}^{-2}$, as shown in Fig. 3, with the highest N_{HI} originating in low z_f halos. Column densities $\leq 10^{14} \text{ cm}^{-2}$ are associated with secondary halos formed at very high redshift, $z_f \sim 20$, where merging – that we have neglected – should be rather important. The redshift distribution of the clouds,

$$\frac{dN}{dz}(z, N_{HI}) = \frac{c}{H_0} (1+z)^{1/2} \int_{v_{c,min}}^{v_{c,max}} dv_c \ n(v_c, z) v_c^{-1} \pi p^2(v_c, N_{HI}), \tag{7}$$

can be very nicely fit by a power law $dN/dz = A_0(1+z)^{\gamma}$ with $\gamma = 2.2$ and $A_0=0.7$, in satisfactory agreement with observational data (Bechtold 1994, Giallongo *et al.* 1996) at this stage.

4. Conclusions and discussion

We have brought arguments suggesting that at least some Ly α QSO absorption lines could originate in secondary halos produced by multiple supernova explosions occurring in Pop III objects. As a first step, we have investigated only a specific CDM model; nevertheless our aim is to extend the analysis to a larger class of theoretical models (CDM+ Λ , HCDM) In this paper we have introduced the main features of our semi-analytical model, deferring a complete exploration of the results, cosmological models, and model details to a forthcoming paper. Yet, two conclusions could be relevant: (i) Ly α clouds should be preferentially found in relatively high pressure regions, as galaxy halos and groups; (ii) HI column densities and redshift distribution are well reproduced by a CDM model if clouds are associated with galaxy halos or groups. We reiterate that predictions for $N_{HI} \lesssim 10^{14}$ cm⁻² might be affected by the simplifications adopted (*i.e.* no merging). In view of the exploration of additional cosmological models, it is important to compare the model predictions on other observables as the dN/dN_{HI} and b distributions, and clustering properties. The metallicity $Z/Z_{\odot} \lesssim 10^{-2}$ found in clouds with $N_{HI} > 3 \times 10^{14}$ cm⁻² by Cowie *et al.* (1995) and Tytler *et al.* (1995) could be explained by primordial enrichment by Pop III objects. A back-on-the envelope calculation using the SN rates from our model and assuming a metal yield of ~ $1M_{\odot}$ per SN, gives $Z/Z_{\odot} \sim 1.2 \times 10^{-4} (v_c/10 \text{ km s}^{-1})^2$ in the secondary halo.

REFERENCES

- Bechtold, J. 1994, ApJS, 91, 1
- Cowie, L.L., Songaila, A., Kim., T. & Hu, E.M. 1995, AJ, 109, 1522
- Efstathiou, G. 1992, MNRAS, 256, 43
- Efstathiou, G., Bond, J.R. & White, S.D.M. 1992, MNRAS, 258, 1P
- Ferrara, A. & Giallongo, E. 1996, MNRAS, 282, 116
- Ferrara, A. & Tolstoy, E. 1997, in preparation
- Giallongo, E. et al., 1996, ApJ, 466, 46
- Gnedin, O.Y. & Ostriker, J.P. 1997, ApJ, 474, 223
- Haardt, F. & Madau, P. 1996, ApJ, 461, 20
- Kompaneets, A.S. 1957, Soviet Phys., 4, 730
- Lacey, C. & Cole, S. 1994, MNRAS, 271, 676
- Lu, L., Sargent, W.L.W., Womble, D.S. & Takada-Hidai, 1996, ApJ, 472, L564
- Press, W.H. & Schechter, P. 1974, ApJ, 187, 425, PS
- Slavin, J.D. & Cox, D.P. 1992, ApJ, 392, 131
- Smith, B.W. 1976, ApJ, 211, 404
- Tegmark, M., Silk, J. & Blanchard, A. 1994, ApJ, 420, 484
- Tegmark, M., Silk, J., Rees, M.J., Blanchard, A., Abel, T. & Palla, F. 1997, ApJ, 474, 1
- Thoul, A.A. & Weinberg, D.H. 1996, ApJ, 465, 608
- Tytler, D., Fan, X.M., Burles, S., Cottrell, L., Davis, C., Kirkman, D. & Zuo, L. 1995, in QSOs Absorption Lines, Proc. ESO Workshop, ed. Meylan, G. (Heidelberg: Springer)
- White, S.D.M. & Frenk, C.S. 1991, ApJ, 379, 52
- Wolfire, M.G., McKee, C.F., Hollenback, D. & Tielens, A.G.G.M. 1995, ApJ, 453, 673

This manuscript was prepared with the AAS IAT_EX macros v4.0.

Fig. 1.— Minimum circular velocity (solid lines) of a DM halo for collapse of the gas as a function of redshift for two values of the UVB power-law spectral index $\alpha = 1.5, 5$. Also shown (dashed line) is the maximum circular velocity required for blow-away to occur. The initial IGM temperature is $T_i = 10^3$ K.

Fig. 2.— Behavior of shell radius, R_s , as a function of z for different values of v_c (from top to bottom) and corresponding DM halo formation redshift z_f ; the evolution is followed up to the stagnation point.

Fig. 3.— N_{HI} of Ly α clouds as a function of v_c for different values of the DM halo formation redshift z_f .





