# Three dimensional, axisymmetric cusps without chaos

S. Sridhar<sup>\*</sup> and J. Touma<sup>†</sup>

\* Inter-University Centre for Astronomy and Astrophysics, Ganeshkhind, Pune 411 007, INDIA

 $^\dagger$  University of Texas, McDonald Observatory, RLM 15.308, Austin, Texas, 78712

IUCAA Preprint 28/97, March 1997

## ABSTRACT

We construct three dimensional axisymmetric, cuspy density distributions, whose potentials are of Stäckel form in parabolic coordinates. As in Sridhar and Touma (1997), a black hole of arbitrary mass may be added at the centre, without destroying the Stäckel form of the potentials. The construction uses a classic method, originally due to Kuzmin (1956), which is here extended to parabolic coordinates. The models are highly oblate, and the cusps are "weak", with the density,  $\rho \propto 1/r^k$ , where 0 < k < 1.

**Key words:** galaxies: elliptical and lenticular, cD—galaxies: kinematic and dynamics—galaxies: structure

## 1 INTRODUCTION

Kuzmin(1956) showed that the density at a general point of an oblate, axisymmetric mass model, whose gravitational potential is of Stäckel form in elliptic coordinates, is simply related to the density on the axis of symmetry. Later, Kuzmin (1973) generalized the theorem to the fully triaxial case. A corollary of Kuzmin's theorem is that the density so constructed is automatically non-negative everywhere as long as it is non-negative on the short axis. The interest in Stäckel models arises from the fact that the motion of point masses in such potentials is completely integrable by the Hamilton-Jacobi method; these potentials are also termed "separable", since the dynamics in three dimensions separates into three independent oscillations (c.f. de Zeeuw 1985a,b for a clear discussion of orbits, and Kuzmin's theorem). Constructing distribution functions for models of galaxies in a steady state (c.f. Binney & Tremaine 1987) is facilitated by the existence of three isolating integrals for three dimensional models. Recently, Sridhar and Touma (1997)—hereafter referred to as ST—presented a family of cuspy, scale-free mass models of non-axisymmetric discs whose potentials are of Stäckel form in parabolic coordinates. The surface density behaves like  $\Sigma \propto 1/r^{\gamma}$ , where  $0 < \gamma < 1$ . Here we generalize Kuzmin (1956) to parabolic coordinates and naturally extend the non-axisymmetric discs of ST to three dimensional, axisymmetric, cuspy density distributions, whose potentials are of Stäckel form in parabolic coordinates. Our main result follows in § 2. Scale–free cusps are presented in § 3, and a complete classification of orbit families is given in § 4. A discussion of the results, in  $\S$  5, completes this paper on mass models.

# 2 KUZMIN'S CONSTRUCTION APPLIED TO PARABOLIC COORDINATES

For axisymmetric configurations, the mass density  $\rho$ , and the gravitational potential U, that it generates are both functions only of  $R = \sqrt{x^2 + y^2}$  and z, the cylindrical radius and the coordinate along the axis of symmetry, respectively. Since the potential is independent of the azimuthal angle ( $\phi$ ), the component of the angular momentum parallel to the z-axis,  $L_z$ , is conserved. Hence the dynamics may be described as two-dimensional motion in a meridional ( $\phi$  = constant) plane, under the influence of an effective potential defined by

$$U_{\rm eff} = U(R, z) + \frac{L_z^2}{2R^2}.$$
 (1)

We study cases when the motion in the meridional plane is separable in parabolic coordinates,  $(\xi, \eta)$ . These may be defined as the roots for  $\tau$  of the equation

$$\tau^2 - 2z\tau - R^2 = 0. \tag{2}$$

Solving this quadratic equation gives

$$\xi = z + \sqrt{z^2 + R^2} 
\eta = z - \sqrt{z^2 + R^2},$$
(3)

where we have chosen  $\eta \leq 0 \leq \xi$ . The inverse relations are

$$R^{2} = -\xi\eta, \qquad z = \frac{1}{2}(\xi + \eta), \qquad r = \frac{\xi - \eta}{2},$$
 (4)

<sup>\*</sup> E-mail: sridhar@iucaa.ernet.in

<sup>&</sup>lt;sup>†</sup> E-mail: touma@harlan.as.utexas.edu

where  $r = \sqrt{R^2 + z^2}$  is the spherical radius. Surfaces of constant  $\xi$  and  $\eta$  are paraboloids of revolution which cut the z-axis at  $\xi/2$  and  $\eta/2$ , and the x-y plane at  $R = \xi$  and  $R = \eta$ respectively. These surfaces are orthogonal to the meridional plane and mutually orthogonal as well; thus  $(\xi, \eta, \phi)$  form an orthogonal coordinate system. Positive and negative ranges of the z-axis are covered by  $\xi$  and  $\eta$  respectively:

$$\xi = 0, \quad z = 2\eta, \text{ for } z \le 0$$
  
 $\eta = 0, \quad z = 2\xi, \text{ for } z \ge 0.$  (5)

The most general axisymmetric potential in parabolic coordinates, for which the Hamilton–Jacobi equation separates, has the Stäckel form, given by

$$U(\xi,\eta) = \frac{F_{+}(\xi)}{\xi - \eta} + \frac{F_{-}(\eta)}{\eta - \xi}.$$
(6)

If  $U(\xi,\eta)$  is of Stäckel form, so is the effective potential,  $U_{\text{eff}}$ , which is defined in equation (1).<sup>‡</sup> The potential due to a point mass is obtained by choosing  $F_- - F_+ = 2GM$ . It is important to note that, to describe an interesting case like the Kepler problem,  $F_+ \neq F_-$ ; in general, we let  $F_+$  and  $F_$ be quite different functions of their arguments. The density corresponding to the point mass assumes the form,

$$\rho_{\bullet} \equiv M\delta(\mathbf{r}) = M \left\{ \frac{2\delta(\xi)\delta(\eta)}{\pi(\xi - \eta)} \right\}$$
(7)

Let  $\rho$  and U be the density and potential, respectively, of a smooth distribution of matter<sup>§</sup>. If we now include a point mass at the origin,

$$\rho_{tot} = \rho + \rho_{\bullet} 
U_{tot} = U - \frac{2GM}{\xi - \eta},$$
(8)

are the total density and potential. These are related to each other through Poisson's equation, or equivalently

$$\nabla^2 U = 4\pi G\rho,\tag{9}$$

where the contribution of the point mass has been dropped from both sides. Expressed in parabolic coordinates, Poisson's equation for (the axisymmetric) potential and density reads

$$\frac{4}{\xi - \eta} \left[ \frac{\partial}{\partial \xi} \xi \frac{\partial}{\partial \xi} - \frac{\partial}{\partial \eta} \eta \frac{\partial}{\partial \eta} \right] U(\xi, \eta) = 4\pi G \rho(\xi, \eta).$$
(10)

We substitute the expression given in equation (6) for U, in equation (10). This results in the following expression for the smoothly distributed density of matter:

$$\pi G \rho(\xi, \eta) = \frac{\xi F_{+}^{''} + \eta F_{-}^{''}}{(\xi - \eta)^2} - \frac{\xi + \eta}{(\xi - \eta)^3} (F_{+}^{'} - F_{-}^{'}), \qquad (11)$$

where the primes denote differentiation with respect to the arguments,  $\xi$  or  $\eta$ , as the case may be. It proves convenient to employ the single variable  $\tau$  in place of the two variables

 $\xi$  and  $\eta.$  Then the density is completely determined by

$$F(\tau) = \begin{cases} F_{+}(\xi) & \text{if } \tau \ge 0\\ F_{-}(\eta) & \text{if } \tau \le 0 \end{cases},$$
(12)

which is evidently one function of one variable. It is also useful to define constants  $C_{\tau}$  and  $D_{\tau}$ , that depend only on the behaviour of F for small  $|\tau|$ :

$$C_{\tau} = \begin{cases} C_{+} \equiv \lim_{\eta \to 0} \eta F_{-}^{\prime} & \text{if } \tau \ge 0 \\ C_{-} \equiv \lim_{\xi \to 0} \xi F_{+}^{\prime} & \text{if } \tau \le 0 \end{cases}$$

$$(13)$$

$$D_{\tau} = \begin{cases} D_{+} \equiv \lim_{\eta \to 0} (\eta F_{-}^{'})^{'} & \text{if } \tau \ge 0 \\ D_{-} \equiv \lim_{\xi \to 0} (\xi F_{+}^{'})^{'} & \text{if } \tau \le 0 \end{cases},$$
(14)

We are now ready to apply Kuzmin (1956)'s method. Let us assume that the density profile on the z-axis,  $\psi(\tau)$ , is specified:

$$\psi(\tau) = \begin{cases} \psi_{+}(\xi) \equiv \rho(\xi, 0) & \text{if } \tau \ge 0\\ \psi_{-}(\eta) \equiv \rho(0, \eta) & \text{if } \tau \le 0 \end{cases},$$
(15)

Then we can use equation (11) to first determine  $F(\tau)$ , and hence  $\rho(\xi, \eta)$ . Restricting equation (11) to the z-axis, results in the following (ordinary) differential equation for  $F(\tau)$ :

$$F^{''} - \frac{F^{'}}{\tau} = \pi G \tau \psi(\tau) - \frac{C_{\tau}}{\tau^2} - \frac{D_{\tau}}{\tau}, \qquad (16)$$

where primes denote differentiation with respect to  $\tau$ . We define  $\Psi(\tau) = \int_0^\tau \psi(s) \, ds$ , and integrate equation (16) twice, to obtain the general solution,

$$F(\tau) = \pi G \int_0^\tau s \Psi(s) ds + A_\tau + B_\tau \frac{\tau^2}{2} + \frac{C_\tau}{2} \ln|\tau| + D_\tau \tau , (17)$$

where  $A_{\tau}$  and  $B_{\tau}$  are constants of integration. We recall, from the discussion following equation (6), that  $A_{\eta} - A_{\xi}$ will merely add to M; to describe a smooth mass distribution, we may, without loss of generality, set  $A_{\eta} = A_{\xi} = 0$ . Substituting this solution for  $F(\tau)$  in equation (11) provides us with a general expression for the Stäckel density (whose z-axis profile is  $\psi(\tau)$ ) in parabolic coordinates:

$$\rho(\xi,\eta) = \frac{\xi^2 \psi_+(\xi) + \eta^2 \psi_-(\eta)}{(\xi-\eta)^2} 
- \frac{2\xi\eta}{(\xi-\eta)^3} [\Psi_+(\xi) - \Psi_-(\eta)] 
+ \rho_{\rm bcd}.$$
(18)

We recall that the Stäckel nature of the potential—hence separability of the Hamilton–Jacobi equation—remains unaffected if a point mass (see eqns. [7] and [8]) with arbitrary M is added at the origin. In equation (18),  $\rho_{\rm bcd}$  is the contribution from the  $B_{\tau}$ ,  $C_{\tau}$ , and  $D_{\tau}$  terms in equation (17). We show in the Appendix that this contribution is generally unphysical, so we ignore  $\rho_{\rm bcd}$  in the rest of the paper. It is evident that the remaining expression for  $\rho(\xi, \eta)$  is non negative.

#### 3 SCALE-FREE CUSPS

A z-axis density profile of the form  $\psi(\tau) = \rho_0 (r_0/|\tau|)^k$ , can give rise to a finite, non-negative Stäckel density only if

<sup>&</sup>lt;sup>‡</sup> To verify this we only need note that adding  $L_z^2/2\xi$  and  $L_z^2/2\eta$  to  $F_+$  and  $F_-$  respectively, adds the centrifugal term to  $U(\xi, \eta)$ . <sup>§</sup> We include density cusps in our "smooth" distribution!

0 < k < 1; this is so because we require cuspy densities to have k > 0, and only when k < 1 is

$$\Psi(\tau) \equiv \int_0^\tau \psi(s) \, ds = \frac{\rho_0 r_0^k}{(1-k)} \Theta(\tau) |\tau|^{1-k} \,, \tag{19}$$

finite. Here,  $\Theta(\tau)$  is  $\pm 1$  accordingly as  $\tau$  is positive, or negative.  $F(\tau)$  is readily computed using equation (17). Scale– free cusps, with a central black hole are described by

$$F_{s+}(\xi) = 2K\xi^{3-k} - GM$$
  

$$F_{s-}(\eta) = -2K|\eta|^{3-k} + GM$$
(20)

where  $K = \pi G \rho_0 r_0^k / 2(1-k)(2-k)$ . The potential-density pairs have simple expressions in parabolic coordinates,

$$U_{s}(\xi,\eta) = \frac{2K[\xi^{3-k} + |\eta|^{3-k}] - 2GM}{\xi - \eta}$$

$$\rho_{s}(\xi,\eta) = \rho_{0}r_{0}^{k}\frac{\xi^{2-k} + |\eta|^{2-k}}{(\xi - \eta)^{2}}$$

$$-\frac{\rho_{0}r_{0}^{k}}{(1-k)}\frac{2\xi\eta}{(\xi - \eta)^{3}}\left[\xi^{1-k} + |\eta|^{1-k}\right]$$

$$+M\delta(\mathbf{r}), \qquad (21)$$

or equally well in spherical polar coordinates:

$$U_{s}(r,\theta) = Kr^{2-k} [(1+\cos\theta)^{3-k} + (1-\cos\theta)^{3-k}] - \frac{GM}{r}$$
  

$$\rho_{s}(r,\theta) = \rho_{0} \left(\frac{r_{0}}{r}\right)^{k} [(2-k-\cos\theta)(1+\cos\theta)^{2-k}]$$

$$+(2-k+\cos\theta)(1-\cos\theta)^{2-k}]$$
$$+M\delta(\mathbf{r}).$$
(22)

Meridional sections of the isocontours of the volume density and potential for k equal to 0.1 and 0.5 are shown in Figure 1. For definiteness, the section may be taken to be the x-z plane. The density (Figures 1a and 1c) is stacked on highly oblate figures of revolution, that display a dip near the z axis. In all figures, the black hole is shown as a dot at the centre. In Figures 1b and 1d, the contribution of the black hole to the potentials is suppressed. As in the corresponding figures of ST, the potential isocontours have k-independent axis ratio of 2.

The motion of point masses in all axisymmetric potential respects two classical, isolating integrals of motion:  $(E, L_z)$ , the energy and the angular momentum about the z-axis. Stäckel potentials are special in that the Hamilton– Jacobi equation separates in special, orthogonal coordinates, giving rise to a third isolating integral  $I_3$  (c.f. Landau & Lifshitz 1976). For motion in the meridional plane,  $L_z$  may be treated as a given constant. We give below, without derivation, expressions for E and  $I_3$ :

$$E = \left(\frac{2\xi}{\xi - \eta}\right) p_{\xi}^2 + \left(\frac{2\eta}{\eta - \xi}\right) p_{\eta}^2 + U_{\text{eff}}(\xi, \eta) , \qquad (23)$$

$$I_{3} = 2\xi p_{\xi}^{2} + F_{+}(\xi) + \frac{L_{z}^{2}}{2\xi} - E\xi$$
$$= 2\eta p_{\eta}^{2} + F_{-}(\eta) + \frac{L_{z}^{2}}{2\eta} - E\eta. \qquad (24)$$

© 0000 RAS, MNRAS 000, 1-5





Figure 1. Isocontours of Density and Potential: Figures (a) and (c) show isocontours of the surface density for two different values of  $\gamma$ . The corresponding potentials are displayed in the panels on the right in (b) and (d). Successive isocontours of the density have ratios of 1.1 and 1.5 in (a) and (c) respectively. Successive potential isocontours in (b) and (d) have ratios of 2. The location of the central black hole is shown as a solid dot in all four figures, although the contribution to the potentials is not included in (b) and (d).

## 4 ORBITS

Stäckel potentials being integrable, orbits are confined to 3– tori in the 6–dimensional phase space. The three isolating integrals, E,  $L_z$  and  $I_3$ , allow us to classify orbit families. When the black hole is present, E can take all real values, whereas only positive values are permited for cusps without black holes For given E, the magnitude of  $L_z$  must be less than, or equal to the angular momentum of the circular, equatorial orbit of the same energy. Having fixed E and  $L_z$ , the third integral,  $I_3$ , will determine the excursions in  $\xi$  and  $\eta$ . From equation (24), the requirement that  $p_{\xi}^2$  and  $p_{\eta}^2$  be non negative gives

$$I_3 \ge g(\xi), \qquad -I_3 \ge g(|\eta|),$$
 (25)

where

$$g(s) = 2Ks^{3-k} + \frac{L_z^2}{2s} - Es - GM; \qquad s \ge 0.$$
(26)

Thus, the range of  $I_3$  is determined by the minimum of the function g(s), which is necessarily non positive. If this minimum value is denoted by  $-I_{\rm m}(E, L_z^2)$ , we obtain the condition  $-I_{\rm m} \leq I_3 \leq I_{\rm m}$ . We note that the dynamics is truly scale–free only in the absence of the black hole.

Given  $(E, L_z, I_3)$ , an orbit belonging to this 3-torus can explore a region of real space, whose boundaries are deter-

 $\P$  No fundamental significance is attached to these positive values of E; they merely reflect the fact that our cuspy potentials have been chosen to be non negative.

mined by the intersections of coordinate surfaces. x-z cuts of these regions, for representative orbit families, are shown shaded in Figures 2a–2d. For definiteness, we have chosen a scale-free potential of the form given in equation (21). When  $I_3$  flips sign, the allowed region for the orbit is reflected about the x-z plane; hence we will consider only  $I_3 \ge 0$ . 1. The  $L_z \neq 0$  orbits are all *loops* about the z-axis. These avoid the z-axis, and generally fill a region of space that is rotationally symmetric about the z-axis; Figure 2a displays the intersection of such a region, for a generic loop, with the x-z plane. When  $I_3 = 0$ , the region is obviously symmetric about the x-y plane, as is evident from Figure 2b. The shaded region now is filled with resonant orbits; the two frequencies, for motion in the meridional plane, are equal. 2. Polar orbits, obtained when  $L_z = 0$ , display greater variety. Motion is restricted to a plane which, for convenience, we think of as the x-z plane, and the orbits can cross the z-axis. In this plane, the orbits are similar to those that live in the phase space of the non-axisymmetric disc potentials considered by ST. To keep the presentation self-contained, we go over the classification here:

(i) E < 0: g(s) increases monotonically with s, reaching a minimum value of -GM for s = 0; thus  $-GM < I_3 < GM$ . For a fixed (say, positive) value of  $I_3$ , the motion is bounded by the coordinate curves

$$\begin{array}{lll}
0 \leq & \xi & \leq \xi_m, \\
0 \leq & |\eta| & \leq \eta_m
\end{array}$$
(27)

where  $\xi_m(E, I_3) > \eta_m(E, I_3) > 0$  are the two roots of  $g(s) = \pm I_3$ . The orbits, named *lenses* by ST, fill a lenticular region bounded by the parabolas  $\xi = \xi_m$  and  $\eta = -\eta_m$ . As the shaded region in Figure 2c indicates, the lens orbits visit the origin. When  $|I_3| = GM$ , its maximum value, the shaded region collapses to an interval of the z-axis,  $0 \le z \le \xi_m/2$ . For negative values of  $I_3$ , the roles of  $\xi$  and  $\eta$  are interchanged.

(ii) E > 0: g(s) is not monotonic in s; a minimum,  $-I_{\rm m} < 0$ , is attained at  $s = [E/2K(3-k)]^{1/(2-k)}$ . Lenses occur for  $|I_3| < GM$ , but a new type of orbits appears when  $GM < |I_3| \leq I_{\rm m}$ : the orbits are centrophobic bananas (they cross the z-axis, but avoid the origin), and the allowed region is shown in Figure 2d. They are parented by a resonant banana orbit which is obtained when  $I_3 = I_{\rm m}$ . When  $M_{\bullet} =$ 0, E is always positive; the lens orbits disappear, leaving the bananas as the only generic family of orbits.

### 5 DISCUSSION

We have generalized Kuzmin's theorem to parabolic coordinates and constructed three dimensional, axisymmetric, cuspy mass models whose potentials are of Stäckel form. A point mass may be added at the centre, while leaving the separable (i.e. Stäckel) nature of the potentials unimpaired. We have also presented a set of scale–free cusps,  $\rho \propto 1/r^k$ , where 0 < k < 1, corresponds to the weak cusps seen in the "core–type" elliptical galaxies of Lauer et. al (1995).

Kuzmin's original construction in elliptic coordinates allows the central axis ratio to be specified as an independent parameter, whereas the very choice of parabolic coordinates forces the isopotential contours (when the contribution of the black hole to the potential is neglected) to



**X X Figure 2.** Orbit Families: Intersection of the region of real space filled by orbits on a 3-torus (in phase space) with the x-z plane is shown as the shaded area in all the figures. (a) and (b) correspond to loop orbits which have  $L_z \neq 0$ . (a) is for a "generic" set of loops, each of which densely fills the shaded region. (b) corresponds to loops with  $I_3 = 0$ ; these resonant, simple, closed (1:1) orbits. Both (c) and (d) are polar orbits with  $L_z = 0$ . These live entirely on one meridional plane, here taken to be the x-z plane. (c) and (d) are filled by centrophilic lenses, and centrophobic bananas, quite similar to those discussed in ST.

be significantly oblate, with axis ratio equal to 2. There are some other essential differences between models in parabolic coordinates on the one hand, and those studied earlier in elliptic/ellipsoidal coordinates. As de Zeeuw, Peletier & Franx (1986) noted, Stäckel models in ellipsoidal coordinates do not give rise to cuspy densities; furthermore, the Stäckel nature is completely destroyed if a point mass is included at the centre. Our models are fortunate to escape these limitations. However, the axisymmetric models we have been able to construct have densities that can be no steeper than 1/r; this not only limits our models to weak cusps, but does not allow construction of finite mass models. On the other hand, the Stäckel models in ellipsoidal coordinates have densities that fall off as steeply as  $1/r^4$  far from the centre.

When the potential of the black hole is neglected, the potential isocontours of our models are similar, concentric ovals. They may be compared with the potentials of the power–law galaxies of Evans (1994), which are stratified on similar, concentric spheroids. However, the dynamics in the power–law models is not integrable (there is no third integral). And like all scale–free models they have infinite mass. Evans' oblate, power–law galaxies also have isodensity contours that are dimpled near the z–axis (see also Richstone 1980, and Toomre 1982 for discussions of scale–free models with flat rotation curves; with k = 2 for these models, the cusps are very strong indeed). Evans notes that there is no real evidence that the density distribution of ellipticals is not dimpled, and further adds that his strongly dimpled

models have projected surface density isocontours that are somewhat boxy. Since our models have only weak cusps, the surface density can be computed only with the specification of a truncation at large r. While it may be tempting to speculate that our Stäckel models capture some essential features of the dynamics in the nuclear regions of giant ellipticals (whose isophotes are known to be boxy), we must be cautious: truncation at large r will likely spoil the exact separability that our models presently enjoy. Whether one can effect a truncation that retains much of the integrable nature is yet to be explored.

While three dimensional, axisymmetric cusps are not expected to be chaotic (c.f. Richstone 1982), it is nevertheless useful to have explicit integrable mass models with simple analytic expressions (for the density and potential) that also include a central black hole. The loops are the only generic orbit family, a fact that was recognised by Ollongren (1962) in his classic study of orbits in galactic potentials. Polar orbits fall in two categories: the centrophobic bananas and the centrophilic lenses. The models presented in this paper are the natural extension of the nonaxisymmetric two dimensional cusps that we have recently presented in ST. Together they exhaust the family of scale– free cusps that can be constructed in parabolic coordinates. Integrable, triaxial cusps still pose a challenge.

#### 6 ACKNOWLEDGMENTS

We thank Dave Syer for comments on the manuscript. JT acknowledges the support of NASA grant NAGW 1477.

### APPENDIX A:

We describe the contribution to  $\rho$  of the  $B_{\tau}$ ,  $C_{\tau}$  and  $D_{\tau}$  terms in Eq. (17). Writing  $\rho_{\rm bcd} = \rho_{\rm b} + \rho c + \rho d$ , the three contributions in order are:

$$\rho_{\rm b}(\xi,\eta) = \frac{B_+ - B_-}{\pi G} \left(\frac{-2\xi\eta}{(\xi-\eta)^3}\right),\tag{A1}$$

which, in the usual  $(r, \theta, \phi)$  polar coordinates, takes the form

$$\rho_{\rm b}(r,\theta) = \left(\frac{B_+ - B_-}{4\pi G}\right) \frac{\sin^2 \theta}{r}.\tag{A2}$$

The density  $\rho_{\rm b}$  is positive for  $B_+ > B_-$  and zero on the z-axis, as is expected of a homogeneous solution of Eq. (16).

$$\rho_{\rm c}(r,\theta) = \left(\frac{C_{-} - C_{+}}{8\pi G}\right) \frac{1}{r^{3}} - \frac{\delta(r)}{16\pi G r^{2}} [C_{+} \ln(1 + \cos\theta) \\ -C_{-} \ln(1 - \cos\theta)],$$
(A3)

$$\rho_{\rm d}(r,\theta) = \left(\frac{D_- - D_+}{4\pi G}\right) \frac{\cos\theta}{r^2}.\tag{A4}$$

The density  $\rho_{\rm d}$  changes sign across the *x-y* plane, unless we require that  $D_+$  or  $D_-$  be discontinuous. All terms in  $\rho_{\rm bcd}$  are unphysical and deserve to be ignored.

© 0000 RAS, MNRAS 000, 1–5

#### REFERENCES

- Binney, J.J., Tremaine, S., 1987, Galactic Dynamics, Princeton University Press, Princeton, Problem 2–13, p. 102
- de Zeeuw, P.T., 1985a, MNRAS, 216, 273
- de Zeeuw, P.T., 1985b, MNRAS, 216, 599
- de Zeeuw, P.T., Peletier, R.F., & Franx, M., 1986, MNRAS, 221, 1001
- Evans, N.W., 1994, MNRAS, 267, 333
- Kuzmin, G.G., 1956, Astr.Zh, 33, 27
- Kuzmin, G.G., 1973, ed. T.B. Omarov, in Dynamics of Galaxies and Clusters (Alma Ata: Akad. Nauk. Kaz. SSR), p. 71 (transl. in IAU Symposium 127, Structure and Dynamics of Elliptical Galaxies, ed. P.T. de Zeeuw (Dordrecht: Reidel), p. 553).
- Landau, L.D., Lifshitz, E.M., 1976, Mechanics, 3rd Edition, Pergamon Press, Oxford
- Lauer, T.R., Ajhar, E.A., Byun, Y.I., Dressler, A., Faber, S.M., Grillmair, C., Kormendy, J., Richstone, D.O., Tremaine, S., 1995, AJ, 110, 2622
- Ollongren, A., 1962, BAN, 16, 241
- Richstone, D.O., 1980, ApJ, 238, 103
- Richstone, D.O., 1982, ApJ, 252, 496
- Sridhar, S., Touma, J., 1997, MNRAS, to appear
- Toomre, A., 1982, ApJ, 259, 535

Having determined the boundary of the region (of the meridional plane) in which the orbit is confined, Ollongren (1962) loses interest in parabolic coordinates as being useful for a description of integrable stellar motion in our Galaxy.