

NEUTRAL POINTS OF OSCILLATION MODES ALONG EQUILIBRIUM SEQUENCES OF RAPIDLY ROTATING POLYTROPES IN GENERAL RELATIVITY

— APPLICATION OF THE COWLING APPROXIMATION —

SHIN'ICHIROU YOSHIDA ¹ AND YOSHIHARU ERIGUCHI

Department of Earth Science and Astronomy, Graduate School of Arts and Sciences, University of Tokyo,
Komaba, Meguro-ku, Tokyo 153, Japan

ABSTRACT

Relativistic Cowling approximation in which all metric perturbations are omitted is applied to non-axisymmetric infinitesimal oscillations of uniformly rotating general relativistic polytropes.

Frequencies of lower order f-modes which are important in analysis of secular instability driven by gravitational radiation are investigated and neutral points of the mode along equilibrium sequences of rotating polytropes are determined. Since this approximation becomes more accurate as stars are more relativistic and/or as they rotate more rapidly, we will be able to analyze how a rotation period of a neutron star may be limited by this instability.

Possible errors in determining neutral points caused by omitting metric perturbations are also estimated.

Subject headings: relativity — stars:rotation — stars:oscillation

1. Introduction

Since the discovery of non-axisymmetric secular instability of rotating stars driven by gravitational radiation (Chandrasekhar 1970, Friedman & Schutz 1978, Friedman 1978), its possibility to limit a rotational period of a neutron star has been pointed out by several authors. As the instability sets in at *neutral points* in the absence of viscosity, it would become the main issue to determine these neutral points for rapidly rotating stars in general relativity. However, thus far, only several investigations in Newtonian gravity (Managan 1985, Imamura, Friedman & Durisen 1985) and one post-Newtonian investigation (Cutler & Lindblom 1992) have been made, because it has been very difficult to manage general relativistic perturbations of rotating stars completely.

In such a situation, Yoshida & Kojima (1997) have examined the accuracy of the relativistic version of the Cowling approximation for oscillations of slowly rotating relativistic stars. Comparing the results by the *relativistic Cowling approximation* (hereafter RCA; for earlier literatures, see e.g. McDermott et al. 1983, Finn 1988, Lindblom & Splinter 1990, Ipser & Lindblom 1992) with ‘exact’ eigenfrequencies obtained by fully relativistic analysis, they argued that the approximation works well for slowly rotating relativistic stars. This suggests that we can determine with a fairly good accuracy the neutral points by this approximation.

¹Research Fellow of the Japan Society for the Promotion of Science

We here investigate the “counter-rotating” f-mode oscillations of rapidly rotating relativistic stellar models by the RCA. Though, as seen in the Newtonian results (Robe 1968), errors of f-mode eigenfrequencies are larger than those of p- or g- modes, absolute values of errors arising from the RCA are likely to become smaller for *relativistic* and *rotating* stars compared to the Newtonian values (see below). Furthermore we will estimate roughly the amount of the corrections to the results of the RCA.

2. Basic Equations and Numerical Method

Here we investigate only *uniformly rotating polytropic* stars. Polytropic relation is defined by $(p/\epsilon_c) = \kappa(\epsilon/\epsilon_c)^{1+1/N}$, where ϵ_c is the central energy density, κ is a measure of strength of gravity of the star and other symbols have their usual meanings.

The stationary axisymmetric equilibrium states are obtained by the KEH scheme (Komatsu, Eriguchi, & Hachisu 1989). The space coordinates (r, θ, φ) and the time coordinate t are used in this paper. As a parameter characterizing the stellar rotation we adopt the ratio of the rotational energy to the gravitational energy $T/|W|$ (see Komatsu et al. 1989).

In the present study, all the Eulerian perturbations of the metric components are omitted. Thus our basic equations consist of (1) the perturbation equation of the baryon number conservation and (2) the perturbation equation of the energy momentum conservation. The perturbed Eulerian variables are assumed to have harmonic dependences on the time and the azimuthal angular coordinate: $\sim e^{-i\sigma t + im\varphi}$, where m is an integer and σ is the frequency. The boundary condition of the problem is that *the Lagrangian perturbation of the pressure vanishes at the original stellar surface*. Since the basic equations closely resemble those of Newtonian gravity, they are solved by the same scheme developed for the Newtonian perturbations (Yoshida & Eriguchi 1995).²

3. Results

3.1. Test Calculations

We have checked our RCA code by applying it to two cases, i.e. spherical stars in general relativity and rotating stars in Newtonian gravity. First our frequencies for relativistic spherical stars agree well with those of other RCA calculations of Yoshida & Kojima (1997). As for rapidly rotating Newtonian stars, by comparing our results with those of the Newtonian Cowling approximation which are obtained by the modified code of Yoshida & Eriguchi (1995), we have found that our RCA code can reproduce the neutral points of f-modes with enough accuracy, i.e. within errors of 2%.

²Although Ipser & Lindblom (1992) have presented an elegant scalar potential formalism, we do not use that formalism here because we prefer a formulation which is parallel to that for the Newtonian case (Yoshida and Eriguchi 1995). We thus derive the relativistic version of the linearized continuity equation and of three components of the linearized Euler equation and solve them directly. See appendix for explicit form of the equations.

3.2. Neutral Points along Sequences

From intersection of $T/|W| - \sigma$ curves of equilibrium sequences and $\sigma = 0$ we can obtain neutral points which are tabulated in Table 1.

In the framework of Newtonian gravity critical values of $T/|W|$ for the $m = 2$ ‘bar’ mode have been considered to be universal, i.e. $T/|W| \sim 0.14$ irrespective of compressibility and/or the rotation laws (Tassoul 1978). However, even for $N = 0.5$ polytropes we could not find a neutral point against this bar mode in weak gravity limit, i.e. for small κ . This may result from the crudeness of the Cowling approximation for lower order modes. In the Newtonian Cowling approximation, smaller frequencies are obtained than those calculated from the full treatment. As a consequence, the critical values of $T/|W|$ at neutral points tend to be raised in this approximation. Thus even if $N < 0.808$ neutral points for the bar mode may not be reached before the termination of equilibrium sequence due to mass-shedding.

On the other hand, neutral points for the bar mode could be found for stronger gravity, although the value of $T/|W|$ does not coincide with the Newtonian universal value 0.14. Remarkable is that even for soft equations of state ($N > 0.808$) with sufficiently strong gravity, neutral points for the bar mode do appear. This is not, however, so surprising because general relativistic effect on equilibrium states becomes significant.³

Our numerical computational results can be summarized as follows: (a) for sequences with N and κ kept fixed, values of $T/|W|$ at neutral points become smaller as the value of m increases; (b) for sequences with N and m kept fixed, values of $T/|W|$ at neutral points become smaller as the value of κ increases; (c) for sequences with m and κ kept fixed, values of $T/|W|$ at neutral points becomes smaller as the value of N increases.

In Newtonian gravity, normalized eigenfrequency scales as: $\frac{\sigma}{\sqrt{4\pi\rho_c}} \sim \sqrt{l} \frac{\bar{\rho}}{\rho_c}$, where ρ_c is the mass density at the centre, $\bar{\rho}$ is the mean density of the star and l is the zenithal quantum number. From this relation, normalized frequencies become smaller for models with higher mass concentration toward the centre. Because of strong gravity, matter distributions are concentrated towards the central region for relativistic models. In other words, relativistic stars can be considered to become softer effectively. Consequently frequencies become lower for larger values of κ .

4. Discussion – How Far is the RCA Reliable ?

Before applying the RCA to realistic neutron stars, it is important to note how far this approximation can be reliably applied. In particular we need to estimate how the strength of relativity and the amount of rotation affect the results.

It should be noted that eigenfunctions of energy density perturbation for f-modes are peaked near the surface. It implies that surface regions are important for f-modes. On the other hand, as is known in Newtonian models, relative importance of gravitational perturbations reduces as the amount of mass participating in oscillations decreases.

Consequently relativistic gravity can be considered to improve the approximation because the effective

³After submitting this paper, we received a preprint of Stergioulas & Friedman (1997), where they also report to have obtained neutral points of $m = 2$ mode for softer equations of state with taking metric perturbations into consideration.

concentration of the matter due to relativity makes the role of metric perturbations less important. Furthermore perturbed density distribution of rotating models is more sharply peaked toward the surface of the star as the stars are spun up. Therefore the accuracy of the Cowling approximation can be relatively increased for rotating models than spherical ones. Thus these two factors can be said qualitatively to have advantages in deciding neutral points.

However, since there have been no investigations on perturbations of rotating stars in full general relativity, the influence of rotation on the errors due to the RCA cannot be quantitatively estimated by direct comparison with the results of full calculations. Thus we need some device to have quantitative estimation about the errors.

Our proposal is as follows. We assume that qualitative dependence of the mode on stellar rotation in general relativity is similar to that of the Newtonian counterpart. In Newtonian theory, differences between the eigenfrequencies obtained by the Cowling approximation and those by the full analysis including gravitational perturbations decrease monotonically as stars are spun up. If this tendency is the same for general relativistic rotating stars, the maximal error of the RCA in determining neutral points can be estimated.

Let us select one $T/|W| - \sigma$ curve for a certain equilibrium sequence computed by the RCA. Since the maximum error of this curve is assumed to be that of the spherical model, the amount of the maximum error can be calculated by comparing our results with exact values of spherical models in general relativity. A parallel displacement of this $T/|W| - \sigma$ curve by this amount of the maximum error yields a curve with the maximal error in determination of neutral points (Fig. 1). Since the amount of displacement required to obtain the correct eigenfrequency from the value by the Cowling approximation decreases as the value of $T/|W|$ is increased, the ‘true eigenfrequency curve’ with gravitational perturbations passes between the curves of the Cowling approximation and of the curve of the maximal error defined above.

Therefore if the error of the eigenfrequency for the spherical model and the inclination of $T/|W| - \sigma$ curve of the RCA at the approximated neutral point are known, the maximal error in determining a neutral point will be determined by:

$$\text{Maximal Error of } \frac{T}{|W|} = \Delta\sigma_0 \cdot \frac{d\left(\frac{T}{|W|}\right)}{d\sigma}, \quad (1)$$

where $\Delta\sigma_0$ is the maximum error of the frequency.

For $N = 1$ polytropes we have obtained the maximal errors of $T/|W|$ at neutral points for several modes (Fig. 2). We also display the post-Newtonian values of $T/|W|$ at neutral points computed by Cutler & Lindblom (1992). Two results agree within the maximal errors estimated here except for $m = 2$ modes which are not available for the post-Newtonian calculations.

5. Concluding Remarks

Here we have shown that the RCA works rather well in evaluating eigenfrequencies of rotating stars in general relativity despite its naiveness. As eigenmode analysis in the RCA is much easier than in the post-Newtonian treatment, this approximation can be extensively used in determining oscillatory frequencies of realistic neutron stars and/or determining their neutral points.

The main drawback of this approximation is that it is hard to include the effect of gravitational radiation

which is responsible for damping or growing of corresponding modes. The post-Newtonian treatment can, in principle, systematically include this effect by increasing further the order of approximation. In contrast the RCA in itself cannot include its effect consistently. Some modification such as an estimation of effective damping due to gravitational radiation by using eigenfunctions obtained by the RCA may be needed.

We are grateful to Dr. Shijun Yoshida for discussions. We would also like to thank the anonymous referee for letting us know the investigation by Stergioulas and Friedman, and Prof. John Friedman for sending us a copy of their preprint.

This paper is presented as a part of the Ph.D. thesis of the first author to the Department of Astronomy, University of Tokyo. This research has been supported by Grant-in-Aid for Scientific Research of the Japanese Ministry of Education, Science and Culture.

REFERENCES

- Chandrasekhar, S. 1970, *Phys.Rev.Lett.*, 24, 611
- Cutler, C., & Lindblom, L. 1992, *ApJ*, 385, 630
- Finn L. S. 1988, *MNRAS*, 101, 367
- Friedman, J. L. 1978, *Comm.Math.Phys.*, 62, 247
- Friedman, J. L., & Schutz, B. F. 1978, *ApJ*, 222, 281
- Imamura, J., Friedman, J.L., & Durisen, R.H. 1985, *ApJ*, 294, 517
- Ipsier, J. R., & Lindblom, L. 1992, *ApJ*, 389, 392
- Komatsu, H., Eriguchi, Y., & Hachisu, I. 1989, *MNRAS*, 239, 153
- Lindblom, L. 1995, *ApJ*, 438, 265
- Lindblom L., & Splinter R. J. 1990, *ApJ*, 348, 198
- Managan, R.A. 1985, *ApJ*, 294, 463
- McDermott P. N., Van Horn H. M., & Scholl J. F. 1983, *ApJ*, 268, 837
- Robe, H. 1968, *Ann.Astrophys.*, 31, 475
- Stergioulas, N., & Friedman, J. L. 1997, *gr-qc/9705056*
- Tassoul, J. L. 1978, *Theory of Rotating Stars*, Princeton University Press, Princeton, New Jersey
- Yoshida, S'i., & Eriguchi, Y. 1995, *ApJ*, 438, 830
- Yoshida, S., & Kojima, Y. 1997, To appear in *MNRAS*

A. Basic Equations of the RCA

A.1. Adiabatic perturbations to stationary axisymmetric configurations

Background spacetime is expressed by the following metric which is used for stationary axisymmetric spacetime:

$$ds^2 = -e^{2\nu(r,\theta)} dt^2 + e^{2\alpha(r,\theta)}(dr^2 + r^2 d\theta^2) + e^{2\beta(r,\theta)} r^2 \sin^2 \theta (d\varphi - \omega(r, \theta) dt)^2. \quad (\text{A1})$$

As we adopt the Cowling approximation here, perturbations to this metric are totally omitted.

As for the matter variables, perfect fluid is assumed:

$$T_{ab} = (p + \epsilon) u_a u_b + p g_{ab}, \quad (\text{A2})$$

where u^a , ϵ , and p are the 4-velocity, the energy density and the pressure of the fluid, respectively. The matter in the equilibrium state is assumed to be polytrope, i.e. $p = K\epsilon^{1+1/N}$ where K is a constant. Perturbations for the matter are assumed to be adiabatic so that the following Lagrangian relation holds for the energy density perturbation $\Delta\epsilon$ and the pressure perturbation Δp :

$$\frac{\Delta p}{p} = \gamma \frac{\Delta\epsilon}{\epsilon}, \quad (\text{A3})$$

where γ denotes the adiabatic exponent. Though this exponent needs not, in general, coincide with the equilibrium polytropic exponent, we assume in this paper that they coincide with each other for simplicity. Under this assumption, this Lagrangian relation is reduced to the Eulerian one for the Euler perturbations for the energy density $\delta\epsilon$ and the pressure δp as follows:

$$\frac{\delta p}{p} = \gamma \frac{\delta\epsilon}{\epsilon}. \quad (\text{A4})$$

Since the background spacetime has timelike and spacelike killing vectors $\frac{\partial}{\partial t}$ and $\frac{\partial}{\partial \varphi}$, all coefficients which appear in the perturbed equations do not depend on neither t nor φ . Consequently solutions of the perturbed equations have dependence of $\sim \exp(-i\sigma t + im\varphi)$ on t and φ variables where m is an azimuthal eigenvalue and integer.

A.2. Introduction of dimensionless variables

We introduce the following dimensionless variables by using ϵ_c and κ (variables underlined are dimensionless):

$$\underline{\epsilon} \equiv \epsilon/\epsilon_c, \quad (\text{A5})$$

$$\underline{p} \equiv p/\epsilon_c = K\epsilon_c^{1/N} \cdot \underline{\epsilon}^{1+1/N} \equiv \kappa \underline{\epsilon}^{1+1/N}, \quad (\text{A6})$$

$$\underline{v}^i \equiv u^i/(u^0 \sqrt{\kappa}), \quad (\text{A7})$$

$$\underline{\sigma} \equiv \sigma/\Omega_c, \quad (\text{A8})$$

$$\underline{\Omega} \equiv \Omega/\Omega_c, \quad (\text{A9})$$

$$\underline{r} = r/r_c, \quad (\text{A10})$$

where Ω is the angular velocity. Parameters r_c and Ω_c are defined by:

$$\Omega_c \equiv \sqrt{4\pi\epsilon_c} \ , \quad (\text{A11})$$

$$r_c \equiv \sqrt{\kappa}/\Omega_c \ . \quad (\text{A12})$$

The quantity $\sqrt{\kappa}$ is a dimensionless sound velocity at the center (normalized by speed of light which is now chosen as $c = 1$). This parameter measures the system's strength of gravity and corresponds to the expansion parameter in weak gravity limit.

Further we introduce the *Emden function* ψ , for it will help us to impose the numerical boundary condition at the stellar surface properly:

$$\underline{\epsilon} \equiv \psi^N \ , \quad \underline{p} = \kappa\psi^{N+1} \ . \quad (\text{A13})$$

A.3. Surface-fitted coordinates

Since a rapidly rotating star deforms from spherical configuration, grid points do not, in general, fall on the surface in the ordinary polar coordinates. This makes it difficult to impose the boundary condition at the stellar surface. To avoid this difficulty, we introduce the *surface-fitted coordinates* as in the Newtonian analysis (see e.g. Yoshida & Eriguchi 1995). For a configuration with the stellar surface $r = R_s(\theta)$ in the polar coordinates (r, θ) , new coordinates $(\tilde{r}, \tilde{\theta})$ are defined by

$$\tilde{r} \equiv r/R_s(\theta) \ , \quad \tilde{\theta} = \theta \ . \quad (\text{A14})$$

Although owing to this coordinate transformation the basic equations become slightly complicated, the boundary condition at the surface, i.e. vanishing of the Lagrangian perturbation of the pressure

$$\Delta p = 0, \quad (\text{A15})$$

can be imposed easily.

Hereafter, for simplicity, 'tilde' and underlines will be omitted in the equations expressed by this new coordinate.

A.4. Hydrodynamic Equations in the Cowling approximation

Basic equations of relativistic hydrodynamics are

$$\nabla \cdot (nu) = 0 \ , \quad (\text{baryon number conservation}) \quad (\text{A16})$$

$$\nabla_b T^{ab} = 0 \ , \quad (\text{energy-momentum conservation}) \quad (\text{A17})$$

$$\tilde{d}\epsilon = \frac{\epsilon + p}{n} \tilde{d}n \ , \quad (\text{the first law of thermodynamics}) \quad (\text{A18})$$

where n is the baryon number density, and differential operators ∇ and \tilde{d} are the covariant derivative and the exterior derivative, respectively.

By introducing 3-velocity components on orthogonal basis,

$$U \equiv -ie^{\alpha-\nu}\delta v^r, \quad (\text{A19})$$

$$V \equiv -ire^{\alpha-\nu}\delta v^\theta, \quad (\text{A20})$$

$$W \equiv e^{\beta-\nu}r \sin \theta \delta v^\varphi, \quad (\text{A21})$$

and rewriting the variable $\delta\psi$ as $\Psi = \delta\psi$, the basic equations for the perturbed quantities are written as follows.

For the baryon number conservation, we have,

$$\begin{aligned} (\sigma - m\Omega)\Psi &+ \left[-\frac{\psi(1+\kappa\psi)}{NR_s(\theta)}e^{\nu-\alpha} \right] \partial_r U + \left[-\frac{e^{\nu-\alpha}\partial_r\psi}{R_s(\theta)} + \frac{\psi(1+\kappa\psi)e^{\nu-\alpha}\partial_r\psi}{2NAR_s(\theta)} \right. \\ &\quad \left. -\psi(1+\kappa\psi)\frac{e^{\nu-\alpha}}{NR_s(\theta)} \left(2\partial_r\nu + \partial_r\alpha + \partial_r\beta + \frac{2}{r} \right) \right] U \\ &+ \left[-\frac{\psi(1+\kappa\psi)e^{\nu-\alpha}}{NR_s(\theta)r} \right] \partial_\theta V + \left[-\frac{\psi(1+\kappa\psi)e^{\nu-\alpha}S}{NR_s(\theta)r} \right] \partial_r V \\ &+ \left[-\frac{e^{\nu-\alpha}(\partial_\theta\psi - rS\partial_r\psi)}{R_s(\theta)r} + \frac{\psi(1+\kappa\psi)e^{\nu-\alpha}(\partial_\theta A - rS\partial_r A)}{2NAR_s(\theta)r} \right. \\ &\quad \left. -\frac{e^{\nu-\alpha}\psi(1+\kappa\psi)}{NR_s(\theta)r} ((\partial_\theta - rS\partial_r)(2\nu + \alpha + \beta) + \cot\theta) \right] V \\ &+ \left[\frac{\psi(1+\kappa\psi)(\omega - \Omega)e^{\nu+\beta}R_s(\theta)r \sin\theta}{NA}(\sigma - m\Omega) - \frac{\psi(1+\kappa\psi)e^{\nu-\beta}m}{NR_s(\theta)r \sin\theta} \right] W = 0. \quad (\text{A22}) \end{aligned}$$

Relativistic version of the Euler equation is obtained by applying the projection tensor $P_{ab} = u_a u_b + g_{ab}$ onto the equation of energy-momentum conservation as follows:

$$\begin{aligned} (\sigma - m\Omega)U &+ [r\partial_r\omega + 2(\omega - \Omega)r\partial_r\beta + 2(\omega - \Omega) + 2\kappa(N+1)Br\partial_r\psi(\omega - \Omega)]e^{\beta-\alpha}\sin\theta W \\ &- (N+1)ABe^{-\alpha-\beta}\frac{\partial_r\Psi}{R_s(\theta)} + \kappa(N+1)AB^2e^{-\alpha-\nu}\frac{\partial_r\psi}{R_s(\theta)}\Psi = 0, \quad (\text{A23}) \end{aligned}$$

$$\begin{aligned} (\sigma - m\Omega)V &+ [(\partial_\theta - rS\partial_r)\omega + 2(\omega - \Omega)(\partial_\theta - rS\partial_r)\beta + 2(\omega - \Omega)\cot\theta \\ &\quad + 2\kappa(N+1)B(\partial_\theta - rS\partial_r)\psi(\omega - \Omega)]e^{\beta-\alpha}\sin\theta W \\ &- (N+1)AB\frac{e^{-\nu-\alpha}}{R_s(\theta)r}(\partial_\theta - rS\partial_r)\Psi + \kappa(N+1)AB^2\frac{e^{-\nu-\alpha}}{R_s(\theta)r}(\partial_\theta - rS\partial_r)\Psi = 0, \quad (\text{A24}) \end{aligned}$$

$$\begin{aligned} (\sigma - m\Omega)W &- \left[2(\omega - \Omega)\partial_r\nu - \kappa(\omega - \Omega)^2e^{2\beta-2\nu}R_s(\theta)^2r^2\sin^2\theta - \partial_r\omega - 2(\omega - \Omega)\left(\partial_r\beta + \frac{1}{r}\right) \right] e^{\beta-\alpha}r \sin\theta U \\ &- \left[2(\omega - \Omega)(\partial_\theta - rS\partial_r)\nu - \kappa(\omega - \Omega)^2e^{2\beta-2\nu}R_s(\theta)^2r^2\sin^2\theta(\partial_\theta - rS\partial_r)\omega \right. \\ &\quad \left. -(\partial_\theta - rS\partial_r)\omega - 2(\omega - \Omega)(\partial_\theta\beta - rS\partial_r\beta + \cot\theta) \right] e^{\beta-\alpha}\sin\theta V \\ &+ \kappa\sigma(N+1)AB(\sigma - \Omega)e^{\beta-3\nu}R_s(\theta)r \sin\theta\Psi \\ &+ m(N+1)AB \left[\frac{e^{-\beta-\nu}}{R_s(\theta)r \sin\theta} - \kappa\omega(\omega - \Omega)e^{\beta-3\nu}R_s(\theta)r \sin\theta \right] \Psi = 0, \quad (\text{A25}) \end{aligned}$$

where functions A , B and S are defined as,

$$A \equiv g_{00} + 2g_{0j}v^j + g_{jk}v^jv^k = -e^{2\nu} + e^{2\beta}r^2\sin^2\theta(\omega - \Omega)^2, \quad (\text{A26})$$

$$B \equiv \frac{1}{1 + \kappa\psi}, \quad (\text{A27})$$

$$S(\theta) \equiv \frac{1}{R_s(\theta)} \frac{dR_s(\theta)}{d\theta}. \quad (\text{A28})$$


Fig. 1.— A schematic behaviour of an eigenfrequency near the neutral point. ‘Full calculation’ means values obtained from full inclusion of gravitational perturbations. ‘Correction with maximal error’ line is the line with the maximal error denoted by vertical arrows and extrapolated from a spherical model. The open circle and square denote approximated neutral points.

Fig. 2.— Critical values of $T/|W|$ at neutral points for several orders of f-modes for $N = 1$ polytrope. In this figure, as a measure of strength of gravity, the ratio of the mass to the radius in Schwarzschild coordinate M/R evaluated for spherical models is adopted. Error bars with open squares for both sides show maximal errors expected for the RCA. Integers m are the orders of the mode. Crosses are values of post-Newtonian results. Horizontal lines in the upper region denote critical values where equilibrium sequences terminate due to mass-shedding.

Table 1. $T/|W|$ Values at Neutral Points

κ	$N = 0.5$			$N = 1.0$			$N = 1.5$		
	$m = 2$	$m = 3$	$m = 4$	$m = 2$	$m = 3$	$m = 4$	$m = 2$	$m = 3$	$m = 4$
0.001	...	$1.36E - 1$	$9.06E - 2$...	$9.31E - 2$	$7.90E - 2$...	$5.69E - 2$	$4.92E - 2$
0.01	...	$1.28E - 1$	$8.30E - 2$...	$9.02E - 2$	$7.39E - 2$...	$5.66E - 2$	$4.84E - 2$
0.1	$1.47E - 1$	$9.25E - 2$	$6.17E - 2$	$9.70E - 2$	$6.95E - 2$	$4.91E - 2$	$5.65E - 2$	$4.77E - 2$	$3.44E - 2$
0.5	$7.92E - 2$	$5.11E - 2$	$3.75E - 2$	$7.53E - 2$	$4.11E - 2$	$2.93E - 2$	$4.14E - 2$	$2.92E - 2$	$1.98E - 2$

Schematic behaviour of an eigenfrequency near a neutral point



T/IWI

