

Signatures of Intergalactic Dust From the First Supernovae

Abraham Loeb and Zoltán Haiman

Astronomy Department, Harvard University, 60 Garden Street, Cambridge, MA 02138

ABSTRACT

We quantify the consequences of intergalactic dust produced by the first Type II supernovae in the universe. The fraction of gas converted into stars is calibrated based on the observed C/H ratio in the intergalactic medium at $z = 3$, assuming a Scalo mass function for the stars. The associated dust absorbs starlight energy and emits it at longer wavelengths. For a uniform mix of metals and dust with the intergalactic gas, we find that the dust distorts the microwave background spectrum by a y -parameter in the range $(0.02\text{--}2) \times 10^{-5}(M_{\text{SN}}/0.3M_{\odot})$, where M_{SN} is the average mass of dust produced per supernova. The opacity of intergalactic dust to infrared sources at redshifts $z \gtrsim 10$ is significant, $\tau_{\text{dust}} = (0.1\text{--}1) \times (M_{\text{SN}}/0.3M_{\odot})$, and could be detected with the Next Generation Space Telescope. Although dust suppresses the Ly α emission from early sources, the redshifts of star clusters at $z = 10\text{--}35$ can be easily inferred from the Lyman-limit break in their infrared spectrum between $1\text{--}3.5\mu\text{m}$.

Subject headings: cosmology: theory – cosmic background radiation

submitted to *ApJ Letters*, April 1997

1. Introduction

Recent spectroscopic observations of the Ly α forest at $z \gtrsim 3$ show evidence for a metallicity $\sim 1\%Z_{\odot}$ in absorbers with HI column densities as low as 10^{15} cm^{-2} (Cowie 1996; Songaila & Cowie 1996; Tytler et al. 1995). Numerical simulations identify such absorbers with mildly overdense regions in the intergalactic medium, out of which nonlinear objects such as galaxies condense (Hellsten et al. 1997, and references therein). Indeed, damped Ly α absorbers, which are thought to be the progenitors of present-day galaxies, show similar metallicities during the early phase of their formation at $3.5 \lesssim z \lesssim 4.5$ (Lu, Sargent, & Barlow 1996). The universality of this mean metal abundance in absorbers spanning a range of six orders of magnitude in HI column density, indicates that an early phase of metal enrichment occurred throughout the universe at $z \gtrsim 5$, before galaxies were assembled.

The formation of an early population of star clusters at redshifts $\sim 10\text{--}30$ is a natural consequence of hierarchical structure formation in Cold Dark Matter (CDM) cosmologies (see, e.g.

Haiman & Loeb 1997, hereafter HL97; Ostriker & Gnedin 1996, and references therein). The first star clusters that form in high abundance have baryonic masses $\sim 10^8 M_\odot$ and virial temperatures $\sim 10^4$ K, which allow their gas to cool via atomic transitions. Less massive objects with lower virial temperatures are unable to cool and fragment into stars due to the rapid photo-dissociation of their molecular hydrogen (Haiman, Rees, & Loeb 1996). Since the potential wells of the first clusters are relatively shallow (~ 10 km s $^{-1}$), supernova-driven winds might have expelled the metal-rich gas out of these systems and mixed it with the intergalactic medium. Incomplete mixing could have led to the observed order-of-magnitude scatter in the C/H ratio along lines-of-sight to different quasars (Rauch, Haehnelt, & Steinmetz 1996; Hellsten et al. 1997). It is an interesting coincidence that the supernova energy output associated with a metal enrichment of $\sim 1\% Z_\odot$ corresponds to ~ 10 eV per hydrogen atom, which is just above the binding energy of these early star clusters. Supernova feedback in these objects could have therefore determined the average metallicity observed in the Ly α forest. Direct observations of these supernovae might be feasible in the future (Miralda-Escudé & Rees 1997).

The measured C/H ratio can be used to calibrate the net fraction of gas which is converted into stars by a redshift $z \approx 3$. In HL97, we have used this fraction and an extension of the Press–Schechter formalism to calculate the early star formation history in a variety of CDM cosmologies. For a large range of models we have found that the universe is reionized by a redshift $z = 10$ – 20 , and that the resulting optical depth to electron scattering, ~ 0.1 , is detectable with future microwave anisotropy experiments. In addition, deep imaging with future infrared telescopes, such as the Space Infrared Telescope Facility (SIRTF) or the Next Generation Space Telescope (NGST), would be able to detect pre-galactic star clusters. In particular, NGST should find $\gtrsim 10^3$ star clusters at $z > 10$ within its field of view of $4' \times 4'$, given its planned detection threshold of 1 nJy in the wavelength range of 1– $3.5 \mu\text{m}$ (Mather & Stockman 1996).

The early epoch of star formation and metal enrichment is inevitably accompanied by the formation of dust in supernova shells. This dust has two important observational signatures. First, the absorption of starlight energy and its re-emission at long wavelengths distorts the spectrum of the cosmic microwave background (CMB) radiation (Wright et al. 1994; Bond, Carr & Hogan 1991; Adams et al. 1989; Wright 1981). Second, the opacity of the intergalactic medium to infrared sources at redshifts $z \gtrsim 10$ could be significant. For these redshifts, infrared in the observer frame corresponds to UV in the source frame—a spectral regime in which dust absorption peaks.¹ Dust obscuration must therefore be considered when predicting the performance of future infrared telescopes such as NGST.

In this *Letter*, we quantify the magnitude of the above signatures of intergalactic dust. In § 2 we describe our method of calculation and in § 3 we compute the distortion of the microwave

¹The effect of intergalactic dust at much lower redshifts was discussed in the context of the reddening (Cheney & Rowan–Robinson 1981; Wright 1981, 1990) and the number count (Ostriker & Heisler 1984; Ostriker, Vogeley, & York 1990) of quasars.

background spectrum and compare it to the current COBE limit. In § 4 we evaluate the opacity of the intergalactic medium to infrared sources. Finally, § 5 summarizes the implications of this work for future microwave-background and infrared observations. Throughout the paper, we adopt a Hubble constant $H_0 = 50 \text{ km sec}^{-1} \text{ Mpc}^{-1}$ and a cosmological density parameter $\Omega = 1$.

2. Method of Calculation

Our first goal is to calculate the distortion of the CMB spectrum due to the production of radiation by stars and its processing through dust. The CMB spectrum is initially a pure black-body with a temperature $T_{\text{CMB}} = 2.728(1+z)$ K (Fixsen et al. 1996). We assume that stars are born with the Galactic initial mass function (IMF) as parameterized by Scalo (1986), and compute their time-dependent spectra using standard spectral atlases (Kurucz 1993) and evolutionary tracks (Schaller et al. 1992). Our stellar modeling is described in detail in HL97. In computing the amount of dust produced by the stars, we assume that each type II supernova yields $0.3M_{\odot}$ of dust which gets uniformly distributed in the intergalactic medium (see discussion in § 3), and that the dust absorption follows the wavelength-dependent opacity of Galactic dust.

To describe the effect of dust on the CMB we follow Wright (1981), and define the comoving number density of photons with a comoving frequency ν as

$$N_{\nu}(z) \equiv \frac{4\pi}{hc(1+z)^3} J_{\nu(1+z)}(z), \quad (1)$$

where $J_{\nu}(z)$ is the specific intensity in $\text{erg cm}^{-2} \text{ s}^{-1} \text{ Hz}^{-1} \text{ sr}^{-1}$. The evolution of $N_{\nu}(z)$ is determined by the radiative transfer equation:

$$-\frac{dN_{\nu}(z)}{dz} = \frac{cdt}{dz} \left[j_{\nu}(z) - \alpha_{\nu(1+z)}(z)N_{\nu}(z) \right], \quad (2)$$

where in an $\Omega = 1$ universe, $(cdt/dz) = (c/H_0)(1+z)^{-5/2}$. Here, $j_{\nu}(z)$ is the comoving emission coefficient due to stars and dust,

$$j_{\nu}(z) = \frac{4\pi}{hc(1+z)^3} J_{\nu(1+z)}^*(z) + \frac{8\pi}{c^3} \left\{ \frac{\nu^3}{\exp[h\nu(1+z)/k_{\text{B}}T_{\text{dust}}] - 1} \right\} \alpha_{\nu(1+z)}(z), \quad (3)$$

and $\alpha_{\nu}(z)$ is the dust absorption coefficient,

$$\alpha_{\nu}(z) = \rho_{\text{dust}}(z)\kappa_{\nu}, \quad (4)$$

for a mass density, ρ_{dust} , in dust. We assume that the dust is in thermal equilibrium with the background radiation field and equate the power it absorbs in the UV to the power it emits in the infrared. The dust temperature T_{dust} as a function of redshift is then derived from the implicit equation,

$$\int_0^{\infty} d\nu \kappa_{\nu(1+z)} \left\{ N_{\nu}(z) - \frac{8\pi}{c^3} \frac{\nu^3}{\exp[h\nu(1+z)/k_{\text{B}}T_{\text{dust}}] - 1} \right\} = 0. \quad (5)$$

We adopt the dust opacity κ_ν based on the mean Galactic extinction law $A(\lambda)/A(J)$ of Mathis (1990),

$$\kappa_\nu = 8.3 \times 10^3 \left[\frac{A(\lambda)}{A(J)} \right] \left(1 - \frac{0.084}{0.12 + (\lambda(\mu\text{m}) - 0.4)^2} \right) \text{ cm}^2 \text{ g}^{-1}, \quad (6)$$

where $\lambda = c/\nu$ is the wavelength; the numerical coefficient converts the extinction per hydrogen atom, with the assumed gas-to-dust ratio of 100, to extinction per gram of dust; and the last term is the albedo term given by Wright (1981) for converting the extinction (absorption plus scattering) to pure absorption.

Given a specific star-formation history, stellar spectra, and dust-yields, one can integrate numerically equations (3)–(5) and get the dust temperature and the net spectrum of the radiation background at $z=0$. As in HL97, we express the star-formation history in terms of the mass fraction of all baryons which are assembled into collapsed objects, $F_{\text{coll}}(z)$, and the fraction of these collapsed baryons which get incorporated into stars, f_{star} . The density parameter of stars is then, $\Omega_{\text{star}}(z) = f_{\text{star}}F_{\text{coll}}(z)\Omega_{\text{b}}$, where Ω_{b} is the total baryonic density parameter ($= 0.05$ in our standard model). We adopt the values of both $F_{\text{coll}}(z)$ and f_{star} from HL97; $F_{\text{coll}}(z)$ is calculated in a standard CDM cosmology based on an extension of the Press-Schechter formalism that takes into account pressure and photo-dissociation of H_2 , and $f_{\text{star}} \sim 4\%$ is found from the requirement that the C/H ratio be 1% of the solar value at $z = 3$. The stellar emission coefficient is found at each redshift through a convolution of the time-dependent emission from all stars born prior to that redshift

$$j_\nu^*(z) = f_{\text{star}}\Omega_{\text{b}}\rho_{\text{c0}}(1+z)^3 \int_z^\infty dz' \frac{dF_{\text{coll}}}{dz'} \epsilon^*[\nu(1+z'), t_{z,z'}], \quad (7)$$

where ρ_{c0} is the current critical density of the universe, $t_{z,z'}$ is the time interval between redshifts z' and z , and $\epsilon^*(\nu, t)$ is the composite physical emissivity (in $\text{erg s}^{-1} \text{ Hz}^{-1} \text{ M}_\odot^{-1}$) of a cluster of stars with the Scalo IMF at time t after its initial starburst (see HL97 for details). We have assumed that all stellar photons escape their parent clouds and are ejected into the IGM. In reality, a fraction of the ionizing photons will get converted into line photons with energies $E < 13.6\text{eV}$ by case B recombinations inside the clouds, but this will only have a minor effect on the total amount of energy absorbed by dust. Free-free emission by the ionized gas in the cluster can be neglected relative to the stellar emission. The mass density of dust at each redshift is given by

$$\rho_{\text{dust}}(z) = f_{\text{dust}}f_{\text{star}}\Omega_{\text{b}}\rho_{\text{c0}}(1+z)^3 \int_z^\infty dz' \frac{dF_{\text{coll}}}{dz'} f_{\text{dep}}(t_{z,z'}), \quad (8)$$

where f_{dust} is the mass fraction of stars which gets converted into dust, and $0 \leq f_{\text{dep}}(t) \leq 1$ is the fraction of the total mass in dust which is deposited after a time t following the starburst (i.e. the mass fraction of the $\geq 8\text{M}_\odot$ stars which have completed their main sequence lifetime). Assuming that each supernova in a Scalo mass function produces 0.3M_\odot of dust leads to $f_{\text{dust}} = 0.001$. The uncertainties in this value will be discussed in the next section.

3. Distortion of the Microwave Background Spectrum due to Dust Emission

To parameterize the deviation of the spectrum at $z=0$ from a $T_{\text{CMB}}=2.728$ K black-body, we compute the Compton y -parameter y_c , defined as

$$y_c \equiv \frac{1}{4} \left[\frac{\int d\nu N_\nu}{\int d\nu N_{0,\nu}} - 1 \right], \quad (9)$$

where the integrals are evaluated over the FIRAS frequency range of 60–600 GHz (Fixsen et al. 1996), and

$$N_{0,\nu} = \frac{8\pi}{c^3} \frac{\nu^3}{\exp(h\nu/k_B T_{\text{CMB}}) - 1}. \quad (10)$$

The y -parameter we derive is subject to a number of uncertainties associated with the formation of dust. Each of these uncertainties can be parameterized by a separate factor that multiplies our result for y_c . The first uncertainty involves the average fraction of the total stellar mass which gets converted into dust, f_{dust} . The value $f_{\text{dust}} = 0.001$ adopted in § 2 is equivalent to assuming that each star with mass above $8M_\odot$ explodes as a supernova and produces $0.3M_\odot$ of dust. This mass is consistent with the values inferred from observations of the depletion of the Si I line in SN 1987A (Lucy et al. 1991), and from the infrared echo of SN 1979C and SN 1980K (Dwek 1983). Theoretical estimates with 100% condensation efficiency predict values as high as $1M_\odot$ of dust per supernova (Dwek 1988; Weaver & Woosley 1980). If the dust yield of supernovae is supplemented by a significant amount of dust produced in stars, the value of f_{dust} could increase even further. On the other hand, a substantial fraction of the dust may be destroyed by sputtering inside the supernova envelope, or by shock waves from other supernovae (Dwek 1997); the y -parameter scales linearly with the survival fraction of dust, f_{surv} . The y -parameter would also be affected by a spatial correlation between the distributions of dust and stellar sources. If dust is correlated with star forming regions, the radiation flux probed by the dust could be enhanced by a factor f_{flux} relative to the average flux in equation (7). For small values of f_{flux} , y_c scales linearly with f_{flux} . The final uncertainty involves the mixing efficiency of carbon with the intergalactic gas. The star-formation efficiency, f_{star} , which is required in order to get the observed metallicity of the Ly α forest is inversely proportional to this mixing efficiency. From equations (2), (3) and (7) we get $y_c \propto f_{\text{star}}^2$, since the radiation emitted by dust is proportional to its opacity and the stellar radiation background – both of which are independently proportional to f_{star} . All of the above uncertainties can be combined through the definition of an overall efficiency parameter η_y :

$$\eta_y \equiv f_{\text{surv}} f_{\text{flux}} \left(\frac{f_{\text{dust}}}{0.001} \right) \left(\frac{f_{\text{star}}}{0.04} \right)^2 \quad (11)$$

so that $y_c \propto \eta_y$, and $\eta_y \equiv 1$ in our standard model.

In addition to the uncertainties about the dust and the star-formation efficiency, we note that if an early generation of low-luminosity quasars of the type discussed by Eisenstein & Loeb (1995) exists, these quasars could have produced UV radiation exceeding the stellar output. The

UV light from these quasars would be reprocessed by dust in much the same way as the stellar radiation is, and could substantially increase the spectral distortion derived from the stars alone. The possible effects of early quasars are ignored in this *Letter* and will be quantified in a future paper.

Figure 1 shows the full spectrum of the background radiation for our standard model, whose parameters are summarized in Table 1. The solid line depicts the total spectrum, which is a sum over the contributions from the original CMB, the stars, and the dust, shown separately by the dotted and dashed lines. The net contribution from dust is negative (absorption) for $\lambda \lesssim 0.015$ cm. The dust emission at longer wavelengths peaks around $\lambda \approx 0.1$ cm, within the COBE range. The y -parameter derived from this spectrum is 0.46×10^{-5} , a factor of 3 below the upper limit of 1.5×10^{-5} established by COBE (Fixsen et al. 1996).

For reference, the bottom panel of Figure 1 shows the star-formation history, i.e. the fraction of baryons confined in stars as a function of redshift, for our standard model. The same panel also shows the histories of the dust temperature and the y -parameter. The dust temperature starts to deviate noticeably from the CMB temperature (dashed line) around $z \approx 10$, when the fraction of baryons in stars $\Omega_{\text{star}}/\Omega_{\text{b}} = f_{\text{star}}F_{\text{coll}} \sim 1\%$. Note that we ignore any contribution to the distortion from redshifts $z \leq 3$. Our calculation is inadequate at low redshifts, since it ignores the enhanced production of metals and dust inside galaxies and the contribution of quasars to the background radiation. The spectral distortion we obtain should therefore be regarded as a lower bound; inclusion of the contribution from lower redshifts could only increase its magnitude.

Table 1 shows the range of values obtained for the y -parameter as one deviates from the input parameters assumed in our standard model. In particular, if the normalization of the standard CDM power-spectrum $\sigma_{8h^{-1}}$ is increased, then star-formation starts earlier and spreads over a wider range of redshifts. Although the optical depth to dust is increased by 70%, the corresponding change in the y -parameter is only 10%, since the average redshift delay between the production of radiation and dust is increased. Tilting the power spectrum to have a lower power-law index n has the opposite effect; stars are produced later and over a narrower range of redshifts. In this case, the optical depth is reduced by 50% but the y -parameter decreases only by 10%. Since we normalize the star-formation efficiency by fixing the total carbon yield of the stars, any change to the slope of the IMF has little effect on the net production of dust. Adding a constant $-0.7 \leq \beta \leq 1$ to the power-law index of the IMF changes the y -parameter in the range $0.17 \leq (y_c/10^{-5}\eta_y) \leq 0.54$. When the photo-dissociation of H_2 is ignored (e.g., due to the renewed formation of H_2 on dust grains), the threshold for fragmentation and star formation in virialized objects is lowered from virial temperatures $T_{\text{vir}} \gtrsim 10^4\text{K}$ (where atomic line cooling is effective) to $T_{\text{vir}} \gtrsim 10^2\text{K}$ (where H_2 cooling operates). In this case, star-formation starts earlier and leads to a slight increase in the y -parameter. The input parameter that affects y_c the most is the baryon density, $y_c \propto \Omega_{\text{b}}^2$; high values of Ω_{b} (such as $\Omega_{\text{b}} \gtrsim 0.09$, if all other parameters are held fixed) violate the existing COBE constraint.

4. Optical Depth to Infrared Sources

Next, we evaluate the optical depth to absorption and scattering by dust along the line of sight to a source at a redshift $z_s > 3$,

$$\tau_{\text{dust}}(\nu) = \frac{c}{H_0} \int_3^{z_s} \frac{dz}{(1+z)^{5/2}} \alpha_{\nu(1+z)}(z). \quad (12)$$

The absorption coefficient $\alpha_{\nu(1+z)}(z)$ is given in equations (4) and (6). Since we are now interested in dust obscuration along a specific line of sight, both absorption and scattering must be included, and so we omit the last term in equation (6). The uncertainties discussed in § 2 concerning the formation and survival of dust, as well as the fraction of the baryons converted into stars, affect the optical depth through the parameter η_τ ,

$$\eta_\tau \equiv f_{\text{surv}} f_{\text{flux}} \left(\frac{f_{\text{dust}}}{0.001} \right) \left(\frac{f_{\text{star}}}{0.04} \right), \quad (13)$$

so that τ_{dust} obtains the value predicted by our standard model times η_τ .

Figure 2 shows the resulting optical depth in our standard model as a function of observed wavelength λ , for three different source redshifts. The optical depth peaks at $\lambda \sim 1\mu\text{m}$, where it reaches a value $\tau_{\text{dust}} = 0.4$. This high value could suppress the Ly α emission peak in the spectra of early star clusters (see Charlot & Fall 1991); attempts to detect this emission line from galaxies at $z \sim 2\text{--}4$ proved difficult in the past (Thompson, Djorgovski & Trauger 1995; Lowenthal et al. 1997). Since the total amount of dust is the same in all models with the same IMF, all other variations in the peak optical depths given in Table 1 arise from differences in the production history of dust. In particular, the optical depth is high when stars form early (e.g., as a result of a high σ_{8h-1} or the lack of H₂ destruction), and low when stars form late (low n). We note that the optical depths we derive are below the values that could cause a measurable reddening in the spectra of quasars at $z \lesssim 5$ (Rowan–Robinson 1995).

To illustrate the potential effect of dust on future observations with NGST, we show the composite spectrum of two star-clusters, located at $z_s=10$ and 20, and containing $10^8 M_\odot$ and $4.6 \times 10^8 M_\odot$ in stars, respectively (bottom panel). The stars are assumed to be distributed on the main sequence according to the Scalo IMF. In both cases the top curve shows the original unprocessed spectrum, and the bottom curve shows the obscured spectrum due to the intervening dust in the standard model (with $\eta_\nu = 1$). These results imply that the performance of NGST near its planned detection threshold of 1 nJy in the wavelength range 1–3.5 μm , could be affected by intergalactic dust.

Another important conclusion that can be drawn from Figure 2 is that despite the obscuration by dust, the Lyman-limit break could easily be identified in the emission spectrum of star clusters at high-redshifts. Additional absorption by neutral hydrogen in the IGM would only make the break appear sharper. For NGST observations between 1–3.5 μm , the detection of this break can identify source redshifts in the range $z_s = 10\text{--}35$. The feasibility of this *UV dropout* technique was

recently demonstrated through the successful photometric identification of galaxy redshifts in the interval $2.5 \lesssim z_s \lesssim 3.5$ (Steidel et al. 1996; Madau et al. 1996).

5. Summary

We have quantified the imprint of intergalactic dust due to the first stars on the y -parameter of the CMB spectrum (in the COBE wavelength range), y_c , and on the opacity to absorption and scattering of radiation from sources at high redshifts, $\tau_{\text{dust}}(z_s)$. We normalized the star-formation efficiency so as to get the observed $\sim 1\%$ solar C/H ratio in Ly α absorbers at $z=3$. With this normalization, we find $y_c \sim 10^{-5}$ and $\tau_{\text{dust}}(\infty) \sim 1$ for plausible choices of the cosmological parameters and stellar properties at high redshifts (see Table 1). Uncertainties due to the mixing efficiency of metals with the IGM, the stellar IMF, the destruction of dust, the spatial correlations between dust and stellar sources, or the cosmological parameters, could change these estimates by an order of magnitude. If early quasars exist, their additional UV radiation would enhance this spectral distortion.

The current COBE limit, $y_c < 1.5 \times 10^{-5}$, already places interesting constraints on the dust formation history of the universe at $z \gtrsim 10$. Complementary microwave anisotropy limits on the scale of $\sim 1''$ would constrain the degree of patchiness in the dust distribution. But most importantly, the future detection of source reddening at $z_s \gtrsim 10$ by NGST, would be instrumental in establishing the properties of the first stars in the universe. Despite the significant dust opacity, Figure 2 shows that infrared photometry by NGST could identify the redshift of sources at $10 \lesssim z_s \lesssim 35$, based on their pronounced Lyman-limit break at observed wavelengths of 1–3.5 μm .

We thank Dimitar Sasselov and Eli Dwek for useful discussions. This work was supported in part by the NASA ATP grant NAG5-3085 and the Harvard Milton fund.

REFERENCES

- Adams, F. C., Freese, K., Levin, J., McDowell, J. 1989, ApJ, 344, 24
 Bond, J. R., Carr, B. J., & Hogan, C. J. 1991, ApJ, 367, 420
 Charlot, S., & Fall, M. S. 1991, ApJ, 378, 471
 Cheney, J. E., & Rowan–Robinson, M. 1981, MNRAS, 195, 831
 Cowie, L. 1996, in HST and the High Redshift Universe, Proc. 37th Herstmonceux conference, eds. Tanvir, N. R., Aragón-Salamanca, A., Wall, J.V. (World Scientific:Singapore), in press, (preprint astro-ph/9609158)
 Dwek, E. 1983, ApJ, 274, 175
 Dwek, E. 1988, ApJ, 329, 814

- Dwek, E. 1997, in Science with the Next Generation Space Telescope, Astr. Soc. of the Pacific Conference Series, in press
- Eisenstein, D.J., & Loeb, A. 1995, ApJ, 443, 11
- Fixsen, D. J., et al. 1996, ApJ, 473, 576
- Haiman, Z., & Loeb, A. 1997, ApJ, in press, preprint astro-ph/9611028
- Haiman, Z., Rees, M. J., & Loeb, A. 1996, ApJ, 476, 458
- Hellsten, U., Dave, R., Hernquist, L., Weinberg, D. H., & Katz, N. 1997, preprint astro-ph/9701043
- Kurucz, R. 1993, CD-ROM No. 13, ATLAS9 Stellar Atmosphere Programs
- Lowenthal, J. D., et al. 1997, ApJ, in press, preprint astro-ph/9612239
- Lucy, L. B., Danziger, I. J., Gouiffes, C., & Bouchet, P. 1991, in Supernovae, ed. S. E. Woosley (New York: Springer), 82
- Madau, P., Ferguson, H. C., Dickinson, M., Giavalisco, M., Steidel, C. C., & Fruchter, A. 1996, MNRAS, 283, 1388
- Mather, J., & Stockman, P. 1996, STSci Newsletter, v. 13, no. 2, p. 15
- Mathis, J. S. 1990, ARA&A, 28, 37
- Miralda-Escudé, J., & Rees, M. J. 1997, ApJ, 478, L57
- Ostriker, J. P., & Gnedin, N. 1997, ApJ, 472, 630
- Ostriker, J. P., & Heisler, J. 1984, ApJ, 278, 1
- Ostriker, J. P., Vogeley, M. S., & York, D. G. 1990, ApJ, 364, 405
- Rauch, M., Haehnelt, M. G., & Steinmetz, M. 1996, ApJ, in press, preprint astro-ph/9609083
- Rowan–Robinson, M. 1995, MNRAS, 272, 737
- Scalo, J. M. 1986, Fundamentals of Cosmic Physics, vol. 11, p. 1-278
- Schaller, G., Schaerer, D., Meynet, G., & Maeder, A. 1992, A&ASS, 96, 269
- Songaila, A., & Cowie, L. L. 1996, AJ, 112, 335
- Steidel, C. C., Giavalisco, M., Dickinson, M., & Adelberger, K. L., 1996, AJ, 112, 352
- Thompson, D., Djorgovski, S., & Trauger, J. 1995, AJ, 110, 963
- Tytler, D. et al. 1995, in QSO Absorption Lines, ESO Astrophysics Symposia, ed. G. Meylan (Heidelberg: Springer), p.289
- Weaver, T. A., & Woosley, S. E. 1980, Ann. N.Y. Ac. Sci., 336, 335
- Wright, E. L. 1981, ApJ, 250, 1
- Wright, E. L. 1990, ApJ, 353, 411
- Wright, E. L., et al. 1994, ApJ, 420, 450

Table 1: Parameter values in our standard model and its variants, and their effect on the expected values of the y -parameter and peak opacity of the intergalactic dust to a source at $z = \infty$. The efficiency parameters η_y and η_τ are defined in equations (11) and (13). Models with a predicted y -parameter in excess of the COBE upper limit, $y_c < 1.5 \times 10^{-5}$, are ruled out. The optical depth of intergalactic dust is significant and could affect observations with NGST.

Parameter	Standard	Range Considered	Compton $y_c/10^{-5}\eta_y$	Opt. Depth $\tau_{\text{dust}}/\eta_\tau$
$\sigma_{8h^{-1}}$	0.67	0.67–1.0	0.42–0.46	0.40–0.69
n	1.0	0.8–1.0	0.38–0.42	0.21–0.40
IMF tilt (β)	0	–0.7–1.0	0.17–0.54	0.16–0.52
H ₂ feedback	yes	yes/no	0.42–0.46	0.40–0.77
Ω_b	0.05	0.01–0.1	0.02–1.71	0.08–0.80




Fig. 1.— **Top panel:** the full spectrum at $z=0$ in our standard model for the star-formation history. The solid line shows the total spectrum, the dotted and dashed lines show the separate contributions of the original CMB, the stars, and the dust. The net contribution of dust results in absorption at $\lambda \lesssim 0.015$ cm, and emission at longer wavelengths. **Bottom panel:** the redshift evolution of the fraction of baryons in stars, the dust temperature, and the y -parameter. Note that $f_{\text{star}} \equiv F_{\text{coll}}(z)^{-1} \Omega_{\text{star}} / \Omega_b$. The dashed curve shows the history of the CMB temperature.




Fig. 2.— **Top panel:** optical depth in our standard model to absorption and scattering by dust at $z \geq 3$, as a function of observed wavelength. The source is located at redshifts $z_s=10$, 20, or 100. **Bottom panel:** the expected flux with (lower curve) and without (upper curve) intervening dust, from star-clusters at $z_s=10$ and 20, containing $10^8 M_\odot$ and $4.6 \times 10^8 M_\odot$ in stars. The stars are assumed to be distributed on the main sequence according to a Scalo IMF. The break in the spectra at the redshifted Lyman limit wavelength includes only the absorption by neutral hydrogen in the stellar atmospheres.