EGRET Gamma-Ray Observations of the Crab P2/P1 Ratio

W. F. Tompkins¹, B.B. Jones, P.L. Nolan

W. W. Hansen Experimental Physics Laboratory, Stanford University, Stanford CA 94305

G. Kanbach

Max Planck Institut für Extraterrestische Physik, D85748 Garching, Germany

and

P. V. Ramanamurthy, D. J. Thompson NASA/Goddard Space Flight Center, Code 661, Greenbelt MD 20771

ABSTRACT

Recent observations of the Crab pulsar by the Energetic Gamma-Ray Experiment Telescope (EGRET) on the Compton Gamma Ray Observatory show that the highenergy gamma-ray light curve has changed little over the lifetime of the instrument. Previous data collected by SAS-2 and COS-B in the years 1972-82, along with earlier EGRET data, suggested a 14 year sinusoidal variation in the flux ratio between the first and second peaks. The new data from EGRET indicate that the flux ratio is constant.

Subject headings: gamma-rays: observations – pulsars : individual (Crab)

Accepted Astrophysical Journal. Scheduled 20 Sep 1997.

¹billt@egret0.stanford.edu

1. Introduction

High energy gamma ray emission from the Crab pulsar was observed by satellite-borne telescopes for 15 years: in 1972-73 by SAS-2 (Kniffen et al. 1974), from 1975-82 by COS-B (Clear et al. 1987), and since 1991 by EGRET (Nolan et al. 1993; Ramanamurthy et al. 1995). Early observations showed possible sinusoidal variation in the relative intensities of the two peaks (Wills et al. 1982) with a time scale of ~ 14 years, and it was suggested that this variation might be due to the precession or free nutation of the neutron star (Kanbach 1990, Özel 1991). An apparent confirmation of the sinusoidal signal was seen in the low energy gamma ray emission (Ulmer et al. 1994), matched in phase and period with the high energy results, but with a smaller amplitude. EGRET data from 1991 through early 1994 were consistent with the expected variation (Nolan et al. 1993, Ramanamurthy et al. 1995), although these observations spanned a time when the ratio of the peaks was predicted to be fairly constant, near the minimum of the sinusoid.

EGRET observations have now extended the available data by over two years. The most recent data were expected to be $4-6\sigma$ from the average of the previous values if the sinusoidal model is correct.

2. Observations and Analysis

The EGRET instrument is a spark chamber gamma-ray telescope with an energy range of 30 MeV - 30 GeV. Details of the instrument design, calibration, and standard analysis software are given in Thompson et al. (1993).

All viewing periods where EGRET was pointed within 20° of the Crab were analyzed, with the exception of viewing period 0021 (1991 Jul 8– 15), in which there was a large solar flare. The Compton Observatory viewing period numbers and dates for these observations are shown in Table 1. The eight viewings numbered 4120 through 5280 (1995 Feb. through 1996 Aug.) have been completed since the time of the previous work of Ramanamurthy et al. (1995).

All photons with measured energy above 50 MeV, which were within an energy-dependent cone of half-angle θ_{max} were used in this analysis. The angle θ_{max} , chosen such that 68% of the photons originating from the pulsar are within the acceptance cone, is given by (Thompson et al. 1993)

$$\theta_{max} = 5^{\circ}.85 \times (E/100 MeV)^{-0.534}$$

The arrival time of each detected photon was transformed to Solar System Barycentric Time using the DE200 ephemeris, then binned according to the pulsar phase at that time, determined from the Princeton Pulsar Timing Database (Arzoumanian et al. 1992). This analysis was performed using the PULSAR program (Fierro 1995).

As seen in Figure 1, the light curve was divided into several sections, including Peak 1 (phase .94) -.04), Peak 2 (phase .32 - .46), and the off-pulse Background (phase .46 - .94). These definitions follow those used in the COS-B analysis (Wills et al. 1982) and are similar to those used by Nolan et al. (1993) and Ramanamurthy et al. (1995). The background (from the Crab nebula, nearby sources, and the diffuse Galactic radiation) was assumed constant as a function of pulsar phase. The off-pulse count rate was then used to find the background-subtracted counts estimates of the two peaks (P1 and P2). In order to avoid any effects of changes in instrument performance, the evolution of the ratio P2/P1 was examined (as in previous analyses).

The two peaks have slightly different energy spectra (Nolan et al. 1993), and the EGRET response at different energies has changed at different rates (Esposito et al. 1997). Thus the ratio P2/P1 is affected by the changes in instrument performance over time. Calculations of this effect, however, indicate that it is an order of magnitude smaller than the errors in P2/P1 due to Poisson fluctuations.



Fig. 1.— Gamma-ray phase histograms of photons with E > 50 MeV for EGRET observations of the Crab in 1991-2 (a) and 1995-6 (b). The vertical dashed lines indicate phase boundaries used in the peak height analysis. The horizontal dashed lines indicate the background level as determined from the data.

The differences in the energy responses of the SAS-2, COS-B, and EGRET instruments are larger than the variation in the EGRET response. However, statistical errors in the previous instruments' data are larger as well. Thus the value of P2/P1 obtained from EGRET data should be comparable to that obtained with SAS-2 and COS-B.

3. Results

The values of P2/P1 obtained, together with the 1σ errors, are shown in Table 1. A reduced data set, where nearby points are joined for clarity, is shown in Figure 2. The data were fit with a constant, yielding $P2/P1 = .54 \pm .03$, with $\chi^2 = 6.01$ with 20 degrees of freedom (DOF). Such a low value of χ^2 might imply that the errors in the data were over-estimated. In this case, however, the errors arise purely from statistical Poisson fluctuations, and the low value must occur purely by chance. The data are very consistent with a constant value of P2/P1. The data were also fit with a straight line ($\chi^2 = 5.96$ with



Fig. 2.— Variation in the ratio of the two peaks in the Crab light curve for E > 50 MeV (from Table 1). For clarity, the 21 observations are grouped into 6 data points, where each point represents the average of several nearby observations. Error bars are 1σ . The dashed line is the average of all EGRET observations.

19 DOF), and a quadratic ($\chi^2 = 5.92$ with 18 DOF). Neither result gives a significantly better fit: the EGRET data are most consistent with no variation.

The EGRET data were also analyzed in conjunction with the SAS-2 (Kanbach 1990) and COS-B (Clear et al. 1987) data. The best fit sinusoid to the previous instruments' data,

$$P2/P1 = 0.85 - 0.56 \sin \left(2\pi \left(T - 1975.67\right)/13.3\right),$$

where T is the year of the observation, and to the combined data set,

$$P2/P1 = 0.544 - 0.060 \sin (2\pi (T - 1976.48) / 11.55)$$

are shown together with the data and the average value of P2/P1 in Figure 3.

As can be seen, the most recent EGRET observations (the last data point) are much less consistent with the large amplitude sinusoid. The constant value of 0.528 ± 0.027 gives a $\chi^2 = 24.0$ (with 27 DOF), indicating a good fit. The sinusoid fit to the combined data gives a period of 11.6 years with $\chi^2 = 21.2$ (with 24 DOF), which does not represent a significant improvement. Thus, using the combined data sets, the



Fig. 3.— Variation in the ratio of the two peaks in the Crab light curve from SAS-2 (1973), COS-B (1975-1983), and EGRET (1991-96), where the EGRET data set has been reduced as in Fig 2. The dotted line is the best fit sinusoid to the pre-EGRET data. The solid line is the best fit sinusoid to all the data. The dashed line is the average of all the data.

data are most consistent with no variation in P2/P1.

The light curves obtained from Phase 1 data (Apr 91–Sept 92) and from Phases 4 and 5 (Feb 95–Aug 96) are shown in Figure 1. The overall shape seems to have changed little, in contrast with the expected change if the 14 year cycle were correct. A χ^2 test was performed to compare the two light curves in a quantitative way. In order to take out possible systematic effects, both an additive offset and a multiplicative factor for the second light curve were fit to the data. The resulting $\chi^2 = 58.5$ (with 48 DOF) is consistent with no change in the light curve. Without the offset, $\chi^2 = 87.9$ (with 49 DOF) was obtained, indicating an inconsistency at the 99.95% confidence level. The offset required indicates a lower background level in the later observations. This could be due to a change in the nebular emission (de Jager et al. 1996), or could be a result of changes in the performance of EGRET. As the gas in the spark chamber ages, the sensitive area at low energies decreases, which decreases the width of the average point spread function. This effect might lower the background in the later observations.

4. Summary and Conclusion

Recent observations with EGRET have provided data relevant to the reported variation in the P2/P1 ratio of the Crab pulsar. Data from SAS-2 and COS-B suggested a sinusoidal variation in this ratio. The EGRET data, both taken alone and in conjunction with the data from previous instruments, are most consistent with a constant value of P2/P1. Examination of the light curves from early and later observations shows no distinct changes in the pulsar's light curve. The EGRET data cannot, of course, rule out past variability in the P2/P1 ratio. Future observations by EGRET and its successors may allow a more precise characterization of the long term behavior of the Crab's light curve.

The EGRET team gratefully acknowledges support from the following: The ARCS Foundation (WFT), Bundesministerium für Bildung, Wissenschaft, Forschung und Technologie grant 50 QV 9095 (MPE), NASA Cooperative Agreement NCC 5-95 (HSC), NASA Grant NAG5-1605(SU), and NASA Contract NAS5-96051 (NGC).

REFERENCES

- Arzoumanian. Z., Nice, D., & Taylor, J.H. 1992, GRO/radio timing data base, Princeton University
- Clear, J. et al. 1987, A&A, 174, 85
- Esposito, J.A. et al. 1997, in preparation.
- de Jager, O.C. et al. 1996, ApJ, 457, 253
- Fierro, J. M. 1995, Ph.D. Thesis, Stanford University
- Kanbach, G. 1990, in The EGRET Scinece Symposium: NASA Conference Publication No. 3071 ed. Fichtel, C.E. et al. 101
- Kniffen, D.A. et al. 1974 Nature, 251 397
- Nolan, P.L. et al. 1993, ApJ, 409, 697
- Özel, M.E. 1991, Europhysics Letters, 14, 3
- Ramanamurthy, P.V. et al, 1995, ApJ, 450, 791
- Thompson, D.J. et al. 1993, ApJS, 86, 629
- Ulmer, M.P. et al. 1994, ApJ, 432, 228
- Wills, R.D. et al. 1982, Nature, 296, 723

This 2-column preprint was prepared with the AAS LATEX macros v4.0.

Viewing Period	Dates	P2/P1
0002	Apr 22–28 1991	0.57 ± 0.11
0003	Apr 28–May 1 1991	0.54 ± 0.15
0004	May 1–4 1991	0.65 ± 0.18
0005	May 4–7 1991	0.55 ± 0.14
0010	May 16–30 1991	0.52 ± 0.07
0360	Aug 11–12 1992	0.66 ± 0.34
0365	Aug 12–20 1992	0.37 ± 0.15
0390	Sep 1–17 1992	0.65 ± 0.17
2130	May 23–29 1993	0.45 ± 0.19
2210	May 13–24 1993	0.49 ± 0.12
3100	Dec 1–13 1993	0.55 ± 0.16
3211	Feb 8–15 1994	0.59 ± 0.11
3215	Feb 15–17 1994	0.61 ± 0.31
4120	Feb 28–Mar 7 1995	0.52 ± 0.15
4130	Mar 7–21 1995	0.58 ± 0.12
4200	May 23–Jun 6 1995	0.45 ± 0.15
4260	Aug 8–22 1995	0.50 ± 0.20
5020	Oct 17–31 1995	0.73 ± 0.14
5260	Jul 30–Aug 13 1996	0.50 ± 0.12
5270	Aug 13–20 1996	0.69 ± 0.21
5280	Aug 20–27 1996	0.48 ± 0.22

TABLE 1 CRAB P2/P1 FOR EACH EGRET Observation