A new measurement of the baryonic fraction using the sparse NGC 3258 group of galaxies

Kristian Pedersen¹

Niels Bohr Institute, Blegdamsvej 17, DK-2100 Copenhagen, Denmark; kristian@dsri.dk

Yuzuru Yoshii

Institute of Astronomy, University of Tokyo, Mitaka, Tokyo 181, Japan; yoshii@omega.mtk.ioa.s.u-tokyo.ac

Research Centre for the Early Universe, University of Tokyo, Bunkyo-ku, Tokyo 113, Japan

and

Jesper Sommer–Larsen

Theoretical Astrophysics Centre, Juliane Maries Vej 30, DK-2100 Copenhagen, Denmark; jslarsen@tac.dk

Received 18 March 1997; accepted 26 May 1997

To appear in Astrophysical Journal Letters

¹Present address: Danish Space Research Institute, Juliane Maries Vej 30, DK-2100 Copenhagen, Denmark

ABSTRACT

New X-ray observations of the sparse NGC 3258 group of galaxies made by the ASCA satellite with good spectral and spatial resolution has revealed that this group has a gravitational potential deep enough to prohibit significant mass removal from the system. The baryonic fraction within $240h_{50}^{-1}$ kpc is found to be $0.065_{-0.020}^{+0.051}$ for $h_{50} = 1$, where $h_{50} \equiv H_0/50 \text{ km s}^{-1} \text{Mpc}^{-1}$, in good agreement with the universal value of 0.05 ± 0.01 predicted by standard Big Bang nucleosynthesis for a Universe with $\Omega_0 = 1$ and $h_{50} = 1$. Since the deep potential of the NGC 3258 group ensures that all pristine intragroup gas has been retained, the baryonic fraction of the NGC 3258 group is indicative of the universal value. Consequently it seems premature to rule out a critical Universe.

Subject headings: dark matter — galaxies: individual (NGC 3258) intergalactic medium — X-rays: galaxies

1. Introduction

A key issue in cosmology is to determine the baryonic fraction in the Universe and X-ray observations are particularly well suited for this task. They have revealed that the main component of light-emitting matter in clusters of galaxies is a $T \sim 10^7$ degree hot gas trapped in the cluster potential([White & Fabian 1995](#page-14-0), [David et al. 1995\)](#page-13-0). Recent X-ray studies with ROSAT have shown that the most abundant galaxy associations([Tully 1987,](#page-14-0) [Nolthenius 1993](#page-13-0)) groups of galaxies with only a handful of galaxies, are also massive enough to retain hot intergalactic gas([Ponman et al. 1993, Mulchaey et al. 1993](#page-13-0)). The X-ray emitting hot gas seems to be rather smoothly and spherically distributed, lending support to the assumption that the gas is in hydrostatic equilibrium with the underlying potential. With this assumption, and since the hot gas is detected far beyond the optical extent of the group, the gravitating mass distribution can be determined to several hundred kiloparsecs from the group center. The hot gas in groups is therefore a much more secure tracer of the gravitational field than the galaxies which do not even assure the physical reality of their grouping on the sky.

The mass fraction of baryons consisting of light-emitting matter (galaxies and hot gas) in the few groups with well-determined masses is around $f_b \approx 0.15$ for $h_{50} = 1$ [\(Mulchaey et](#page-13-0) [al. 1996\)](#page-13-0) which is not too dissimilar to $f_b \simeq 0.20 - 0.25 h_{50}^{-3/2}$ ([White & Fabian 1995,](#page-14-0) [David](#page-13-0) [et al. 1995\)](#page-13-0) in clusters of galaxies. If groups or clusters are representative of the Universe as a whole, the above baryonic fractions may be identified with the ratio of baryon to total mass densities Ω_b/Ω_0 in the Universe. Neither the baryonic fraction of the groups studied so far nor the baryonic fraction of clusters can be reconciled with the standard Big Bang nucleosynthesis (SBBN) prediction $\Omega_b h_{50}^2 = 0.05 \pm 0.01$ for a critical $\Omega_0 = 1$ Universe if $h_{50} = 1$ $h_{50} = 1$ $h_{50} = 1$ ([White et al. 1993\)](#page-14-0).

One approach for narrowing in on the universal baryonic fraction is to obtain deep X-ray observations of galaxy groups with a sufficiently deep potential, to ensure that de-gassing has not been important. Galaxy groups are much more abundant than galaxy clusters and those groups having kept all their primordial gas should give a relatively fair estimate of the baryonic fraction in the Universe.

The Advanced Satellite for Cosmology and Astrophysics (ASCA) provides superior energy coverage and spectral resolution compared to previous X-ray observatories, yielding more accurate determination of temperature and metal abundance of hot intergalactic gas in galaxy associations. In order to study the hot gas content of a deep potential group and determine its content of light-emitting matter and dark matter, we have obtained new ASCA observations of the NGC 3258 group for which the optical parameters are listed in Table 1. The four mass estimators of [Heisler et al. 1985](#page-13-0) yield masses in the range $0.9 - 2.4 \times 10^{13} M_{\odot}$ with a mean of $M_{vir} = 1.7 \times 10^{13} M_{\odot}$ being uncertain within a factor of about two. The large virial mass of the NGC 3258 group indicates that it has a deep gravitational potential. Furthermore the short crossing time and short collapse time point to a bound and relaxed system.

EDITOR: PLACE TABLE [1](#page-11-0) HERE.

2. X-ray studies of the NGC 3258 group of galaxies

We obtained a 22400 sec exposure of the NGC 3258 group of galaxies with ASCA in May 1994. More than 4000 photons from an extended source centered on the NGC

3258 group were detected with each of the two Gas Imaging Spectrometers (GISs). One of the Solid-state Imaging Spectrometers (SIS0) detected more than 1800 photons whereas the other (SIS1) were so badly affected by "hot and flickering" pixels that the NGC 3258 group emission from this detector could not be recovered. Consequently the SIS1 data were excluded from the analysis. The ASCA data were screened for local contaminating sources and background emissions from the detectors, and the sky were subtracted using the publically available long exposures at high Galactic latitude.

EDITOR: PLACE FIGURE [1](#page-15-0) HERE.

Diffuse emission centered on the group is clearly detected by each of the GIS detectors and by the SIS0 detector (Fig. [1](#page-15-0)). In the SIS0 image there is a clear peak situated at the centroid of the diffuse emission. Furthermore a point-like source North-West of the group center is detected and diffuse emission is extending North-East-wards beyond the GIS field of view, towards the poor cluster around NGC 3268.

The point-like source is $11''$ off-set a background Seyfert galaxy located at a redshift of $z = 0.0636$ (M.I. Andersen, private communication). The X-ray emission is well fitted as a point source shining through the intragroup gas. No other obvious optical counterparts are found within the 40′′ pointing error circle (90% confidence) of ASCA. In the analysis of the diffuse group emission a circular region centered on the X-ray point source was masked out. For the GIS data, a radius of 4' circle was chosen, and for the SIS0 data a radius of 3'.

In order not to be biased by the emission extending towards the NGC 3268 cluster, only photons within 10′ of the optical axis in this quadrant of the detector were included in the model fits to the GIS images. The contribution of "stray photons" from the NGC 3258 cluster outside the field of view were estimated from ray tracing simulations, kindly provided by the astrophysics group of Nagoya University. At most, a few percent of the

photons within any region in the GIS field of view originates outside the field of view.

The peak of the diffuse emission is off-set NGC 3258 by 1.1' and off-set NGC 3260 by 1.8 ′ , making it most likely that the peak is attributed to the minimum of the group potential. We analysed the surface brightness distribution of the diffuse intragroup emission using a model-fitting procedure [\(Pedersen 1996](#page-13-0)). Given the spherically symmetric appearance of the group X-ray emission and the number of detected photons, we generated azimuthally symmetric model images of the group emission characterised by the so-called β-model, $S(r) = S_0(1 + (r/r_c))^{-3\beta/2}$, generally used to fit the X-ray surface brightness profile of clusters of galaxies [\(Jones & Forman 1984](#page-13-0)). Vignetting of the X-ray telescopes was applied and the model images were folded with the point spread function of the X-ray telescopes and the GIS detectors. We found from the spectra that most of the photons were in the range of 1–2 keV, so that we adopted the vignetting and point spread functions at an energy of 2 keV. We checked that using the telescope and detector responses at an energy by 1 keV lower or higher does not change the conclusion of our analysis. Model images were then fitted individually to each GIS image and the SIS0 image with the slope of the surface brightness profile and the core radius as free parameters. Due to the low number of counts per pixel element a maximum-likelihood statistic was minimised rather than χ^2 .

Model fits to each of the GIS images were mutually consistent and simultaneous fit to both GIS images yielded a slope of $\beta = 0.6 \pm 0.2$ and a core radius of $r_c = 112 \pm 50 h_{50}^{-1}$ kpc. These two parameters are tightly linked in a way that larger β results in larger r_c . The quoted uncertainties arise from varying the level of the background emission by 20% which dominates over the statistical errors for a low surface brightness source like the NGC 3258 group. The intragroup emission is detected out to at least $240 h_{50}^{-1}$ kpc.

The diffuse emission extends all the way to the edge of the SIS0 field of view, making it difficult to constrain the surface brightness profile from this detector. However, the best-fit

profile from the SIS0 image is entirely consistent with the best-fit profile from simultaneous fit to both GIS images. The three independent ASCA images give tight constraints on the distribution of intragroup gas in the NGC 3258 group.

EDITOR: PLACE FIGURE [2](#page-15-0) HERE.

Integrated spectra of the masked region within 15' of the centroid of the group peak for the GIS detectors and within 14′ for the SIS0 detector were accumulated. The Fe-L complex around 1 keV, which is typical of a $1-2$ keV plasma, is visible in the SIS0 spectrum (Fig. [2\)](#page-15-0). This enables reliable determination of metal abundance in the intragroup gas. Plasma emission models were folded with the instrument responses and fitted to the spectra, using the XSPEC version 9.00 software. MEKAL models [\(Mewe et al. 1996\)](#page-13-0) as wellas Raymond-Smith models ([Raymond & Smith 1977](#page-13-0)) provide good fits to all three spectra. Simultaneous fit of MEKAL models to the GIS2 and the GIS3 spectra yields a temperature of $kT = 1.90_{-0.18}^{+0.21}$ keV and a metal abundance of $Z = 0.11_{-0.10}^{+0.15}$ solar (1 sigma errors) with $\chi^2 = 117$ for 141 degrees of freedom. Best-fit values of these parameters for the SIS0 spectrum are $kT = 1.58^{+0.16}_{-0.13}$ keV and $Z = 0.09^{+0.06}_{-0.05}$ solar ($\chi^2 = 59.3$ for 61 degrees of freedom). Fully consistent results are obtained when Raymond-Smith models are used. While the metal abundance of the NGC 3258 group is very low in concordance with the abundances of other groups deduced from lower resolution ROSAT spectra([Mulchaey et](#page-13-0) [al. 1996\)](#page-13-0), the temperature $kT = 1.7 \pm 0.2$ keV is significantly higher than those in the few other groups where the intragroup gas has been detected [\(Mulchaey et al. 1996](#page-13-0)). We also extracted spectra from three annuli centered on the centroid of the diffuse emission. In the process of spectral fitting, we took the mixing of photons between annuli into account, but for the three wide annuli chosen a maximum of 25% of the photons originated outside the annulus itself. No significant variations of kT and Z were seen out to 15'.

3. Mass determinations of the NGC 3258 group

From the profiles of gas density and temperature, we estimated the intragroup gas mass and the gravitating mass in the NGC 3258 group. Integrating the bestfit β model out to the detected extent, the gas mass inside 240 h_{50}^{-1} kpc is directly obtained as $M_{gas} = 7.8 \times 10^{11} h_{50}^{-5/2} M_{\odot}$. On the other hand, the mass of galaxies is $M_{gal} = 7.1 \times 10^{11} h_{50}^{-1} M_{\odot}$, assuming a standard mass to light ratio of $M/L_B = 8h_{50}(M/L_B)_{\odot}$ for NGC 3258 and NGC 3260 and $M/L_B = 3h_{50}(M/L_B)_{\odot}$ for NGC 3257.

The relatively high temperature and steep density profile of the intragroup gas implies that the gravitational potential well is deeper than in the few other sparse groups which have been subject to detailed X-rays studies. It should be safe to assume that no gas has been expelled from the group by galactic winds([Dekel & Silk 1986,](#page-13-0) [Yoshii & Arimoto](#page-14-0) [1987\)](#page-14-0). The smooth appearance and short sound crossing time of the intragroup gas ensure that the gas is approximately in hydrostatic equilibrium with the group potential, allowing us to estimate the gravitating mass. In concordance with the temperature profile deduced from the spectral fits, we model the intragroup gas as isothermal gas, giving the following expression for the gravitating mass([Cowie et al. 1987\)](#page-13-0)

$$
M_{grav}(r) = 1.8 \times 10^{12} h_{50}^{-1} M_{\odot} (kT/1 \,\text{keV}) (r/1') \beta \frac{(r/r_c)^2}{1 + (r/r_c)^2} \quad . \tag{1}
$$

Numerical simulations of clusters of galaxies show that this mass estimator is unbiased and accurate to better than 30%, even when significant substructure is present [\(Evrard et al.](#page-13-0) [1996\)](#page-13-0). Using $\beta = 0.6$, $r_c = 112h_{50}^{-1}$ kpc, and $kT = 1.7$ keV, the gravitating mass inside 240 h_{50}^{-1} kpc is obtained as $M_{grav} = 2.3 \times 10^{13} h_{50}^{-1} M_{\odot}$. Since M_{grav} far exceeds M_{gas} and M_{gal} , we conclude that most of mass in the NGC 3258 group is dark which is reflected in a large mass to light ratio $M_{grav}/L_B = 237 h_{50}(M/L_B)_{\odot}$. The virial mass estimate is a bit lower than the X–ray determined mass. This is to be expected since the galaxy motions only probe the inner parts of the group whereas the X–ray data trace mass contributions

out to $240h_{50}^{-1}$ kpc from the group center.

The baryonic fraction or the mass ratio of galaxies and gas to the total gravitating matter inside $240h_{50}^{-1}$ kpc is given by

$$
f_b = \frac{M_{gal} + M_{gas}}{M_{grav}} = 0.031 + 0.034 h_{50}^{-3/2} \quad . \tag{2}
$$

The baryonic fraction at large radii is nearly independent of the radius at which it is evaluated (Fig. [3](#page-15-0)), decreasing from $0.092_{-0.016}^{+0.041}$ ($h_{50} = 1$) at $r = 150h_{50}^{-1}$ kpc to $0.065_{-0.020}^{+0.051}$ $(h_{50} = 1)$ at $r = 240h_{50}^{-1}$ kpc.

EDITOR: PLACE FIGURE [3](#page-15-0) HERE.

The low baryonic fraction relative to the few other groups with well-determined gas fractions is due to a relatively steep gas density profile and a high intragroup gas temperature, yielding a low gas mass and a high gravitating mass. A summary of the results is given in Table 2.

EDITOR: PLACE TABLE [2](#page-12-0) HERE.

Cosmological implications

Our deep ASCA pointing of the sparse NGC 3258 group has enabled a precise determination of the baryonic fraction in this system. The derived value of $f_b \approx 0.065$ is comparable to the SBBN prediction of the universal baryonic density $\Omega_b h_{50}^2 = 0.05 \pm 0.01$ with $\Omega_0 = 1$ and $h_{50} \approx 1$.

Our discovery of a sparse galaxy group having a low baryonic fraction is interesting to discuss in light of the cosmological implications associated with recent measurements of primordial deuterium in high-redshift quasar absorption line systems([Rugers & Hogan](#page-14-0) [1996, Tytler et al. 1996, Songaila et al. 1997](#page-14-0)). Although the reported D/H abundance is still controversial, a range of $4 \times 10^{-5} < D/H < 2.4 \times 10^{-4}$ seems to persist [\(Songaila et al. 1997](#page-14-0)), yieldinga universal baryonic density of $0.022 < \Omega_b h_{50}^2 < 0.064$ from SBBN theory ([Hogan](#page-13-0) [1995\)](#page-13-0) where the higher D/H corresponds to the lower Ω_b . If the baryonic fraction derived in clusters([White & Fabian 1995,](#page-14-0) [David et al. 1995](#page-13-0)) of $f_b \simeq 0.20 - 0.25 h_{50}^{-3/2}$ is representative of the Universe, the total matter density in the Universe would be $0.09 < \Omega_0 h_{50}^{-1/2} < 0.32$. However, taking the baryonic fraction of the NGC 3258 group as representative, motivated by the ability of its deep potential to retain pristine gas, the estimated total matter density is $0.34_{-0.16}^{+0.15} < \Omega_0 < 0.99_{-0.45}^{+0.44}$ $(h_{50} = 1)$ or $0.20_{-0.05}^{+0.07} < \Omega_0 < 0.57_{-0.20}^{+0.17}$ $(h_{50} = 1.5)$. We note in passing, that these bounds are in accord with constraints on Ω_0 from large-scale flows ([Dekel et al. 1997](#page-13-0)) and from the first acoustic peak in the cosmic microwave background ([Hancock & Rocha 1996](#page-13-0)).

In conclusion, our measured low f_b value removes much of the prevailing argument that the baryonic fraction in galaxy systems only leaves room for a low-density Universe. Specifically, it is premature to rule out a critical Universe, and we conjecture that such a Universe is still a viable option in modern cosmology.

We are grateful to the ASCA Team for assistance with the analysis of ASCA data. Y. Tawara, M. Watanabe, A. Furuzawa and F. Akimoto of Nagoya University are gratefully acknowledged for providing ray tracing simulations of the stray light effects from the NGC 3268 cluster. We thank Hans Ulrik Nørgaard–Nielsen for enlightning discussions and useful suggestions to the manuscript. This work has been supported in part by the Grant-in-Aid

for Center-of-Exellence Research (07CE2002) of the Ministry of Education, Science, and Culture, Japan.

TABLE 1

NGC 3258 GROUP PROPERTIES

		L_B	v_r
Object	Type	$(10^{10} h_{50}^{-2} L_{B0})$	$(km s^{-1})$
NGC 3258 E1		6.2	2792
NGC 3260 E_p		2.1	2416
NGC 3257 $SAB(s)0$		1.4	3172

Parameters for individual member galaxies^{*a*}

Parameters for the NGC 3258 group b </sup>

σ	t_{cross}	t_{coll}	M_{vir}	M_{gal}
			(km s^{-1}) (H_0^{-1}) (H_0^{-1}) $(10^{12} h_{50}^{-1} M_{\odot})$ $(10^{12} h_{50}^{-1} M_{\odot})$	
400	0.07	0.13	17	(0.71)

Table 1:

^aGalaxy type, blue luminosity L_B , heliocentric radial velocity v_r are drawn from RC3. The blue luminosity L_B has been corrected for internal and Galactic absorption assuming a distance of $56 h_{50}^{-1}$ Mpc.

^bGalaxy positions are drawn from RC3. The velocity dispersion σ has been corrected for measurement errors. The crossing time t_{cross} is evaluated as the inertial radius [\(Jackson](#page-13-0) [1975\)](#page-13-0) divided by the velocity dispersion. The collapse time t_{coll} is estimated assuming a homogenous sphere with the group virial mass M_{vir} which initially expands with the universe and then collapses under its own gravity [\(Gunn & Gott 1972](#page-13-0)). The mass of galaxies M_{gal} is estimated from their blue luminosities assuming $M/L_B = 8 h_{50}(M/L_B)_{\odot}$ for NGC 3258 and NGC 3260 and $M/L_B = 3 h_{50} (M/L_B)_{\odot}$ for NGC 3257.

NGC 3258 GROUP X-RAY RESULTS^a

r_c	kT	L_X	M_{gas}	M_{grav}
$(h_{50}^{-1}$ kpc)	(keV)	(Z_{\odot}) $(10^{42} h_{50}^{-2} \text{ erg/s})$ $(10^{12} h_{50}^{-5/2} M_{\odot})$ $(10^{12} h_{50}^{-5/2} M_{\odot})$		
0.6 ± 0.2 110 ± 50 1.7 ± 0.2 0.1 ± 0.1		4.3 ± 0.3	$0.78^{+0.32}_{-0.24}$	23^{+6}_{-8}

Note – Quoted errors on derived X-ray parameters are those from the systematics which dominate over the statistical errors.

Table 2:

^aThe temperature kT and metal abundance Z are the best-fit values when combining integrated spectra within 14′ of the group center for SIS0 and within 15′ of the group center for GIS2 and GIS3. The X-ray luminosity L_X in the 0.5–5 keV band is estimated from the half of intragroup emission which is not contaminated with the background Seyfert emission or the extension towards NGC 3268 assuming a spherically symmetric surface brightness distribution. The gas mass M_{gas} and the gravitating mass M_{grav} are those integrated out to $r = 240 h_{50}^{-1}$ kpc from the group center assuming that the intragroup gas is isothermal and in hydrostatic equilibrium.

REFERENCES

- Cowie, L., Henriksen, M. & Mushotzky, R. 1987, ApJ, 317, 593
- David, L.P., Jones, C. & Forman, W. 1995, ApJ, 445, 578
- Dekel, A. & Silk, J. 1986, ApJ, 303, 39
- Dekel, A., Burstein, D. & White, S.D.M. 1997, *Critical Dialogues in Cosmology*, in press
- Evrard, A.E., Metzler, C.A. & Navarro, J.F. 1996, ApJ, 469, 494
- Gunn, J.E. & Gott, J.R. 1972, ApJ, 176, 1
- Hancock, S. & Rocha, G. 1996, *astro-ph* 9612016
- Heisler, J., Tremaine, S. & Bahcall, J.N. 1985, ApJ, 298, 8
- Hogan, C.J. 1995, ApJ, 441, L17
- Jackson, J. 1975, MNRAS, 173, 41P
- Jones, C. & Forman, W. 1984, ApJ, 276, 38
- Mewe, R., Kaastra, J. & Liedahl, D. 1996, *Legacy*, 6, 16
- Mulchaey, J.S., Davis, D.S., Mushotzky, R.F. & Burstein, D. 1993, ApJ, 404, L9
- Mulchaey, J.S., Davis, D.S., Mushotzky, R.F. & Burstein, D. 1993, ApJ, 456, 80
- Nolthenius, R. 1993, ApJS, 85, 1
- Pedersen, K. 1996, *Ph.D. Thesis*, Niels Bohr Institute, Copenhagen
- Ponman, T.J. & Bertram, D. 1993, Nature, 363, 51
- Raymond, J. & Smith, B. 1977, ApJS, 35, 419
- Rugers, M. & Hogan, C.J., ApJ, 459, L1
- Songaila, A., Wampler, E.J. & Cowie, L.L. 1997, Nature, 384, 137
- Tully, R.B. 1987, ApJ, 321, 280
- Tytler, D., Fan, X.-M. & Burles, S. 1996, Nature, 381, 207
- White, S.D.M., Navarro, J. Evrard, A. & Frenk, C. 1993, Nature, 366, 429
- White, D.A. & Fabian, A.C. 1995, MNRAS, 273, 72
- Yoshii, Y. & Arimoto, N. 1987, A&A, 188, 13

This manuscript was prepared with the AAS IATEX macros v4.0. $\,$

Fig. 1.— Optical image from the Digitized Sky Survey over-layed with ASCA GIS2 contours. The GIS2 contours were generated from the background subtracted image smoothed to 2^{\prime} A point-source model of the emission from the background Seyfert was subtracted. No correction for vignetting was applied to the image. The field is 30['] by 30′ with North up and East to the left. The contour scale is linear, ranging from 8.9×10^{-5} counts/arcmin²/sec to 7.6×10^{-4} counts/arcmin²/sec. In the North-Eastern corner, contour lines are crowding at the edge of the detector boundary.

Fig. 2.— Integrated SIS0 spectrum within 14′ of the group center. Detections from a 3′ circular region around the Seyfert galaxy were excluded. The histogram is the best fit bremsstrahlung model with absorption from Galactic hydrogen, $kT = 1.65^{+0.16}_{-0.15}$ keV. Below the residuals are shown in units of standard deviations. The model is acceptable $(\chi^2 = 49)$ for 53 degrees of freedom), but systematic residuals due to the Fe-L line complex are seen around 1 keV.

Fig. 3.— Baryonic fraction of the NGC 3258 group versus radial distance from the group center for $h_{50} = 1$. The solid line is the baryonic fraction from the best fit intragroup gas temperature ($kT = 1.7$ keV) and the best fit intragroup gas density distribution ($\beta = 0.6$, $r_c = 112h_{50}^{-1}$ kpc). The dotted lines show extreme values of the baryonic fraction when combining errors on the gas temperature with errors on the gas density distribution. The dashed line marks the standard big bang nucleosynthesis prediction for a critical $\Omega_0 = 1$ Universe (Hogan 95) with $h_{50} = 1$.

