

VLBA Imaging of the OH Maser in IIZw35

Adam S. Trotter, James M. Moran, & Lincoln J. Greenhill
Harvard-Smithsonian Center for Astrophysics, MS 42, 60 Garden Street, Cambridge, MA
02138

Xing-Wu Zheng
Department of Astronomy, Nanjing University, Nanjing, Jiangsu, China 210093

Carl R. Gwinn
Department of Physics, University of California, Santa Barbara, CA 93106

Submitted to *The Astrophysical Journal Letters*, 14 April 1997

Accepted 6 June 1997

ABSTRACT

We present a parsec-scale image of the OH maser in the nucleus of the active galaxy IIZw35, made using the Very Long Baseline Array at a wavelength of 18 cm. We detected two distinct components, with a projected separation of 50 pc (for $D = 110$ Mpc) and a separation in Doppler velocity of 70 km s^{-1} , which contain 50% of the total maser flux. Velocity gradients within these components could indicate rotation of clouds with binding mass densities of $\sim 7000 \text{ M}_{\odot} \text{ pc}^{-3}$ and total masses $\gtrsim 5 \times 10^5 \text{ M}_{\odot}$. Emission in the 1665-MHz OH line is roughly coincident in position with that in the 1667-MHz line, although the lines peak at different Doppler velocities. We detected no $\lambda = 18$ cm continuum emission; our upper limit implies a peak apparent optical depth $|\tau_{\text{OH}}| > 3.4$, assuming the maser is an unsaturated amplifier of continuum radiation.

Subject headings: masers — galaxies: active — galaxies: nuclei — galaxies: individual (IIZw35)

1. Introduction

IIIZw35 is a pair of galaxies catalogued in Zwicky (1971) as a 16th-magnitude compact blue double. Its systemic velocity, based on observations of CO emission (Mirabel & Sanders 1987), is 8289 km s^{-1} (heliocentric, optical definition), corresponding to a distance of 110 Mpc ($H_0 = 75 \text{ km s}^{-1} \text{ Mpc}^{-1}$; 1 mas is equivalent to a projected separation of 0.53 pc). Optical and IR images (Chapman et al. 1990) show two components separated by $9''$ along a position angle of 20° . In optical images, the northern galaxy is brighter and redder than the southern, and emits the majority of its energy in the far infrared; *IRAS* measured its $10 - 120 \mu\text{m}$ luminosity to be $3.5 \times 10^{11} L_\odot$, with a spectral energy distribution consistent with thermal emission from dust at 65 K. Chapman et al. (1990) classify the northern galaxy as either a LINER or Seyfert 2, based on optical emission line ratios.

The nucleus of the northern component of IIIZw35 is the site of strong OH maser emission. The total luminosity in the 1667-MHz transition, assuming isotropic emission, is $530 L_\odot$, with a ratio of flux density in the 1667 and 1665-MHz spectral peaks of 9:1 (Staveley-Smith et al. 1987). The 1667-MHz spectrum has two peaks at approximately $\pm 35 \text{ km s}^{-1}$ from the systemic velocity. The maser extends in Doppler velocity over about 100 km s^{-1} , but there is also faint ($< 10 \text{ mJy}$) blueshifted emission extending more than 500 km s^{-1} from v_{sys} . Baan, Haschick, & Henkel (1989) interpret the blueshifted wings in this and other OH maser galaxies to be due to molecular outflows.

Chapman et al. (1990) imaged the $\lambda = 18 \text{ cm}$ continuum emission from IIIZw35 with the VLA and MERLIN and measured a flux density of $30 \pm 10 \text{ mJy}$ and an angular size of $\lesssim 300 \text{ mas}$ ($T_B > 10^3 \text{ K}$). The centroid of the maser emission was observed to coincide with that of the 18 cm continuum to within $0.3''$. Assuming the maser acts as an unsaturated amplifier and completely covers the continuum, they inferred an apparent peak OH optical depth $|\tau_{\text{OH}}| \sim 1.7$. Montgomery & Cohen (1992) mapped the OH maser emission with MERLIN and observed an extended ($\sim 80 \text{ mas}$) structure with an apparent velocity gradient of $2.5 \text{ km s}^{-1} \text{ pc}^{-1}$ along a position angle of -20° . They interpreted the gradient to be the signature of a rotating molecular disk in the nucleus of IIIZw35.

2. Observations

We observed III Zw 35 on 25 March 1995 at $\lambda = 18$ cm with the NRAO Very Long Baseline Array (VLBA)¹ in spectral–line mode. Only the stations at Fort Davis, Kitt Peak, Los Alamos, North Liberty, Owens Valley and Pie Town provided useful data. The baseline fringe spacings ranged from 17 to 200 mas. The data were recorded in both left and right circular polarization, in two 8–MHz passbands centered on $\nu = 8438$ and 7000 km s^{−1}. The former covers both the 1667–MHz and the 1665–MHz transitions of OH, while the latter was tuned to lie outside the known range of maser emission. The data were correlated at NRAO; each passband was divided into 512 spectral channels, yielding a Doppler velocity spacing of 2.8 km s^{−1}. All further processing was undertaken with AIPS at the CfA. After initial calibration of the amplitudes, fringe rates, and delays, the visibilities were phase–referenced to the channel of the peak line emission, at 8245 km s^{−1}. Analysis of residual fringe rates established the position of the maser peak: $\alpha_{1950} = 01^{\text{h}}41^{\text{m}}47^{\text{s}}.925 \pm 0^{\text{s}}.007$, $\delta_{1950} = 16^{\circ}51'05''.8 \pm 0''.2$. An error in this reference position during processing resulted in a systematic error in the positions of spectral features that scales linearly with frequency offset from the maser peak. The magnitude of this error is 1.7 mas MHz^{−1}; the resulting systematic errors are < 1 mas in the 1667–MHz line profile, and < 4 mas in the 1665–MHz line profile. The 1– σ noise level in a uniformly–weighted spectral channel image was 3.8 mJy/beam. Spectral channels in the lower–frequency passband were averaged to produce continuum visibilities. The 1– σ noise level in a uniformly–weighted continuum image was 0.18 mJy/beam.

3. Results

Figure 1 shows the distribution of 1667–MHz OH maser emission in the nucleus of III Zw 35. We detected emission in two clumps, with a projected separation of 90 mas (50 pc) along an axis with a position angle of -10° . The spectral peak of the maser emission is in the southern clump. Figure 2 shows expanded views of the northern and southern clumps. The positions of the 1665–MHz line peaks are also plotted. In the single–channel images, the maser features were unresolved; we place an upper limit of 17 mas on their angular sizes, which implies a peak brightness temperature $T_{\text{OH}} > 3 \times 10^6$ K. Figure 3 compares the imaged flux density in both clumps with the single–dish spectrum of Staveley–Smith et al. (1987). Note that $\sim 50\%$ of the total maser flux density was resolved in our VLBA images.

¹The National Radio Astronomy Observatory is a facility of the National Science Foundation operated under cooperative agreement with Associated Universities, Inc.

Sensitivity limitations did not permit us to image the full range of blueshifted emission, which extends $\sim 500 \text{ km s}^{-1}$ from the systemic velocity at a level of less than 10 mJy (Baan, Haschick, & Henkel 1989).

The northern clump has a projected extent of $\sim 10 \text{ pc}$ (Figure 2a). It consists of multiple spectral features, over a velocity range of 110 km s^{-1} (Figure 3). We observed an east-to-west velocity gradient of $11 \text{ km s}^{-1} \text{ pc}^{-1}$ over $\sim 5 \text{ pc}$ in the velocity range $8290 < v < 8340 \text{ km s}^{-1}$ (Figure 1, *inset*). Between 8230 and 8290 km s^{-1} , we did not detect a gradient; all of the emission in this range lies in a compact ($\lesssim 2 \text{ pc}$) cluster at the SE end of the clump (Figure 2a). A recent high-resolution intercontinental VLBI observation revealed a north-south gradient in this cluster of $30 \text{ km s}^{-1} \text{ pc}^{-1}$ (P. Diamond, private communication). Lower sensitivity and angular resolution prevented us from detecting a gradient of this magnitude.

The brightest emission is in the southern clump. The imaged line profile is well fit by a Gaussian function with a peak of 130 mJy at 8245 km s^{-1} and a FWHM of 30 km s^{-1} , with line wings extending $\pm 50 \text{ km s}^{-1}$ from the peak at a level of $< 50 \text{ mJy}$. Most of the emission in the southern clump is confined to a region approximately 3 pc across (Figure 2b); however, emission in the line wings extends $\sim 5 \text{ pc}$ SE and NW of the maser peak. Evidence for a velocity gradient is ambiguous within the compact part of the southern clump, but emission from the line wings suggests a velocity gradient of about $10 \text{ km s}^{-1} \text{ pc}^{-1}$, increasing along a position angle of 110° . This gradient is of roughly the same magnitude as that in the northern clump, but in the opposite direction (Figure 1).

We imaged 1665-MHz maser emission in both the northern and southern clumps. The southern clump has two spectral peaks, at 8232 and 8258 km s^{-1} , both with flux densities of $\sim 10 \text{ mJy}$. These two peaks straddle the 1667-MHz peak in Doppler velocity (Figure 3). Both of the southern 1665-MHz features are coincident with the compact component of the 1667-MHz clump within the position errors (Figure 2b). In the northern clump, we detected one 1665-MHz feature, also with a peak flux density of $\sim 10 \text{ mJy}$. The velocity of this feature is 8353 km s^{-1} , which is near the red edge of the 1667-MHz profile. Moreover, while the velocity of the 1667-MHz emission increases to the west in the northern clump, this 1665-MHz feature lies at its eastern end (Figure 2a). It appears that the two OH maser clumps differ in their relative distributions of 1667 and 1665-MHz emission, both in angle and in velocity. Note that the features we attribute to the 1665-MHz transition could be redshifted ($\sim 8600 \text{ km s}^{-1}$) 1667-MHz features. Single-dish spectra (*e.g.*, Mirabel & Sanders 1987) do show faint emission filling the gap between the main 1667-MHz profile and the putative 1665-MHz profile. However, the near angular and Doppler velocity coincidence of emission in the two profiles suggests we are detecting the 1665-MHz transition in both

clumps, rather than highly redshifted 1667-MHz features.

We did not detect any 18 cm continuum emission in the nucleus of IIZw35 with the VLBA. We imaged various subsets of the data with a variety of weighting schemes to confirm this non-detection. We place a 5σ upper limit on the continuum flux density of 1 mJy/beam. The visibility amplitudes, averaged over the entire observation, were consistent with $\sigma = 1$ mJy noise on all baselines (180–2200 km).

4. Discussion

The concentration of maser emission into two compact clumps is likely a result of inhomogeneity in the distribution and excitation of OH in the inner 100 pc of IIZw35, and not to compact structure in the background continuum. Chapman et al. (1990) observed a $\lambda = 18$ cm continuum source with a flux density of ~ 30 mJy with MERLIN. The non-detection of the continuum on our shortest VLBA baselines places a lower limit on the angular size of a 30 mJy Gaussian source of 170 mas, or an upper limit on the continuum brightness temperature, $T_C < 10^5$ K. This suggests that the 18 cm nuclear continuum structure is smooth on the angular scale of the maser clump separation. It is also possible that filamentary compact continuum sources of < 1 mJy are being amplified by the northern and southern maser clumps. However, the components of the northern clump show a clear gradient of velocity with position, suggesting that they sample a coherent, possibly rotating, region of molecular gas (Figure 1, *inset*). 50% of the total maser flux density was resolved in our VLBA images (Figure 3), which may indicate an extended halo component of maser emission. Smoothly distributed molecular gas could act as a low-gain amplifier of an extended continuum source. Alternatively, there may exist a population of compact maser clumps, each with a flux density below our detection limit.

If the velocity gradient of $11 \text{ km s}^{-1} \text{ pc}^{-1}$ that we observe over 5 pc in the northern clump (Figure 1; *inset*) is due to rotation of a homogeneous, gravitationally bound cloud, the implied mass density is $\sim 7000 M_\odot \text{ pc}^{-3}$ (total enclosed mass $\gtrsim 5 \times 10^5 M_\odot$). The gradient suggested in the line wings of the southern clump could be produced by rotation of a cloud of similar density. A difficulty in interpreting gradients in spectral images arises when two or more distinct components are close in angle (less than a beamwidth) and velocity (less than a linewidth). Blending of the individual components can simulate a smooth velocity gradient in an extended feature. The MERLIN observations of Montgomery & Cohen 1992 had a beamwidth of 280 mas and a channel spacing of 14 km s^{-1} ; at this resolution, spectral blending between the two distinct maser clumps appeared as a gradient. True gradients can only be identified with adequate resolution. With this caution in mind,

we note that the S-shaped structure in the northern maser clump (Figure 2a) could be produced by four or more blended, unresolved features. Even so, these features would still display a correlation between velocity and position.

Baan, Haschick, & Henkel (1989) noted that the apparent luminosity of extragalactic OH masers scales approximately as L_{FIR}^2 . They interpret this correlation to imply that pumping by nuclear FIR radiation produces a column density of inverted OH that is proportional to L_{FIR} , which then acts as a low-gain amplifier of the 18 cm radio continuum, itself proportional to L_{FIR} (Norris, Allen, & Roche 1988). The apparent optical depth estimated for the OH maser in III Zw 35, $|\tau_{\text{OH}}| \sim 1.7$, was based on the assumption that the maser completely covers the background continuum (Chapman et al. 1990). Our observation indicates that the true covering factor is significantly less than unity. We place an upper limit on the brightness temperature of the 18 cm background continuum, $T_C \lesssim 10^5$ K, and a lower limit on the peak brightness temperature of the maser, $T_{\text{OH}} > 3 \times 10^6$ K. Assuming the maser is unsaturated, we derive a lower limit on the apparent optical depth, $|\tau_{\text{OH}}| > 3.4$.

If we assume that the maser is an unsaturated amplifier of background continuum emission, that the excitation temperatures of the 1667 and 1665-MHz transitions are equal, and an LTE optical depth ratio $\tau_{1667}/\tau_{1665} = 1.8$, then the ratio of 1667 to 1665-MHz line peaks in single-dish spectra of 9:1 implies $|\tau_{1667}| \sim 5$ (Henkel & Wilson 1990). However, the two transitions do not peak at exactly the same velocities, and the resulting variation in line ratios suggests $3 \lesssim |\tau_{1667}| \lesssim 5$ across the emission profile. Randell *et al.* (1995) compared the observed main-line and satellite-line ratios with models of extragalactic OH emission regions, and found that the OH line ratios are sensitive to a number of properties of the molecular gas, including density, dust temperature and the presence of velocity gradients. The observed range of 1667 to 1665-MHz line ratios may be due to variations of these quantities within the maser clumps. Our lower limit on the peak apparent optical depth, $|\tau_{1667}| > 3.4$, is consistent with the range of optical depths implied by the mainline flux ratios. It therefore appears that low-gain models of extragalactic OH masers (*e.g.* Baan, Haschick, & Henkel 1989) do not apply in III Zw 35.

One possible explanation for the angular structure of this OH maser is that the clumps trace edge-on segments of molecular shocks. Shocks are expected to be associated with energetic molecular outflows, such as those inferred from the blueshifted line wings in several OH maser galaxies, including III Zw 35 (Baan, Haschick, & Henkel 1989). The positions of OH masers in several Galactic star-forming regions, *e.g.*, W3(OH) (Bloemhof, Reid, & Moran 1992), W75N (Baart et al. 1986) and G5.89–0.39 (Wood 1993), have been observed to coincide with shock fronts in molecular outflows. The abundance of OH can be significantly enhanced in a post-shock medium (see, *e.g.*, Elitzur & De Jong

1978; Hollenbach & McKee 1989). However, unlike most extragalactic OH masers, those in Galactic star-forming regions typically have 1667 to 1665-MHz line ratios less than one (Caswell, Haynes, & Goss 1980). Mirabel, Rodríguez, & Ruiz (1989) have observed extended (pc-scale), weak OH maser emission in a number of star-forming regions that exhibits line ratios closer to those seen in extragalactic masers, and suggest that the two phenomena may be related; if so, then a shock origin for the observed maser structure would appear less likely. Detailed modelling (Randell et al. 1995) also suggests that the brightest extragalactic OH masers may arise in pc-scale clouds with H₂ densities on the order of 10⁴ cm⁻³. We note that images of OH absorption toward the center of our Galaxy (Boyce & Cohen 1994) reveal a number of “wideline clouds” that exhibit velocity widths in excess of 100 km s⁻¹, and intrinsic velocity gradients. However, the typical size of these clouds is rather larger (~ 100 pc), and their velocity gradients at least an order of magnitude smaller, than in the maser clumps in III Zw 35.

Although the continuous structure inferred by Montgomery & Cohen (1992) was probably a result of spectral blending, it is still possible that the two maser clumps lie in a rotating disk. We cannot rule out the presence of emission in the region between the two clumps, given that 50% of the total maser flux density was resolved. The velocity separation of the maser peaks could be indicative of rotation about a central mass of $2 \times 10^7 M_{\odot}$. A difficulty with this model is that the line profiles of the two clumps overlap in velocity by more than 50 km s⁻¹, which would imply superimposed non-rotational velocities comparable to the disk rotational velocity. Furthermore, the velocity gradients we observe are more than an order of magnitude greater than what we would expect from differential rotation alone. More elaborate models, such as a warped disk, or emission on the surface of a thick torus, cannot be ruled out by this observation, but are difficult to quantify. It is possible, despite the similar angular and velocity structure of the clumps, that they are physically unconnected structures, *e.g.*, counter-rotating, gravitationally bound molecular clouds, or shocks associated with a molecular outflow.

5. Conclusion

We have used the VLBA to image the OH maser in the nucleus of III Zw 35. We detect maser emission in two clumps, with a projected separation of 50 pc along a position angle of -10°. The northern clump is extended, and exhibits a west-to-east projected velocity gradient of ~ 11 km s⁻¹ pc⁻¹. The brightest maser emission arises in the southern clump, which is mostly compact (< 3 pc); however, faint emission from the line wings suggests a velocity gradient of similar magnitude to that in the northern clump, but in the opposite

direction. We detected no 18 cm continuum emission in the nucleus, which places a lower limit on the apparent maser optical depth, $|\tau_{\text{OH}}| > 3.4$, assuming the maser is unsaturated and is amplifying continuum radiation. This is the highest apparent gain yet determined for an extragalactic OH maser. The observed flux ratios of the 1667 to 1665-MHz OH maser lines are consistent with optical depths of this magnitude or greater. Variations of the line ratio with angular position and velocity suggest that physical conditions are not uniform within the maser clumps. We did not find evidence for a 100-pc rotating, edge-on molecular structure, as reported by Montgomery & Cohen (1992); however, since 50% of the total maser flux was resolved, there is likely an extended or clumpy component of maser emission in III Zw 35 that does not appear in our images. Velocity gradients within the clumps may indicate rotation on smaller scales (3–10 pc). The observed gradients could be produced by two counter-rotating, gravitationally bound molecular clouds, each with an average density of roughly $7000 \text{ M}_{\odot} \text{ pc}^{-3}$. Alternatively, the maser clumps could be segments of molecular shocks viewed edge-on.

REFERENCES

- Baan, W. A., Haschick, A. D., & Henkel, C. 1989, *ApJ*, 346, 680
- Baart, E. E., Cohen, R. J., Davies, R. D., Rowland, P. R., & Norris, R. P. 1986, *MNRAS*, 219, 145
- Bloemhof, E. E., Reid, M. J., & Moran, J. M. 1992, *ApJ*, 397, 500
- Boyce, P. J. & Cohen, R. J. 1994, *A&AS*, 107, 563
- Caswell, J. L., Haynes, R. F., & Goss, W. M. 1980, *Australian J. Phys.*, 33, 639
- Chapman, J. M., Staveley-Smith, L., Axon, D. J., Unger, S. W., Cohen, R. J., Pedlar, A., & Davies, R. D. 1990, *MNRAS*, 244, 281
- Elitzur, M. & De Jong, T. 1978, *A&A*, 67, 323
- Henkel, C. & Wilson, T. L. 1990, *A&A*, 229, 431
- Hollenbach, D. J. & McKee, C. F. 1989, *ApJ*, 342, 306
- Mirabel, I. F. & Sanders, D. B. 1987, *ApJ*, 322, 688
- Mirabel, I. F., Rodríguez, L. F., & Ruiz, A. 1989, *ApJ*, 346, 180
- Montgomery, A. S. & Cohen, R. J. 1992, *MNRAS*, 254, 23P

- Norris, R. P., Allen, D. A., & Roche, P. F. 1988, MNRAS, 234, 773
- Randell, J., Field, D., Jones, K. N., Yates, J. A., & Gray, M. D. 1995, A&A, 300, 659
- Staveley-Smith, L., Cohen, R. J., Chapman, J. M., Pointon, L., & Unger, S. W. 1987, MNRAS, 226, 689
- Wood, D. O. S. 1993, in *Astrophysical Masers*, ed. A. W. Clegg & G. E. Nedoluha (Berlin: Springer-Verlag), 141
- Zwicky, F., 1971, *Catalogue of Selected Compact and Post Eruptive Galaxies* (Zurich: Speich)

Fig. 1.— Distribution of 1667-MHz OH maser emission in the nucleus of III Zw 35. The angular position offsets are referenced to the maser peak. Features greater than 5σ (10 mJy) in the spectral images are indicated by circles; the diameter of each circle is proportional to the flux density of the feature. The heliocentric velocity of the spectral peak in each clump is indicated (optical definition; $v(\text{Heliocentric, Optical}) = v(\text{LSR, Radio}) + 235 \text{ km s}^{-1}$ at v_{sys}). The directions and ranges of the velocity gradients observed in each clump are indicated with arrows. The synthesized beam is $25 \times 12 \text{ mas}$ with a position angle of -27° . *Inset*: A plot of velocity vs. R.A. offset in the northern maser clump. The solid line corresponds to the best-fit velocity gradient (over the range $8290 < v < 8340 \text{ km s}^{-1}$) of $5.6 \pm 0.4 \text{ km s}^{-1} \text{ mas}^{-1}$, or $11 \pm 1 \text{ km s}^{-1} \text{ pc}^{-1}$.

Fig. 2.— Expanded views of the OH maser clumps: (a) The northern clump; (b) The southern clump. The diameter of each circle is proportional to the flux density of the feature. Selected features are labelled with their heliocentric velocities. The filled squares indicate the positions of the 1665-MHz emission peaks; their heliocentric velocities are labelled in italics.

Fig. 3.— The 1667-MHz OH maser spectrum of III Zw 35. The heavy solid curve is a single-dish spectrum taken from Staveley-Smith *et al.* (1987). The two solid curves show the total imaged power as a function of heliocentric velocity in each maser clump; the dotted line is the sum of both clumps. Arrows mark the velocities of the 1665-MHz emission peaks. The marker at 8289 km s^{-1} indicates the systemic velocity. Note that maser emission in single-dish spectra extends to velocities beyond those plotted here.





