X-RAY OBSERVATIONS AND THE STRUCTURE OF ELLIPTICAL GALAXIES¹

Fabrizio Brighenti^{2,3} and William G. Mathews³

²Dipartimento di Astronomia, Università di Bologna, via Zamboni 33, Bologna 40126, Italy brighenti@astbo3.bo.astro.it

³University of California Observatories/Lick Observatory, Board of Studies in Astronomy and Astrophysics, University of California, Santa Cruz, CA 95064

mathews@lick.ucsc.edu

ABSTRACT

We compare optical and high quality x-ray data for three bright ellipticals in the Virgo cluster, NGC 4472, 4649, and 4636. The distribution of total mass in NGC 4472 and 4649 determined from x-ray data is sensitive to the stellar mass over a considerable range in galactic radius extending to $r \approx r_e$, the effective radius. The agreement of x-ray and optically-determined stellar masses provides a unique verification of the stellar mass to light ratio which is essentially constant over the range $0.1 \leq r/r_e \leq 1$. However, for NGC 4636 the dark matter is important at all radii $\geq 0.35r_e$. Evidently the dark to stellar mass ratio varies in quite different ways in ellipticals of comparable optical luminosity, implying that the radial structure of dark halos may not be universal. There is some evidence in NGC 4636 for additional support of the hot interstellar gas at $r \leq 0.35r_e$; either a field $B \sim 10^{-4}$ G or a small (mechanically unstable) central region of high gas temperature ($T \sim 10^7$ K) is required. The global temperature structure in the hot interstellar medium of many recently observed ellipticals is very similar, reaching a maximum near $3 - 4r_e$. This feature, which may suggest a new structural scale in these galaxies, is inconsistent with current theoretical gas dynamical models.

Subject headings: galaxies: elliptical and lenticular – galaxies: cooling flows – galaxies: individual(NGC4472, NGC4636, NGC4649) – xrays: galaxies

1. INTRODUCTION

Elliptical galaxies are large, bright stellar systems with considerable structural regularity. The stellar light profiles are well fit by the de Vaucouleurs law (Burkert 1993) and the stellar velocity dispersion, surface brightness, and effective radius r_e are constrained to a fundamental plane (Djorgovski & Davis 1987; Dressler et al. 1987). By contrast, the x-ray emission from hot interstellar gas in ellipticals exhibits enormous variations in L_x/L_B (Eskridge et al 1995) and, to a lesser extent, in gas temperature (Davis & White 1996).

We have re-examined optical and x-ray observations of ellipticals in preparation for a new series of gas dynamical studies of the evolution of the interstellar gas. Although current x-ray observations are sparse, we have found that they reveal more useful information about elliptical galaxies than is currently realized. Gas temperature profiles T(R) for elliptical galaxies recently determined with ROSAT and ASCA reveal a surprising uniformity for radii $r \leq 10r_e$. The total mass and density of all gravitating matter, $M_{tot}(r)$ and $\rho_{tot}(r)$, can be found from the variation of temperature and density in the hot interstellar gas by assuming that the gas is in hydrostatic equilibrium in the galactic potential.

We show here that the total mass determined from the x-ray gas for two bright Virgo ellipticals (NGC 4472 and 4649) is in excellent agreement with a de Vaucouleurs profile for $0.1r_e \leq r \leq r_e$ using optically determined mass to light ratios. Remarkably, in another bright Virgo elliptical NGC 4636 dark matter contributes substantially to the total mass within r_e and the total mass found from x-ray observations is *lower* than the known stellar mass for $r \leq 0.35r_e$, implying that thermal pressure is not the only support for the hot gas. These results suggest that AXAF x-ray observations of elliptical galaxies will provide powerful new constraints on both the interstellar physics and the nature of stellar populations in these galaxies.

2. SCALE OF THERMAL STRUCTURE IN ELLIPTICALS

Figure 1 shows a plot of the radial variation of gas temperature with projected galactic radius R/r_e for six early type galaxies. The remarkable feature that all galaxies share is a positive temperature gradient out to about ~ $3r_e$ followed by a leveling off or gradual decrease toward larger radii. Data for Figure 1 are taken from the following sources: NGC 1399: Jones et al. (1997); NGC 5044: David et al. (1994); NGC 4636: Trinchieri et al. (1994); NGC 507: Kim & Fabbiano (1995); NGC 4472: Irwin & Sarazin (1996); NGC 4649: Trinchieri et al. (1997). The temperature profile observed with the ASCA satellite by Mushotsky et al. (1994) for NGC 4636 is in excellent agreement with that of Trinchieri et al. (1994) for 20 percent solar abundance. Additional properties of these galaxies are listed in Table 1. It is remarkable that T(R) is so consistent in spite of the wide range of luminosities and cosmic environments among these galaxies. NGC 4636, 4649 and 4472 are in the Virgo cluster while NGC 1399, 5044, and 507 are the brightest galaxies in small groups or clusters. NGC 507 has an exponential surface brightness profile that is consistent with a very massive, face-on S0 galaxy (Magrelli, et al. 1992).

The temperature maximum observed in ellipticals at $\sim 3r_e$ suggests a new previously unrecognized structural scale length. This cooling of gas within $3 - 4r_e$ is not a natural result of galactic cooling flows as many have suggested. In theoretical models, the radial gas velocity in the hot interstellar gas is very subsonic since otherwise the gas replenishment time would be impossibly short and L_x would be much too low. Straightforward spherical models of the hot interstellar gas (e.g. Fig. 3 of Brighenti & Mathews 1996) typically result in subsonic inflow with dT/dr < 0 for $r \gtrsim 0.1r_e$. Although the gas loses thermal energy by emitting the x-rays observed, it is heated by compression as it descends deeper into the galactic potential, i. e. dT/dr < 0. In this sense the term "galactic cooling flow" is a misnomer. In "mass drop-out" models, in which gas is removed from the inner flow, the remaining gas is actually heated since the Pdv work done by gravity must raise its temperature even higher to provide hydrostatic support for gas at larger radii. However, this rise in T(r) is fully composated when radiation from the cooler, dropping-out gas is included (e.g. White & Sarazin 1987). In any case, mass dropout is usually invoked at small galactic radii, less than the scale of dT(r)/dr < 0 in Figure 1. The gas temperature is also sensitive to heating by supernova explosions, but this heating should be proportional to the stellar density so dT/dr < 0 at $r \lesssim r_e$ would again be expected. To our knowledge no theoretical cooling flow model can produce temperature profiles resembling those in Figure 1. Fortunately, regardless of the possibly complex thermal history of the interstellar gas, the observed variation of T and ρ can be used to determine the underlying mass distribution as we now discuss.

3. COMPARISON OF X-RAY AND OPTICAL DATA FOR THREE VIRGO GALAXIES

NGC 4472, 4649 and 4636 listed in Table 1 are all members of the Virgo cluster. We adopt a distance of D = 17 Mpc for all three galaxies, similar to the Malmquist-corrected value $D = 17.2 \pm 1.9$ Mpc of Gonzalez & Faber (1997). For any particular Virgo galaxy the distance is uncertain by ~ 20 percent due to the unknown location of the galaxy along the line of sight through the cluster.

The total mass $M_{tot}(r)$ in these galaxies can be found from the condition for hydrostatic equilibrium:

$$M_{tot}(r) = -\frac{kT(r)r}{G\mu m_p} \left(\frac{d\log\rho}{d\log r} + \frac{d\log T}{d\log r} + \phi_m \frac{d\log P_m}{d\log r}\right)$$

where the last term allows for the possibility of magnetic pressure $P_m = B^2/8\pi$ and $\phi_m = P_m/P$ is the ratio of magnetic to gas pressure. The proton mass is m_p and we assume $\mu = 0.63$. The strong negative density gradient $d \log \rho/d \log r$ is expected to be the largest of the three derivatives. Only the shape of $\rho(r)$, not its absolute normalization, influences $M_{tot}(r)$. The density distribution $\rho(r)$ in the hot interstellar gas can be found from the x-ray surface brightness distribution. T(r) and possibly also $\phi_m(r)$ must be known to some precision in order to determine reliable total masses. The total mass density is $\rho_{tot}(r) = dM_{tot}/4\pi r^2 dr$.

Density profiles are available for all three ellipticals from *Einstein* HRI observations (Trinchieri, Fabbiano & Canizares 1986) and from ROSAT PSPC (Trinchieri et al. 1994; Irwin & Sarazin 1996; Trinchieri, Fabbiano, & Kim 1997). Fortunately there is a substantial range of angular scale over which these data sets overlap so values of $\rho(r)$ from *Einstein* and ROSAT data can be renormalized to agree. Having done this we fit $n(r) = \rho(r)/m_p$ with a sum of functions $n(r) = \sum n_i(r)$ where $n_i(r) = n_o(i)[1 + (r/r_o(i))^{p(i)}]^{-1}$ and the temperature is fit with $T(r) = 2T_m [r_m/(r+r_{ot}) + (r/r_m)^q]^{-1}$. The temperature T(R) observed at any projected radius R is an average along the line of sight weighted by ρ^2 and differs in principle from the temperature T(r) at physical radius r. However, ρ^2 is a very steep function of galactic radius and we find $T(r) \approx T(R)$ within 10 percent, sufficient for our purposes here considering the observational uncertainties involved (Fig. 1). Figures 2a, 2d and 2g show our fits to the x-ray data. (The parameters for these fits are: NGC 4472: $n_o(i) = 0.095, 0.00597, -0.0004; r_o(i) = 0.107, 0.95, 10; p(i) = 2.0, 1.14, 1.19; T_m = 0.75; r_m = 0.75; r_m = 0.0004; r_m = 0.$ 0.5; $r_{ot} = 0.75$; q = 0; NGC 4636: $n_o = 0.151$; $r_o = 0.172$; p = 1.57; $T_m = 0.5375$; $r_m = 0.475$; $r_{ot} = 0.6$; $q = 0.0; \text{ NGC 4649: } n_o(i) = .1, \ .0014; \ r_o(i) = .15, \ 3.0; \ p(i) = 1.8, \ 3.0; \ T_m = 0.9; \ r_m = 4.0; \ r_{ot} = 4.0; \ q = 1.0; \ r_{ot} = 1.0; \ q = 1.$ 0.0 with radii in r_e , densities in cm⁻³ and temperatures in 10⁷ K.) Total masses $M_{tot}(r)$ (Figs. 2b, 2e and 2h) and the corresponding total mass densities $\rho_{tot}(r)$ (Figs. 2c, 2f, and 2i) are determined with $\phi_m(r) = 0$. Our values for $M_{tot}(r)$ in Figure 2b are in satisfactory agreement with those of Irwin and Sarazin (1996) (scaled to D = 17 Mpc) which are based only on ROSAT data; our $M_{tot}(r)$ in Figure 2h is in excellent agreement with Mushotsky et al. (1994).

It is of particular interest to compare $M_{tot}(r)$ and $\rho_{tot}(r)$ with corresponding stellar values. The total mass M_{*t} is found from L_B using "stellar" mass to light ratios of van der Marel (1991), determined by comparing stellar velocities measured out to $r_{obs} \approx 0.5r_e$ with solutions of the 2D Jeans equations in slowly rotating model galaxies. This "stellar" M_{*t}/L_B is sensitive to all mass within r_{obs} , and may contain a component of dark matter which can be checked a posteriori (see below). In general mass to light values determined only in the core region (e.g. Faber et al. 1997) are larger; central black holes may account for this increase but typical holes may not be sufficiently massive (Kormendy, J. & Richstone, D. 1995; Faber et al. 1997). $M_*(r)$ and $\rho_*(r)$ in Figure 2 are evaluated with de Vaucouleurs profiles (Young 1976).

The Structure of NGC 4472 and 4649: We find it quite remarkable that the total mass and density indicated by the x-ray observations agree almost exactly with stellar values in the range $0.1 \leq r \leq 1r_e$

where the observational data is of high quality (Figs 2b and 2e). Discounting a conspiracy of compensating errors, this superb agreement is possible only (i) if the stellar mass to light values determined by van der Marel (1991) are essentially correct and constant for stars out to ~ $1r_e$, and (ii) if magnetic pressure or rotation contributes little or nothing to the total pressure support of gas in $0.1 \leq r \leq 1r_e$. The total masses of dark and stellar matter are equal at ~ $2.5r_e$ which includes r_{obs} so van der Marel's M_*/L_B are likely to be totally stellar. Dark matter dominates at $r/r_e \geq 2.5$. At $r = 10r_e$ the total mass to light ratio is ~ 78 (NGC 4472) and ~ 110 (NGC 4649) and M_{tot}/L_B is undoubtedly higher at $r/r_e \gtrsim 10$. The stellar to dark halo transition is particularly striking in the density plots (Figs 2c and 2f). None of the qualitative features in Figure 2 are changed if $(M_*/L_B)_{core}$ values are used from Faber et al. (1997), but the overall fit in $0.1 \leq r/r_e \leq 1$ is of lower quality. The discrepancy $M_{tot} < M_*$ at $r \leq 0.1r_e$ may result from observational inaccuracies in this region although $M_{tot}(r)$ would be underestimated if a large magnetic field were present.

The Structure of NGC 4636: Our total mass $M_{tot}(r)$ for NGC 4636 (Fig. 2h) is in excellent agreement with the mass determined by Mushotsky et al (1994) who also assumed D = 17 Mpc. A de Vaucouleurs profile is used to determine $M_*(r)$ and $\rho_*(r)$; we have verified that NGC 4636 satisfies a de Vaucouleurs profile by plotting the surface brightness data of Peletier et al (1990) against $R^{1/4}$.

The most surprising result evident from Figures 2h and 2i is the relative dominance of dark matter in NGC 4636 as compared to NGC 4472 and 4649. The dark mass $M_{dark} \equiv M_{tot} - M_*$ in NGC 4636 becomes equal to the stellar mass M_* at about $r = 1.2r_e$ and $M_{dark} \approx 0.83M_*$ at $r = r_e$ where the total mass to light ratio is 19. This large amount of dark matter may be unusual since it has been difficult until recently to find evidence of dark matter in ellipticals from stellar velocities measured within r_e (e.g. Carollo et al. 1995). The "stellar" mass to light ratio M_*/L_B from van der Marel is determined from velocity observations only out to $r_{obs}/r_e = 0.45$, where $M_* \gg M_{dark}$ (Fig. 2h). It is remarkable that there is no change in slope in Figure 2h as $M_{tot}(r)$ crosses below $M_*(r)$ at $r \approx 0.33 r_e$ since all observations should be reliable at this radius. This failure of the x-ray determined mass to detect the stellar mass suggests large magnetic fields, unusually high gas temperatures or rotational support in this region. Magnetic fields $B \sim 9 \times 10^{-5}$ G would be required at $r = 0.1 r_e$ to support the hot gas against the stellar mass within (see Mathews & Brighenti 1997). Or a curious central temperature inversion may exist: a mean temperature within 1 kpc (0.20') of 0.9×10^7 K is implied. This inversion is constrained so that the mean projected gas temperature within 1', 0.75×10^7 , is within the observed limits $0.667 + 0.029 / -0.041 \times 10^7$ K. However, the high temperature interpretation for $M_{tot} < M_*$ in NGC 4636 is unlikely since it is buoyantly unstable; dT/dr is superadiabatic for $r \lesssim 0.08 r_e \approx 0.7$ kpc. Alternatively, M_*/L_B could decrease with galactic radius so that M_* is lower than we think near $r/r_e \approx 0.1$ in Figure 2h, but this seems unlikely in view of the uniform M_*/L_B implied by Figures 2b and 2e. None of these conclusions are changed if the core value $(M_*/L_B)_{core} = 12.69$ is used instead. The different distribution of dark matter at $r \lesssim r_e$ in NGC 4636 may provide evidence against a universal dark halo structure as proposed by Navarro, Frenk & White (1996), but for $r \gtrsim r_e$ the dark halos are all very similar having slope $\rho_{dark} \propto r^{-1.9}$ at $r \sim 10r_e$. The total mass to light ratio near the outer limit of the x-ray observations shown in Figure 2g, $r = 12.5r_e$, is about 125.

How do these Virgo ellipticals differ in other respects? NGC 4636 is a relatively isolated galaxy quite far ($\gtrsim 3$ Mpc) from the core of Virgo. Nevertheless, its x-ray image shows some azimuthal asymmetry at $r \gtrsim 5r_e$, but this emission may arise from a different source (Trinchieri et al. 1994). It also has a broad, faint stellar distribution characteristic of CD galaxies. NGC 4472 is optically the brightest galaxy in the Virgo cluster, lies within a small subgroup, and is interacting with a nearby dwarf irregular galaxy, UGC 7636. At radii $\gtrsim 3r_e$ the x-ray isophotes show a tail-like structure which may result from the motion of NGC 4772 through the Virgo cluster medium (Forman et al 1985; Irwin and Sarazin 1996). NGC 4472 has a small kinematically decoupled (non-rotating) core (Davies 1989) within ~ $0.096r_e$. NGC 4649 appears to be close to the center of Virgo. However, none of these galaxies shows evidence of unusual x-ray or optical structure at $0.1 \leq r/r_e \leq 3$ where the difference in dark to stellar mass is so apparent in Figure 2.

4. CONCLUSIONS

(1) A previously unrecognized spatial scale at ~ $3r_e$ is suggested by the similarity of gas temperature profiles in many recently observed ellipticals. (2) The stellar mass to light ratios for NGC 4472 and 4649 have been verified by x-ray observations; the interstellar gas temperature is also correct. (3) The mass to light ratio in NGC 4472 and 4649 determined from stellar motions are constant over the range $0.1r_e \leq r \leq r_e$. (4) In NGC 4472 and 4649 the hot interstellar gas is supported out to $r \sim r_e$ by thermal gas pressure, other means of support such as magnetic pressure or rotation are not evident. (5) Hot interstellar gas in the center of NGC 4636 may be supported by magnetic stresses ($B \sim 10^{-4}$ G), a transient unstable region of unexpectedly high temperatures ($T \sim 10^7$ K), or rotation. (6) Elliptical galaxies with comparable luminosities can have quite different distributions of dark matter relative to stellar matter. (7) In NGC 4636 the mass of dark matter is comparable to that in stars at $r \sim r_e$; in NGC 4472 and 4649 the mass of dark halo matter is negligible at $r \sim r_e$. (8) The "stellar" mass to light ratios for these three ellipticals as determined by van der Marel (1991) are unlikely to be contaminated by dark matter from the galactic halos. AXAF observations will be useful in interpreting the variation of M_*/L_B along the fundamental plane (Pahre & Djorgovski 1997) and in verifying central magnetically supported regions.

Thanks to Michael Loewenstein and Sandra Faber for enlightenment and encouragement. Our work on the evolution of hot gas in ellipticals is supported by NASA grant NAG 5-3060 for which we are very grateful. In addition WGM is supported by a UCSC Faculty Research Grant and FB is supported in part by Grant ASI-95-RS-152 from the Agenzia Spaziale Italiana.

REFERENCES

- Brighenti, F. & Mathews, W. G. 1996, ApJ, 470, 747
- Burkert, A., 1993, A&A, 278, 23
- Carollo, C. M., de Zeeuw, P. T., van der Marel, R. P., Danziger, I. J., & Qian, E. E. 1995, ApJ, 441, L25
- David, L. P., Jones, C., Forman, W., & Daines, S. 1994, ApJ, 428, 544
- Djorgovski, S. & Davies, M. 1987, ApJ, 313, 59

Dressler, A., et al. 1987, ApJ, 313, 42

Eskridge, P. B., Fabbiano, G., & Kim, D-W. 1995, ApJ, 441, 182

- Faber, S. M., Tremaine, S., Ajhar, E. A., Byun, Y-I., Dressler, A, Gebhardt, K., Grillmair, C., Kormendy, J., Lauer, T. R., & Richstone, D. 1997, ApJ (in press)
- Faber, S. M., Wegner, G., Burstein, D., Davies, R. L., Dressler, A., Lynden-Bell, D., & Terlevich, R. J. 1989, ApJS, 69, 763
- Forman, W., Jones, C., & Tucker, W. 1985, ApJ, 293, 102
- Grillmair, C., & Tremaine, S. 1996, AJ, 112, 105

- Gonzalez, A. H. & Faber, S. M. 1997, ApJ, (in press)
- Jones, C., Stern, C., Forman, W., Breen, J., David, L., & Tucker, W. 1997, ApJ, (in press)
- Irwin, J. A. & Sarazin, C. L. 1996, ApJ, 471, 683
- Kim, D-W. & Fabbiano, G. 1995, ApJ, 441, 182
- Kormendy, J. & Richstone, D. 1995, ARA&A, 33, 581
- Magrelli, G., Bettoni, D., & Galletta, G.1992, MNRAS, 256, 500
- Mathews, W. G. & Brighenti, F. 1997, ApJ, (in press).
- Mushotzky, R., Loewenstein, M., Awaki, H, Makishima, K, Matsushita, K., & Matsumoto, H. 1994, ApJ, 436, L79
- Navarro, J. F., Frenk, C. S., & White, S. D. M. 1996, ApJ, 462, 563
- Pahre, M. A. & Djorgovski, S. G. 1997, in *The Second Stromlo Symposium: The Nature of Elliptical Galaxies*, eds. M. Arnaboldi, G. S. Da Costa, & P. Saha, ASP Conf. Ser. 116, 154
- Peletier, R. F., Davies, R. L., Illingworth, G. D., Davis, L. E., & Cawson, M. 1990, AJ, 100, 1091
- Trinchieri, G., Fabbiano, G., & Canizares, C. R. 1996, ApJ, 310, 637
- Trinchieri, G., Fabbiano, G., & Kim, D-W. 1997, A&A, 318, 361
- Trinchieri, G., Kim, D-W., Fabbiano, G., & Canizares, C. R. C. 1994, ApJ, 428, 555
- van der Marel, R. P. 1991, MNRAS, 253, 710
- White, R. E. & Sarazin, C. L. 1987, ApJ, 318, 621

Young, P. J. 1976, AJ, 81, 807

Fig. 1.— Radial dependence of gas temperature with projected radius in five early type galaxies all showing maxima at $\sim 3r_e$.

Fig. 2.— (a,d,g) Observations and our fits to gas temperature (top) and density (bottom) for Einstein HRI (filled circles) and ROSAT (open circles) data; (b,e,h) solid curves: M_{tot} ; points are values of M_{tot} from Irwin and Sarazin (1996) for 4472 and Mushotsky et al. (1994) for 4636; long-dashed curves: stellar mass M_* ; short-dashed curves: dark mass $M_{dark} \equiv M_{tot} - M_*$ which loses accuracy when $M_* \geq M_{tot}$; dot-dashed curves: stellar mass density ρ_* with a horizontal cut at the core or break radius r_b ; short-dashed curves: dark mass density.

This preprint was prepared with the AAS LATEX macros v4.0.

NGC	Type ^a	D^{b} (Mpc)	$\frac{L_B{}^c}{(10^{10} L_{B\odot})}$	$M_{*t}/L_B^{\mathrm{c,d}}$	$\frac{L_x^{\rm c}}{(10^{41} {\rm e/s})}$	$r_e{}^c$ (kpc)	r_e (')	r_b^{e} (")
507	SO	72.2	19.81		59.3	26.94	1.285	
1399	E1	17.9	2.80	10.85	8.25	3.63	0.706	3.14
4472	E2	17	7.89	9.20	4.54	8.57	1.733	2.41
4649	E2	17	5.35	9.03	1.61	6.07	1.227	3.58
4636	EO+	17	3.54	10.74	3.79	8.32	1.683	3.21
5044	E0	37.3	7.11		52.6	14.24	1.315	

Table 1. OPTICAL AND X-RAY PROPERTIES OF SIX ELLIPTICALS

^aType from RC3.

^bDistances NGC 507 & 5044 from Faber et al. (1989); NGC 1399 from Faber et al. (1997); NGC 4636 and 4472 from Gonzalez & Faber (1997).

^cScaled to the distance in the second column.

^dMass to light ratios from van der Marel (1991).

^eBreak radii into core region from Faber et al.(1997).



