

Thin Layer Approximation in 3-D

G. S. Bisnovatyi-Kogan ^{*} S. A. Silich [†]

August 4, 2021

Abstract

Equations are derived which describe a propagation of strong shocks in the interstellar matter, without any demands for a symmetry, in a thin layer approximation (2.5 dimensions). Using these equations permits to calculate a propagation of shock waves from nonsymmetric supernovae explosions in the medium with arbitrary density distribuion, formation of superbubbles in galaxies.

1 Thin layer approximation

A thin shell approximation for discription of strong shocks is based on two simplifications. First, it is assumed that all swept-up intercloud gas accumulates into the thin shell just behind the shock front and moves with the velocity u . Second, the pressure distribution inside the cavity $P(r, t)$ is taken to be uniform. The equations of mass and momentum conservation may be expressed as follows (Cherny, 1957) :

$$M = M_0 + 4\pi \int_0^R \rho(r)r^2 dr, \quad \frac{d(Mu)}{dt} = 4\pi R^2(P_{\text{in}} - P) + Mg, \quad (1)$$

where M is the mass of the shell, M_0 is the ejected mass, R is the shock radius and u is the gas velocity behind the shock; $\rho(r)$ and P are the density and the pressure of the ambient gas, g is the external gravitational field. For the adiabatic blastwave without gravity

$$1) \frac{dR}{dt} = \frac{\gamma + 1}{2} u \quad \text{or} \quad 2) \frac{dR}{dt} = u; \quad E_0 = E_{\text{th}} + \frac{1}{2}Mu^2, \quad (2)$$

where $E_0 = \text{const}$ is the energy of the explosion, $E_{\text{th}} = \frac{4\pi}{3(\gamma-1)}P_{\text{in}}R^3$ is the thermal energy of the blastwave, and γ is the adiabatic power. Equations (1)- (2) have a simple solution for the homogeneous case if the swept-up mass is

^{*}Space Research Institute, 84/32 Profsoyuznaya, Moscow, 117810, Russia

[†]Main Astronomical Observatory of the National Academy of Sciences of Ukraine 252650, Kiev, Golosiiv, Ukraine

much greater than the ejected one $R = \left[\frac{\xi_0 E_0}{\rho_0} \right]^{1/5} t^{2/5}$, where (Chernyi, 1957, Bisnovaty-Kogan and Blinnikov, 1982)

$$1) \xi_0 = \frac{75(\gamma - 1)(\gamma + 1)^2}{16\pi(3\gamma - 1)}, \quad 2) \xi_0 = \left[\frac{75(\gamma - 1)}{8\pi} \right]^{1/5}. \quad (3)$$

Comparison with an exact self-similar (SS) solution (Sedov, 1946) shows, that the case 2) gives much better precision, $\xi_0^{1/5} = 1.033, 1.014, 1.036$ for (SS), (1), (2) cases respectively at $\gamma = 1.4$; and similarly $\xi_0^{1/5} = 1.15, 1.12, 1.15$ for $\gamma = 5/3$.

2 Three-dimensional shocks

Introduce following Bisnovaty-Kogan and Silich (1991), Silich (1992) (see also Palouš, 1990, Bisnovaty-Kogan and Silich, 1995) Cartesian coordinate system (x, y, z) . Let m is the mass, \mathbf{r} is the radius-vector, \mathbf{u} is the velocity of the particular Lagrangian element of the shock, $\rho(x, y, z) = \rho_0 f(x, y, z)$ is the ambient gas density, \mathbf{n} is the normal to the shock front unit vector, \mathbf{g} is the acceleration of the external gravitational field, \mathbf{V} is the velocity field of the undisturbed gas flow, Σ is the surface area of the Lagrangian element, $m = \sigma \Sigma$, σ is the surface density, $\Delta P = P_{\text{in}} - P$ is the pressure difference between the hot interior and warm (cold) external gas. The pressure $P_{\text{in}} = (\gamma - 1)E_{\text{th}}/\Omega$ of the hot tenuous gas within the cavity is a function of the bubble thermal energy E_{th} and volume Ω . To describe shock expansion we must introduce element of the surface area and define volume of any closed three-dimensional region. It is well known from the differential geometry that any surface may be specified parametrically, with the Cartesian coordinates at any point on the surface being function of two parameters, λ_1 and λ_2 : $x = x(\lambda_1, \lambda_2)$, $y = y(\lambda_1, \lambda_2)$, $z = z(\lambda_1, \lambda_2)$. Then the element of the surface area may be defined by the expression (Budak and Fomin, 1965)

$$d\Sigma = S(\lambda_1, \lambda_2) d\lambda_1 d\lambda_2, \quad (4)$$

$$S(\lambda_1, \lambda_2) = \left\{ \left[\frac{\partial(y, z)}{\partial(\lambda_1, \lambda_2)} \right]^2 + \left[\frac{\partial(z, x)}{\partial(\lambda_1, \lambda_2)} \right]^2 + \left[\frac{\partial(x, y)}{\partial(\lambda_1, \lambda_2)} \right]^2 \right\}^{1/2}, \quad (5)$$

where $\partial(x_i, x_j)/\partial(\lambda_1, \lambda_2) \equiv \det(\partial(x_i, x_j)/\partial(\lambda_1, \lambda_2))$ are the appropriate Jacobians. Similarly could be obtained expressions for Ω and \mathbf{n}

If parameters λ_1 and λ_2 are considered as the Lagrangian coordinates of the shock front, then the equations for 3-D shock propagation may be written for the mass $\mu = \sigma S(\lambda_1, \lambda_2)$ per unit of Lagrangian square, in a compact form, convenient for numerical integration

$$\frac{d\mu}{dt} = \rho\chi, \quad (6)$$

$$\frac{du_x}{dt} = \frac{\Delta P}{\mu} \frac{\partial(y, z)}{\partial(\lambda_1, \lambda_2)} - \frac{u_x - V_x}{\mu} \rho\chi + g_x, \quad (7)$$

$$\frac{du_y}{dt} = \frac{\Delta P}{\mu} \frac{\partial(z, x)}{\partial(\lambda_1, \lambda_2)} - \frac{u_y - V_y}{\mu} \rho\chi + g_y, \quad (8)$$

$$\frac{du_z}{dt} = \frac{\Delta P}{\mu} \frac{\partial(x, y)}{\partial(\lambda_1, \lambda_2)} - \frac{u_z - V_z}{\mu} \rho\chi + g_z, \quad (9)$$

$$\frac{dx}{dt} = u_x, \quad \frac{dy}{dt} = u_y, \quad \frac{dz}{dt} = u_z. \quad (10)$$

Here function χ is defined as follows:

$$\chi = (u_x - V_x) \frac{\partial(y, z)}{\partial(\lambda_1, \lambda_2)} + (u_y - V_y) \frac{\partial(z, x)}{\partial(\lambda_1, \lambda_2)} + (u_z - V_z) \frac{\partial(x, y)}{\partial(\lambda_1, \lambda_2)}. \quad (11)$$

Approximating the shock front by a number N of Lagrangian elements one get a system of $7N$ differential equations of the mass and momentum conservation. This set of equations is coupled with the equation for the gas pressure within the cavity, and the equation of energy for $E_{\text{tot}} = E_{\text{th}} + E_{\text{k}} + E_{\text{g}}$, consisting from the thermal energy of the hot bubble interior, kinetic and gravitational energies of the shell. The kinetic and gravitational energies of the shell are determined by corresponding surface integrals. Variations of the total energy E_{tot} of the remnant throughout the adiabatic stage of evolution are defined by the energy input rate $L(t)$, kinetic and thermal energies of the swept-up interstellar gas with temperature $T(x, y, z)$:

$$E_{\text{tot}} = E_0 + \int_0^t [L(t) + \frac{1}{2} \int_{\lambda_{1,\text{min}}}^{\lambda_{1,\text{max}}} \int_{\lambda_{2,\text{min}}}^{\lambda_{2,\text{max}}} \dot{\mu}(V^2 + 3kT/\eta) d\lambda_1 d\lambda_2] dt, \quad (12)$$

where E_0 is the initially deposited energy, η is the mean mass per particle. On the radiative phase of expansion the gas behind the shock front cools so quickly that does not add to the total energy of the remnant. Rarefied hot gas inside the cavity expands adiabatically and accelerates the surrounding dense shell. Time-derivative of the thermal energy of the remnant is defined then by the equation, which is used instead of (12):

$$\frac{dE_{\text{th}}}{dt} = L(t) - \int_{\lambda_{1,\text{min}}}^{\lambda_{1,\text{max}}} \int_{\lambda_{2,\text{min}}}^{\lambda_{2,\text{max}}} P_{\text{in}} u_n S(\lambda_1, \lambda_2) d\lambda_1 d\lambda_2, \quad (13)$$


where u_n is the velocity component normal to the shock front. Calculations of supershell formation in the plane-stratified differentially rotating Galactic disk have been performed by Silich *et al.* (1994) and are represented in the Fig. 1. An hourglass remnant with a noticeable degree of deformation by the Galactic shear have appeared. Formation of elongated superbubbles due to differential galactic rotation gives a possibility to determine unambiguously a direction of the galactic rotation (Mashchenko and Silich, 1995). The means of this determination is shown in the Fig. 2.

Figure 1: Galactic superbubble morphology for different locations of the parent OB-association relative to the galactic plane. Left: the OB-association is at the midplane of the Galaxy. Right: the OB-association is 50pc above the Galactic plane.

Figure 2: Scheme of the HI holes orientation in the plane of view.

References

- [1] Bisnovaty-Kogan G. S. and Blinnikov S.I. 1982, *Astron. Zh.* **59**, 876 [*Sov. Astron.* **26**, 530 (1982)].
- [2] Bisnovaty-Kogan, G. S. and Silich S.A., 1991, *Astron. Zh.* **68**, 749 [*Sov. Astron.* **35**, 370 (1991)].
- [3] Bisnovaty-Kogan, G. S. and Silich S.A., 1995, *Rev. Mod. Phys.* **67**, 661.
- [4] Budak B. M. and Fomin S.V. 1967, *Multiple integrals and series* (Nauka, Moscow).
- [5] Chernyi G. G., 1957, *Doklady Acad. Sci. USSR* **112**, 213.
- [6] Mashchenko S. Ya. and Silich S.A. 1995, *Astron. Zh.* **72**, 660.
- [7] Palouš J., 1990, in *The Interstellar Disk-Halo Connection in Galaxies*, edited by H. Bloemen (Srerrewacht, Leiden) p. 101.
- [8] Sedov L. P., 1946, *Dok. Akad. Nauk SSSR*, **42**, 17.
- [9] Silich, S. A., 1992, *Astrophys. Space. Sci.* **195**, 317.
- [10] Silich S. A., Franco J. Palouš, J. and Tenorio-Tagle G. 1994, in *Violent Star Formation from 30 Doradus to QSOs*, edited by G. Tenorio-Tagle (Cambridge Univ. Press, Cambridge) p. 162.



This figure "FIG2.GIF" is available in "GIF" format from:

<http://arxiv.org/ps/astro-ph/9707188v1>