A High Deuterium Abundance at $z=0.7^{1}$

J. K. Webb^{2,3}, R. F. Carswell⁴, K. M. Lanzetta⁵, R. Ferlet⁶, M. Lemoine⁷, A. Vidal-Madjar⁶, D. V. Bowen⁸

¹ Based on observations with the NASA/ESA Hubble Space Telescope, obtained at the Space Telescope Science Institute, which is operated by the Association of Universities for Research in Astronomy, Inc., under NASA contract NAS5–26555.

² School of Physics, University of New South Wales, Sydney, NSW 2052, Australia

³ Astronomy Centre, Sussex University, Falmer, Brighton BN1 9QH, England

⁴ Institute of Astronomy, Madingley Road, Cambridge, CB3 0HA, England

⁵ Astronomy Program, Department of Earth and Space Sciences, State University of New York at Stony Brook, Stony Brook, NY 11794–2100, USA

⁶ Institut d'Astrophysique, 98 bis Boulevard Arago, Paris 75014, France

⁷ Department of Astronomy & Astrophysics, Enrico Fermi Institute, The University of Chicago, Chicago IL 60637–1433, USA

⁸ Royal Observatory Edinburgh, Blackford Hill, Edinburgh EH9 3HJ, Scotland

Of the light elements, the primordial abundance of deuterium, $(D/H)_p$, provides the most sensitive diagnostic¹ for the cosmological mass density parameter Ω_B . Recent high redshift D/H measurements are highly discrepant^{2,3,4,5,6}, although this may reflect observational uncertainties^{7,8}. The larger D/H values, which imply a low Ω_B and require the Universe to be dominated by non-baryonic matter (dynamical studies indicate a higher total density parameter), cause problems for galactic chemical evolution models since they have difficulty in reproducing the large decline down to the lower presentday D/H. Conversely, low D/H values imply an Ω_B greater than derived from ⁷Li and ⁴He abundance measurements, and may require a deuterium abundance evolution that is too low to easily explain. Here we report the first measurement at intermediate redshift, where the observational difficulties are smaller, of a gas cloud with ideal characteristics for this experiment. Our analysis of the z = 0.7010 absorber toward 1718+4807 indicates D/H = $2.0 \pm 0.5 \times 10^{-4}$ which is in the high range. This and other independent observations suggests there may be a cosmological inhomogeneity in (D/H)_p of at least a factor of ten.

Measurements can be achieved using absorption line spectroscopy of gas clouds intersecting the sight-lines to distant, background quasars by measuring the strengths of the absorption lines of the H I and D I Lyman series. The D I transitions occur at an isotopic shift of -81 km s^{-1} with respect to the H I series. Such a measurement is observationally difficult for a number of reasons^{9,10}: (1) ill-placed H I Ly α -forest absorption lines can masquerade as deuterium and (2) most QSO absorption

systems exhibit complex internal velocity structure, which may render parameter estimation for individual components of interest unreliable or impossible. To overcome the first difficulty, we searched for suitable absorbers at intermediate redshifts (i.e. at z < 1), where the number density of Ly α -forest systems is far lower than at high redshifts. To overcome the second difficulty, we searched for suitable absorbers with apparently simple velocity structure. We found one object which appears optimal for a D/H analysis; only one absorbing component is revealed by the data and the velocity dispersion in that component is small enough to easily detect and accurately measure the deuterium abundance.

The absorber at redshift z = 0.7010 toward the QSO 1718+4807 ($z_{\rm em} = 1.084$, $m_V = 15.3$) was selected on the basis of a remarkably abrupt partial Lyman-limit discontinuity in the International Ultraviolet Explorer (IUE) spectrum¹¹. The extreme sharpness of the Lyman break clearly indicates simple velocity structure and low velocity dispersion parameter. This simple velocity structure contrasts with the complex nature of the higher redshift systems reported so far, both those which give low D/H measurements^{2,3,7} and those which give high values^{4,5,6} so the measured column densities (based on Ly α and the Lyman limit alone) are likely to be more reliable than in those cases. Hubble Space Telescope (HST) spectra were obtained of the spectral region covering the Lyman- α and Si III transitions using the Goddard High Resolution Spectrograph and the G270M grating. All spectra were extracted, binned to linear wavelength scales and corrected to vacuum, heliocentric wavelengths using standard procedures. Absorption-line parameter estimates were derived using VPFIT¹², a computer program based on Gauss-Newton unconstrained optimisation. Parameter errors were derived from the diagonal terms of the parameter covariance matrix computed at the best fit.

The D/H we derive from the combined data is $D/H = 2.0\pm0.5\times10^{-4}$ which is approximately an order of magnitude larger than recent ground-based measurements of high-redshift absorbers^{2,3,7} (although the reliability of these measurements is presently a matter of debate⁸) and an order of magnitude larger than a recent intermediate redshift measurement¹³ at z=0.5. It is of course possible that our high value is caused by a weak H I line which just happens to fall very close to the D I wavelength. Unfortunately, *a posteriori* statistics do not provide a reliable means of assessing the probability of this because the absorption cloud we have studied was specifically selected as having uniquely simple velocity structure, a property which itself implies a low probability of an H I interloper successfully mimicking D I. We note however that the probability of a *randomly placed* interloper mimicking deuterium is extremely small; using the observed number density of absorbers in the spectrum and adopting a generous 4-sigma tolerance on the line position, it is less than 1 percent.

To examine this possibility further, we re-fitted the data replacing D I with H I (so that the redshift of the putative interloper was a free parameter). The best fit resulted in the new H I line falling at -86 ± 5 km s⁻¹ from the strong component, implying that the detected absorption feature *is* deuterium and not an interloper. Furthermore, the Doppler parameter of this putative H I interloper ($b = 21 \pm 4$ km s⁻¹) lies between that of the strong H I component but greater than

the Si III value, which is again consistent with the expectation for D I. Taken together, these points indicate that the most reasonable and likely interpretation of the data is that we have detected D I.

A further possibility is that some other transition from an absorbing cloud at some other redshift just happens to coincide with the D I wavelength at $z_{abs} = 0.701024$. The closest candidate is ISM Cr II $\lambda 2066.16$. The D I feature falls at 2067.31 Å. This corresponds to a shift of ~ 170 km s⁻¹, thus so Cr II $\lambda 2066.16$ is not a serious contender for D I contamination. Furthermore, Cr II $\lambda 2056$ is not observed and yet has an oscillator strength 2 times higher than Cr II $\lambda 2066.16$. We conclude that no Cr II contamination occurs.

We cannot reliably estimate the heavy element abundances of the $z_{abs} = 0.701024$ system, since the only species in the observed range is Si III (log $N(\text{Si III}) = 12.81 \pm 0.04 \text{ cm}^{-2}$). An upper limit can be derived, but this only constrains the abundances to be less than solar; they could easily be very much lower. We note that it would be astonishing if the $z_{abs} = 0.701024$ system turned out to have already undergone substantial chemical evolution and yet exhibit such a high D/H. Future observations of any unsaturated absorption lines in this system (potentially Mg II or Fe II, for example) should yield reliable heavy element abundance constraints. Astration (deuterium destruction in stars) means that any particular D/H detection provides a *lower* limit to the primordial value. Therefore, because our measured D/H is high, the lack of information about the heavy element abundances does not alter the interpretation of the data described below.

In standard models of galactic chemical evolution, the mass fraction of deuterium decreases steadily with time, by a factor ~ 2 – 3 over 10 Gyr, so that when plotted versus metallicity, it appears constant until a metallicity 1/10 solar and drops abruptly thereafter¹⁴. The two extremes of the recent high redshift D/H observations imply radically different consequences for the baryonic density parameter, Ω_B , and its cosmological significance^{15,16}. A high (D/H)_p would be in excellent agreement with standard homogeneous BBN and the observed "primordial" abundances of ⁴He and ⁷Li, but observations of D/H in the interstellar medium then require destruction of deuterium by a factor ~ 10 up to the present epoch. Whether this much astration can be easily explained by chemical evolution models is unclear, although some recent models may succeed¹⁷.

A ratio D/H = 2.0×10^{-4} , as obtained here toward Q1718+4807, corresponds to a mass fraction of ⁴He $Y \simeq 0.233 \pm 0.002$, ⁷Li/H $\simeq 1.8^{0.7}_{0.6} \times 10^{-10}$ and $\Omega_B \simeq 0.006 \pm 0.003 \ h^{-2}$, where hdenotes the value of the Hubble constant in units of 100 km s⁻¹ Mpc⁻¹ and the errors only include 1σ errors on the nuclear cross-sections of BBN (values were derived using useful fitting formulae¹⁸). Remarkably, these figures agree precisely with the measured abundances, $Y \simeq 0.233 \pm 0.005$ and ⁷Li/H $\simeq 1.8^{+0.5}_{-0.3} \times 10^{-10}$. The above baryonic mass density parameter may be compared with the measured amount of visible matter in galaxies, $\Omega_{\rm vis} \sim 0.002 - 0.005 \ h^{-1}$. This shows that a high primordial deuterium abundance leaves little room for *baryonic* dark matter and supports the view that the missing mass inferred from dynamical studies must be non-baryonic. On the other hand, a low (D/H)_p implies BBN values of ⁴He and ⁷Li/H which differ from the measured values by amounts corresponding to 2σ observational limits (although it avoids the difficulty in explaining such rapid deuterium destruction, but if too low, may still be problematic if the implied astration is too small compared to predictions).

Several explanations have been put forward which could explain discrepant values of the D/H ratio. Post-BBN chemical evolution processes might result in strong deuterium depletion in some gas clouds, giving rise to low observed abundances. By redshifts $z \approx 3 - 4$, stars down to $\sim 2M_{\odot}$ have had time to eject deuterium-poor gas. However, the observational upper limits on C/H and Si/H in clouds at high redshift where D/H $\sim 2.5 \times 10^{-5}$ imply metal abundances less than $\sim 10^{-2}$ solar, constraining the fraction of gas which has been cycled through stars to be less than about 5%. Such models appear to offer reasonable compatibility with light element abundance observations only by admitting unreasonable stellar populations, such as a primordial population of supermassive stars of mass $M \gtrsim 1000M_{\odot}$, or an initial mass function strongly peaked around $M \approx 6M_{\odot}$.

Alternatively, primordial isocurvature baryon fluctuations¹⁹ could account for a variation of the $(D/H)_p$ ratio by a factor ~ 10 but only on scales corresponding to a Jeans mass of $M_J \sim 10^5 - 10^6 M_{\odot}$. For a specific class of these models, there are opposing views as to whether or not the observed isotropy of the cosmic microwave background rules out substantial *intrinsic* D/H fluctuations¹⁹⁻²². We note, however, that homogeneous, critical-density models with a low baryonic component, $\Omega_B < 0.01 h^{-2}$, appear to predict degree–scale microwave background fluctuations which are significantly below those actually observed²³. Also, assuming no segregation between baryons and dark matter²⁴, X-ray observations of galaxy clusters support this point²³ since for critical-density models they independently suggest baryonic density parameters of $\Omega_B > 0.02 h^{-3/2}$. Both of these types of observation therefore appear to be in conflict with a baryonic density parameter for a homogeneous universe as low as that derived from our results for D/H in the $z_{\rm abs} = 0.701024$ gas cloud reported here. Therefore our results indicate that either the universe does not have a critical total density, or, if it does, Big-Bang nucleosynthesis must have occurred inhomogeneously.

1. Epstein, R.I., Lattimer, J.M., Schramm, D.N., 1976, Nature, 263, 198-202. The Origin of Deuterium.

2. Tytler, D., Fan, X-M., Burles, S., 1996, Nature, 381, 207-209. Cosmological baryon density derived from the deuterium abundance at redshift z = 3.57.

3. Burles, S., Tytler, D., 1996, Science, submitted. Cosmological deuterium abundance and the baryon density of the universe.

4. Rugers, M., & Hogan, C.J., 1996, Astrophys. J., 469, L1-L4. Confirmation of High Deuterium Abundance in Quasar Absorbers

5. Carswell, Rauch, Weymann, Cooke, Webb, 1994, Mon. Not. R. Astron. Soc., 268, L1-L4. Is there deuterium in the z = 3.32 complex in the spectrum of 0014+813?

6. Rugers, M., Hogan, C.J., 1996, Astron. J., 111(6), 2135-2140. High Deuterium Abundance in a New Quasar Absorber

7. Tytler, D., Burles, S., Kirkman, D., 1996, preprint astro-ph/9612121. New Keck spectra of Q0014+813:

annulling the case for high deuterium abundances.

8. Songaila, A., Wampler, E.J., Cowie, L.L, 1997, *Nature*, 385, 137-139. A high deuterium abundance in the early Universe.

9. Webb, J.K., Carswell, R.F., Irwin, M.J., Penston, M.V., *Mon. Not. R. Astron. Soc.*, **250**, 657-665, 1991. On measuring the deuterium abundance in QSO absorption systems

10. Wampler, E.J., Nature, 383, 308, 1996. Alternative hydrogen cloud models.

11. Lanzetta, K.M., Turnshek, D., Sandoval, J., 1993, Astrophys. J. Suppl., 84, 109-184. Ultraviolet spectra of QSOs, BL Lacertae objects, and Seyfert galaxies

12. Webb, J.K, PhD thesis, Cambridge University, 1987. QSO absorption lines.

13. Vidal-Madjar, A., Ferlet, R., Lemoine, M, 1996, preprint astro-ph/9612020. Deuterium abundance and cosmology.

14. Pagel, B.E.J., Nucleosynthesis and Chemical Evolution of Galaxies, Cambridge University Press 1997.

15. Schramm, D.S., Turner, M.S., Nature, 1996, 381, 193-194. Deuteronomy and numbers.

16. Hata, N., Steigman, G., Bludman, S., Langacker, P., 1996, preprint astro-ph/9603087. Cosmological implications of two conflicting deuterium abundances.

17. Vangioni-Flam, E., Cassé, M., 1995, *Astrophys. J.* **441**, 471-476. Cosmological and astrophysical consequences of a high primordial deuterium abundance

18. Hogan, C.J., preprint, astro-ph/9609138. Big bang nucleosynthesis and the observed abundances of light elements.

19. Copi, C.J., Olive, K.A., Schramm, D.N., 1996, preprint astro-ph/9606156. Implications of a primordial origin for the dispersion in D/H in quasar absorption systems.

20. Copi, C.J., Schramm, D.N., Turner, M.S., 1995, *Science*, **267**, 192. Big-Bang Nucleosynthesis and the Baryon Density of the Universe.

21. Jedamzik, K., Fuller, G.M., 1995, *Astrophys. J.*, **452**, 33. Nucleosynthesis in the Presence of Primordial Isocurvature Baryon Fluctuations.

22. Jedamzik, K., Fuller, G.M., 1996, preprint astro-ph/9609103. Is the deuterium in high redshift Lyman limit systems primordial?

23. White, M., Viana, P.T.P., Liddle, A.R., Scott, D., 1996, Mon. Not. R. Astron. Soc., 283, 107. Cold dark matter models with high baryon content

24. White, S.D.M., Navaro, J.F., Evrard, A.E., Frenk, C.S., 1993, *Nature*, **366**, 429. The Baryon Content of Galaxy Clusters: A Challenge to Cosmological Orthodoxy.

ACKNOWLEDGEMENTS. KML acknowledges support by NASA, STScI, and NSF. ML acknowledges support by the NASA, DoE, and NSF at the University of Chicago. JKW is grateful for valuable discussions with M. Ashley, J. Barrow, A. Liddle, and R.J. Tayler and to SUN Microsystems Australia Pty Ltd for providing computing facilities for this work.

Fig. 1.— Data and fits for 1718+4807. The top panel illustrates the HST H I and D I profile, showing the model profiles for H I (dotted line), D I (dashed line) and H I+D I (solid line). The middle panel illustrates the HST Si III λ 1206.51 profile and fit. The HST spectrum was observed on 3/3/1995 (total integration time 282 minutes). The spectral resolution is 20,000. The bottom panel illustrates the IUE spectrum and fit. The pixel size is 1.18Å and the adopted spectral resolution is 2.5 pixels. To minimise the number of free parameters, we adopted a single redshift for H I, D I and Si III lines and the Lyman limit. The parameter constraints arise as follows. The IUE spectrum provides an accurate determination of the H I column density $(\log N(\text{H I}) = 17.24 \pm 0.01)$ $\rm cm^{-2}$). The Si III λ 1206 absorption line supports the single component velocity structure implied by the IUE Lyman limit and determines the cloud redshift precisely $(z_{abs} = 0.701024 \pm 0.00007)$. The Ly α absorption is clearly asymmetric, showing additional absorption in the blue wing at the position corresponding to D I. Since Si III accurately constrains the D I position and because the Doppler parameter (which is equal to $\sqrt{2}$ times the RMS velocity dispersion) b(D I) is constrained by the overall fit, only one free parameter, N(D I), is required to fit the excess absorption seen at the D position. The D I column density is thus accurately determined and is $\log N(\text{D I}) =$ 13.57 ± 0.06 cm⁻². The Doppler parameters are dominated by non-thermal broadening with an inferred temperature of 1.9×10^4 K and b(H I) = 25.5 km s⁻¹ and b(D I) = 22.2 km s⁻¹ and $b(\text{Si III}) = 18.7 \text{ km s}^{-1}$, with the same error on each of $\sigma = 0.5 \text{ km s}^{-1}$. From the above, we derive $D/H = 2.0 \pm 0.5 \times 10^{-4}.$

