A unified model for the spectral variability in GRS 1915+105

Τ.	Belloni ¹ ,	Μ.	$Méndez^{1,2}$, A.R.	$King^3$,	Μ.	van	der	$Klis^1$,	J.	van	Para	$\mathrm{dijs}^{1,4}$

accepted.

ApJ Letters, accepted for publication, 1997 August 12th

¹Astronomical Institute "Anton Pannekoek", University of Amsterdam and Center for High-Energy Astrophysics, Kruislaan 403, NL-1098 SJ Amsterdam, the Netherlands

²Facultad de Ciencias Astronómicas y Geofísicas, Universidad Nacional de La Plata, Paseo del Bosque S/N, 1900 La Plata, Argentina

³Astronomy Group, University of Leicester, Leicester LE1 7RH, United Kingdom

⁴Physics Department, University of Alabama in Huntsville, Huntsville, AL 35899, USA

ABSTRACT

We have analyzed the spectral variations of the superluminal black-hole X-ray binary GRS 1915+105 by using data obtained with the PCA on the Rossi XTE. We find that, despite the marked differences in the structure and the time scale of variability, all spectral changes can be attributed to the rapid disappearing of the inner region of an accretion disk, followed by a slower re-filling of the emptied region. The time scale for each event is determined by the extent of the missing part of the disk. The observed relation between the duration of an event and the radius of the disappearing region matches remarkably well the expected radius dependence of the viscous time scale for the radiation-pressure dominated region of an accretion disk.

Subject headings: accretion, accretion disks — binaries: close — black hole physics — instabilities — X-rays: stars — stars: individual GRS 1915+105

1. INTRODUCTION

The X-ray source GRS 1915+105 is the best of the two only examples of galactic objects that show superluminal expansion in radio observations (Mirabel & Rodríguez 1994). It is located at a distance of 12.5 kpc and its relativistic jets are directed at 70° from the line of sight (Mirabel & Rodríguez 1994). After its discovery as an X-ray transient in 1992 (Castro-Tirado, Brandt, & Lund 1992), a number of outbursts has been reported, although it is possible that the source has never completely switched off in between. The Rossi X-ray Timing Explorer (RXTE) started observing GRS 1915+105 in a bright and variable state in April 1996 and continued to observe it regularly at least once per week since then. During this period, the source has displayed a remarkable richness in variability, ranging from quasi-periodic burst-like events, deep regular dips and strong quasi-periodic oscillations, alternated with quiescent periods (Greiner, Morgan & Remillard 1996,Morgan, Remillard, & Greiner 1997,Chen, Swank & Taam 1997,Belloni et al. 1997,Taam, Chen & Swank 1997). Because of its similarities with the other galactic superluminal source GRO 1655-40 (Zhang et al. 1994), whose dynamical mass estimate implies the presence of a black hole (Bailyn et al. 1995), and because of its high luminosity exceeding the Eddington limit for a neutron star, the source is suspected to host a black hole.

In a previous paper (Belloni et al. 1997, hereafter Paper I) we showed that the complicated light curve of GRS 1915+105 can be described by the rapid (\sim 1 s) appearance and disappearance of emission from an optically thick inner accretion disk. In this paper we report on the results of the analysis of a later observation which allowed a more detailed investigation of the spectral variability of the source. We find that the variation time scale $t_{\rm rec}$ of the disk source is completely specified by the maximum size $R_{\rm max}$ of it inner radius. The relation between these quantities is precisely as expected for the viscous evolution of a radiation-pressure dominated accretion disk.

2. X-RAY OBSERVATIONS

Among the many PCA observations of GRS 1915+105, that of 1997 June 18th deserves particular attention, since it reproduces within one day most of the timing behaviors observed from the source. The data consist of two segments of roughly one hour each. Since the two segments are essentially similar, we will restrict ourselves to the second one, which starts at 14:36 UT and ends at 15:35 UT. A 1200 s representative stretch of the 2.0-40.0 keV light curve, binned to 1 second, can be seen in the top panel of Figure 1. It consists of a sequence of 'bursts' of different duration with quiescent intervals in between. All bursts start with a well-defined sharp peak and decay faster than they rise. The longer bursts show oscillation (or sub-bursts) towards the end. In the following we define an event as a full cycle of one quiescent interval followed by one burst. With this definition there is no interval between events. In order to maintain precision. separate events are considered only as those whose quiescent count rate goes below 10000 cts/s in the total 2.0-40.0 keV band. We measured the start of an event (corresponding to the end of the previous one) as the time of the small dip at the end of the decay (see Figure 1). Since all the bursts start with a sharp peak, the time of the peak can be taken as the separation between the quiescent phase and the burst. Finally, the bottom level of each quiescent part has been measured as the lowest level excluding the small dip mentioned above. With these numbers it is possible to associate four quantities to an event: total length, length of the quiescent part, length of the outburst and level of the bottom. Two strong correlations emerge: (i) the longer the event, the lower the bottom level (see below); and (ii) a one-to-one relation between quiescent and burst duration (Figure 2). No correlation is seen between a burst duration and the following quiescent interval.

In order to follow the spectral evolution, we produced two X-ray colors: HR1 = B/A and HR2 = C/(A+B), where A, B and C are the counts in the 2.0–5.0 keV, 5.0–13.0 keV and 13.0–40.0 keV intervals respectively. We produced color-color diagrams by selecting events according to their total length. It turns out that there is a very strong correlation between the pattern traced out in the CC diagram by each event, and its duration. In Figure 3 one can see two extreme cases and

an intermediate one. All other events follow a similar pattern. During short events (top panel) the quiescent period is at the bottom left in the C-C diagram, then the source moves up to the top left during peak, then down and right to the dip. In longer events the sequence is the same, with the only difference that the quiescent part is located farther to the right the longer the event is. The small oscillations at the end of long bursts show up as oscillations around the peak points in the C-C diagram (mid and bottom panels in Figure 3).

The spectral evolution is extremely reproducible throughout the different events. We therefore divided the C-C diagram in regions 0.01 wide in HR2 and 0.1 wide in HR1 and accumulated energy spectra for all the (74) populated regions of the diagram. The spectra were accumulated in 48 energy bins with 1 second time resolution. The background subtraction was performed with a background spectrum extracted from a blank sky observation and normalized to the highest channels where the source signal is absent. We used the latest detector response matrix available (April 1997, Version 2.1.2). A systematic error of 2% has been added to the data (see Paper I). In order to avoid possible problems due to the background subtraction, we limited our fits to energies below 30.0 keV. Following paper I, for each of the regions we fitted the data with a "standard" spectral model for black-hole candidates, consisting of a disk-blackbody model (Mitsuda et al. 1984) plus a power law. A broad Gaussian line was added at a fixed energy of 6.3 keV as in Paper I. We found the interstellar absorption to be stable around 4.5×10^{22} cm⁻², so it was fixed to that value. Notice that this is lower than the value found in paper I because of the improved calibration. It is slightly lower than the best-fit ROSAT value (Greiner, Morgan & Remillard 1996). The resulting parameters for the power law component depend on the inclusion of the Gaussian line mentioned above, but the disk-blackbody parameters are independent of this. In the following we will restrict ourselves to the discussion of the latter. Since both distance and inclination of the system are known (the latter is assumed to be equal to the observed inclination of the radio jet, see Mirabel & Rodríguez 1994), it is possible to derive the inner radius of the accretion disk directly from the model without further assumptions.

The fitting procedure yielded spectral parameters for each of the 74 regions. By identifying

the region of each 1-s bin, it is possible to produce a time history of relevant spectral parameters with a resolution of 1 second. The inner radius and temperature are shown in the bottom two panels of Figure 1. As it can be seen, during bursts the temperature is above 2 keV and the radius stable around 20 km, parameters consistent with those reported in Paper I. During quiescent phases, the temperature drops to lower values and the radius increases. There is a strong correlation between the length of the event and the largest radius reached (Figure 4). In order to check the results of our 'grid' procedure, we extracted spectra for the longest event in 16 s bins from the Standard 2 data, which have a much better energy resolution. We obtained similar results. Since the spectral parameters coming from these data are more accurate, we used those for the estimate of the values of the accretion rate. Notice that the radii determined for the disk hole are smaller than that reported in Paper I, primarily due to the new PCA calibrations. Because of these differences, the behavior of the accretion rate derived from the model fits (which depends on the third power of the inner radius) is now radically different: the value of M during the quiescent phase is around $1.7-10\times10^{-8}\mathrm{M}_{\odot}/\mathrm{yr}$ (the range being determined by the spin of the central black hole, which determines the innermost stable orbit), while in the burst phase it is around a factor of two higher and variable.

We produced similar color-color diagrams for a number of other RXTE observations of GRS 1915+105. All observations we analyzed follow the pattern outlined above, except that of 1996 June 16th (Greiner, Morgan & Remillard 1996). All the quasi-periodic bursts observed in many observations (see Taam, Chen & Swank 1997) are consistent with repetitive short events like the ones described here.

3. DISCUSSION

The results described in the previous section can be interpreted within the model presented in Paper I, providing a unified picture of the variability observed in GRS 1915. In Paper I, we modeled the large amplitude changes as emptying and replenishing of the inner accretion disk caused by a viscous-thermal instability. The small radius observed during the quiescent period

was identified with the innermost stable orbit around the black hole, while the large radius during the burst phase was the radius of the emptied section of the disk. The smaller oscillations were interpreted as failed attempts to empty the inner disk. As it can be seen from Figure 1, from this observation we find that all variations, from major events like the ones described in Paper I to small oscillations observed at the end of a large event, can be modeled in exactly the same fashion (see Figure 1). In this scenario, the "flare state" presented in Paper I is simply a sequence of small events following a big one, similar to the small oscillations in Figure 1. Both the spectral evolution and the duration of the event are determined by one parameter only, namely the radius of the missing inner section of the accretion disk. For a large radius, the drop in flux will be large and the time needed for re-fill will be long. Following Paper I, it is natural to associate the length of the quiescent part of an event t_q to the viscous time scale of the radiation-pressure dominated part of the accretion disk. This can be expressed as $t_{\text{visc}} = R^2/\nu$, where $\nu = \alpha c_S H$. From Frank, King & Raine (1992) the scale-height H and sound speed c_S can be found, leading to

$$t_{\text{visc}} = 30\alpha_2^{-1} M_1^{-1/2} R_7^{7/2} \dot{M}_{18}^{-2}$$
 (1)

where $\alpha_2 = \alpha/0.01$, R_7 is the radius in units of 10^7 cm, M_1 the central object mass in solar masses, and \dot{M}_{18} is the accretion rate in units of 10^{18} g/s. Notice that even the largest radii derived here are well within the radiation-pressure dominated part of the disk (see Equation 2 in Paper I). The line in Figure 4 represents the best fit to the data with a relation of the form $t_{\rm q} \propto R^{7/2}$. The fit is excellent, with the exception of the point corresponding to the longest event. The qualitative agreement with the theoretical expectation is striking, although by substituting the appropriate values for the mass and accretion rate we find that our best fit predicts rather small values of α_2 (0.004 and 0.05 for the Schwarzschild and extreme Kerr cases respectively). This indicates a small viscosity in the disk, although we stress that t_{visc} is only a time scale, so that additional corrections might be necessary in order to allow a precise quantitative comparison.

An event can therefore be pictured in the following way. At the start of a quiescent period, the disk has a central hole, whose radius is R_{max} . The hole is either empty or filled with gas whose radiation is too soft to be detected. Slowly the disk is re-filled by a steady accretion rate

 \dot{M}_0 from outside. Each annulus of the disk will move along the lower branch of its S-curve in the $\dot{M}-\Sigma$ plane trying to stabilize at \dot{M}_0 (see Paper I). The surface gravity increases as the annulus moves towards the unstable point at a speed determined by the local viscous time scale. During this period, no changes are observed in the radius of the hole, since all the matter inside does not radiate in the PCA band. The observed accretion rate is \dot{M}_0 . At some point, one of the annuli will reach the unstable point and switch to the high- \dot{M} state, where the accretion rate is larger than \dot{M}_0 , causing a chain-reaction that will "switch on" the inner disk. The observed accretion rate is now higher than the external value \dot{M}_0 . A smaller, hot radius is now observed. At the end of the outburst, the inner disk runs out of fuel and switches off, either jumping back to the $\dot{M} < \dot{M}_0$ state or emptying completely. A new hole is formed and a new cycle starts. Notice that in this scenario the more "normal" state for the source is the one at high count rates, where the disk extends all the way to the innermost stable orbit: in this state the energy spectrum is similar to that of conventional black-hole candidates (see Tanaka & Lewin 1995).

Not only the start and end of a major burst, but also all the small amplitude oscillations within a burst show the same timing signature of decaying faster than they rise. This is in agreement with what was already noticed in Paper I: the rise time is determined by the speed at which a heating wave moves through the central disk, while the faster decay time is due to the rapid fall of matter into the black hole (or into a relativistic jet).

Chen, Swank & Taam (1997) found that when HR2 exceeds 0.1, the power density spectrum is similar to the one observed in black-hole transient systems during the Very High State (see Van der Klis 1995). In these occasions, a strong 1-6 Hz QPO peak was found, positively correlated with the count rate. Notice that the limit HR2>0.1 is an indication that the source was in a quiescent state. Our spectral results show that the fast timing features (both QPO and band-limited noise, see Morgan, Remillard, & Greiner 1997) cannot originate from the innermost regions of the optically thick accretion disk, since those are missing during the quiescent phases. The fact that the QPO frequency increases with count rate is in qualitative agreement with the model, since a higher count rate indicates a smaller inner disk radius, and therefore shorter time scales.

The radii for the disappearing region of the disk found here are considerably smaller than that reported in Paper I. This is entirely due to the improved knowledge of the spectral response of the PCA. During the burst phase the accretion rate through the optically thick accretion disk is found to be higher than in the quiescent phase, contrary to what we reported in Paper I.

Although we can model the spectral evolution and the time scale of the events, most of the variability is yet to be explained. The main question to be answered is: what determines the length of the following outburst? Or, in more physical terms, what determines how large the next missing section of the disk will be? In some observations the events are very regular, in some others they are extremely random, and in some others no events are observed at all. In the observation reported here a striking one to one relation between quiescent and burst time is observed, a relation which applies to the "outburst" and "quiescent" states of the observation presented in Paper I but is obviously not satisfied in other observations (see e.g. Taam, Chen & Swank 1997) nor during the "flare" state of Paper I. Moreover, as already mentioned, one observation among the ones we analyzed does not fit this pattern and requires a different interpretation (1996 June 16th). Nevertheless, the model sketched here provides a satisfactory interpretation of the cause of the changes in the X-ray emission. It fits not only the energy spectral distribution, but also the complex variability shown by GRS 1915+105.

MM is a fellow of the Consejo Nacional de Investigaciones Científicas y Técnicas de la República Argentina. This work was supported in part by the Netherlands Organization for Scientific Research (NWO) under grant PGS 78-277 and the Netherlands Foundation for Research in Astronomy (ASTRON) under grant 781-76-017. ARK gratefully acknowledges a PPARC Senior Fellowship. JVP acknowledges support from NASA under contract NAG 5-3003. We thank Keith Jahoda for helping with the calibration of the PCA data, and the RXTE team for making these data publicly available. We thank D. Battacharya for his careful reading of the manuscript.

REFERENCES

Bailyn, C. D., Orosz, J. A., McClintock, J. E., & Remillard, R. A. 1995, Nature, 378, 157

Belloni, T., Méndez, M., King, A.R., van der Klis, M, & van Paradijs, J., 1997, ApJ, 479, L145

Castro-Tirado, A. J., Brandt, S., & Lund, S. 1992, IAU Circ. 5590

Chen, X., Swank, J.H., & Taam, R.E., 1997, ApJ, 477, L41

Frank, J., King, A., & Raine, D. 1992, "Accretion power in Astrophysics", Cambridge University Press, Cambridge

Greiner, J., Morgan, E., & Remillard, R.A. 1996, ApJ, 473, L107

Mitsuda, K., et al. 1984, PASJ, 36, 741

Mirabel, I. F., & Rodríguez, L. F. 1994, Nature, 371, 46

Morgan, E., Remillard, R. A., & Greiner, J. 1997, ApJ, 482, 993

Taam, R.E., Chen, X., & Swank, J.H., 1997, ApJ, 485, L83

Tanaka, Y., & Lewin, W.H.G., 1995, in "X-ray binaries", eds. Lewin, W.H.G., Van Paradijs, J., & Van den Heuvel, E.P.J., Cambridge Univ. Press., Cambridge, p2.

Van der Klis, M., 1995, in "X-ray binaries", eds. Lewin, W.H.G., Van Paradijs, J., & Van den Heuvel, E.P.J., Cambridge Univ. Press., Cambridge, p252.

Zhang, S. N., Wilson, C. A., Harmon, B. A., Fishman, G. J., Wilson, R. B., Paciesas, W. S., Scott, M., & Rubin, B. C. 1994, IAU Circ. 6046

This manuscript was prepared with the AAS LATEX macros v4.0.

- Fig. 1.— Upper panel: 2.0–40.0 keV PCA light curve. Time zero corresponds to 1997 June 18th 14:36 UT. Middle and lower panels: corresponding inner radius and temperature (see text).
- Fig. 2.— Correlation between burst and quiescent time (as defined in the text). The line is a y=x relation.
- Fig. 3.— Three color-color diagrams for events grouped by total length.
- Fig. 4.— Correlation between total length of the event and maximum inner radius of the disk. The line is the best fit with a power law with fixed index $\gamma = 3.5$ and normalization 1.44×10^{-5} . The last point has been excluded from the fit.







