$Ly\alpha$ absorbers arising in galaxy clusters

A. Ortiz-Gil^{1,2}, K.M. Lanzetta², J.K. Webb¹, X. Barcons³

¹Dept. of Astrophysics & Optics, UNSW, Sydney 2052, NSW, Australia

²Dept. of Physics & Astronomy, SUNY, Stony Brook, NY 11794-3800, USA

³IFCA, CSIC-UC, Avda. de los Castros s/n, E-39005, Santander, Spain

Abstract. We present here new GHRS observations of $Ly\alpha$ absorption lines associated with groups or clusters toward the QSOs 1545+2101 and 0850+4400. In the first case we have identified at least eight distinct $Ly\alpha$ absorption features, with a mean redshift of $\langle z \rangle = 0.2648 \pm 0.0002$ and a velocity dispersion of 163 ± 57 km s⁻¹. We have also identified a group or cluster of galaxies in the vicinity of this QSO with a mean redshift of $\langle z \rangle = 0.2643 \pm 0.0004$ and velocity dispersion 223 ± 91 km s⁻¹. The spectrum of QSO 0850+4400, of poorer quality, reveals two Ly α absorption features at $z = 0.0909500 \pm 0.0000070$ (which is just resolved) and $z = 0.0948215 \pm 0.0000090$, separated by ~ 1060 km s⁻¹. A group or cluster of galaxies is also present in the vicinity of the QSO line-of-sight with a mean redshift $\langle z \rangle = 0.0901 \pm 0.0007$ and velocity dispersion of 530 ± 200 km s⁻¹ The results of this work establish that $Ly\alpha$ absorption can occur in denser than average galaxy environments, and that it arises in discrete objects spanning a velocity range similar to that of the cluster galaxies. Although a one-to-one relationship between absorbers and galaxies is difficult to establish in such a dense environment, the results obtained here are indeed consistent with the $Ly\alpha$ absorption lines being associated with individual galaxies also in groups and clusters. Moreover, the data shows clearly that line clustering takes place in the $Ly\alpha$ forest.

1 Absorption systems & groups of galaxies

Details about the GHRS spectra of 1545+2101 and 0850+4400 (see Fig. 1) will be presented elsewhere [3]. A standard Voigt profile fitting [1] to individual lines (or groups of them) shows the following features in each field: (a) 1545+2101: two lines are found at $z = 0.2504707 \pm 0.0000030$ and $z = 0.2522505 \pm 0.0000016$, with a velocity separation of ~ 427 km s⁻¹. Also present is a group of eight lines, whose redshift centroid is $\langle z \rangle = 0.2648 \pm 0.0002$ with a velocity dispersion of 163 ± 57 km s⁻¹. (b) 0850+4400: two lines are detected at $z = 0.0909500\pm 0.0000070$ and $z = 0.0948215\pm 0.0000090$, with a velocity separation of 1062 ± 3 km s⁻¹.

A number of galaxies in these two fields have been observed spectroscopically [2] to determine their possible connection to the absorption features detected on the QSOs spectra: (a) 1545+2101: a galaxy is found at z = 0.2510, its impact parameter value being $\rho = 306.4 \ h^{-1}$ kpc. Moreover, there is a group or cluster of six galaxies with redshift centroid $\langle z \rangle = 0.2643 \pm 0.0004$ and velocity dispersion of $223 \pm 91 \ \text{km s}^{-1}$, whose impact parameters range from 47.6 to $456.4h^{-1}$ kpc. (b) 0850+4400: there is a group or cluster of seven galaxies with redshift centroid $\langle z \rangle = 0.0901 \pm 0.0007$ and velocity dispersion of $530 \pm 200 \ \text{km s}^{-1}$. Impact parameter range is $38.5 - 117.8 \ h^{-1}$ kpc



Figure 1: Spectrum of 1545+2101 (left) and enlarged extract from spectrum of 0850+4400 (right). Tick marks in the upper pannel indicate the predicted wavelengths of Ly α at the redshifts of the galaxies. Tick marks in the lower pannel show the positions of the detected absorption lines

 $(h = H_0/(100 \text{ km s}^{-1}\text{Mpc}^{-1}), q_0 = 0.5).$

The fact that we observe a number of resolved lines shows that the absorptions arise in discrete clouds of cold gas instead of arising in a large single diffuse gas component. The comparison between the cross-correlation function (CF) between the galaxies and absorbers in the data with the CF corresponding to a random case shows that there is a non-random connection between the absorbers and the galaxies in our data. If we assume that galaxies within the real group or cluster are distributed in velocity space according to a gaussian distribution, a one-to-one match between the absorbers and the galaxies cannot be established with the present data.

2 Conclusions

From these results we can conclude the following: (1) The detection of groups of $Ly\alpha$ absorption lines implies that some of the $Ly\alpha$ absorbers do actually cluster. (2) The velocity spanned by the $Ly\alpha$ absorption lines arising in groups or clusters is consistent with the velocity dispersion of the corresponding group or cluster of galaxies. This implies that the $Ly\alpha$ absorbers arising in those clusters occupy the same region in space than the galaxies themselves. (3) There is no strong preference for $Ly\alpha$ absorbers to avoid overdense environments. (4) $Ly\alpha$ absorption caused by galaxy groups or clusters arises in discrete clouds of cold gas, rather than in some hypothetical smoothly distributed cold phase in the intracluster medium.

References

- [1] Lanzetta K.M., & Bowen D.V., 1992, ApJ , 391, 48
- [2] Lanzetta K.M., Bowen D.V., Tytler D., & Webb J.K., 1995, ApJ , 442, 538
- [3] Ortiz-Gil A., Lanzetta K.M., Webb J.K., Barcons X., 1997, in preparation