

The Absorption Spectrum of the QSO PKS 2126-158 ($z_{\text{em}} = 3.27$) and the Clustering Properties of Metal Systems

Valentina D’Odorico¹

¹*International School for Advanced Studies, SISSA, via Beirut 2-4, I-34014 Trieste, Italy*

Abstract. We report the detection of a significant clustering signal on scales $\Delta v \lesssim 300 \text{ km s}^{-1}$, for a sample of 71 C IV absorbers. This result confirms previous findings and, together with data at lower column density, it points to a trend of decreasing clustering amplitude with decreasing column density, analogous to that observed for Ly α lines.

1 Observations and data analysis

Spectra of the $z_{\text{em}} = 3.268$ quasar PKS 2126–158 have been obtained at the ESO NTT telescope in the range $\lambda\lambda 4300 - 6620 \text{ \AA}$ with a resolution $R \simeq 27000$ and an average s/n ratio $\simeq 25$ per resolution element. 12 metal systems have been identified, two of which were previously unknown. An enlarged sample of C IV absorptions (71 doublets) has been used to analyze the statistical properties of this class of absorbers. A full report on the data and an extended version of the discussion can be found in the paper [2].

2 Clustering of C IV clouds

We have adopted as a statistical tool the two-point correlation function (TPCF) computed in the velocity space. The reference Poissonian distribution of pairs is obtained by averaging 5000 numerical simulations of the observed number of redshifts, randomly generated in the same redshift range as the data, according to the cosmological distribution $\propto (1+z)^\gamma$ where $\gamma = -1.2$ ([5]).

The TPCF for the C IV clouds, in the range $30 \leq \Delta v \leq 10000$ shows a strong clustering signal at small velocity separations (Fig. 1) while, at larger scales no significant signal is found. In the velocity interval $30 \leq \Delta v \leq 500 \text{ km s}^{-1}$, 66 pairs are observed, while ~ 6 are predicted for a homogeneous distribution, $\xi(30 < \Delta v < 500 \text{ km s}^{-1}) = 10 \pm 1$. Similar results, with a significant correlation on scales up to $\sim 300 \text{ km s}^{-1}$, have been obtained in the works by [4], [7] and [6], carried out at comparable resolution. Degrading the resolution to that of previous works based on larger samples ([5]; [3]) a compatible result is obtained.




Figure 1: TPCF at small velocity separations for the C IV systems (binned at 60 km s^{-1}). The dotted line shows the 95% confidence level.

3 Correlations with the column density

Correlations between total equivalent widths and number of components and between total equivalent widths and velocity spread of the individual C IV systems are observed. The TPCF for lower column density C IV absorption systems shows a weaker signal than the present data ([7]; [6]). These three pieces of evidence suggest a trend of decreasing clustering amplitude with decreasing column density. An analogous behaviour has been observed for Ly α lines by [1].

Acknowledgements. I would like to thank S. Cristiani, S. D’Odorico, E. Giallongo and A. Fontana for their support and useful discussions.

References

- [1] Cristiani S., D’Odorico S., D’Odorico V., Fontana A., Giallongo E., Savaglio S., 1997, MNRAS 285, 209
- [2] D’Odorico V., Cristiani S., D’Odorico S., Fontana A., Giallongo E., 1997, A&A Suppl. in press
- [3] Heisler J., Hogan C., White S.D.M., 1989, ApJ 347, 52
- [4] Petitjean P., Bergeron J., 1994, A&A 283, 759
- [5] Sargent W.L.W., Boksenberg A., Steidel C.C., 1988, ApJS 68, 539
- [6] Songaila A., Cowie L.L., 1996, AJ 112, 335
- [7] Womble W.S., Sargent W.L.W., Lyons R.S., 1996, eds Bremer M.N., in *Cold Gas at High Redshift*, Proc. of the Kluwer Conference