Constraints on Torque-Reversing Accretion-Powered X-ray Pulsars

Insu Yi¹ and J. Craig Wheeler^{2,3}

¹Institute for Advanced Study, Princeton, NJ 08540; yi@sns.ias.edu

²Department of Astronomy, University of Texas, Austin, Texas 78712; wheel@astro.as.utexas.edu

³Institute for Theoretical Physics, University of California, Santa Barbara, 93106-4030

ABSTRACT

The observed abrupt torque reversals in X-ray pulsars, 4U 1626-67, GX 1+4, and OAO 1657-415, can be explained by transition in accretion flow rotation from Keplerian to sub-Keplerian, which takes place at a critical accretion rate, $\sim 10^{16} - 10^{17}g/s$. When a pulsar system spins up near equilibrium spin before the transition, the system goes into spin-down after transition to sub-Keplerian. If a system is well into the spin-up regime, the transition can cause a sharp decrease in spin-up rate but not a sudden spin-down. These observable types of abrupt torque change are distinguished from the smooth torque variation caused by change of \dot{M} in the Keplerian flow. The observed abrupt torque reversal is expected when the pulsar magnetic field $B_* \sim 5 \times 10^{11} b_p^{-1/2} L_{x,36}^{1/2} P_{*,10}^{1/2} G$ where the magnetic pitch parameter $b_p \sim$ a few, $L_{x,36}$ is the X-ray luminosity in $10^{36} erg/s$, and $P_{*,10}$ is the pulsar spin period in 10s. Observed quasi-periodic oscillation (QPO) periods tightly constrain the model. For 4U 1626-67, $\dot{M} \approx 2.7 \times 10^{16} g/s$ with $b_p^{1/2} B_* \approx 2 \times 10^{12} G$. We estimate $\dot{M} \sim 6 \times 10^{16} g/s$ and $b_p^{1/2} B_* \sim 5 \times 10^{13} G$ for GX 1+4, and $\dot{M} \sim 1 \times 10^{17} g/s$ and $b_p^{1/2} B_* \sim 2 \times 10^{13} G$ for OAO 1657-415. Reliable detection of QPOs before and after torque reversal could directly test the model.

Subject headings: accretion, accretion disks – pulsars: general – stars: magnetic fields – X-rays: stars

1. Introduction

Sudden torque reversal events in some accretion-powered X-ray pulsars such as 4U 1626-67, GX 1+4, and OAO 1657-415 have recently been detected (e.g. Chakrabarty et al. 1993,1996,1997ab). The spin-up and spin-down rates are puzzlingly similar despite abrupt torque reversal. The torques remain largely steady before and after reversal, which plausibly indicates the existence of an ordered, stable accretion flow. These systems are distinguished from those showing random torque fluctuations seen in some wind-fed pulsar systems (e.g. Nagase 1989, Anzer & Börner 1995). In the torque-reversing systems, the mass accretion rate \dot{M} appears gradually modulated with a small amplitude on a time scale $\gtrsim yr$ which is much longer than the typical reversal time

scale $\gtrsim days$. The recently reported flux and spectral changes around the time of the reversal in 4U 1626-67 shows that the \dot{M} seems to change by about a few 10% in the X-ray emitting region close to the neutron star (Vaughn & Kitamoto 1997).

Recently, Yi, Wheeler, & Vishniac (1997)¹ have suggested an explanation for the torque reversal phenomenon. The reversal is triggered by transition of the accretion flow from Keplerian rotation to sub-Keplerian rotation. The transition occurs when M crosses the critical rate $\dot{M}_c \sim 10^{16} - 10^{17} g/s$. For magnetized pulsar systems, the inner region of accretion flow lies roughly at a radius $R_o \sim 5 \times 10^8 B_{*,12}^{4/7} \dot{M}_{16}^{-2/7} cm$, where $\dot{M}_{16} = \dot{M}/10^{16} g/s$ and $B_{*,12}$ is the stellar field strength in units of $10^{12}G$. This radius is similar to the size of a white dwarf, which strongly suggests that the transition could be similar to that of cataclysmic variables (e.g. Patterson & Raymond 1985, Narayan & Popham 1993). The details of the transition process remain unknown. The relevant time scale for the transition is likely to be the thermal time scale $t_{th} \sim (\alpha \Omega_K)^{-1}$ or the viscous-thermal time scale $t_{vt} \sim (R/H)t_{th}$, where α is the conventional α viscosity coefficient, H is the thickness of the accretion disk at radius R from the star, and $\Omega_K = (GM_*/R^3)^{1/2}$ is the Keplerian rotation rate for the stellar mass M_* . The time scales $t_{th} < t_{vt} \sim 10^3 s$ for $\alpha \sim 0.3$, $R \sim 10^9 cm$, and $\dot{M}_{16} = 1$ (Frank, King, & Raine 1992). After the transition, the rotation of the accretion flow becomes sub-Keplerian due to large internal pressure (cf. Narayan & Yi 1995). The change of rotation induces an abrupt ($\leq day$) decrease of torque exerted on the star by the accretion flow. The observed X-ray emission is not expected to be significantly affected since Mchange is small.

We show how the proposed model tightly constrains the pulsar system parameters. The estimated parameters are largely consistent with other available estimates. We also derive a simple criterion which identifies candidate pulsar systems by their observable parameters. Combining this with an additional constraint from the quasi-periodic oscillation (QPO), we suggest a possible test of the model. We also discuss different types of torque variation and how they differ observationally from each other. The observed reversal events are likely to occur in pulsar systems near spin equilibrium with $\dot{M} \sim \dot{M}_c \sim a$ few $\times 10^{16} g/s$. We take the neutron star moment of inertia $I_* = 10^{45} g \ cm^2$, radius $R_* = 10^6 cm$, and $M_* = 1.4 M_{\odot}$. The magnetic field is assumed to be a dipole configuration (Frank et al. 1992). The angular velocity of the star $\Omega_* = 2\pi/P_*$.

2. Accretion Flow Transition and Torque Reversal

When the accretion flow rotation is Keplerian, the corotating stellar magnetic field lines interact with the rotating accretion flow with $\Omega_K(R)$ (e.g. Yi et al. 1997, Wang 1995 and references therein). In a steady state, the pitch of the azimuthally stretched magnetic field is given

¹An erratum is to be published in ApJL due to errors in the quoted parameters. The corrected parameters include substantially higher magnetic fields and mass accretion rates which are consistent with those in this work.

by

$$\frac{B_{\phi}(R)}{B_{z}(R)} = \frac{\gamma}{\alpha} \frac{\Omega_{*} - \Omega_{K}(R)}{\Omega_{K}(R)},$$
(2-1)

where $\gamma \leq 1$ measures the vertical velocity shear length scale. The dimensionless ratio $b_p = \gamma/\alpha$, which determines the magnetic pitch, is not well constrained, but is likely to be order unity (e.g. Wang 1995 and reference therein). The magnetically disrupted accretion flow gives an inner edge at $R = R_o$ determined by

$$\delta = \left(\frac{R_o}{R_c}\right)^{7/2} \left[1 - \left(\frac{R_o}{R_c}\right)^{3/2}\right]^{-1}$$
(2-2)

where $R_c = (GM_*P_*^2/4\pi^2)^{1/3}$ is the Keplerian corotation radius and

$$\delta = \frac{2(2\pi)^{7/3} R_*^5}{(GM_*)^{2/3}} \frac{(\gamma/\alpha) B_*^2}{P_*^{7/3} L_x} \approx 2.1 \times 10^{-2} B_{eff,11}^2 P_{*,10}^{-7/3} L_{x,36}^{-1}.$$
(2-3)

We have defined $B_{eff,11} = b_p^{1/2} B_*/10^{11} G$ and $L_x = GM_* \dot{M}/R_*$. The torque exerted on the star is

$$N = \frac{7}{6} N_0 \frac{1 - (8/7)(R_o/R_c)^{3/2}}{1 - (R_o/R_c)^{3/2}}$$
(2-4)

where $N_0 = \dot{M}(GM_*R_o)^{1/2}$ (Yi et al. 1997, Wang 1995). The spin equilibrium N = 0 is achieved when $R_o/R_c = x_{eq} = (7/8)^{2/3}$. The pulsar spin evolution follows $\dot{P}_* = -P_*^2 N/2\pi I_*$ as \dot{M} varies, which we model as a linear variation determined simply by $d\dot{M}/dt$. If the accretion flow remains Keplerian with gradually varying \dot{M} , the disruption radius and the resulting torque N varies according to eqs. (2-2),(2-4). If a pulsar system evolves from a spin-up (spin-down) state and \dot{M} decreases (increases) gradually, it is possible that $R_o/R_c = x_{eq}$ is reached at a certain \dot{M} and then moves to $R_o/R_c > x_{eq}$ ($< x_{eq}$) and hence spin-down (spin-up). As long as \dot{M} variation is smooth and gradual the torque reversal event is expected to be smooth. Figure 1 shows an example of the smooth transition for 4U 1626-67, which is clearly distinguished from the observed abrupt reversal.

When the decreasing \dot{M} reaches \dot{M}_c below which the accretion flow becomes sub-Keplerian (Yi et al. 1997), the accretion flow-magnetic field interaction would change on a time scale $\leq day$. For sub-Keplerian rotation, $\Omega/\Omega_K = A < 1$, the new corotation radius $R'_c = A^{2/3}R_c$ and the new inner edge R'_o is determined by

$$\delta' = \left(\frac{R'_o}{R'_c}\right)^{7/2} \left[1 - \left(\frac{R'_o}{R'_c}\right)^{3/2}\right]^{-1}$$
(2-5)

where $\delta' = \delta A^{-7/3}$. The new torque on the star after the transition is

$$N' = \frac{7}{6} N'_0 \frac{1 - (8/7) (R'_o/R'_c)^{3/2}}{1 - (R'_o/R'_c)^{3/2}},$$
(2-6)

where $N'_0 = A\dot{M}(GM_*R'_o)^{1/2}$. As in the Keplerian flow, N' = 0 would occur when $R'_o/R'_c = x_{eq}$. In a sub-Keplerian flow supported by internal pressure, $\Omega = A\Omega_K$ is largely determined by the ratio of magnetic to gas pressure in the accreted plasma. For equipartition between magnetic and gas pressures, with the pressure ratio, say, 0.05 - 1, we expect $A \approx 0.14 - 0.4$ for all $\alpha = 0.01 - 0.3$ (e.g. Narayan & Yi 1995). For the Keplerian flow, the equilibrium spin period is $P_{eq} \approx 13(b_p/10)^{3/7} B_{*,12}^{6/7} \dot{M}_{16}^{-3/7} s$. For the sub-Keplerian rotation, the new equilibrium spin period $P'_{eq} = P_{eq}/A > P_{eq}$. A system in a spin-up state with a restricted period ratio given by $1 \leq P_*/P_{eq} \leq A^{-1}$ before transition, would evolve toward $P'_{eq} \gtrsim P_*$ after transition, i.e. a sudden torque reversal. Figure 1 shows a fit to the observed reversal event in 4U 1626-67 (cf. Yi et al. 1997). Vaughn and Kitamoto (1997) report that \dot{M} changes by ~ 20% before and after the the observed reversal. The model in Figure 1 is consistent with this finding. On the other hand, if a system has a pre-transition $P_*/P_{eq} > A^{-1}$ that places it well into the spin-up regime, the transition could merely lead to a sudden decrease of spin-up rate, but not to spin-down (i.e. $P_* > P'_{eq}$), a behavior most likely in systems with smaller B_* . Figure 1 shows a hypothetical pulsar similar to 4U 1626-67 except with a smaller B_* . Observations of this type of torque change could support the present accretion flow transition model.

Quasi-periodic oscillation (QPO), if observed, could provide an additional constraint. Such a constraint would depend critically on the nature of the QPO mechanism. Adopting the widely used beat frequency model (e.g. Lamb et al. 1985), we get the QPO periods

$$P_{QPO} = P_* \left[(R_o/R_c)^{-3/2} - 1 \right]^{-1}, \qquad P'_{QPO} = P_* \left[(R'_o/R'_c)^{-3/2} - 1 \right]^{-1}$$
(2-7)

before and after transition, respectively. It remains unclear whether the conventional beat frequency model can explain the recently observed kHz QPOs in some X-ray pulsar systems. The QPO periods we use in the following section are much longer than than the kHz QPO periods. There exists no indication that the long period QPOs arise from other physical processes.

3. Application to Observed X-ray Pulsar Systems

The expressions derived so far can be readily applied to the observed systems.

4U 1626-67 has $P_* \approx 7.660s$ at reversal and the observed torques for the adopted I_* are $N \approx 5.37 \times 10^{33} gcm^2/s^2$ and $N' \approx -4.51 \times 10^{33} gcm^2/s^2$, or the observed torque ratio $(N'/N)_{obs} \approx -0.840$ (Chakrabarty 1996, Chakrabarty et al. 1997a). Using the observed $P_{QPO} \approx 25s$ during spin-up (Shinoda et al. 1990), eq. (2-7) gives $R_o/R_c \approx 0.84$ or $\delta \approx 2.3$ from eq. (2-2). Using the observed N and the derived R_o/R_c in eq. (2-4), we get $\dot{M} \approx 2.7 \times 10^{16} g/s$. Making use of the observed N' and $\delta' = \delta A^{-7/3}$, we solve eqs. (2-5), (2-6) to obtain $A \approx 0.46$, which suggests that the accreted plasma has an equipartition strength magnetic field. From eq. (2-3), we then find $B_{eff} = b_p^{1/2} B_* \approx 1.7 \times 10^{12} G$. This field strength is smaller than the estimates $B_* \sim (6-8) \times 10^{12} G$ obtained by Pravado et al. (1979) and Kii et al. (1986) based on the energy cutoff in X-ray spectrum and the energy dependence of the pulse shape. Although uncertainties in the latter estimates of B_* are not clear, a simple comparison suggests that $b_p^{1/2}$ is at most order unity (cf. Wang 1995). The observed 0.7-60 keV flux $F_x \sim 2.4 \times 10^{-9} erg/s/cm^2$ (Pravado et al. 1979) gives the distance estimate $(L_x/4\pi F_x)^{1/2} \sim 4.2$ kpc, which is consistent with the previous estimates (e.g. Chakrabarty 1996). Eq. (2-7) suggests a possible $P'_{QPO} \sim 120s$ after transition. Detection of such a QPO could directly test the model.

GX 1+4 reversed torque from spin-down to spin-up around the spin period $P_* \approx 122.15s$ (Chakrabarty 1995, Chakrabarty et al. 1997b) with the measured torques $N \approx 3.77 \times 10^{34} g cm^2/s^2$ and $N' \approx -3.14 \times 10^{34} g cm^2/s^2$, which give $(N'/N)_{obs} \approx -0.833$. A QPO of period $P_{QPO} \sim 250s$ was reported during the 1976 spin-up (Doty et al. 1981), but none near the recent reversal episode. The applicability of this QPO is therefore questionable. Nevertheless, assuming that the 1976 spin-up state is similar to the recent spin-up, we can adopt this QPO to make estimates for GX 1+4. The spin period around 1976 is close to the recent spin period, which may help to justify our use of the 1976 QPO data. Following the same procedure as for 4U 1626-67, we estimate that $R_o/R_c \sim 0.77$, $\delta \sim 1.2$, and $\dot{M} \sim 2.9 \times 10^{16} g/s$. We get $A \sim 0.12$, which again indicates a magnetic field strength near equipartition. Finally from eq. (2-3), $B_{eff} \sim 3 \times 10^{13} G$ or $B_* \sim 10^{13} G$ for $b_p \sim 10$. The observed flux $F_x \sim 2 \times 10^{-10} erg/s/cm^2$ in the range of 20-60 keV (Chakrabarty et al. 1997b) gives a distance estimate of ~ 15kpc for the derived \dot{M} whereas for the Doty et al. (1981) value of $F_x \sim 8 \times 10^{-9} erg/s/cm^2$ in the 1.5-55 keV range gives a distance estimate of ~ 2.4 kpc. During the recent spin-down, $P'_{QPO} \sim 6160s$ would be predicted. This prediction, however, is much less certain than that for 4U 1626-67 due to the lack of reliable P_{OPO} measurement near the recent spin-up. We note that the recent GX 1+4 reversal is considerably more gradual than the 4U 1626-67 event. It is possible that the smooth transition type shown in Figure 1 could be relevant in this case (Yi et al. 1997).

OAO 1657-415 has an observed pulse period $P_* \approx 37.665s$ at the time of reversal. Recent observations give $N \approx 4.40 \times 10^{34} gcm^2/s^2$ and $N' \approx -1.06 \times 10^{34} gcm^2/s^2$ (Chakrabarty et al. 1993) or $(N'/N)_{obs} \approx -0.241$. There exists no reported QPO for this system. If we take A = 0.4 based on our estimates in 4U 1626-67 and GX 1+4, the observed torque ratio $(N'/N)_{obs}$ corresponds to $\delta \sim 1.2$ or $R_o/R_c \sim 0.77$ as in GX 1+4, which gives $\dot{M} \sim 1.0 \times 10^{17} g/s$. Therefore, we get $B_{eff} \sim 2 \times 10^{13} G$ or $B_* \sim 10^{13} G$ for $b_p \sim 10$. For $P_* \sim 37s$, $R_o/R_c \sim 0.77$, and A = 0.4, we expect $P_{QPO} \sim 75s$ and $P'_{QPO} \sim 430s$ before and after reversal, respectively. For a detected flux of $\sim 10^{-9} erg/s/cm^2$ (White & Pravado 1979, Mereghetti et al. 1991) and the estimated $\dot{M} \sim 1 \times 10^{17} g/s$, the distance ~ 12 kpc.

We caution that the predicted QPO frequencies may not be easily detectable. During spin-down, the accretion flow thickens considerably and the pulsed polar emission may be significantly diluted by scattering (Yi et al. 1997). The long period QPOs may appear as occasional flares with repetition time scales of $\sim a$ few $10^3 s$.

4. Necessary Condition for Torque Reversal in Systems Near Spin Equilibrium

The parameters derived above can be understood by a simple expression that constrains the pulsar parameters and essentially determines the pulsar magnetic field. We assume that the system reverses its torque from spin-up to spin-down. Just before reversal, we take $R_o/R_c = x_{eq} - \epsilon$ with $\epsilon < x_{eq}$. Expanding eqs. (2-2),(2-4) to first order in ϵ/x_{eq} and combining we get

$$N = \frac{7N_0}{4} \frac{\epsilon/x_{eq}}{1 - x_{eq}^{3/2}} = \frac{7}{2(7 - 4x_{eq}^{3/2})} \left[1 - \frac{1 - x_{eq}^{3/2}}{x_{eq}^{7/2}} \delta \right].$$
 (4-1)

After transition to the sub-Keplerian state, we take $R'_o/R'_c = x_{eq} + \epsilon'$ also with $\epsilon' < x_{eq}$. We expand eqs. (2-5),(2-6) to first order in ϵ'/x_{eq} and combine them to derive

$$N' = -\frac{7N'_0}{4} \frac{1}{1 - x_{eq}^{3/2}} \frac{\epsilon'}{x_{eq}} = -\frac{7N'_0}{2(7 - 4x_{eq}^{3/2})} \left[\frac{1 - x_{eq}^{3/2}}{x_{eq}^{7/2}} \delta' - 1\right].$$
(4-2)

The ratio of the sub-Keplerian spin-down torque (N') to the Keplerian spin-up torque (N) is then

$$\frac{N'}{N} = -A^{4/3} \left[1 + \frac{g(x_{eq})(1 - x_{eq}^{3/2})}{4(7/4 - x_{eq}^{3/2})} (\delta' - \delta) \right] \frac{g(x_{eq})\delta' - 1}{1 - g(x_{eq})\delta}$$
(4-3)

where $g(x_{eq}) = (1 - x_{eq}^{3/2})/x_{eq}^{7/2}$. We have made use of the first order expression

$$\frac{N_0'}{N_0} = A^{4/3} \left(1 + \frac{\epsilon'}{x_{eq}} + \frac{\epsilon}{x_{eq}} \right) = A^{4/3} \left[1 + \frac{g(x_{eq})(1 - x_{eq}^{3/2})}{4(7/4 - x_{eq}^{3/2})} (\delta' - \delta) \right].$$
(4-4)

For $x_{eq} = (7/8)^{2/3}$, we get

$$\frac{N'}{N} = -A^{4/3} \left[1 + 6.10 \times 10^{-3} (\delta' - \delta) \right] \frac{0.171\delta' - 1}{1 - 0.171\delta}.$$
(4-5)

Given the observational constraint $N' \leq N$, eq. (4-5) indicates that small values of $A \sim 0.1$ are unlikely, which is consistent with the result $A \geq 0.3$ in the previous section. This in turn suggests that the magnetic to gas pressure ratio is roughly at the level of equipartition (cf. Narayan & Yi 1995). From eqs. (4-1),(4-2), the first necessary condition for the torque reversal is N'/N < 0 or

$$A^{7/3} \frac{x_{eq}^{7/2}}{1 - x_{eq}^{3/2}} < \delta < \frac{x_{eq}^{7/2}}{1 - x_{eq}^{3/2}}$$
(4-6)

or for $x_{eq} = (7/8)^{2/3}$

$$\delta_{min} = 5.86 A^{7/3} < \delta < 5.86. \tag{4-7}$$

The second condition comes from the observational requirement, $|N'/N|_{obs} < 1$, which gives (cf. eq. 4-5)

$$\delta < 5.86 \frac{1 + A^{4/3}}{1 + A^{-1}}.\tag{4-8}$$

Finally, the observed continuous X-ray emission throughout the whole reversal process requires the accretion to be continuous. This translates into the condition $R_o/R_c < 1$ and $R'_o/R'_c < 1$ in order to avoid an angular momentum barrier at the corotation radius. By expressing ϵ' in terms of δ , we get

$$\delta < 13.39 A^{7/3} = \delta_{max}. \tag{4-9}$$

For $A \leq 0.51$, δ_{max} replaces the upper bounds in eqs. (4-7),(4-8). For $A \geq 0.51$, the upper bound on δ is given by eq. (4-8) as shown in Figure 2. As $A \to 1$, the allowed parameter space shrinks rapidly, which simply indicates that the rotation gets closer to the Keplerian and the spin-down becomes more difficult when the mass accretion rate is fixed. Therefore, the rotation needs to be substantially sub-Keplerian after transition in order for the reversal to occur. For most of the Avalues, δ 's range is limited to be within a factor of ~ 2 . Therefore the condition for torque reversal becomes $\delta_{min} < \delta < \delta_{max}$ and the magnetic field should be close to

$$B_{eff} \approx 7 \times 10^{11} \delta^{1/2} L_{x,36}^{1/2} P_{*,10}^{7/6} G \tag{4-10}$$

where $\delta_{\min}^{1/2} = 2.4A^{7/6} < \delta^{1/2} < 3.7A^{7/6} = \delta_{\max}^{1/2}$. If the accreted plasma is magnetized with near equipartition between magnetic pressure and gas pressure, the sub-Keplerian flow with $A \sim 0.1 - 0.4$ (Narayan & Yi 1995) suggests that $\delta^{1/2}$ is essentially a constant of order unity (cf. Figure 2). Therefore, eq. (4-10) provides an estimate of the magnetic field within a factor of ~ 2 . If the distance to the pulsar and an independent estimate on B_* (say through X-ray spectrum and emission) are known, this relation can constrain $b_p = (B_{eff}/B_*)^2$. The pulsar parameters derived in the previous section are consistent with eq. (4-10). This can be checked as follows. Using eqs. (4-5), we derive

$$\delta = \frac{A^{4/3}}{0.1707} \frac{1 - A^{-4/3} (N'/N)_{obs}}{A^{-1} - (N'/N)_{obs}}.$$
(4-11)

For a given A, we confirm that the observed torque ratio gives δ 's which are close enough to the derived values in the previous section. Therefore we conclude that eq. (4-10) can be reliably used to identify the characteristics of the accreting pulsar systems that should show torque reversals.

The condition for the torque reversal, $5.86A^{7/3} < \delta < 13.39A^{7/3}$, for $A \leq 0.51$ allows a very tightly constrained physical parameters of the pulsar systems. Based on the results in the previous section, our model suggests $A \approx 0.4$ and therefore $0.69 < \delta < 1.58$. Then, in eq. (4-10), $0.8 < \delta^{1/2} < 1.3$ is determined accurately. If this is the case, we can derive $B_{eff} \approx 7 \times 10^{11} L_{x,36}^{1/2} P_{*,10}^{7/6} G$ essentially without uncertainties in δ . Since $\delta \propto A^{7/3}$, $B_{eff} \propto A^{7/6}$ has a crucial dependence on A.

5. Discussion

The current analysis and the study of Yi et al. (1997) suggest that the observed torque reversals occur when the accretion flow makes a transition at a critical rate $\dot{M}_c \sim 10^{16} - 10^{17} g/s$.

This points to an interesting connection between X-ray pulsar systems and other compact accretion systems such as cataclysmic variables and black hole transients. We speculate that the spectral transitions seen in the latter systems may be due to a common accretion flow transition mechanism (cf. Narayan & Yi 1995). The observed torque reversal could be a signature of a pulsar system near spin equilibrium with \dot{M} gradually varying near \dot{M}_c . The proposed mechanism tightly constrains the pulsar field strength. If a pulsar does not have an appropriate field strength (eq. 4-10), it would not undergo a torque reversal even if \dot{M} passes through \dot{M}_c causing the transition. If all systems have a similar \dot{M}_c (Narayan & Yi 1995), the pulsar magnetic field strength can be determined despite a major uncertainty in the distance. It is possible that some systems with $P_* > P_{eq}$ abruptly change their torques without reversing sign at $\dot{M} \sim \dot{M}_c$. These types of sudden torque change are distinguishable from the smooth torque variation in a Keplerian accretion flow. If QPOs can be detected near torque reversal, the proposed model could be further tested in detail.

Although the spectral and flux changes are generally small in the torque reversing systems (Chakrabarty 1996), which indicates the small changes in \dot{M} , at least in 4U 1626-67, some spectral and flux changes have been reported (Vaughn & Kitamoto 1997). The hardening of the spectrum after the transition to spin-down could be attributed to the accretion flow transition itself (Yi et al. 1997). It is however unclear if the hot accretion flow with a large vertical thickness could directly affect the X-ray spectrum through scattering since the expected scattering depth through the hot flow is at most a few $\times 10^{-2}$ for $\dot{M} \sim 2 \times 10^{16} g/s$ (Narayan & Yi 1995) appropriate for 4U 1626-67.

Gradual M modulation on a time scale ranging from $\sim yr$ to a few decades is required for the proposed torque reversal. Such a modulation can be driven by several known mechanisms. (i) In the case of GX 1+4, the binary orbital motion on a time scale $\sim yr$ is plausible (Chakrabarty 1996), which could cause \dot{M} variation through orbital modulation of mass transfer. In 4U 1626-67, an orbital time scale longer than a few years is unlikely (Rappaport et al. 1977, Joss et al. 1978, Shinoda et al. 1990). OAO 1657-415 has an orbital time scale of $\sim 10d$ (Chakrabarty et al. 1993). (ii) Several precession time scales involving the pulsar and the secondary remain viable (e.g. Thorne et al. 1986). (iii) X-ray irradiation-induced mass flow oscillation (Meyer & Meyer-Hofmeister 1990) could provide long time scale (> month) oscillations when certain conditions such as disk size and α are met. Interestingly, in 4U 1626-67 the observed optical pulsation frequency is the same as that in X-rays, which has been attributed to the reprocessing of X-rays by the accretion disk (Ilovaisky et al. 1978, Chester 1979). (iv) The disk instability model (e.g. Smak 1984) has not been applied to neutron star systems and long term \dot{M} variation due to the disk instability needs further investigation.

We acknowledge receiving useful references from L. Bildsten and D. Chakrabarty. JCW appreciates the hospitality of the Institute for Theoretical Physics which is supported by NSF grant No. PHY94-07194. This work has been supported in part by SUAM Foundation (IY) and

by NSF grant AST95-28110 (JCW).

REFERENCES

Anzer, U. & Börner, G. 1995, A&A, 299, 62

- Chakrabarty, D. 1996, Ph.D. Thesis, California Institute of Technology
- Chakrabarty, D. et al. 1993, ApJ, 403, L33
- Chakrabarty, D. et al. 1997a, ApJ, 474, 414
- Chakrabarty, D. et al. 1997b, ApJ, 481, L101
- Chester, T. J. 1979, ApJ, 227, 569
- Doty, J. P., Hoffman, J. A., & Lewin, W. H. G. 1981, ApJ, 243, 257
- Frank, J., King, A. R., & Raine, D. 1992, Accretion Power in Astrophysics (Cambridge: Cambridge University Press)
- Ilovaisky, S. A., Motch, C., & Chevalier, C. 1978, A&A, 70, L19
- Joss, P. C., Avni, Y., & Rappaport, S. 1978, ApJ, 221, 645
- Kii, T., Hayakawa, S., Nagase, F., Ikegami, T., & Kawai, N. 1986, PASJ, 38, 751
- Lamb, F. K., Shibazaki, N., Alpar, M. A., & Shaham, J. 1985, Nature, 316, 239
- Mereghetti, S. et al. 1991, ApJ, 366, L23
- Meyer, F. & Meyer-Hofmeister, E. 1990, A&A, 239, 214
- Nagase, F. 1989, PASJ, 41, 1
- Narayan, R. & Popham, R. 1993, Nature, 362, 820
- Narayan, R. & Yi, I. 1995, ApJ, 452, 710
- Patterson, J. & Raymond, J. C. 1985, ApJ, 292, 535
- Pravado, S. H. et al. 1979, ApJ, 231, 912
- Rappaport, S. et al. 1977, ApJ, 217, L29
- Shinoda, K. et al. 1990, PASJ, 42, L27
- Smak, J. 1984, PASP, 96, 5
- Thorne, K. S., Price, R. H., & MacDonald, D. A. 1986, Black Holes: The Membrane Paradigm (New Haven: Yale University Press)
- Vaughn, B. A. & Kitamoto, S. 1997, preprint (astro-ph/9707105)
- Wang, Y.-M. 1995, ApJ, 449, L153
- White, N. E. & Pravado, S. H. 1979, ApJ, 233, L121

Yi, I., Wheeler, J. C., & Vishniac, E. T. 1997, ApJ, 481, L51 (erratum to appear in ApJL)

This preprint was prepared with the AAS ${\rm IAT}_{\rm E}{\rm X}$ macros v4.0.

Figure Caption

Figure 1: Left columns: Torque reversal event of 4U 1626-67. The solid line is the combined BATSE and pre-BATSE spin period history adopted from Chakrabarty (1996). The dotted lines correspond to the model with the accretion flow transition where $B_* = 5.4 \times 10^{11}$ G, the mass accretion rate decreases from an initial rate of $\dot{M}_i = 3.2 \times 10^{16} g/s$ at a rate $d\dot{M}/dt = -6 \times 10^{14} g/s/yr$. The accretion flow makes a transition at $\dot{M}_c = 2.4 \times 10^{16} g/s$. The dashed line corresponds to the Keplerian model without accretion flow transition where $B_* = 10^{12}$ G, $\dot{M}_i = 5.9 \times 10^{16} g/s$, and $d\dot{M}/dt = -2.35 \times 10^{15} g/s/yr$. The observed event is noticeably more sudden than the latter model. Right Column: A hypothetical pulsar similar to 4U 1626-67 (with abrupt reversal) except $B_* = 3 \times 10^{11}G$. When $P_* > P_{eq}/A$ (i.e. well into spin-up), the accretion flow transition results in abrupt decrease of torque without reversal (dotted line). The dashed line is the corresponding Keplerian model without transition. The parameters are $b_p = 10$ and A = 0.46 (see text).

Figure 2: Allowed region in the $\delta - A$ parameter space for the torque reversal. The region between the two lines are allowed for the torque reversal.



