# On the robustness of the microlensing optical depth as a measure of MACHO density

E. J. Kerins

URA CNRS 1280, Observatoire de Strasbourg, 11 Rue de l'Université, F-67000 Strasbourg,

France

\_;

Received \_\_\_\_\_

accepted \_\_\_\_\_

# ABSTRACT

The optical depth is widely used in Galactic microlensing studies as a means to determine the density of MACHOs, since in theory it depends only upon their spatial distribution and is therefore less model-dependent than other microlensing observables. However, since the measured optical depth is restricted to that of events with measurable timescales, inferences of total MACHO density are dependent on the assumed timescale distribution. Using the 2-year LMC results of the MACHO collaboration, we illustrate this point by showing how, for an assumed isothermal halo, the inferred MACHO halo fraction as determined from optical depth estimates depends upon MACHO mass.

The analysis highlights the following conclusions: (1) The MACHO density inferred from optical depth measurements depends on the assumed MACHO mass function for a given Galactic distribution function; (2) without extra information on the MACHO mass function, such as can be obtained from the rate-timescale distribution, optical depth measurements can provide a lower limit but not an upper limit on the MACHO density for a given Galactic distribution function; (3) a comparison between the inferred total optical depth of different Galactic models, or of different components in a multi-component Galactic model, requires knowledge of the underlying timescale distribution for each model or component.

For our assumed Galactic halo model we find excellent agreement between our model-dependent lower-limit halo fraction of f > 0.3 (84% confidence) or f > 0.15 (97.5% confidence), and the MACHO collaboration's own modelindependent optical depth lower-limit estimates. MACHO's optical depth upper limits are consistent with the minimum value of the upper limit derived for our assumed model.

Subject headings: Dark matter — gravitational lensing — Galaxy: halo — Galaxy: stellar content — Galaxy: structure

#### 1. Introduction

Several microlensing experiments are undertaking searches to detect compact halo dark matter (MACHOs), as well as other low-luminosity stellar populations, following the suggestion of Paczyński (1986). One of the principal quantities which characterises the microlensing properties of a particular MACHO population is the optical depth  $\tau$ . This quantity determines the average number of microlensing events in progress at any instant in time per background source star. In its simplest form it is given by

$$\tau = \int_0^L \frac{\pi R_{\rm e}}{m} \rho(x) \,\mathrm{d}x,\tag{1}$$

where x is the distance along the line of sight between the observer and MACHO, L is the observer–source distance, m is the MACHO mass,  $\rho$  is the MACHO mass density at x and

$$R_{\rm e} = \sqrt{\frac{4Gmx(L-x)}{c^2L}} \tag{2}$$

is the Einstein radius. Equation (1) is valid if all sources are at distance L, otherwise one needs to further integrate equation (1) over the spatial distribution of sources (Kiraga & Paczyński 1994).

So far, theoretical predictions have been compared to measures of the optical depth obtained from observations towards the LMC and Galactic bulge (Alcock et al. 1997a; Renault et al. 1997; Udalski et al. 1994; Alard et al. 1995; Alcock et al. 1997b), with further tentative comparisons also starting to emerge from observations towards SMC (Alcock et al. 1997c) and from pixel experiments directed towards M31 (Ansari et al. 1997; Crotts & Tomaney 1997). For non-pixel based experiments, observational determinations of the optical depth are based on the model-independent estimate

$$\tau_{\text{meas}} = \frac{1}{E} \frac{\pi}{4} \sum_{i=1}^{N_{\text{obs}}} \frac{t_{\text{e},i}}{\mathcal{E}(t_{\text{e},i})},\tag{3}$$

(e.g. Udalski et al. 1994; Alcock et al. 1997a; Alcock et al. 1997b), where  $t_{e,i}$   $(i = 1 \dots N_{obs})$ are the measured event timescales (we define  $t_e \equiv 2R_e/V_T$ , with  $V_T$  the MACHO velocity across the observer-source line of sight),  $\mathcal{E}$  is the efficiency with which timescales  $t_{\rm e}$  are detected and E is the "effective exposure"; that is the average observation time per source star multiplied by the total number of stars observed. Equation (3) basically measures the fraction of the total observing time for which microlensing events are in progress. Errors are typically determined by a boot-strap method in which the range in optical depth is estimated from random timescale realisations generated from the observed  $t_{{\rm e},i}$  (e.g. Alcock et al. 1997b; Alcock et al. 1997a). Han & Gould (1995) have shown that error estimates based on naive Poisson statistics can significantly underestimate the true error.

Alcock et al. (1997a) point out that equation (3) is *not* a measure of the total optical depth, but only of the optical depth of events which fall within a particular range of timescales (those for which  $\mathcal{E} > 0$ ). They stress that an estimate of the total optical depth requires one to input a timescale distribution.

In this study we define the concept of *observable* optical depth and use it, together with the MACHO collaboration's 2-year LMC results, to place limits on the total optical depth, and hence MACHO halo fraction, for the "standard" isothermal halo model analysed by MACHO. The analysis highlights the dependency of the results on MACHO mass for a given Galactic model. Whilst optical depth measurements allow one to determine a lower limit on the density of MACHOs for a given Galactic distribution function, no upper limit can be obtained without using extra information, such as can be obtained from a rate-timescale analysis.

#### 2. Optical depth-timescale distribution

Because of the timescale dependence of equation (3), its theoretical analogue is *not* equation (1), since this is an implicit integral over all event timescales. Instead, one can use

the following expression:

$$\tau = \frac{\pi}{4} \int_{t_{\rm e}(\mathcal{E}>0)} t_{\rm e} \frac{\mathrm{d}\Gamma}{\mathrm{d}t_{\rm e}} \,\mathrm{d}t_{\rm e},\tag{4}$$

where  $\Gamma$  is the event rate. Equation (4) is almost a re-statement, in somewhat expanded form, of the relation  $\tau = (\pi/4)\langle t_e \rangle \Gamma$ , where  $\langle t_e \rangle$  is the average event duration. The one difference is the restriction to timescales  $t_e(\mathcal{E} > 0)$ .

Equation (4) points to an expression for the differential contribution to the optical depth from events of duration  $t_e$ :

$$\frac{\mathrm{d}\tau}{\mathrm{d}t_{\mathrm{e}}} = \frac{\pi}{4} t_{\mathrm{e}} \frac{\mathrm{d}\Gamma}{\mathrm{d}t_{\mathrm{e}}}.$$
(5)

Evidently, just as the rate-timescale distribution  $d\Gamma/dt_e$  depends upon the spatial, velocity and MACHO mass distributions, so too must the optical depth-timescale distribution  $d\tau/dt_e$ . Its integral, if performed over a restricted range of timescales, is similarly model dependent, unlike the expression in equation (1). Hence, to evaluate  $\tau$  within a certain range of event durations  $t_e$  one must specify both the full Galactic distribution function and the MACHO mass function.

As an example, we employ the cored isothermal halo model originally analysed by Griest (1991), and denoted model S in the MACHO collaboration's halo analyses. For the case of a discrete mass function and stationary line of sight, the rate-timescale distribution towards the LMC for this model is

$$\frac{\mathrm{d}\Gamma}{\mathrm{d}t_{\mathrm{e}}} = \frac{2\,V_{\mathrm{c}}^2\rho_0}{m}(a^2 + R_0^2)\int_0^L \frac{\beta(m,x)^2\exp[-\beta(m,x)]}{(x^2 - 2xR_0\cos b\cos l + a^2 + R_0^2)}\,\mathrm{d}x\tag{6}$$

(c.f. Griest 1991), where  $V_c = 220 \text{ km s}^{-1}$  is the halo velocity normalisation,  $\rho_0 = 0.0079 \text{ M}_{\odot} \text{ pc}^{-3}$  is the local halo density, a = 5 kpc is the halo core radius,  $R_0 = 8.5 \text{ kpc}$  is the Sun's Galactocentric distance,  $(l = 280^\circ, b = -33^\circ, L = 50 \text{ kpc})$  is the LMC position in Galactic coordinates and  $\beta \equiv (2R_e/V_c t_e)^2$ . From equations (5) and (6), the optical depth-timescale distribution becomes

$$\frac{\mathrm{d}\tau}{\mathrm{d}t_{\mathrm{e}}} = \frac{\pi V_{\mathrm{c}}^2 \rho_0 t_{\mathrm{e}}}{2m} (a^2 + R_0^2) \int_0^L \frac{\beta(m, x)^2 \exp[-\beta(m, x)]}{(x^2 - 2xR_0 \cos b \cos l + a^2 + R_0^2)} \,\mathrm{d}x.$$
(7)

Both distributions are plotted in Figure (1) for the LMC direction, assuming a MACHO mass  $m = 1 \text{ M}_{\odot}$ . To ease comparison, the distributions are both normalised such that their peak values are unity. It is evident from the figure just how much more sensitive the optical depth is than the rate to longer duration events.

# 3. Observable and observed optical depths

Having defined the optical depth-timescale distribution we can now compare theoretical prediction with observation in either of two ways. The first method would be to directly apply equations (3) and (4) to microlensing data. Instead, we proceed by defining the *observable* optical depth

$$\tau_{\rm oble} \equiv \int_{t_{\rm e}(\mathcal{E}>0)} \mathcal{E}(t_{\rm e}) \frac{\mathrm{d}\tau}{\mathrm{d}t_{\rm e}} \,\mathrm{d}t_{\rm e}.\tag{8}$$

This quantity represents the total optical depth which is potentially observable to a microlensing experiment with detection efficiency  $\mathcal{E}$ . Since it already incorporates the detection efficiency, it should be compared not to equation (3) but to the directly *observed* optical depth in the absence of efficiency corrections:

$$\tau_{\rm obsd} = \frac{1}{E} \frac{\pi}{4} \sum_{i=1}^{N_{\rm obs}} t_{{\rm e},i}.$$
(9)

In the limit of low-number statistics, which is presently the case for searches towards the LMC and SMC, comparison of the quantities  $\tau_{oble}$  and  $\tau_{obsd}$  should provide a more stable estimate than can be obtained from equations (3) and (4), since it is less dependent on the efficiency estimates for particular timescales.



Fig. 1.— The optical depth-timescale and rate-timescale distributions towards the LMC for a standard isothermal halo and a lens mass  $m = 1 \text{ M}_{\odot}$ . The line of sight is assumed to be stationary with respect to the Galactic rest frame. The peak of both distributions is normalised to unity for ease of comparison.

An evaluation of the quantities  $\tau_{oble}$  and  $\tau_{obsd}$  allows a straightforward estimate of the halo fraction  $f = \tau_{obsd}/\tau_{oble}$  for our adopted halo model. This fraction represents the total MACHO fraction, not just the fraction within some timescale range. The first 2 years of MACHO observations towards the LMC have uncovered 8 microlensing candidates (Alcock et al. 1997a), including a likely binary event (Bennett et al. 1996). MACHO analyses both this sample and a six-event sub-sample which excludes the binary event, since it seems likely that the event originates from within the LMC, and also excludes one of the weaker of the other candidates to preserve the overall average event duration. We restrict our analysis to this 6-event sub-sample. The timescales of these 6 events yield an observed optical depth  $\tau_{obsd} = 5.7 \times 10^{-8}$  for an effective exposure  $E = 1.82 \times 10^7$  star-years. Note that this is smaller than the value quoted by MACHO since it does not compensate for the effect of efficiencies. Instead, it represents the optical depth actually measured by the experiment.

As mentioned previously, the potentially observable optical depth  $\tau_{oble}$  is sensitive not only to the halo model but to the assumed MACHO mass function. We calculate inferred halo fractions for discrete MACHO mass functions with masses ranging from  $0.01 - 10 \text{ M}_{\odot}$ . This range includes the  $0.1 - 1 \text{ M}_{\odot}$  range favoured by the MACHO collaboration's maximum-likelihood analysis of the rate-timescale distribution (Alcock et al. 1997a). The MACHO 2-year detection efficiencies are incorporated as required in equation (8).

The solid line in Figure (2) shows the preferred halo fraction f as a function of assumed MACHO mass m. The errors on f are determined by Monte-Carlo simulation and are discussed in the following section. The most obvious point to note is the mass dependency of f. Whilst the preferred value for f is quite stable between  $\sim 0.1 - 1 \,\mathrm{M}_{\odot}$  at  $f \simeq 0.5$ , it becomes arbitrarily large for small and large MACHO masses, with a minimum preferred value of f = 0.47 occurring for  $m = 0.23 \,\mathrm{M}_{\odot}$ . This variation is directly related to the timescale correspondence between the efficiency and optical depth distributions. For very



Fig. 2.— Halo fraction determinations resulting from the 2-year MACHO LMC 6-event sample. A discrete MACHO mass function is assumed. The solid line denotes the preferred value as a function of m. Also shown are 68% and 95% confidence-level regions bounded by dashed and dotted lines, respectively. The horizontal dashed line at f = 1 demarcates a full MACHO halo, whilst the cross denotes the MACHO collaboration's 2-D maximum likelihood solution based on the rate-timescale distribution ( $m = 0.41 \text{ M}_{\odot}$ , f = 0.51).

high- or low-mass MACHOs the peak in the optical depth-timescale distribution occurs at relatively long and short timescales, respectively, where the efficiency is low. Hence,  $\tau_{oble}$  is very small for these cases, and so  $f \propto \tau_{oble}^{-1}$  becomes large for a given  $\tau_{obsd}$ . Hence, for a particular Galactic distribution function, optical depth measurements alone can specify fonly as a function of m.

#### 4. Error estimates

To obtain error estimates on our inferred halo fraction f we use a Monte-Carlo procedure in which 10000 microlensing "experiments" are conducted for each assumed MACHO mass m and halo fraction f. The errors are obtained from the resulting distribution of "observed" optical depths.

The calculation procedure is as follows: for each assumed mass m the efficiencycorrected rate-timescale distribution  $\mathcal{E}d\Gamma/dt_e$  and optical depth-timescale distribution  $\mathcal{E}d\tau/dt_e$  are calculated for a full MACHO halo (f = 1), together with their integrals  $\Gamma_{oble}$ and  $\tau_{oble}$ . Comparison of  $\tau_{oble}$  with  $\tau_{obsd} = 5.7 \times 10^{-8}$ , as computed from the MACHO 2-year LMC results, gives the preferred estimate of f(m) discussed in the previous section. For a range of assumed f, the expected number of detectable events for the model is computed as  $N_{exp}(m) = f E \Gamma_{oble}$ . This number is used to generate a Poisson realisation  $N_{obs,j}$  for mass m, fraction f and experiment j. Event durations  $t_{e,i}$   $(i = 1 \dots N_{obs,j})$  are generated from the distribution  $\mathcal{E}d\Gamma/dt_e$ . The optical depth observed in experiment  $j, \tau_j$ , is then computed from equation (9). Repeating this process for each experiment results in the distribution  $P_{\tau_j}(f,m)$ , of which some fraction  $F_{\tau_j > \tau_{obsd}}(f,m)$  give  $\tau_j$  in excess of the measured value of  $5.7 \times 10^{-8}$ . For a given (f,m), the quantity  $\langle \tau_j \rangle$ , the average over all experiments, is found to be typically in agreement to within 0.5% of the quantity  $f\tau_{oble}$  as obtained by direct integration over  $\mathcal{E}d\tau/dt_e$ . For an assumed m, the values of f for which  $F_{\tau_j > \tau_{obsd}}$  is 0.16 and 0.84 (0.025 and 0.975) bound a 68% (95%) confidence region in f. These confidence intervals are bracketed by the dashed (dotted) lines in Figure (2). It is clear that these intervals are larger for larger lens masses m. This is because a measured  $\tau_{obsd}$  implies a fixed total time  $\sum_i t_{e,i}$  for the summed event durations. The number of events required to produce this summed duration is inevitably larger for low-mass MACHOs than for high-mass MACHOs, due to the smaller Einstein radius of low-mass MACHOs [c.f. equation (2)]. As a result,  $N_{exp}$  is required to be large for low-mass MACHOs so Poisson fluctuations about  $N_{exp}$  are small. The converse is true for very massive MACHOs, which give rise to inherently longer durations. Since  $N_{exp}$ for these objects is typically smaller, larger Poisson fluctuations can arise, which in turn produce larger errors in f.

The actual number of detected events,  $N_{obs}$ , is of course a known quantity but is not directly used in the optical depth analysis. This is because the analysis is concerned only with the sum of the event timescales  $\sum_i t_{e,i}$  not the number of events, and this is essentially why the analysis places no constraint on the MACHO mass m. Rate-timescale analyses use the observed number of events together with the probability of observing each event duration (for an assumed underlying timescale distribution) to constrain both the MACHO density and mass.

Figure (2) shows that, using optical depth measurements alone, one can obtain firm mass-independent lower limits on f for the assumed halo model. The measured value for  $\tau_{obsd} = 5.7 \times 10^{-8}$  implies lower limits of f > 0.3 (84% confidence) and f > 0.15 (97.5% confidence) for the model adopted here. Both of these estimates are in excellent agreement with the MACHO collaboration's model-independent boot-strap determination (Alcock et al. 1997a), and thus support MACHO's assertion that the lower limits it derives ought to be robust. Figure (2) also highlights the fact that upper limits cannot be placed without extra information on the MACHO mass. The MACHO collaboration's boot-strap analysis gives upper limits which are consistent with the minimum derived upper-limit values in Figure (2). These minimum values are f < 0.7 (84% confidence) and f < 0.9 (97.5% confidence). Note that these upper limits apply only to specific MACHO masses, and that the mass at which the upper limit minimises is a function of the required confidence level. The comparison of upper-limit estimates confirms the MACHO collaboration's concerns that its model-independent boot-strap method may tend to underestimate upper limits, since it does not take account of the possible contribution of events with durations exceeding the experiment lifetime.

Plotted in Figure (2) is the MACHO collaboration's preferred value for f and mbased on a 2-D maximum likelihood analysis of the rate-timescale distribution (Alcock et al. 1997a). The correspondence between this estimate  $(f = 0.51, m = 0.41 \text{ M}_{\odot})$  and the optical depth f(m) constraint for the same mass is reassuring and shows that optical depth measures can also serve as important consistency checks on rate-timescale analyses. One would not have the right to expect such good agreement in cases where the assumed model is a poor approximation of the actual Galactic distribution function, or where the underlying timescale distribution is poorly sampled by the experiment. For  $m = 0.41 \text{ M}_{\odot}$ we obtain  $f = 0.48^{+0.62}_{-0.31}$ , where the quoted errors bound a 95% confidence interval. This implies a *total* halo optical depth  $\tau = 2.24^{+2.91}_{-1.45} \times 10^{-7}$ , which is to be compared to the MACHO collaboration's own model-independent estimate of  $\tau_2^{200} = 2.06^{+2.38}_{-1.29} \times 10^{-7}$  for events with durations between 2 and 200 days (Alcock et al. 1997a). Comparison of the two values shows that the MACHO collaboration's estimate is slightly lower than ours, which is to be expected since our estimate is not restricted to a certain timescale range. However, the MACHO estimate is not *much* lower because, for the assumed Galactic model, the timescales produced by 0.41  $M_{\odot}$  lenses typically fall within the MACHO efficiency range and are thus relatively well sampled.

### 5. Discussion

The optical depth is a familiar concept in gravitational microlensing studies and is widely used as a means to determine relatively model-independent constraints on the density of MACHOs in our Galaxy. In this study we emphasise that a comparison between the observed optical depth and the predicted total optical depth is a model-dependent procedure and that failure to take proper account of this may give rise to misleading results.

We have shown that even if one specifies a distribution function for the MACHO population, optical depth constraints on their density will still be a function of the MACHO mass. Whilst optical depth measurements can be used to place firm mass-independent lower limits on the MACHO density they cannot be used to place upper limits. The rate-timescale distribution, however, can provide both lower and upper limits on the MACHO density because of its ability to simultaneously constrain both the MACHO density and typical mass.

In cases where more than one Galactic component contributes significantly to the observed lensing rate, as is believed to be the case for searches directed towards the Galactic bulge, a proper calculation of the relative contribution of each component to the observed optical depth requires one to assume both distribution functions and mass functions for each component. Such assumptions are implicit in calculations which simply compare observed and theoretical optical depths without regard to timescale distributions. This is also relevant to LMC searches if the LMC or an intervening structure is contributing significantly to the microlensing statistics (Sahu 1994; Zhao 1997; Zaritsky & Lin 1997). Similarly, if one wishes to make a comparison between the halo fractions of different Galactic models, as inferred from optical depth measurements, on must specify full distribution functions and mass functions for each model.

In summary, optical depth measurements can provide firm lower limits on the density

in MACHOs and can serve as useful consistency checks for rate-timescale analyses. By themselves, optical depth measurements cannot provide upper limits on the MACHO density. Comparisons between the preferred MACHO density of different Galactic components, or of different Galactic models, are meaningful only if one assumes both a full distribution function and MACHO mass function for each component or model.

The author wishes to thank Will Sutherland for providing the MACHO 2-year LMC efficiencies and David Bennett for pointing out errors in an earlier version of this work. This research is supported by an EU Marie Curie TMR postdoctoral fellowship.

## REFERENCES

- Alard, C., et al., 1995, ESO Messenger, 80, 31
- Alcock, C., et al., 1997a, ApJ, 486, 697
- Alcock, C., et al., 1997b, ApJ, 479, 119
- Alcock, C., et al., 1997c, ApJ, submitted
- Ansari, R., et al., 1997, A&A, 324, 843
- Bennett, D. P., et al., 1996, Nucl. Phys. B (Proc, Suppl.), 5113, 152
- Crotts, A. P. S., Tomaney, A. B., 1997, ApJ, 473, L87
- Griest, K., 1991, ApJ, 366, 412
- Han, C., Gould, A., 1995, ApJ, 449, 521
- Kiraga, M., Paczyński, B., 1994, ApJ, 430, L101
- Paczyński, B., 1986, ApJ, 304, 1
- Renault, C., et al., 1997, A&A, in press
- Sahu, 1994, Nature, 370, 275
- Udalski, A. et al., 1994, Acta Astron., 44, 165
- Zaritsky, D., Lin, D. N. C., 1997, AJ, in press
- Zhao, H. S., 1997, MNRAS, submitted

This manuscript was prepared with the AAS  ${\rm L\!A}T_{\rm E\!X}$  macros v4.0.