Diffractive Scintillation of the Pulsar PSR B1259-63

N. M. McClure-Griffiths^{1,2}, Simon Johnston³, Daniel R. Stinebring¹, and Luciano Nicastro⁴

Received _________________; accepted

¹Department of Physics, Oberlin College, Oberlin, OH 44074

²Present address: Department of Astronomy, University of Minnesota, Minneapolis, MN 55455

³Research Centre for Theoretical Astrophysics, University of Sydney, NSW 2006, Australia

⁴ Instituto TeSRE – CNR, Via Gobetti 101, 40129 Bologna, Italy

ABSTRACT

PSR B1259–63 is in a highly eccentric 3.4 yr orbit around the Be-star SS2883. The system is located in the direction of the Sagittarius-Carina spiral arm at a distance of ∼1.5 kpc. We have made scintillation observations of the pulsar far from periastron at 4.8 GHz and 8.4 GHz, determining the diffractive bandwidth, $\Delta \nu_{\rm d}$, and timescale, $\Delta t_{\rm d}$, at both frequencies. We find no dependency on orbital phase until within 30 days of periastron. The data indicate that the scintillation is caused, not by the circumstellar environment, but by an Hii region within the Sagittarius-Carina spiral arm located at least three-quarters of the way to the pulsar. Close to periastron, when the line-of-sight to the pulsar intersects the disk of the Be star, the electron densities within the disk are sufficient to overcome the 'lever-arm effect' and produce a reduction in the scintillation bandwidth by six orders of magnitude.

Subject headings: ISM: structure — pulsars: individual (PSR B1259-63)

1. Introduction

Scintillation studies provide us with a powerful tool for determining conditions along the line-of-sight to a particular pulsar. They have the potential to determine both the strength of the scattering process and the location of the dominant scattering screen and have been extremely successful in determining some properties of the interstellar medium (Cordes et al. 1991; Gupta 1995). In binary pulsars which are interacting with their companion stars, such as the ablating millisecond pulsars, scatter broadening provides us with a vital clue in understanding the eclipse mechanisms (Luo $\&$ Melrose 1995). We present observations and interpretation of the scintillation parameters for the binary pulsar PSR B1259–63, the only known radio pulsar with a Be star companion.

PSR B1259–63 was identified as a pulsar in a highly eccentric ($e \sim 0.87$), 3.4 yr orbit with the 10 M_{\odot} Be-star, SS2883 (Johnston et al. 1992). Optical observations determined that SS2883 is of spectral type B2e, and also suggested that the system is at a distance $D \sim 1.5$ kpc, placing the pulsar in or just beyond the Sagittarius-Carina spiral arm (Johnston et al. 1994). Be stars are characterized by a circumstellar disk of cool, dense material, which extends outward from the equator to a few tens of stellar radii, and a hot, low-density polar wind.

Observations of the pulsar have shown that dispersion measure (DM) variations around the orbit are ≤ 0.2 cm⁻³pc, except close to periastron where the DM increases by ~15 cm⁻³pc. At this time the pulsar undergoes extreme scatter broadening at frequencies up to at least 5 GHz (Johnston et al. 1996). The increased DM and the scatter broadening can be explained by the line-of-sight to the pulsar traversing the Be star's wind and disk (Melatos, Johnston, & Melrose 1995). PSR B1259–63 thus potentially provides us with a probe of the wind structure of a Be star (far from periastron) and the disk structure (close to periastron).

A priori, one can think of four potential locations for the scattering material: the general interstellar medium, dense material within the spiral arm, an Hii bubble around the hot star, or the wind/disc of the Be star. In this paper we examine each of these possibilities in turn.

2. Observations and Results

We made a total of 46 independent observations of PSR B1259–63 between 1993 August and 1997 February using the Parkes 64-m radio telescope. This excludes data taken in 1993 December and 1994 January during the periastron passage of the pulsar (Johnston et al. 1996). The observations were made at 4.8 GHz and 8.4 GHz using dual-channel, cryogenically cooled systems sensitive to two orthogonal linear polarizations. The signals from the receivers were down-converted to an intermediate frequency and then passed into a back-end filterbank which consisted of a number of contiguous frequency channels. Prior to 1996 October the filterbank consisted of 64 channels, each 5 MHz wide for a total bandwidth of 320 MHz. Since 1996 October we have used an alternative back-end system consisting of 128 channels, each 3 MHz wide (total bandwidth 384 MHz). The signal was detected, high and low-pass filtered, and sampled at 0.6 ms. The observations were typically between 30 min and 100 min in duration.

The off-line data analysis was performed in two stages. First, a dynamic spectrum was constructed by forming pulse profiles for each frequency channel over a small length of time, typically 10 s. In order to maximize the signal-to-noise ratio, we often summed consecutive frequency channels and time-intervals, making sure that we retained adequate resolution across individual scintillation maxima. Subsequently, a 2-dimensional autocorrelation was formed of the summed dynamic spectrum, and 1-dimensional Gaussians fitted to the zero lag in time and frequency to find the diffractive bandwidth, $\Delta\nu_{\rm d}$, and diffractive time scale,

 $\Delta t_{\rm d}$, respectively. No significant tilt of the autocorrelation function was observed, and we computed fractional uncertainties of the values based on the number of scintles present in the dynamic spectrum following Cordes (1986). The scintillation parameters for each observation and their associated errors are plotted as a function of days from periastron in Figure [1.](#page-12-0)

The main finding from these observations is that there is little or no change in the scintillation parameters with orbital phase from 25 days after periastron until 100 days before periastron. The apparent decrease in the scintillation bandwidth at 4.8 GHz in late 1996 and early 1997 is most likely due to the improved spectral resolution. The standard deviation of the values throughout the orbit is ∼30%, consistent with the individual uncertainties at both 4.8 GHz and 8.4 GHz. Averaging all data at 4.8 GHz, we find that $\Delta \nu_{\rm d} = 8.5 \pm 0.9$ MHz and $\Delta t_{\rm d} = 190 \pm 10$ s. At 8.4 GHz, we find that $\Delta \nu_{\rm d} = 70 \pm 20$ MHz and $\Delta t_d = 360 \pm 100$ s. The contrast of these values far from periastron with the ∼20 Hz scintillation bandwidth at 4.8 GHz just prior to periastron is startling.

3. Discussion

Scattering in the interstellar medium (ISM) is due to electron density inhomogeneities which are thought to be distributed as a power-law, $P_{3N} = C_N^2 q^{-\beta}$. P_{3N} is the 3-dimensional electron density power spectrum, C_N^2 gives the averaged level of turbulence along a particular line-of-sight, q is the spatial wavenumber, and β is the spectral index of the power spectrum (e.g. Kaspi & Stinebring 1992). In the discussion below we assume Kolmogorov turbulence such that $\beta = 11/3$.

Using the definition for C_N^2 given in Cordes (1986) and the parameters for PSR B1259– 63, we find $C_N^2 = 10^{-1.3}$ m^{-20/3}. This value places PSR B1259-63 in the top five pulsars

ranked by C_N^2 and is more than two orders of magnitude greater than the 'canonical' value of 10^{-3.5} m^{-20/3} for the warm, ionized (ISM). Further evidence for the enhanced density towards the pulsar comes from the pulsar's DM. At a distance of 1.5 kpc, the Taylor & Cordes (1993) model yields an expected DM of \sim 50 pc cm⁻³, in contrast to the measured value of 147 pc cm^{-3} .

The scintillation velocity, V_{iss} , can be computed from

$$
V_{iss} = 3.85 \times 10^4 \frac{\sqrt{\Delta \nu_{\text{d,MHz}} x D_{\text{kpc}}}}{\nu_{\text{GHz}} \Delta t_d} \text{ km s}^{-1}.
$$
 (1)

where ν is the observing frequency and x is the ratio between the screen-observer distance and the screen-pulsar distance (Gupta et al. 1994; Gupta 1995). Assuming a mid-placed $(x = 1)$ scattering screen we derive a scintillation velocity of ~150 km s⁻¹. We note that the scintillation velocity, as a function of orbital phase, is constant within the errors. The small and slowly changing component of the pulsar's orbital velocity cannot be detected in the data. The proper motion of PSR B1259–63 has not been measured, although Johnston et al. (1994) estimate a recessional velocity of 80 km s⁻¹ based on the shift of the H α line. Theoretical considerations of the evolution of systems like PSR B1259–63 give an upper limit of 90 km s⁻¹ for the space velocity (Brandt & Podsiadlowski 1995). To reconcile this with the measured scintillation velocity implies $x \ge 3$, placing the screen closer to the pulsar than the observer.

Evidently, then, there is good evidence for enhanced-density scattering material located significantly closer to the pulsar than mid-way from the observer. In the following discussion we derive an expression linking the scintillation bandwidth with the total electron density variance along the line-of-sight, δn_e^2 , the outer scale of density inhomogeneities, l_o , and the thickness of the scattering screen, ΔL . We then consider three possible locations of the scattering screen: the wind of the Be star, a possible Strömgren sphere surrounding the system, and the Sagittarius-Carina spiral arm, all three of which satisfy the condition on x .

 χ From geometric ray optics, we can compute the time delay, τ , of scattered rays and using the 'uncertainty relationship', $2\pi\tau\Delta\nu_{\rm d}=1$, we have

$$
\Delta\nu_{\rm d} = \frac{c}{\pi} \frac{(1+x)^2}{x} \frac{1}{D\theta_{\rm scatt}^2},\tag{2}
$$

where c is the speed of light and θ_{scatt} is the scattering angle at the screen. (Note that for a mid-placed screen, $x = 1$, the angular broadening, θ , is $\theta_{\text{scatt}}/2$ and equation 2 becomes the familiar $\Delta \nu_{\rm d} = c/(\pi D \theta^2)$.) By relating the electron density inhomogeneities to the coherence length of the electromagnetic phase, θ_{scatt} can be expressed as

$$
\theta_{\text{scatt}} = \frac{c}{2\pi\nu} \left[\frac{A_{\beta}\,\nu^2}{2\pi r_e^2 c^2} \frac{1}{\Delta L \, C_N^2} \right]^{-3/5} \tag{3}
$$

(Cordes et al. 1985), where r_e is the classical electron radius. C_N^2 is given by

$$
C_N^2 = C_{\rm SM} \frac{\delta n_e^2}{l_o^{2/3}}.
$$
 (4)

Both A_β and C_{SM} are dimensionless constants which depend only on β and have values of 2.49 and 0.18, respectively, for $\beta = 11/3$ (Cordes et al. 1985, Cordes et al. 1991). Combining these equations and putting the parameters in convenient astronomical units we find that -10^{16}

$$
\delta n_e^2 = 4.0 \,\text{cm}^{-6} \quad \Delta \nu_{\text{d,MHz}}^{-5/6} \quad D_{\text{kpc}}^{-5/6} \quad \nu_{\text{GHz}}^{11/3} \quad \frac{l_{o,\text{pc}}^{2/3}}{\Delta L_{\text{pc}}} \quad \left[\frac{(1+x)^2}{x} \right]^{5/6} . \tag{5}
$$

The wind of the Be star near apastron is a possible scattering site. However, if this were the case, one would expect a modulation in the scintillation parameters as a function of orbital phase. Our observations cover a period in which the distance of the pulsar from the Be star varies by a factor of 3; assuming a wind profile density which varies as r^{-2} , this implies a change in electron density by a factor of ∼10. Given that the scintillation bandwidth scales as $\Delta n_e^{-12/5}$ (from equation [5]), if $\delta n_e^2/n_e^2$ and l_o remain constant we should observe a change in $\Delta \nu_{\rm d}$ by a factor ~250. Furthermore, the wind velocity is as high as \sim 1000 km s⁻¹ and this would then be the dominant velocity in the system, causing a diffractive timescale much shorter than observed. Finally, it is unlikely that the electron density within the wind is large enough to cause the observed scintillation until very close to periastron. Assuming the scattering screen is located only a few hundred light seconds from the pulsar and that $l_o \approx \Delta L$ is also of this order, then $\delta n_e^2 \sim 10^{11}$ cm⁻⁶. For even a fully modulated wind with $\delta n_e^2 \approx n_e^2$ the implied electron densities are only encountered within ± 30 days of periastron.

The UV flux from hot, high-mass stars can form an HII bubble (or Strömgren sphere) within the ISM. The interface between these two regions can be a source of turbulence and is a possible site for scattering. We can put limits on the emission measure of the Strömgren sphere from a 12-hr synthesis image of the pulsar made using the Australia Telescope Compact Array. The observations were made at frequencies of 1.4 and 2.4 GHz and were gated into 8 bins synchronous with the pulsar period. After removal of the pulsed component from the image, the upper limit on the brightness temperature of any residual emission is ∼1 K. Using standard equations for free-free emission from Hii regions and assuming a temperature of 10^4 K, the upper limit on the emission measure is 650 cm⁻⁶pc. The radius of a Strömgren sphere for a B2 star with effective temperature of 2×10^4 K is related to the electron density within the surrounding medium by $R_s = 13n_e^{-2/3}$ pc (Prentice & ter Haar 1969). Hence, we see that $n_e \le 20$ cm⁻³ and $R_s > 1.8$ pc for the Strömgren sphere to be undetectable. Applying equation [\(5\)](#page-6-0), assuming $x \sim 750$ and $l_o \approx \Delta L \approx 1$ pc, we see that δn_e^2 must be larger than $\sim 3.7 \times 10^4$ cm⁻⁶ to yield the observed scintillation bandwidth. Such a large value is inconsistent with the upper limit on the electron density imposed by the lack of radio emission. Thus, the Strömgren sphere cannot be the source of the observed scattering.

The Sagittarius-Carina arm intersects the line-of-sight at a distance of ∼1 kpc. The spiral arm likely contains numerous, high-density Hii regions which can cause both excess

DM and scattering. If we estimate $l_o \sim 1$ pc for an H_{II} region and use $x = 3$, equation [\(5](#page-6-0)) yields $\delta n_e^2 \Delta L = 600 \text{ cm}^{-6} \text{pc}$. From the excess DM we have that $n_e \Delta L = 100 \text{ cm}^{-3} \text{pc}$ and assuming that the condition $\delta n_e^2 \approx n_e^2$ holds within the HII region then we find $n_e \sim 6 \text{ cm}^{-3}$ and $\Delta L \sim 10$ pc. The derived electron density is consistent with ~ 6 cm⁻³ and ~ 3 cm⁻³ obtained by Heiles, Reach, & Koo (1996) and McKee & Williams (1997), respectively, for 'typical' Hii regions. We thus conclude that the source of the enhanced scattering is one (or several) H_{II} regions within the Sagittarius-Carina spiral arm.

A number of other factors point to the Sagittarius-Carina spiral arm as a dominant source of scattering. PSR J1243–6423, only 2.2° distant from PSR B1259–63 on the sky, has very similar scintillation parameters to those of PSR B1259–63 (Johnston et al. 1997). Its DM-derived distance is 12 kpc, yet indications from Hi absorption suggest a distance as low as 2.2 kpc (Frail & Weisberg 1990). It seems evident that the spiral arm contributes both to its large DM and relatively small scintillation bandwidth. Nicastro & Johnston (1997) recently discovered a pulsar only 7′ from PSR B1259–63 with an exceedingly high DM of 875 pc cm⁻³. Such a large DM is not expected in this part of the Galactic plane, and this again points to the spiral arm as a significant source of excess electrons.

Near periastron we conclude that the disk of the Be star must dominate the scattering. Twenty days from periastron $\Delta \nu_{\rm d}$ is 30 Hz at 1.4 GHz. We assume that the proximity of the scattering screen implies $x \sim 5 \times 10^8$. We take $\Delta L \approx l_o \sim 100$ light seconds and hence obtain δn_e^2 of ~ 10¹⁴ cm⁻⁶. Again, assuming $\delta n_e^2 \approx n_e^2$, an electron density of ~ 10⁷ cm⁻³ is required. Modeling of the disk of the Be star (Melatos et al. 1995) shows that the maximum density encountered along the line of sight through the disk reaches 10⁷ cm⁻³ some 25 days prior to periastron and rises by an order of magnitude at periastron itself. Hence, near to periastron, there is sufficient density in the Be star's disk to produce the observed scattering. Because the scattering angle is $\sim 0.5^{\circ}$, the angular broadening produces an

image size of 8×10^{10} cm on the screen. This value is significantly larger than the diffractive coherence scale of $\sim 10^9$ cm for scattering in the spiral arm. We thus expect a quenching of the scintillation seen through the spiral arm, which is confirmed by the observations.

4. Conclusions

We have determined the scintillation parameters of the binary pulsar PSR B1259–63 as a function of orbital phase and conclude that there is no change in these parameters except during a small portion of the orbit around the time of periastron. The scintillation bandwidth is much less than expected if the scintillation were solely due to the background ISM. We have demonstrated that the electron densities are not sufficiently large in either the wind of the Be star far from periastron or a putative Strömgren sphere to produce the observed scattering. The Sagittarius-Carina spiral arm is therefore the most likely origin of the scintillation far from periastron. At periastron, however, the electron density in the Be star disk is sufficiently large to counterbalance the 'lever-arm effect' of having the screen so close to the pulsar. This produces extremely strong scattering, and the resultant broadening of the pulsar image is sufficient to quench further scintillation in the ISM.

The back-end system used in this work was designed and built by A. Lyne at the University of Manchester. We thank R. Manchester for help with the observations, B. Rickett for fruitful discussions and H. Fagg and M. McColl for receiver installation and maintenance. NMM-G and DRS gratefully acknowledge research support from NSF grant AST-9317605 and hospitality and support from the Australia Telescope National Facility. The Parkes radio telescope is part of the Australia Telescope which is funded by the Commonwealth of Australia for operation as a National Facility managed by CSIRO.

REFERENCES

- Brandt, N. Podsiadlowski, P. 1995, MNRAS, 274, 461.
- Cordes, J. M., Weisberg, J. M., Frail, D. A., Spangler, S. R., & Ryan, M. 1991, Nature, 354, 121.
- Cordes, J. M. 1986, ApJ, 311, 183.
- Cordes, J. M., Weisberg, J. M., Boriakoff, V. 1985, ApJ, 288, 221.
- Frail, D. A., & Weisberg, J. M. 1990, AJ, 100, 743.
- Gupta, Y. 1995, ApJ, 451, 717.
- Gupta, Y., Rickett, B. J., & Lyne, A. G. 1994, MNRAS, 269, 1035.
- Heiles, C., Reach, W. T., & Koo, B.-C. 1996, ApJ, 466, 191.
- Johnston, S., Manchester, R. N., Lyne, A. G., Bailes, M., Kaspi, V.M., Guojun Qiao, & D'Amico, N. 1992, ApJ, 387, L37.
- Johnston, S., Manchester, R. N., Lyne, A. G., Nicastro, L., & Spryromilio, J. 1994, MNRAS, 268, 430.
- Johnston, S., Manchester, R. N., Lyne, A. G., D'Amico, N., Bailes, M., Gaensler, B. M., & Nicastro, L. 1996, MNRAS, 279, 1026.
- Johnston, S., Nicastro, L., & Koribalski, B. 1997, In Preparation.
- Kaspi, V. M., & Stinebring, D. R. 1992, ApJ, 392, 530
- Luo, Q., & Melrose, D. B. 1995, ApJ, 452, 346
- McKee, C. F., & Williams, J. P. 1997, ApJ, 476, 144.
- Melatos, A., Johnston, S., & Melrose, D. B. 1995, MNRAS, 275, 381.
- Nicastro, L., & Johnston, S. 1997, MNRAS, Submitted.

Prentice, A. J. R. & ter Haar, D. 1969, MNRAS, 146, 423.

Taylor, J. H., & Cordes, J. M. 1993, ApJ 411, 674.

This manuscript was prepared with the AAS IATEX macros v4.0. $\,$

Fig. 1.— Diffractive scintillation parameters as a function of orbital phase. Periastron is represented as day 0 in all plots. Figures (a) and (c) show diffractive bandwidth with the associated uncertainties at 4.8 GHz and 8.4 GHz, respectively. Figures (b) and (d) show diffractive timescale and associated uncertainties at 4.8 GHz and 8.4 GHz, respectively.