Radio-Luminous Southern Seyfert Galaxies. I. Radio Images and Selected Optical/Near-Infrared Spectroscopy

M. A. Bransford^{1,2}, P. N. Appleton^{1,2}

Erwin W. Fick Observatory and Department of Physics and Astronomy, Iowa State University, Ames, IA 50011

C. A. Heisler¹

Mount Stromlo and Siding Spring Observatory, ANU, Private Bag, Weston Creek Post Office, ACT 2611, Australia

R. P. $Norris^1$

CSIRO Australia Telescope National Facility, P. O. Box 76, Epping, NSW, 2121, Australia

A.P. Marston³

Dept. of Physics and Astronomy, Drake University, Des Moines, IA 50311

Send off-print requests to M. A. Bransford e-mail: mabransf@iastate.edu. Higher quality figures (.ps) are available via anonymous ftp, ftp pv7408.vincent.iastate.edu, login: anonymous.mabransf, password: yourusername@domain, cd outgoing, binary, get figures.tar (approx. 1 Meg). Updated version includes revised coordinates of galaxies in Table 1.

Received _____; accepted Astrophysical Journal, October 1997

¹Visiting astronomer at the Anglo-Australian Observatory

²Visiting astronomer at the Australia Telescope National Facility. The Australia telescope is funded by the Commonwealth of Australia for operation as a National Facility managed by CSIRO.

³Visiting astronomer at CTIO, CTIO is operated by Associated Universities for Research in Astronomy (AURA) under contract to the NSF

ABSTRACT

This is the first of two papers in which a study is made of a sample of 12 southern radio-luminous Seyfert galaxies. Our aim is to investigate possible correlations between radio morphology and nuclear/circumnuclear emission-line properties. In this paper we present radio images at $\lambda = 13$, 6, and 3 cm taken with the Australia Telescope Compact Array (ATCA), global far-infrared (FIR) properties for the whole sample, and optical and near-infrared (NIR) spectroscopy of an interesting subset. We find a mixture of radio morphologies, including linear, diffuse and compact sources. When the FIR colors of the galaxies are considered there is an indication that the compact radio sources have warmer FIR colors than the diffuse sources, whereas the linear sources span a wide range of FIR colors. There is a wide variation in radio spectral-indices, suggesting that free-free absorption is significant in some systems, particularly IRAS 11249-2859, NGC 4507, and NGC 7213.

Detailed emission-line studies are presented of 4 galaxies IC 3639, NGC 5135, NGC 3393 & IRAS 11249-2859. In IC 3639 we present evidence of vigorous, compact star formation enclosed by very extended [OI] λ 6300 emission, suggestive of the boundary between a diffuse outflow and the surrounding ISM. In another galaxy, IC 5063, we see evidence for the possible interaction of a highly collimated outflow and the surrounding rotating inner disk. Of the 5 galaxies which show compact radio emission, 4 have radio/FIR flux ratios consistent with an energetically dominant AGN, whereas IC 4995 exhibits evidence for a very compact starburst.

1. Introduction

Indications that Sevfert galaxies may contain a significant starburst component come from several different directions, although it is not yet clear how such star formation might affect the properties of an active galactic nucleus (AGN). Norris, Allen, & Roche (1988) and Roy et al. (1997) showed that most Seyfert galaxies, unlike radio quasars, roughly follow the radio-far-infrared correlation, which suggests that their bolometric luminosity is dominated by star formation activity. Extended minor-axis radio emission from Seyferts (Baum et al. 1993) suggests that many Seyfert galaxies may contain powerful circumnuclear starbursts. The inner few hundred parsecs of Mrk 477 have recently been shown to contain an extremely luminous and compact starburst (Heckman et al. 1997). In NGC 1068 an extended luminous starburst region is observed 3 kpc from the central AGN (e.g. Telesco & Decher 1988). Genzel et al. (1995) made high-resolution NIR spectroscopic images of NGC 7469, which show a circumnuclear ring of star formation 480 pc in radius. A blueshifted ridge of emission between the ring and the nucleus is interpreted as infalling gas that may provide fuel for the AGN. The star-formation in NGC 1068 and NGC 7469 may result from a stellar bar which is both triggering extended star-formation and funneling material inwards to the AGN. Previous investigations have shown that half of the total luminosity of NGC 1068 (Telesco et al. 1984; Lester et al. 1987) and Mrk 477 (Heckman et al. 1997), and nearly two-thirds of that of NGC 7469 (Genzel et al. 1995), arises from starburst activity.

This study is aimed at looking for possible correlations between arcsec radio structure⁴ and optical–IR emission-line properties in a sample of southern radio-luminous Seyfert galaxies (see Table 1). The mapping of arcsec radio structure in southern Seyfert galaxies has only become feasible recently with the advent of the Australia Telescope National Facility and many of these radio sources have not yet been mapped. Our pilot study focuses on one aspect of the properties of AGNs, namely the possible relationship between AGN or starburst activity and the morphology of the arcsec-scale radio emission. For example, we wish to investigate the possibility that

⁴At the distance of our sample galaxies this corresponds to linear scales of a few hundred parsecs.

radio-emitting plasma ejected from an AGN might interact and trigger star formation in the inner regions of galaxies. Alternatively, in other systems, the radio emission itself may originate in circumnuclear star formation and this will be investigated via spectroscopic observation.

In this paper we present the radio images, at three wavelengths, of the galaxies in our sample, as well as optical and NIR spectra for an interesting subset. In addition we will compare the thermal properties of the dust emission (as measured by global FIR properties of the galaxies, e.g. Heisler & Vader 1994; Condon et al. 1991) with the radio properties. We show that for a subset of these data, the FIR color temperatures seem to correlate with radio morphology, suggesting that in some cases the FIR colors are intimately connected with the nuclear dust properties of Seyferts. (see §4.3). In a second paper (Paper II) we will complete the spectroscopic observations and draw more general conclusions about the sample.

In §2 we present the sample selection, in §3 the details of the observations, in §4 the radio data, and in §5 we present optical and NIR spectra of some of the galaxies. In §6 we present our summary and conclusions.

2. The Sample

The sample galaxies listed in Table 1 are a radio-bright subset of a 2.4 GHz Australia Telescope National Facility (ATNF) survey of Seyfert galaxies by Roy et al. (1997).⁵ Galaxies from this list were selected if they were more luminous than $L_{2.4GHz} > 10^{21.7}$ W Hz⁻¹, with declinations $< -20^{\circ}$, and redshifts z < 0.04. One galaxy that met these criteria, IC 4329A, was not included in our sample since it had been imaged in the radio by A.L. Roy (private communication) and radio maps of this source will be published elsewhere. Our 2.4 GHz observations reveal that the range in luminosity of our sample is $log(L_{2.4GHz}/W-Hz^{-1}) = 21.7-23.2$ (see Figure 1). By way of comparison, Centaurus A, the most radio-luminous Seyfert galaxy in the southern hemisphere,

⁵The Roy et al. sample is drawn from the de Grijp et al. 1992 sample of IRAS selected galaxies.

has $L_{2.7GHz} = 10^{23.8}$ W Hz⁻¹ (Bolton & Shimmins 1973). Approximately 40% of the galaxies lie on the "radio-bright" (see Fig. 1) side of the well-known FIR/radio correlation (Helou, Soifer, & Rowan-Robinson 1985) and were therefore expected to show composite AGN/star formation characteristics suitable for a study of this kind. The galaxies in the sample have been observed as part of various optical emission-line surveys (Phillips, Charles, & Baldwin 1983; van den Broek et al. 1991; de Grijp et al. 1992). In Table 1 we list some of the optical properties of the sample galaxies.

3. The Observations

3.1. Radio Observations

Radio observations were made with the 6 km Compact Array (ATCA) on 1994 July 24 and 25, at a wavelength λ 13 cm and on 1995 November 19-22, at λ 6 and λ 3 cm. Each source was observed for typically six 20-min periods spread over 12 h, in order to provide good coverage of the u-v (Fourier-transform) plane. The total on-source integration times were between 1.29 and 5.66 h. Unresolved phase calibrators were observed before and after every target observation. Data were taken in two independent linear polarizations at frequencies centered on 2.378 and 2.496 GHz with a bandwidth 128 MHz for the 13 cm observations. The 6 and 3 cm observations were made simultaneously, with frequencies centered on 4.798 and 8.638 GHz with a bandwidth of 128 MHz. Flux densities were calibrated assuming a flux density for the standard ATCA calibrator PKS B1934-638 of 13, 5.8, and 2.8 Jy at 13, 6, and 3 cm, respectively.

AIPS was used to perform the standard calibration and data reduction. Images were made and CLEANed using the AIPS routine MX, and then corrected for primary beam effects. The rms flux densities in source-free areas of the CLEANed images ranged from 50 to 720 μ Jy beam⁻¹. Table 2 lists the clean beam dimensions and rms noise levels, for each source observed at all three wavelengths. In the case of brighter sources (≥ 100 mJy) we used "self-calibration" to provide additional phase corrections. The resulting dynamic range achieved in most of the images was typically 50-100 at 13 cm and approximately 20-30 at 3 cm, although in most cases the images were limited by thermal noise ratio rather than dynamic range.

Total fluxes for the sources were calculated by blanking all data below twice the rms noise in the image in a region close to the source. Because these observations are optimized for studying the nuclear region, they are relatively insensitive to extended emission. Thus our images are primarily of the nuclear region and will therefore not show any smooth extended structure on scales of tens of arcsec, and consequently the total radio fluxes quoted may be an underestimate. Table 3 lists the fluxes of the sources at each of the three wavelengths and other radio properties. The images have resolutions in the range 1-8". The galaxy NGC 5135 was observed solely at 13 cm, because published 6 and 3 cm VLA images are available for this object (Ulvestad & Wilson 1989). Spectral indices were calculated, in the core regions, for 13 and 6 cm, 13 and 3 cm, and 3 and 6 cm ($\alpha(6,13)$, $\alpha(3,13)$, and $\alpha(3,6)$ are listed in Table 2), by restoring the higher frequency images with the same beam and cell-size as used at the lower frequencies. To calculate the spectral index accurately, we measured the flux density over similar regions for each source, at both frequencies.

3.2. Infrared and Optical Observations

NIR spectroscopy and imaging were performed at the 3.9 m AAT, in 1994 July and October. The July observations resulted in only 2 h of useful non-photometric K-band imaging data because of bad weather. Direct imaging was done using the HgCdTe 128 x 128 array detector of the Infrared Imaging Spectrograph (IRIS) and a field of view of 1.'3 x 1.'3. Although we obtained K-band images for IRAS 18325-5926, IC 4995, IC 5063 and NGC 7213, only NGC 7130 shows interesting structure (see §5.2).

Attempts to obtain long-slit, NIR spectra during this run were unsuccessful because of bad weather. During part of the AAT Director's discretionary time on 1994 October 21-23, non-photometric conditions allowed us to obtain a series of long-slit, NIR spectra for four galaxies: IRAS 18325-5926, IC 4995, IC 5063 and NGC 7213. At f/15 the image scale was 0.61 pixel⁻¹ and the spectral resolution was ~300. The telescope was programmed to step the slit across the galaxy in 0".6 increments and integration times of 180-220 s were obtained at each position. In this way, a spectral cube was built up giving full two-dimensional spectral information across the inner 60" of each galaxy. Two sets of scans were made, one in each of the H (λ 1.65 μ m) and K (λ 2.2 μ m) NIR bands. Infrared emission was detected from the Br γ , H₂ v=1-0 S(1) 2.121 μ m and [FeII] λ 1.644 μ m lines for all four galaxies observed. Representative K-band spectra are presented in §5.

We obtained long-slit optical spectroscopy at both the AAT and CTIO of four galaxies, to search for extended emission associated with their radio and NIR structure in their nuclear regions and inner disks. The spectroscopic optical observations were made at intermediate dispersion at the AAT in 1995 August and low dispersion at the 1.5 m CTIO telescope in 1995 April.

The AAT observations were made using the RGO spectrograph, in combination with the TEK 1k CCD detector (1024 x 1024) at f/8 with the 25 cm camera and a grating with 270 grooves mm^{-1} in first order to produce a spectral resolution of ~3.3 Å pixel⁻¹ over a wavelength range of 4170-7625 Å. The spatial plate scale was 0".77 pixel⁻¹. The slit was set to a width of 1".66 for galaxy observations, which was comparable to the atmospheric seeing. Total integration times at each position were 1200-1800 s. These observations were made during photometric conditions for only 3 hours on the first night. The standard star LTT 7379 (Stone & Baldwin 1983) was observed at approximately the same airmass as the galaxies, with a wide slit to collect all the light, for purposes of flux calibration. The rest of the night, and the next night were wiped out because of poor weather. Two fully flux-calibrated data sets were obtained for IC 3639 and NGC 5135 for various slit orientations and positions.

CTIO spectra were taken for two galaxies (NGC 3393 and IRAS 11249-2859) in 1995 April with the 1.5 m telescope, in combination with the GEC 10 CCD, at f/7.5. The grating used provided a very low spectral resolution of 8.5 Å pixel⁻¹ and a wavelength coverage of 4325-6750 Å. The plate scale was 1".9 pixel⁻¹. The slit width was 3", which was larger than the atmospheric seeing (typically 1".2). The integrations were 900 s in length. The spectra were obtained before we knew what the radio structure of our sample was, and so all the slit positions were taken

along position angle (PA) 90° (N through E). For both the CTIO and AAT spectra we made no correction for telluric absorption.

The fluxes of the emission-lines were corrected for internal reddening using the observed Balmer decrement and assuming a standard interstellar reddening curve (Savage & Mathis 1979). The relative line emissivities for Case B recombination were given by Osterbrock (1989). We used the SPLOT task in IRAF to measure and de-blend the optical features in our spectra. We extracted spatial information of the emission-line gas from the various slit positions, by binning the flux into $\sim 1.4^{\prime\prime}$ increments for the AAT spectra. For the photometric data, the line flux ratios are accurate to the $\pm 10-15\%$ level in the brighter nuclear regions, decreasing to $\pm 20-30\%$ far from the nucleus. Table 4 presents a complete summary of the observations made on each galaxy at radio, NIR, and optical wavelengths.

4. The Radio and FIR Properties

In this section we present our radio observations. In §4.1, we provide quantitative and qualitative definitions of the L-, D- and C-class radio morphologies, along with the presentation of the full-resolution λ 13, 6, and 3 cm contour maps, which are also shown in Figure 2a,b. In §4.2 we present and discuss the radio spectral indices derived from our radio data, and in §4.3, we discuss the FIR/radio correlation and FIR spectral indices, and their role in distinguishing AGN from starburst activity.

4.1. Radio Morphology

The flux densities, morphology, and radio spectral indices derived from the radio images are presented in Table 3. 50% of the galaxies observed at 13 cm show extended emission, and $\sim 33\%$ show extended emission at both 6 and 3 cm. We find a wide variation in radio spectral indices. Free-free absorption is likely to be important in the cores of two systems: NGC 4507 and NGC 7213, and in the outer radio components of IR 11249-2859 (see §4.2). There is also a wide variation in the radio morphologies of the galaxies, which fall into four main classes (e.g. Wilson 1988). The physical significance of these classes is supported by the fact that different optical emission line ratios are associated with the different morphological radio types and that there appears to be some correlation with IRAS color ratios (see §4.3):

1. **D**-class diffuse radio morphology which is characterized by a low surface brightness component surrounding a compact core. The low surface brightness emission is generally extended on scales $\geq 5''$. D-class sources have been linked with starburst activity in that the brightness temperature and spectral index of the radio emission are consistent with a combination of supernova remnants, diffuse synchrotron emission from supernova-accelerated cosmic-ray electrons, and H II regions (e.g. Ulvestad 1982; Condon et al. 1982).

2. L-class linear structure which straddles a compact core believed to be powered primarily by an outflow from an AGN. The outflow may, however, also arise from a circumnuclear starburst (e.g. Baum et al. 1993). We define the radio emission as "linear" if the axial ratio of the linear structure is ≥ 3 . L-class radio sources often have optical emission line ratios consistent with high ionization, high-velocity gas associated with the narrow-line region (NLR) (Haniff, Wilson, & Ward 1988; Wilson, Ward, & Haniff 1988; Whittle et al. 1988), and as such are intimately connected with an AGN. Whittle et al. (1988) studied the velocity structure of the [OIII] λ 5007 emission line in a sample of L-class sources, and found that in some cases there was evidence for an outflow of ionized gas, suggesting that the radio emission of some L-class sources may be due to an outflow of plasma from an AGN.

3. C-class compact sources, which are either unresolved or only marginally resolved on the arcsec scale by our present observations (see Table 2).

The D and L classifications have to be used with care, as they refer to the dominant source of radio flux which can cover a wide range of scales (arc seconds to arc minutes). For example, a faint L-class source may be contained in D-class emission, as in NGC 1068, where a collimated structure is seen inside a starburst disk (e.g. Wilson & Ulvestad 1982b; Wynn-Williams, Becklin, & Scoville 1985). Other ambiguities may also arise. A D-class source seen edge-on might appear as an L-class source. A linear radio structure could be the result of an outflow from an AGN or circumnuclear starburst, but could also be emission from a bar or an edge-on spiral arm. These ambiguities can sometimes be avoided by comparing the optical and radio emission. For example, radio emission from a spiral arm or a bar will be coincident with the optical structure.

As shown in Table 3, our sample contains two D-class, four L-class, one is a hybrid L+D-class (see below), and five C-class sources. IC 3639 and NGC 5135, the only D-class sources in our sample, are nearly face-on galaxies and their D classification was straightforward and not hampered by any orientation effects.

On the other hand, care must be taken in the interpretation of the L-class objects in the sample. In one case, NGC 7130, the elongated radio structure may be due to the existence of an asymmetric stellar bar in the galaxy. This source is known to contain vigorous circumnuclear star-formation (Shields & Filippenko 1990 - see §5.2), and the long-dimension of its λ 13 cm radio structure is oriented along a position angle (PA) 0° which is close to that of an inner bar. We therefore attribute the radio emission in this galaxy to either shocks or star formation within the bar.

However, radio emission from the other L-class galaxies probably represents an outflow from the nucleus. The PA of the linear structure seen in NGC 3393 at 13 cm is 23° and is not coincident with spiral arms or a bar. The asymmetric double radio source seen at 3 cm is also seen in recent VLA 6 cm observations at higher spatial resolution (R. Morganti, private communication). NGC 4507, while not highly collimated, meets our criterion for an L-class, although the knots seen at 6 cm to the north and the south could be due to a "hot spot" ring, extra-nuclear H II regions, or SN near the nucleus, and not a nuclear outflow. As we discuss in §4.2, the radio spectral index of the nucleus of NGC 4507 is flat, suggesting that there is significant free-free absorption. The position angle of the linear structure seen at 13 cm and 6 cm in IRAS 11249-2859 is 125° and is at an angle of 25° with the major axis of the galaxy. It is therefore unlikely that the emission is from H II regions at the end of an edge on spiral arm or from a bar.

4.2. Radio Spectral Indices

There are four main effects which are likely to influence the radio spectral index of Seyfert and starburst galaxies:

1. the synchrotron emission from the interaction of cosmic rays with the interstellar magnetic field. The cosmic rays are generated and re-accelerated by supernovae and supernova remnants, but have a lifetime which extends their range beyond the supernova remnants themselves. This synchrotron radiation is expected to have a spectral index of \sim -0.7. The emission generally dominates the radio power of starburst galaxies, and is presumably responsible for the tight FIR/radio correlation obeyed by these galaxies;

2. the synchrotron emission from relativistic particles generated by the AGN. This radiation could, in principle, cover a wide range of spectral indices. For example, the cores of radio-loud radio galaxies and quasars typically have a flat-spectrum core (because of synchrotron self-absorption) but steep-spectrum extended radio-lobes (because of the cooling of high-energy electrons). However, most Seyfert galaxies have AGN core spectral indices in the region of -0.7. Condon et al. (1991) argue that, amongst infrared-luminous galaxies, even those that are flat-spectrum do not have a sufficiently high brightness temperature for synchrotron self-absorption to be important;

3. thermal emission from H II regions. Compact H II regions in our galaxy generate free-free emission from hot electrons. Most are optically thin, giving a flat spectrum, but most compact H II regions are optically thick at centimeter wavelengths, giving a spectral index ~ 2 . However, Condon et al. have demonstrated that, at centimeter wavelengths, the integrated flux of such regions is generally small compared to the synchrotron emission, so that star-formation regions in starburst and Seyfert galaxies have a typical synchrotron spectrum;

4. the radio emission from ultra-luminous infrared galaxies is optically thick to free-free absorption, so that the typical synchrotron spectrum of these galaxies is flattened at low frequencies (Condon et al. 1991).

The combined result of these effects in Seyfert and starburst galaxies is to produce a typical

radio spectral index of -0.7 with a flattening at low frequencies in some starburst sources because of free-free absorption. In our sample of galaxies, excluding NGC 4507 and NGC 7213, the spectral indices of the cores lie in the range -0.57 to -1.05, which is consistent with a dominant non-thermal synchrotron process, although the cosmic-ray electrons may be associated either with star formation, or else with an AGN core.

NGC 4507 and NGC 7213 differ from the other Seyferts in our sample, having low spectral indices ($\alpha(3,6) = -0.34$ and -0.25, respectively). Other Seyfert galaxies are known to contain flat-spectrum radio sources (Ulvestad & Wilson 1989). These authors suggested that the flat spectral indices they derived for some of their galaxies could arise from synchrotron self-absorption or free-free absorption from the ionized gas. Given that NGC 7213 is a Seyfert 1 galaxy, it is relevant that Antonucci & Barvainis (1988) reported that 8 out of 17 RQQs and Seyfert 1 galaxies imaged at 20, 6 and 2 cm had flat spectral indices, arising from a flat or rising component which becomes dominant by 2 cm. This 2 cm "excess" could be due to free-free emission either from a starburst (requiring a brightness temperature $\leq 10^4$ K) or from $\sim 10^6$ K nuclear free-free emission (Antonucci & Barvainis 1988).

The core of NGC 7213 is marginally resolved by our 1" beam at 3 cm. Assuming a core size equal to our synthesized beam, the brightness temperature of the core in this source is about 3000 K, which is too low for synchrotron self-absorption, suggesting that free-free absorption is responsible for the low spectral index, although free-free emission is also plausible. The free-free absorption may arise from a vigorous starburst present in the nuclear region of this galaxy, or, perhaps, from a high column density narrow-line region. Interestingly, NGC 7213 contains a ring of H II regions 20-40" away from the nucleus (Evans et al. 1996), which, at the distance of this galaxy (see Table 1), corresponds to a ring 2-3 kpc in radius. Yet, our radio observations reveal that in the inner 1" (or inner 100 pc) star-formation is an important contribution to the radio emission. Radio observations of much higher resolution and dynamic range are important for this source in order to understand the morphology of the radio emission.

Unfortunately, the core of NGC 4507 is unresolved at 3 cm so we cannot calculate the

brightness temperature of the emission for that source, and as a consequence we cannot rule out synchrotron self-absorption as the cause for the flat spectral index.

We have obtained radio spectral indices of the knots in the linear structure imaged at 13 and 6 cm, in IRAS 11249-2859. The spectral index, $\alpha(13,6)$, of the first knot to the south-east (see Fig. 2a,b) is -0.88, consistent with synchrotron emission, whereas the first two knots to the north-west are -0.23 and -0.34, respectively, which is consistent with free-free absorption.

4.3. FIR/Radio Correlation and FIR Spectral Indices

The FIR/radio correlation is thought to arise from a common dependence of the FIR and radio emission on the formation of massive stars (e.g. Lisenfeld, Volk, & Xu 1996), and it is generally believed to hold up to high FIR and radio luminosities. Galaxies, or regions within a galaxy, that adhere to the FIR/radio correlation are likely to be regions of massive star-formation. Our sample contains a mix of galaxies that fell on, or were radio-bright with respect to, the FIR correlation, as illustrated by Figure 1, which is based on our 13 cm flux densities (see Table 3) and published IRAS fluxes (see Table 5).

A comparison between the FIR flux and the radio flux is a useful indicator of the dominant FIR and radio emission mechanisms. For example, Helou et al. (1985) define a parameter q, which is defined as:

$$q \equiv \log[(FIR/3.75 \times 10^{12} \text{ Hz})/S_{1.49GHz}]$$
 (1)

where FIR
$$\equiv 1.26 \ge 10^{-14} (2.58 S_{60\mu} + S_{100\mu}).$$
 (2)

It has been shown that q is an indicator of the relative importance of starburst to AGNdominated activity in the nuclear heating of dust in galaxies. For example, the mean value of q for the IRAS Bright Galaxy Sample starbursts is 2.34 ($\sigma_q \sim 0.19$) and for AGNs q < 2 (see Condon et al. 1991).

We present in Table 5 the q-values for our sample, based on IRAS FIR fluxes and on our

13 cm (2.437 GHz) radio flux densities. We use $q \equiv \log[(FIR/3.75 \times 10^{12} \text{ Hz})/\text{S}_{2.4\text{GHz}}]$, similar⁶ to that of Helou et al. (1985). Also presented in Table 5 are the published IRAS FIR fluxes (de Grijp et al. 1992) and the corresponding FIR spectral indices ($\alpha \equiv d \log \text{S/d } \log \nu$). Dust heated primarily by a luminous buried AGN is found in some galaxies, (i.e., the so called "warmer" galaxies) and Vader et al. (1993) suggested that such sources emit strongly near 60 μ m, whereas the UV emission from a pure starburst population is believed to be re-radiated mostly between 60 and 100 μ m. In particular, galaxies with star-formation tend to have steeper far-infrared spectral indices ($\alpha(25,60) \leq -1.4$ and $\alpha(60,100) < -1.1$), and those systems dominated by an AGN have a relatively flat FIR spectrum.

Of the 5 "compact" (C) radio sources in our sample, 4 (IRAS 11215-2806, IRAS 13059-2407, IRAS 18325-5926, & NGC 7213) have FIR flux ratios and q-values that support AGN-dominated FIR and radio emission. These systems probably have very little nuclear/circumnuclear star-formation.

The one exception to this is IC 4995. This galaxy shows unresolved structure in the 3 and 6 cm radio maps and yet has a q-value suggestive of a star formation-dominated galaxy. Spectra, which will be presented in Paper II, show little evidence for significant powerful large-scale star formation in this galaxy which argues against the FIR flux coming from star formation in the outer disk. Furthermore, the point-like nature of the K-band image of this galaxy (not shown) also argues AGAINST disk-wide star formation contributing to the large FIR flux. We therefore conclude that circumstantial evidence points to IC 4995 containing a very compact starburst region which is unresolved by our present observations. Alternatively, IC 4995 may turn out to be the exception to the rule discovered by Helou et al. (that high q values are indicative of starburst-dominated systems). For this reason it would be highly advantageous to obtain higher

⁶Note that we have used a slightly different definition of q from Helou et al. (1985). Assuming a spectral index between 2.4 and 1.49 GHz of -0.7, the borderline between AGN and Starburst would be shifted by 0.14 in dex between the two definitions, which is a small effect

resolution optical, IR and radio observations of this galaxy.

In Fig. 3, we present a plot of the IRAS S(100)/S(12) flux ratio versus S(60)/S(25) ratio to create a color-color diagram for the sample. This combination of IRAS flux ratios has recently been found to be well correlated in samples of Seyfert 2 galaxies (e.g. Heisler, Lumsden, & Bailey 1997) and is being modeled by Dopita et al. (1997). It appears that our galaxies also show a tight correlation in these colors. A lower limit for the S(100)/S(12) flux ratio is shown for NGC 3393, IRAS 13059-2407, and IC 4995. For IRAS 11215-2806, only the S(60) and S(25) band fluxes are known. Reasonable limits on the S(100)/S(12) flux ratio would imply that it too follows the correlation (see vertical line in Fig. 3).

Figure 3 shows a number of interesting properties for the sample. If the radio morphologies of the galaxies are taken into consideration it is clear that the C-class galaxies occupy the lower-left (warmer FIR colors) part of the diagram, whereas the D-class sources (and the one partially resolved C-class source) lie at the cooler end of the correlation. The L-class sources span the entire range, although the two most highly collimated sources, IC 5063 and IRAS 11249-2859 lie in the warmer region of the diagram. This morphological segregation, based on IRAS colors, suggests that the global FIR properties are linked in some way to the radio morphologies.

To understand how this might be, we will consider what has been learned recently about galaxies with small (i.e. warm) S(60)/S(25) IRAS flux ratios. Heisler et al. (1997) have shown that Seyfert 2 galaxies with warmer S(60)/S(25) colors are most likely to show broad permitted lines in their polarized nuclear light (Hidden Broad-line regions-HBLRs), and lower line-of-sight extinction to the nucleus. Within the context of the unified theory of Seyfert galaxies, Heisler et al. provide evidence that the warmer colors are a result of the viewing orientation of the inner dust torus which is postulated to surround the AGN. A dust torus seen closer to face-on would show warmer IRAS colors than those seen more nearly edge-on because the viewer sees more deeply into the inner regions as a result of a higher viewing inclination.

Can the same argument be applied to our sample? IC 5063 (L+D) shows the warmest colors in Fig. 3 and is already know to be a HBLR galaxy, and so its warm colors are consistent with the Heisler et al. picture. A more challenging problem is to explain why the C-class sources should have warmer IRAS colors. One possibility might be that the compact sources are collimated "jet-like" sources seen end-on. If the jets were emitted roughly orthogonally to the dust torus, then this would provide a natural explanation for the warmer colors since the viewing direction would be favorable for seeing deeper into the dust torus. In at least one of the C-class sources (IR 1832-59) the galaxy exhibits high excitation optical emission lines (Isasawa et al. 1995) and is suspected, from X-ray evidence, to contain a Seyfert 1 nucleus. This would seem to support the view that we can almost view directly the Seyfert 1 core.

Certainly the two galaxies with the coolest colors, NGC 5135 and NGC 7130 show evidence for large-scale extended star formation and their FIR colors are likely to be dominated by heat sources spread over a large scale.

However, we caution that our understanding of correlations like that in Figure 3 is still in its infancy, and any possible correlation between warm IRAS colors and radio morphology may be entirely coincidental. Work such as modeling by Dopita et al. (1997) may help to clarify the nature of such a correlation in terms of optical depth effects in the dusty torus at 12 μ m.

5. Optical and Near-Infrared Data and its Relation to Radio Structure

In §5.1 we present the AAT and CTIO optical spectra, and we discuss the correlations of the line ratios with radio structure for IC 3639, NGC 5135, NGC 3393, and IRAS 11249-2859. In §5.2 we present the AAT NIR spectra and imaging, and in §5.3 we discuss in detail the radio structure of IC 5063 and its relation to published optical imaging and spectroscopic observations.

5.1. Optical Spectroscopy of a Subset of the Sample

5.1.1. IC 3639

IC 3639 has been observed spectroscopically and has been imaged in the the H α line (van den Broek et al. 1991; known as galaxy 24SW in that paper) as part of a study of galaxies with large FIR-to-optical luminosities. This Seyfert 2 galaxy shows considerable extended H α emission over much of its compact, high surface-brightness disk. The overall extent of the disk is only 3 kpc. A nearby companion shows a ring of H II regions (van den Broek et al. 1991; galaxy 24NE). IC 3639 has been shown, from optical spectropolarimetry, to contain a hidden broad line region (Heisler et al. 1997). Our 13 cm contour map of this galaxy shows a very rich structure. Surrounding the central core is a diffuse radio structure with several small unresolved knots of emission, some of which are coincident with H α knots, particularly in the northern half of the galaxy. This galaxy is clearly a D-class radio source. The overall envelope of the radio emission down to the level of 0.16 mJy beam⁻¹ follows closely the boundaries of the optical galaxy, suggesting that the radio emission is intimately related to disk processes, i.e., star-formation and supernova events.

Our spectroscopic observations from the AAT reveal highly extended [OI] $\lambda 6300$ and H α emission. We were able to map spatially the optical emission-lines in and near the Seyfert nucleus. Representative spectra are presented in Figure 4a,b,c. In Figure 5, we plot the line ratios obtained for IC 3639 on the standard optical diagnostic diagrams (Baldwin, Phillips & Terlevich 1981; Veilleux & Osterbrock 1987) [OIII] $\lambda 5007/H\beta$ vs [SII] $\lambda \lambda 6731,6716/H\alpha$, [OI] $\lambda 6300/H\alpha$ and [NII] $\lambda 6584/H\alpha$ for PA 90°, centered on the nucleus. The line ratios change from AGN-dominated emission in the nucleus, to H II-region-like emission at a radius of ~1.5 kpc, which suggests that circumnuclear star-formation is important. Note the very blue continuum in Figure 4c, which suggests a very young stellar population. Tables 6a,b,c contain emission-line fluxes and flux ratios for the detected lines at 1".4 intervals from the center for the three observed slit positions (see below).

It is interesting that in two of the line diagnostic diagrams, those involving ratios of [OI] $\lambda 6300$ and [SII] to hydrogen (Fig. 5b,c), the points representing emission just outside the nucleus encroach significantly into the LINER region. We will argue below that this may be a result of enhanced [OI] and [SII] emission caused by weak (150 km s⁻¹) shocks in the circumnuclear environment (Dopita & Sutherland 1995).

A marked east/west asymmetry exists in both the emission lines and the radio distribution (e.g. in Fig. 5a,c, note how the line ratios show this east/west asymmetry). We will discuss this in relation to a simple model of the galaxy below.

Along PA 0° (see Fig. 6a,c) the line ratios change smoothly from AGN to H II-region-like emission with increasing distance from the nucleus. Again we note that the [OI] emission-line ratios dip significantly into the LINER region. Unlike the spectrum taken at PA 90°, there is no noticeable asymmetry on either side of the nucleus.

We also obtained a spectrum 7^{''}.⁵ N of the nucleus along PA 90° (see Fig. 7a,c for BPT and VO diagrams). The [OI] distribution is remarkably asymmetric, showing strong LINER-type emission in the east, but H II-region-like emission in the west. This mimics the behavior at PA 90° through the nucleus, except that no AGN emission is seen.

The radio emission is highly extended and asymmetric at 13 cm, being stronger and more extensive to the north-west (see Fig. 2a). The two spectra which probed the east-west character of the galaxy also reflect an asymmetry. Strong optical emission on the western side of the nucleus, in the region of the high surface-brightness radio emission, shows H II-region-like spectra, whereas to the east the emission is more characteristic of LINER emission, and this occurs on the edge of the rapidly declining radio emission. This behavior is seen in both east-west spectra which cut through the AGN and to the north of the AGN. We note that LINER characteristics (especially enhanced line ratios of [OI] $\lambda 6300$ and [SII] $\lambda \lambda 6716,6731$ to H α) could arise from either a low-powered AGN illuminating ambient narrow-line region gas or from high-velocity shocks (See Dopita & Sutherland, 1995 for discussion). The fact that the spectra appear H II-region-like over the area of the extended radio emission region but LINER-like at the edges of the radio structure, suggest to us that the LINER emission may be from shocked gas resulting from a tipped outflow from a widespread starburst in the central few kpc. The morphology of the radio emission itself is reminiscent of outflows such as those seen in starburst systems like M82. Unfortunately, because the radio surface brightness of the very extended 13 cm emission is low, it is not possible to determine a spectral index of the plasma, but the radio emission may be from relativistic electrons which have burst out of the disk asymmetrically, perhaps from supernova explosions. In this simple picture, gas from the outflow interacts with ambient ISM material to shock and produce a LINER spectrum. If the LINER spectrum were interpreted as coming predominantly from shock waves, they would imply supersonic flow velocities of the order of 150-200 km s⁻¹ for IC 3639 (Dopita & Sutherland 1995). One can test this model further by obtaining higher dispersion spectra to look for mass-motions in the gas in order to identify conclusively a superwind in this galaxy.

An indication that nuclear star-formation may have been episodic in IC 3639 comes from the marginal detection of Balmer absorption (H β) in a ring at about 7" from the nucleus. Three 1-d spectra, at that angular distance from the nucleus, hint at Balmer absorption. However, it is not seen in any of the other extracted 1-d spectra. If these features are confirmed (e.g. by deeper spectra in the blue) this might suggest evidence for a post-starburst population some 1-3 Gyr old. The H β emission line, with faint absorption superimposed, was not corrected for absorption and we caution that the A_v derived from the Balmer decrement may be unreliable. As a consequence the line flux ratios, in these three spectra, could be in error by ~ 30%. We show this feature in Figure 3b.

5.1.2. NGC 5135

NGC 5135 is a barred spiral. The 13 cm radio image (see Fig. 2a) shows considerable extended structure. Our observations show that, in addition to the central extended emission regions, there are knots of emission on a larger scale. In particular, there are two relatively bright knots of emission, one at a position angle of 10°, 18″.4 (4.7 kpc) distant from the central source, and another at 279°, 13″.8 (3.5 kpc) away from the center of emission. Comparison with the optical image shows that the strongest emission is located at the center of the bright optical bar (see Fig. 2a). Ulvestad & Wilson (1989) obtained 6 and 20 cm images of NGC 5135, providing arcsec resolution with the A and B arrays of the VLA. At both wavelengths they saw an asymmetric structure consisting of a bright core with amorphous, fainter extension to the north-east (D-class source). The extension broke up into several smaller knots giving it a bubble-like appearance. All of this structure lies within the inner contours of our 13 cm map.

We obtained optical spectra at the AAT for this galaxy along PA 19° and 45°. The former intersects the radio knot to the north-east and the latter follows the diffuse radio emission detected by Ulvestad & Wilson. Along both position angles, the emission is extended over $\sim 5''$. The optical diagnostic diagrams (see Fig. 8a,c for representative spectra along PA 19° and Fig. 9a,c for BPT and VO diagrams; also see Table 7 for fluxes and flux ratios) along PA 19° reveal that the emission is that of an AGN to the south-west, but changes in the north-east (where one of the radio knots resides) from AGN to LINER to H II region-like. Although not very extended, the optical emission is consistent with star-formation in the circumnuclear region. Again, this confirms that a D-class source has significant amounts of star-formation. Unfortunately, the observing conditions for NGC 5135 became poor, and it is possible that fainter off-nuclear emission was not detected.

5.1.3. NGC 3393

A CTIO spectrum was obtained for NGC 3393 along a position angle of 90° through the nucleus. Because of the very low dispersion, we were unable to de-blend the H α and [NII] $\lambda\lambda 6548,6584$ lines, and so were unable to construct line ratio diagnostics in the nuclear regions. In both the east and west directions, [OI] $\lambda 6300$ is extended over $\sim 2''$ which, at the distance of this galaxy, is ~ 500 pc. The H α is more extended, covering $\sim 10''$, or ~ 2.6 kpc, to the east and $\sim 6''$, or ~ 1.6 kpc, to the west, with a similar morphology to that of the 6 and 3 cm radio emission. The extracted 1-d spectrum for the nucleus of NGC 3393 is presented in Figure 10a.

5.1.4. IRAS 11249-2859

A spectrum was obtained for IRAS 11249-2859 along a position angle of 90° through the nucleus, cutting through the radio knot to the east. We show in Figure 10b a 1-d spectrum of the

nucleus of IRAS 11249-2859.

It is interesting to note that the H α is extended over $\sim 6''$, or 2.7 kpc, and [OI] $\lambda 6300$ is extended over $\sim 2''$, or ~ 1.4 kpc. Proceeding east, away from the nucleus, the [OI] emission at first declines and then increases in strength $\sim 4''$ from the center and then disappears again at larger distances. This increase in flux at 4'' corresponds to the position of the south-east radio knot. A similar increase in line emission at the position of the knot is seen in the (blended) H α & [NII] $\lambda\lambda 6548,6584$ emission and the [SII] $\lambda\lambda 6731,6716$ lines. The ratio of [SII] to H α is 0.39, [NII] $\lambda 6584$ to H α is 0.48, and [OI]/H α is 0.04. Off the nucleus, H β and [OIII] $\lambda 5007$ disappear completely, thus we cannot construct the $[OIII]/H\beta$ ratio which would better determine if the emission is AGN-like or consistent with a star-formation region. We note that these ratios are also consistent with a low to intermediate velocity shock-wave process (Dopita & Sutherland 1995). Given that the optical emission lines are coincident with one of the components of the linear triple radio source (the SE knot), it is clear that follow-up observations of higher sensitivity along a PA of 125° are highly desirable. We speculate that these observations might reveal that the nuclear outflow is causing wide scale shocks in the disk of this galaxy giving rise to the enhanced [OI] and possibly inducing secondary star-formation, consistent with the flat radio spectral indices of the knots to the NW.

5.2. Near-Infrared Spectra and Imaging

All four galaxies observed spectroscopically in the NIR (IRAS 18325-5926, IC 4995, IC 5063, and NGC 7213; see Table 4) show detectable emission lines of $Br\gamma$, H_2 , and [FeII]. The nuclear spectra are presented in Figure 11.

Unlike the other galaxies observed at NIR wavelengths, IRAS 18325-5926 shows no evidence for an underlying stellar continuum (in particular the CO (2-0) band-head is missing). This is consistent with significant obscuration, even at infrared wavelengths, by dust near the nucleus. Unusually high levels of obscuration in this probable heavily embedded Seyfert 1 nucleus was also inferred from the optical and X-ray observations of Isasawa et al. (1995). We note that the lack of the CO band-head could be due to a stronger continuum from the Seyfert 1 nucleus.

NGC 7130 may be an example of a collisional ring galaxy (e.g. see the review by Appleton & Struck-Marcell 1996) formed as a result of a small companion passing through the disk of a larger system. In the radio, at 13 cm, this galaxy is clearly extended with a linear north-south structure (L-class source). However, the origin of the linear structure is unclear since it is coincident with a bar along the same PA.

This galaxy has been studied in detail by Shields & Filippenko (1990), who spectrally mapped the nuclear regions of the galaxy. Their work clearly shows the composite nature of this galaxy, in that the line intensity ratios vary radially from those indicative of AGNs, close to the nucleus, to that of H II regions at increasing distances from the center.

We obtained a K-band continuum infrared image of NGC 7130 (Fig. 12a,b), but no line data were obtained. In the high-surface-brightness inner regions, two tightly-wrapped spiral arms emanate from a somewhat asymmetric bar. The arms almost form a ring or lens component. The bright knot in the disk of the galaxy is a foreground star. However, the high-contrast, grey-scale image shown in Figure 12b clearly shows the existence of a large faint ring of emission which is confirmed by the Digitized Sky Survey (DSS) plate of the area. A small companion galaxy lies embedded in this outer ring and Figure 12 also shows an additional faint plume to the south-west of the nuclear regions which may be connected to the outer ring.

NGC 7130 has the highest FIR luminosity, $\log[FIR/L\odot]=11.04$, of all the galaxies in our sample. Much of the star-formation that powers ultra-luminous far-infrared galaxies is believed, in part, to be triggered by interactions. While this image alone provides no evidence for the star-formation properties of this galaxy, its morphology alone strongly suggests that it is interacting.

5.3. Radio Structure of IC 5063 and its Relation to Published Optical Spectra and Imaging

IC 5063 is a Seyfert 2 galaxy which is known to be abnormally radio-loud (Danziger, Goss, & Wellington 1981; Roy & Norris 1997). We measure a flux density of 747 mJy at 13 cm. This galaxy, along with IRAS 13059-2407, is the furthest from and thus the most "radio-loud" with respect to, the FIR/radio correlation for our sample. At 3 cm the emission is in the form a linear structure composed of the bright central nucleus with two distinct knots towards the south-east, along PA $115\pm3^{\circ}$. Morganti, Oosterloo, & Tsvetanov (1997) have also imaged this galaxy at 3 cm and our maps agree very well in morphology and flux. At 6 cm, the remnants of the linear structure can be seen, but a faint shell of emission is seen extending from the end of the linear structure, perhaps indicative of a bow-shock. The diffuse emission observed at 6 cm may result from the interaction of the linear structure, i.e. "plasmons" ejected from the AGN, with the ambient ISM in the inner disk. This galaxy is a hybrid L+D-class (see Figure 2b), since it is an L-class source at 3 cm and a D-class source at 6 cm. In Figure 13, we label the bright central source (at 3 cm) as knot A, the first knot to the south-east as knot B, and the second, fainter radio source as knot C. From our K-band spectra (see Fig. 11) we extract a $\log[H_2/Br\gamma]$ ratio of 0.14, which is confirmation of an AGN in the nucleus (Mouri & Taniguchi 1992). AGN-dominated FIR emission is suggested by the FIR flux ratios and the q-value for this galaxy (see Table 5).

IC 5063 is fascinating because it may be a merging system which contains one of the best-defined ionization cones known to date (Colina, Sparks, & Macchetto 1991). It is an S0 galaxy with very peculiar dust lanes and emission-lines which define a cross-shape centered near the AGN. Colina et al. have suggested that the cross-shape has some of the characteristics of an ionization cone illuminated by an AGN. To explain the increase in the excitation of the ionized gas with distance from the center, they suggested that this may be the result of decreasing density and a strong metallicity gradient in the central few arcsec. However, they could not completely rule out a local source of heating in the cone. Of special significance is the fact that the orientation of the linear triple is coincident with the inner edge of the ionization cone reported by Colina et al.

(1991), in just the region where the ionized gas shows enhanced excitation. Knot B (see Fig. 13), in the 3 cm map, seems to coincide exactly with the peak in the brightness profile for both H α and [OI] λ 6300 emission. The fact that the peak in the emission line strength was offset from the "nucleus" was discussed at length by Colina et al. but was not explained. However, these authors did note that the velocity of the peak differed by 80 km s⁻¹ from the surrounding fainter gas. We will argue below that this may be a signature of ionized gas being ejected supersonically into the surrounding ISM in the form of plasmons.

In an H α imaging survey of galaxies with far-infrared spectral energy distributions peaking near 60 μ m, Heisler & Vader (1995) showed that this galaxy has three bright knots along, roughly, but not exactly, the same position angle as the radio knots seen in our 3 cm image. It is interesting to note that radio knots A and B have similar strengths in H α . We have recently obtained new IRIS NIR long-slit spectra along the radio structure seen at 3 cm (to be published later). The knot identified in the radio image as B contains both a NIR continuum source and lines of [FeII] $\lambda 1.644\mu$ m and Br γ , with fainter IR emitting knots on either side (this confirms the existence of the nuclear [FeII] $\lambda 1.644\mu$ m detected in our spectra obtained in 1994 October). Thus, the Heisler & Vader H α imaging and our NIR spectroscopy suggest that the nucleus is knot B, although this is by no means clear. In particular, the absolute positions of the optical and infrared data are uncertain by about an arcsec, and cannot therefore be accurately superimposed on the radio image.

Figures 13a,b show that there is a slight difference in relative position between the H α knots (shown as crosses) and the 3 cm radio knots. Figure 13a assumes that radio knot B is the nucleus and, hence, the H α knot and knot B are made to align, whereas figure 13b assumes that radio knot A is the nucleus. In each case, the other H α knots are offset from the radio knots. We therefore speculate, regardless of whether knot A or knot B is the nucleus, that these offsets are consistent with plasmons being ejected into a counter-clockwise rotating disk. The plasmons ram into the ambient ISM causing enhanced H α emission along their leading edges. We note that if two "extra-nuclear" radio blobs are indeed moving at a minimum of 80 km s⁻¹ relative to

the surrounding gas, they would be highly supersonic relative to their surroundings and might naturally produce bow-shocks on their leading edge. It is therefore significant that the 6 cm map shows a bow-shock-like morphology ahead of the possible ejected plasma (such emission might be produced by electrons trapped in tangled magnetic fields in the bow-shock ahead of the ejected material). Further radio observations of higher dynamic range and spatial resolution may help to define this morphology better. Other evidence to support the view that gas outflow from the AGN is interacting with the ISM of the inner disk are broad (~600 km s⁻¹), blueshifted absorption features seen against a disk of neutral hydrogen emission (Morganti et al. 1997).

We emphasize that we are not suggesting that the overall morphology of the optical emission-line properties of IC 5063 are driven by the radio source, but rather that additional heating is occurring along one edge of the ionization cone that seems to be related to the radio structure.

6. Conclusions

1) Our sample contains a mixture of radio morphologies. There are four L-class sources, two D-class sources, one L+D-class, and five compact (unresolved or marginally resolved on the arcsec scale) sources. Four of the galaxies in the sample (NGC 4507, IC3639, NGC 5135 and NGC 7130) show evidence for being AGN/starburst composites, and a 5th galaxy (IC 4995) is likely to contain a compact starburst core on a scale smaller than our arc second (r < 400 pc) resolution based on its IR and radio characteristics. Further optical properties of the galaxies not studied optically here will be presented in Paper II.

2.) There is a correspondence between D-class radio morphology and circumnuclear star-formation. Our optical spectroscopic observations of IC 3639, luminous in the FIR $(\log[FIR/L\odot]=10.33)$, show clearly the importance of circumnuclear star-formation and the composite AGN/starburst nature of this galaxy. The radio emission has a clear east-west asymmetry that is seen also in the optical spectra, particularly in the (highly extended) [OI] λ 6300

emission line. The radio knots appear to be contained within a region of highly extended optical [OI] $\lambda 6300$ emission. Our BPT diagrams reveal that the emission changes within the inner 1.5 kpc from AGN, through LINER, to H II-region-like spectra radially away from the nucleus. The [OI] emission that is detected to the east of the nucleus is consistent with emission that is being driven by large scale shocks (v ≈ 150 km s⁻¹), perhaps being generated by the outflow of material from a nuclear starburst. In a similar vein, our optical spectroscopy of NGC 5135 also clearly demonstrates the importance of circumnuclear star formation. Along PA 19°, the emission-line ratios change radially away from the nucleus, changing from AGN to LINER to H II-region-like emission.

3.) Some morphological segregation occurs when the galaxies global FIR colors are investigated. For example, on a plot of IRAS S(100)/S(12) versus S(60)/S(25), the C-class sources cluster to warmer IRAS colors than the D-class sources. The L-class sources have a wide range of color temperatures, but the hottest galaxies are those with well defined highly collimated radio emission. It is not clear why the radio morphology and the IRAS colors are related, unless the higher colors relate to the orientation of the dust torus (see for example Heisler et al (1997), and Dopita et al. (1997)), and by implication, the radio jet direction.

4.) Our CTIO spectra for NGC 3393 and IRAS 11249-2859 show that the radio and optical emission-line gas is extended on roughly the same scale. In particular we detected, in [OI] λ 6300, the radio knot to the south-east in IRAS 11249-2859. NGC 7130, another of the L-class sources, has been shown in the work of Shields & Filippenko (1990) to contain a significant amount of circumnuclear star-formation. The L classification of this galaxy is probably due to a bar along the same PA as the radio emission. Because of the lack of optical and NIR spectra we cannot make any conclusions about the correlation of, or lack thereof, radio morphology with circumnuclear star-formation for the L-class sources. Future papers are planned in order to shed light on this;

5.) The radio spectral indices for the core-regions of the sample are generally in a range consistent with synchrotron emission, either from the AGN itself or from cosmic rays accelerated by star-formation activity. Two notable exceptions are NGC 4507 (L-class) and NGC 7213, whose

low spectral indices indicate that they may suffer from free-free absorption of a thermal source. A likely source for the free-free absorption in the marginally resolved radio source NGC 7213 might be circumnuclear star-formation, as suggested by the brightness temperature of the radio emission. Furthermore, the collimated structure seen in IRAS 11249-2859 contains knots that are flat-spectrum sources suggesting that they might be undergoing free-free absorption;

6.) IC 5063, an L+D class radio source, has IRAS fluxes and a q-value consistent with AGN-dominated emission, although spectroscopic confirmation of this statement is pending. The radio emission seen at 6 and 3 cm suggest that "plasmons" are being ejected from the nucleus into the disk of the galaxy, giving rise to extensive shocks, possibly a bow-shock morphology at 6 cm, and significant local heating of one edge of a well-defined ionization cone seen in optical emission lines. In the nucleus NIR spectra revealed $Br\gamma$, H_2 and [FeII] (along with the CO (2-0) bandhead feature);

7.) Our infrared observations of IRAS 18325-5926, IC 4995, and NGC 7213 reveal H₂, Br γ , and [FeII] λ 1.644 μ m. Also, we detect the CO (ν =2-0) band-head absorption feature, longward of 2.3 μ m, in IC 4995, IC 5063 and NGC 7213.

Acknowledgments

We are grateful to Russell Cannon for granting us director's time at the AAT in order to obtain near-infrared spectroscopy presented in this paper. We would also like to thank the staff at both the Paul Wild Observatory and Siding Spring Observatory for their very generous advice and help. M.A. Bransford thanks Sammy for his warm and friendly demeanor during visits to Australia, and to R. Morganti for very valuable comments, which helped to improve the paper. We thank STSci/SERC for use of the digitized sky survey. We would also like to thank the referee for very valuable comments and suggestions that helped to improve the paper. The Australia Telescope is funded by the Commonwealth of Australia for operation as a National Facility managed by CSIRO. This work is partly funded by an NSF/US-Australia Cooperative Research Grant INT-9418142. A. Marston was supported by NASA JOVE grant NAG8-264 and a grant from NASA administered by the American Astronomical Society.

REFERENCES

- Antonucci, R., & Barvainis, R. 1988, ApJ, 332, L13
- Appleton, P.N., & Struck-Marcell, C. 1996, in Fundamentals of Cosmic Physics, 16, 111
- Baldwin, J.A., Phillips, M.M., & Terlevich, R.J. 1981, PASP, 93,5
- Baum, S. A., O'Dea, C. P., Dallacassa, D., de Bruyn, G., & Pedlar, A. ApJ, 1993, 419, 553
- Bolton, J.G., & Shimmins, A.J. 1973 AuJPA, 30, 1
- Colina, L., Sparks, W.B., & Macchetto, F. 1991, ApJ, 370, 102
- Condon, J.J., Huang, Z.P., Yin, Q.F., & Thuan, T.X. 1991, ApJ, 378, 65
- Condon, J.J., Condon, M.A., Gisler G., & Puschell, J.J. 1982, ApJ, 252, 102
- Danziger, I.J., Goss, W.M., & Wellington, K.J. 1981, MNRAS, 196, 845
- de Grijp, M.H.K., Keel, W.C., Miley, G.K., Goudfrooij, P., & Lub, J. 1992, A&AS, 96, 389
- Dopita M.A., & Sutherland, R.S. 1995 ApJ, 455, 468
- Dopita, M.A., Heisler, C., Lumsden, S., & Bailey, J. 1997, in preparation
- Evans, I.N., Koratkar, A.P., Storchi-Bergmann, T., Kirkpatrick, H., Heckman, T.M., & Wilson, A.S. 1996, ApJS
- Genzel, R., Weitzel L., Tacconi-Garman, L.E., Blietz, M., Cameron M., Krabbe, A., Lutz, D., & Sternberg, A. 1995, ApJ, 444, 129
- Haniff, C.A., Wilson, A.S., & Ward, M.J. 1988, ApJ, 334, 104
- Heckman, T.M., Gonzalez-Delgao, R., Leitherer, C., Meurer, G.R., Krolik, J., Wilson, A.S., Koratkar, A., & Kinney, A. 1997, ApJ, 482, 114

- Heisler, C.A., & Vader, J.P. 1995, AJ, 110, 87
- Heisler, C.A., & Vader, J.P. 1994, AJ, 107, 35
- Heisler, C.A., Lumsden, S.L., & Bailey, J.A. 1997, Nature, 385, 700
- Helou, G., Soifer, B.T., & Rowan-Robinson, M. 1985, ApJ, 298, L7
- Isasawa, K., Kunieda, H., Tawara, Y., Awaki, H., Koyamo, K., Murayama, T., & Taniguchi, Y. 1995, AJ, 110, 551
- Lester, D.F., Joy, M., Harvey, P.M., Ellis, H.B., & Parmer, P.S. 1987, ApJ, 321, 755
- Lisenfeld, U., Volk, H.J., & Xu, C. 1996, A&A, 314, 745
- Morganti, R., Oosterloo, T., & Tsvetanov, Z. 1997, in Emission Lines in Active Galaxies: New Methods and Techniques, PSAP Conference Series, vol. 113, edited by B.M. Peterson, F.-Z. Chang, & A.S. Wilson, pg. 310
- Mouri, H. & Taniguchi, Y. 1992, ApJ, 386, 68
- Norris, R. P., Allen, D. A., & Roche, P. F. 1988, MNRAS, 234, 773
- Osterbrock, D.E. 1989, in Astrophysics of Gaseous Nebulae and Active Galactic Nuclei, University Science Books
- Phillips, M.M., Charles, P.A., & Baldwin, J.A. 1983, ApJ, 266, 485
- Roy, A.L., & Norris, R.P. 1997, MNRAS, in press
- Roy, A.L., Norris, R.P., Kesteven, M.J., Troup, E.R., & Reynolds, J.E. 1997, in preparation
- Savage, B.D., & Mathis, J.S. 1979, ARA&A, 17, 73
- Shields, J.C., & Filippenko, A.V. 1990, AJ, 100, 1034
- Stone, R.P.S., & Baldwin, J.A. 1983, MNRAS, 204, 347

- Telesco, C. M., & Decher, R. 1988, ApJ, 334, 573
- Telesco, C.M., Becklin, E.E., Wynn-Williams, C.G., & Harper, D.A. 1984, ApJ, 282, 427
- Ulvestad, J.S., & Wilson, A.S. 1989, ApJ, 343, 659
- Ulvestad, J.S. 1982, ApJ, 259, 96
- Vader, J.P., Frogel, J.A., Terndrup, D.M., & Heisler, C.A. 1993, AJ, 106, 1743
- van den Broek, A.C., van Driel, W., de Jong, T., Lub, J., de Grijp, M.H.K., & Goudfrooij, P. 1991, A&AS, 91, 61
- Veilleux, S., & Osterbrock, D.E. 1987 ApJS, 63, 295
- Whittle, M., Pedlar, A., Meurs, E.J.A., Unger, S.W., Axon, D.J., & Ward, M. 1988, ApJ, 326, 125
- Wilson, A.S., & Ulvestad, J.S. 1982b, ApJ, 263, 576
- Wilson, A.S. 1988, A&A, 206, 41
- Wilson, A.S., Ward, M.J., & Haniff, C.A. 1988, ApJ, 334, 121
- Wynn-Williams, C.G., Becklin, E.E., Scoville, N.Z. 1985, ApJ, 297, 607

This manuscript was prepared with the AAS LAT_EX macros v4.0.

Table 1. The Sample

Name	$\begin{array}{c} \mathrm{RA}(\mathrm{J2000})^{\mathrm{a}}\\ \mathrm{(h\ m\ s)} \end{array}$	DEC(J2000) ^a (d m s)	$m_{ m b}{}^{ m b}$ (mag)	Redshift ^c	Distance ^d (Mpc)	$Scale^d$ (kpc arcsec ⁻¹)	Type ^e	Diameter ^b (arcmin)
NGC 3393	10 48 23.4	$-25 \ 09 \ 44.0$	13.09	0.0137	54.8	0.266	Sy 2	$2.2 \ge 2.0$
IRAS 11215-2806	$11\ 24\ 02.7$	$-28 \ 23 \ 15.3$	13.00	0.0135	54.0	0.262	Sy 2	— x —
IRAS 11249-2859	$11\ 27\ 23.4$	$-29\ 15\ 27.5$	14.71	0.0234	93.6	0.454	Sy 2	$1.6 \ge 0.4$
NGC 4507	$12 \ 35 \ 36.7$	-39 54 34.5	12.92	0.0132	52.8	0.256	Sy 2	$1.7 \ge 1.3$
IC 3639	$12 \ 40 \ 52.8$	$-36\ 45\ 22.4$	13.00	0.0125	50.0	0.242	Sy 2	$1.2 \ge 1.2$
IRAS 13059-2407	$13\ 08\ 42.0$	$-24 \ 22 \ 58.0$	16.00	0.0141	56.4	0.273	Sy 2	— x —
NGC 5135	$13\ 25\ 44.0$	-29 50 00.2	12.88	0.0132	52.9	0.256	Sy 2	$2.6 \ge 1.8$
IRAS 18325-5926	$18 \ 36 \ 58.3$	$-59 \ 24 \ 08.2$	13.20	0.0192	76.8	0.372	Sy 2	— x —
IC 4995	$20\ 19\ 59.0$	$-52 \ 37 \ 18.6$	14.28	0.0163	65.2	0.316	Sy 2	$1.2\ge 0.8$
IC 5063	$20\ 52\ 02.1$	$-57 \ 04 \ 06.7$	12.89	0.0113	45.2	0.219	Sy 2	$2.1 \ge 1.4$
NGC 7130	$21 \ 48 \ 19.5$	-34 57 04.0	12.98	0.0159	63.6	0.308	Sy 2	$1.5 \ge 1.4$
NGC 7213	22 09 16.2	$-47 \ 10 \ 00.2$	11.01	0.0050	20.0	0.097	Sy 1	$3.1 \ge 2.8$

 $^{\rm a}3$ cm radio positions; except NGC 5135 which is 13 cm

^bTaken from the NED database

^ctaken from de Grijp et al. (1992), except NGC 7130 which is taken from van den Broek et al. (1991) and NGC 5135 which is taken from Phillips et al. (1983)

 $^{\rm d} {\rm Based}$ on a Hubble constant of $H_{\rm o} = 75 \ {\rm km \ s^{-1} \ Mpc^{-1}}$

^eTypically Seyferts are typed as follows: emission line galaxies which contain significantly broader (by $\sim 500 \text{ km s}^{-1}$) permitted lines than the forbidden narrow lines are classified as Seyfert 1 (Sy1), narrow line objects that have [OIII] $\lambda 5007/\text{H}\beta > 3$ and [NII] $\lambda 6584/\text{H}\alpha > 0.5$ are classified as Seyfert 2 (Sy 2)

Name	wavelength	Beam	rms noise
	(cm)	(arcsec x arcsec)	$(mJy beam^{-1})$
NGC 3393	13	7.22 x 3.95	0.09
	6	$4.07 \ge 1.91$	0.10
	3	$2.46 \ge 1.04$	0.19
RAS 11215-2806	13	8.00 x 3.42	0.14
	6	$3.85 \ge 1.78$	0.11
	3	$2.12 \ge 0.99$	0.19
RAS 11249-2859	13	7.85 x 3.17	0.04
	6	$3.58 \ge 1.86$	0.11
	3	$2.00 \ge 1.01$	0.19
NGC 4507	13	$5.24 \ge 3.93$	0.17
	6	$2.58 \ge 1.98$	0.10
	3	$1.76 \ge 1.01$	0.16
C 3639	13	$5.33 \ge 3.86$	0.05
	6	$2.78 \ge 2.02$	0.07
	3	$1.55 \ge 1.15$	0.13
RAS 13059-2407	13	$7.95 \ge 3.74$	0.22
	6	4.35 x 1.93	0.27
	3	$2.38 \ge 1.08$	0.20
NGC 5135	13	$6.83 \ge 3.72$	0.09
RAS 18325-5926	13	$4.75 \ge 3.28$	0.07
	6	$2.35 \ge 1.55$	0.08
	3	$1.30 \ge 0.87$	0.10
C 4995	13	$5.01 \ge 3.16$	0.07
	6	$2.44 \ge 1.80$	0.13
	3	$1.34 \ge 0.98$	0.18
C 5063	13	4.23 x 3.57	0.72
	6	$2.32 \ge 1.74$	0.51
	3	$1.31 \ge 0.97$	0.16
IGC 7130	13	$6.19 \ge 3.51$	0.15
	6	$3.17 \ge 1.86$	0.10
	3	1.71 x 1.01	0.11
NGC 7213	13	4.93 x 3.78	0.12
	6	$2.27 \ge 1.94$	0.15
	3	$1.26 \ge 1.08$	0.16

Table 2. Clean beam parameters and rms noise

Table 3. Radio Properties

Name	S_{13cm}^{a} (mJy)	S _{6cm} ^a (mJy)	S _{3cm} ^a (mJy)	$ Log(L_{13cm}) W Hz^{-1} $	Morphology ^b	$\alpha(6,13)^{\rm c}$	$\alpha(3,13)^{d}$	$\alpha(3,6)^{\mathrm{e}}$
NGC 3393	40.9 ± 1.1	20.5 ± 0.4	13.3 ± 1.0	22.14	L	-0.91 ± 0.03	$-0.79 {\pm} 0.02$	-0.62 ± 0.09
IRAS 11215-2806	33.0 ± 0.7	20.0 ± 0.1 20.0 ± 0.4	12.0 ± 0.8	22.04	C	-0.72 ± 0.03	-0.67 ± 0.02	-0.75 ± 0.07
IRAS 11249-2859	$33.4 {\pm} 0.7$	$22.3 {\pm} 0.5$	$8.8 {\pm} 1.0$	22.52	\mathbf{L}	$-0.62 {\pm} 0.04$	$-0.73 {\pm} 0.03$	$-0.79 {\pm} 0.12$
NGC 4507	$23.4{\pm}2.0$	$10.8 {\pm} 0.4$	$8.8{\pm}0.6$	21.87	\mathbf{L}	$-0.84{\pm}0.09$	$-0.59 {\pm} 0.05$	$-0.34{\pm}0.11$
IC 3639	$42.0{\pm}0.5$	$18.2{\pm}0.3$	$9.5{\pm}0.6$	22.07	D	$-0.88 {\pm} 0.03$	$-0.92{\pm}0.01$	$-1.01{\pm}0.06$
IRAS 13059-2407	346 ± 2	215 ± 1	108 ± 1	23.10	С	$-0.71 {\pm} 0.01$	$-0.93 {\pm} 0.01$	$-1.17 {\pm} 0.02$
NGC 5135	112 ± 2	—	_	22.55	D	—	—	—
IRAS 18325-5926	123 ± 1	$67.2{\pm}0.4$	35.5 ± 0.3	22.91	С	$-0.88 {\pm} 0.01$	$-0.97{\pm}0.01$	$-1.05 {\pm} 0.02$
IC 4995	$8.2{\pm}0.5$	$5.2{\pm}0.5$	$3.1{\pm}0.7$	21.70	\mathbf{C}	$-0.61 {\pm} 0.08$	$-0.69 {\pm} 0.05$	$-0.71 {\pm} 0.16$
IC 5063	747 ± 12	389 ± 5	$214{\pm}10$	23.24	L+D	$-0.96 {\pm} 0.02$	$-0.98 {\pm} 0.01$	$-1.05 {\pm} 0.05$
NGC 7130	$97.7 {\pm} 0.7$	$59.0{\pm}0.8$	$30.8 {\pm} 0.4$	22.65	\mathbf{L}	$-0.67 {\pm} 0.01$	$-0.84{\pm}0.01$	$-0.98 {\pm} 0.01$
NGC 7213	168 ± 1	207 ± 1	176 ± 1	21.88	С	$+0.30{\pm}0.01$	$+0.02{\pm}0.01$	$-0.26 {\pm} 0.02$

^aExtended + core flux from BLANKed images

 $^{\rm b}{\rm L}$ = Linear Class, D = Diffuse Class, C = Compact Class

 $^{\rm c}{\rm Core}$ spectral index for 13 and 6 cm

 $^{\rm d}{\rm Core}$ spectral index for 13 and 3 cm

 $^{\rm e}{\rm Core}$ spectral index for 6 and 3 cm

Name	Imaged in radio?	Optical long-slit spectra?	NIR long-slit spectra?	NIR imaging?	
NGC 3393	yes	yes	no	no	
IRAS 11215-2806	yes	no	no	no	
IRAS 11249-2859	yes	yes	no	no	
NGC 4507	yes	no	no	no	
IC 3639	yes	yes	no	no	
IRAS 13059-2407	yes	no	no	no	
NGC 5135	yes	yes	no	no	
IRAS 18325-5926	yes	no	yes	yes	
IC 4995	yes	no	yes	yes	
IC 5063	yes	no	yes	yes	
NGC 7130	yes	no	no	yes	
NGC 7213	yes	no	yes	yes	

Table 4. Observations made on the sample

Table 5. Published IRAS Fluxes, and computed q-values

Name	$12 \mu \mathrm{m}$ (Jy)	$25 \mu \mathrm{m}$ (Jy)	$60 \mu m$ (Jy)	$100 \mu { m m}$ (Jy)	α (12,25)	α (25,60)	α (60,100)	$\log [FIR/L\odot]$	q	AGN/starburst composite
NGC 3393	< 0.25	0.71	2.38	3.94	< -1.42	-1.38	-0.99	10.07	1.92	
IRAS 11215-2806	< 0.44	0.31	0.59	<1.00	< 0.48	-0.74	> -1.03	< 9.45	<1.41	
IRAS 11249-2859	0.11	0.34	0.60	0.94	-1.54	-0.65	-0.88	9.92	1.40	
NGC 4507	0.46	1.42	4.52	5.40	-1.54	-1.32	-0.35	10.16	2.39	Х
IC 3639	0.66	2.32	7.08	10.77	-1.71	-1.27	-0.82	10.33	2.37	Х
IRAS 13059-2407	< 0.25	0.71	1.41	1.75	< -1.42	-0.78	-0.41	9.82	0.72	
NGC 5135	0.64	2.40	16.91	28.64	-1.81	-2.23	-1.03	10.92	2.34	Х
IRAS 18325-5926	0.60	1.39	3.17	4.09	-1.14	-0.94	-0.50	10.49	1.53	
IC 4995	< 0.25	0.36	0.90	1.28	< -0.50	-1.05	-0.69	9.77	2.17	X? ^a
IC 5063	1.17	3.87	5.89	4.26	-1.63	-0.48	0.63	10.19	0.94	
NGC 7130	0.59	2.12	16.48	25.57	-1.75	-2.34	-0.86	11.04	2.37	Х
NGC 7213	0.63	0.74	2.56	8.63	-0.22	-1.42	-2.38	9.52	1.43	х

^aInferred from high q-value–see text

Table 6a. IC 3639 reddening-corrected emission-line ratios, relative to $H\beta = 100$, at P.A. 90° centered on nucleus

distance ^a (arcsec)	${\rm H}\beta^{\rm b}$ (x 10 ⁻¹⁵ ergs s ⁻¹ cm ⁻²)	A_v^c (mag)	$[OIII]$ $\lambda 4959$	$[OIII]$ $\lambda 5007$	[OI] λ6300	$[\rm NII] \\ \lambda 6548$	$[NII] \\ \lambda 6583$	$[SII] \\ \lambda 6717$	$[SII] \\ \lambda 6731$
10.0	1.0	1.04	69.9	0.45	20.0	22.0	105	20.0	54.0
-19.6	1.3	1.64	63.3	245	30.2	23.0	105	80.9	54.0
-18.2	5.0	2.74	161	361	62.4	27.1	117	64.7	55.2
-16.8	61.0	1.31	241	693	50.8	82.1	239	57.2	63.4
-15.4	5.3	2.04	110	286	29.0	36.0	110	66.1	49.0
-14.0	5.5	2.05	107	290	23.3	39.7	121	75.6	50.3
-12.6	3.6	1.92	99.4	305	26.4	43.8	140	78.9	50.4
-11.2	2.3	1.69	127	346		48.9	163	76.6	63.4
-9.8	6.5	2.71	194	502	34.0	50.4	159	82.5	49.3
-8.4	2.5	1.55	115	302	34.1	48.0	156	74.2	48.4
-7.0	4.1	1.36	95.3	275	23.5	47.4	145	60.0	50.4
-5.6	6.8	1.23	96.8	485	29.1	64.0	166	67.4	57.9
-4.2	13.1	0.87	235	656	44.0	69.5	219	56.6	54.9
-2.8	61.0	1.31	241	693	50.8	82.1	239	57.2	63.4
-1.4	372	2.20	259	745	50.5	89.0	240	54.0	59.4
0.0	863	1.35	336	891	55.4	97.3	238	56.8	69.7
+1.4	54.1	3.29	201	399	47.7	88.0	210	64.3	52.1
+2.8	16.9	2.45	68.6	153	17.9	50.2	138	57.6	42.3
+4.2	18.1	2.49	60.2	134	23.8	46.3	128	59.1	42.0
+5.6	7.3	2.27	83.6	132	19.6	43.0	141	67.8	51.0
+7.0	1.9	1.24	68.6	124	20.9	49.3	134	80.9	57.2
+8.4	2.1	0.68	38.9	90.4	12.5	36.3	109	55.8	41.0
+9.8	4.8	1.01	35.4	90.0	6.2	33.9	96.2	47.1	36.2
+11.2	10.6	1.85	36.1	104		31.8	90.5	45.5	35.3
+12.6	6.0	1.85	44.8	105	15.4	31.4	98.3	53.1	37.0
+14.0	4.8	1.91	40.3	111		29.8	93.7	57.1	40.1
+15.4	3.4	1.78	63.2	145	21.9	28.4	98.2	61.2	44.1
+16.8	4.3	2.22	64.2	233		27.0	97.9	59.5	47.2
+18.2	2.2	1.94	101	199	22.5	40.9	107	70.1	46.4

 a Negative angular distances are to the east of the nucleus, positive angular distances are west of the nucleus, and 0.0 indicates the nucleus

 $^{\rm b}{\rm H}\alpha$ fluxes may be obtained by multiplying these fluxes by 2.85

^ccalculated from the Balmer decrement assuming Case B recombination

$distance^{a}$	${ m H}eta^{ m b}$	$A_v{}^c$	[OIII]	[OIII]	[OI]	[NII]	[NII]	[SII]	[SII]
(arcsec)	$(x \ 10^{-15} ergs \ s^{-1} \ cm^{-2})$	(mag)	$\lambda 4959$	$\lambda 5007$	$\lambda 6300$	$\lambda 6548$	$\lambda 6583$	$\lambda 6717$	$\lambda 6731$
-15.4	4.3	1.22	49.4	136	9.7	23.8	76.2	45.7	36.4
-14.0	4.9	1.30	48.2	122	16.5	26.4	81.7	48.8	36.6
-12.6	3.3	0.94	34.0	105	14.1	29.2	87.5	51.7	38.9
-11.2	3.1	0.83	33.7	91.0	14.6	29.3	91.7	55.4	37.2
-9.8	3.6	0.72	28.7	84.4	19.2	30.6	91.1	51.8	34.5
-8.4	4.5	0.53	28.5	77.1	12.9	30.5	90.1	47.5	33.6
-7.0	5.9	0.49	31.1	81.0	9.1	28.5	85.9	44.8	31.9
-5.6	10.6	0.91	31.0	82.2	6.4	27.4	83.8	41.3	29.2
-4.2	10.0	1.12	31.4	74.1	6.8	30.9	91.5	45.4	33.9
-2.8	8.1	1.06	29.4	80.6	9.9	33.7	96.2	49.7	35.7
-1.4	7.9	1.00	26.9	83.6	8.2	32.8	96.3	47.7	35.8
0.0	14.9	1.49	30.3	72.5	5.8	31.5	100	48.9	35.6
+1.4	21.1	2.07	28.8	69.7	9.3	33.2	105	52.6	38.4
+2.8	13.3	2.11	37.1	75.9		36.5	108	56.7	42.3
+4.2	6.1	1.61	36.9	77.5		33.0	113	57.5	46.2
+5.6	4.1	1.28	29.1	67.7		35.7	110	61.2	43.7
+7.0	6.4	1.86	36.0	85.3	10.2	31.6	107	60.0	44.2
+8.4	4.5	1.76	34.4	71.1		25.6	102	65.6	41.8
+9.8	4.7	2.12		123		30.9	106	65.2	48.5
+11.2	5.4	2.35		124		30.9	89.3	66.2	46.4
+12.6	4.5	2.28		158		32.2	94.7	64.0	56.0

Table 6b. IC 3639 reddening-corrected emission-line ratios, relative to $H\beta = 100$, at P.A. 90°,

7".5 North of nucleus

^aNegative angular distances are to the east, positive angular distances are west, and 0.0 indicates the point 7...⁵ directly north of the nucleus

 $^{\rm b}{\rm H}\alpha$ fluxes may be obtained by multiplying these fluxes by 2.85

^ccalculated from the Balmer decrement assuming Case B recombination

Table 6c. IC 3639 reddening-corrected emission-line ratios, relative to $H\beta = 100$, at P.A. 0° centered on nucleus

distance ^a (arcsec)	${\rm H}\beta^{\rm b}$ (x 10 ⁻¹⁵ ergs s ⁻¹ cm ⁻²)	A_v^c (mag)	$[OIII] \\ \lambda 4959$	$[OIII]$ $\lambda 5007$	$[OI]$ $\lambda 6300$	$[NII]$ $\lambda 6548$	$[NII]$ $\lambda 6583$	$[SII] \\ \lambda 6717$	$[SII] \\ \lambda 6731$
-19.6	1.1	1.78	65.1	151		35.7	90.4	90.4	41.7
-13.0 -18.2	1.5	1.44	59.2	191	18.3	35.6	96.6	82.0	49.3
-16.2	2.8	1.44	61.1	132	11.4	27.7	94.9	56.7	45.8
-15.4	3.8	1.63	55.0	98.4	13.3	29.4	94.7	57.7	38.6
-14.0	4.9	1.83	45.6	115	9.8	23.4 27.7	100	57.0	41.5
-12.6	4.6	1.61	40.0 34.7	97.8	12.1	30.0	100	55.8	42.7
-11.2	4.6	1.47	49.9	84.3	12.1	33.1	101	51.6	37.8
-9.8	8.0	1.98	10.0	84.8	8.7	29.2	102	52.8	34.7
-8.4	17.0	2.50		112	20.9	37.4	122	47.4	38.1
-7.0	20.6	2.43	58.3	140	21.5	40.3	131	55.8	41.4
-5.6	25.4	2.55	164	422	31.4	56.3	173	58.3	51.2
-4.2	59.9	2.46	302	810	50.3	86.5	228	60.3	58.4
-2.8	111	1.88	262	734	50.8	93.7	243	57.6	59.5
-1.4	317	2.02	263	737	51.1	92.4	241	57.7	59.6
0.0	189	2.72	297	848	58.2	89.9	237	54.9	57.3
+1.4	60.3	2.80	191	608	49.1	60.4	188	50.4	47.9
+2.8	22.5	2.00	60.4	139	15.1	39.0	127	46.2	35.4
+4.2	17.9	1.50	24.7	74.9	13.1	34.2	105	45.4	33.8
+5.6	19.8	1.58	26.0	72.7	6.7	30.9	97.0	44.1	33.7
+7.0	20.5	1.92	30.2	91.2	9.3	30.1	92.2	44.1	34.0
+8.4	11.4	1.84	36.1	108	32.8	31.3	92.1	47.2	34.6
+9.8	8.4	1.54	39.0	101	12.2	28.5	87.4	45.4	34.7
+11.2	7.3	1.16	40.9	118	8.0	26.7	82.6	43.5	33.3
+12.6	9.6	1.23	49.9	135	8.5	23.5	75.9	45.2	32.7
+14.0	8.2	1.16	44.3	131	13.2	21.4	76.5	43.8	31.6
+15.4	4.9	1.34	34.6	133	18.8	23.1	77.5	51.3	34.2
+16.8	2.2	1.68	47.3	132		20.6	84.6	58.9	38.9

 a Negative angular distances are to the south of the nucleus, positive angular distances are north of the nucleus, and 0.0 indicates the nucleus

 $^{\mathrm{b}}\mathrm{H}\alpha$ fluxes may be obtained by multiplying these fluxes by 2.85

^ccalculated from the Balmer decrement assuming Case B recombination

-39-

Table 7. NGC 5135 reddening-corrected emission-line ratios, relative to $H\beta = 100$, at P.A. 19° centered on nucleus

distance ^a (arcsec)	${ m H}\beta^{ m b}$ (x 10 ⁻¹⁵ ergs s ⁻¹ cm ⁻²)	A_v^c (mag)	[OIII] $\lambda 4959$	$[OIII]$ $\lambda 5007$	$[OI]$ $\lambda 6300$	$[NII] \\ \lambda 6548$	$[NII]$ $\lambda 6583$	$[SII] \\ \lambda 6717$	$[SII] \\ \lambda 6731$
-2.8	394	4.10	268	647	18.3	73.4	198	40.9	36.8
-1.4	203	2.68	205	572	16.9	86.2	222	43.9	38.3
0.0	140	1.96	154	409	22.4	97.9	253	50.9	41.7
+1.4	125	1.87	146	408	24.0	110	281	61.5	48.2
+2.8	89.6	1.85	116	324	27.5	110	287	66.5	49.3
+4.2	37.6	1.76	85.9	232	29.5	43.5	114	27.5	21.0
+5.6	23.6	2.19	61.0	167	14.5	64.4	169	47.0	40.7

 a Negative angular distances are to the south-west of the nucleus, positive angular distances are north-east of the nucleus, and 0.0 indicates the nucleus

 $^{\rm b}{\rm H}\alpha$ fluxes may be obtained by multiplying these fluxes by 2.85

^ccalculated from the Balmer decrement assuming Case B recombination

Legends for Figures

Fig. 1.— FIR-radio correlation for sample. Note how the galaxies lie on the "radio-bright" side of the correlation. NGC 4507 has only upper limits for its FIR luminosity.

Fig. 2.— (a) 13 cm radio - optical overlays. The optical images were obtained from the Digitized Sky Survey (J plates). The numbers at the top of the overlays are thousands of counts, representing the stretch for the grey-scale. At the lower right is an inset showing more clearly the radio structure.
(b) 6 and 3 cm radio maps. The galaxies are presented two by two, from left to right are the lower and higher frequencies, respectively. In each case, at all frequencies, the contours are at (-2.5, 2.5, 4, 8, 16, 32, 64, 128, 256, 512, 1024) times the rms noise (see Table 3 for rms values) measured in a source-free area of the map.

Fig. 3.— A FIR color-color plot of the IRAS $S(100 \ \mu m)/S(12 \ \mu m)$ flux ratio versus the $S(60 \ \mu m)/S(25 \ \mu m)$ ratio. Solid squares represent the C-class sources, the open square with the circle inside represents the marginally resolved source (NGC 7213), the crosses represent the L-class sources, and the open circles represent the D-class sources. In the case of IRAS 11215-2806, only the S(60) and S(25) fluxes are known, and the unconstrained S(100)/S(12) ratio is denoted by an open square with a vertical line running through it. Interestingly, there is a segregation of the morphological classes in the color-color plot: the compact sources lie in the lower left (warm FIR colors) and the diffuse sources (and the marginally resolved source) lie from the middle to the upper right (cool FIR colors). The L-class sources span the whole range, but the most highly collimated sources reside in the lower left.

Fig. 4.— Spectra of IC 3639, PA 90°, going from AGN (a) to LINER (b) to H II-region-like (c). Note in that in (b) we marginally detect an H β absorption feature. Confirmation of this feature is pending. The vertical scale is F_{λ} (ergs cm⁻² s⁻¹ Å⁻¹) and the wavelengths are in the rest frame of the galaxy. Fig. 5.— BPT and VO diagrams for IC 3639, PA 90°. For this figure, and for the following BPT and VO diagrams: (a) is $\log([NII]/H\alpha)$ vs $\log([OIII]/H\beta)$, (b) is $\log([SII]/H\alpha)$ vs $\log([OIII]/H\beta)$, (c) is $\log([OI]/H\alpha$ vs $\log([OIII]/H\beta)$, and the points proceed radially away from the nucleus APPROXIMATELY from upper right to lower left with 1".4 increments. Of interest in these figures is the east-west asymmetry. To the east of the nucleus, where there is a rapidly declining radio emission, the optical emission is consistent with a LINER and to the west, the emission is more like that of H II-regions. This effect is seen in all three diagnostics.

Fig. 6.— BPT and VO diagrams for IC 3639, PA 0°. Again, the optical emission proceeds, radially away from the nucleus, from AGN to H II-region-like. Note that south of the nucleus the emission proceeds to H II-region-like at a larger relative distance than to the north. The emission to the south dips into the LINER region (at ~ 5 ."6), but only significantly in the [OI] diagnostic.

Fig. 7.— BPT and VO diagrams for IC 3639, PA 90°, 7^{\prime}.⁵ North of the nucleus. This emission is consistent with H II-regions. Of interest is (c), which mimics the east-west asymmetry seen in Figure 5a,c. To the east, just north of the nucleus, the emission is consistent with a LINER, and not until ~5^{\prime}.⁶ east does it dip into the H II-region portion of the graph.

Fig. 8.— Spectra of NGC 5135, along PA 19°, going from AGN (a) to LINER (b) to H II-region-like (c). The vertical scale is F_{λ} (ergs cm⁻² s⁻¹ Å⁻¹) and the wavelengths are in the rest frame of the galaxy.

Fig. 9.— BPT and VO diagrams for NGC 5135, PA 19°. Of interest is a north-east south-west asymmetry. To the north-east (there is a radio knot at 13 cm along this position angle, see Fig. 2a) the emission goes from AGN to LINER to H II-region-like, whereas to the south-west the emission remains consistent with an AGN.

Fig. 10.— CTIO spectra. (a) Nuclear spectrum for NGC 3393, along PA 90°. (b) Nuclear spectrum for IRAS 11249-2859 along PA 90°. The vertical scale is F_{λ} (ergs cm⁻² s⁻¹ Å⁻¹) and the wavelengths are in the rest frame of the galaxy.

Fig. 11.— K-band IR spectra for IRAS 18325-5926, IC 5063, NGC 7213, and NGC 4995. Note that we have detected H_2 and $Br\gamma$ in all four cases. Also detected in all cases, except IRAS 1832, is the CO 2-0 band-head absorption feature.

Fig. 12.— K-band continuum IR images of NGC 7130. Figure 12a shows a contour plot of the intensity of the K-band continuum emission. It clearly shows the two tightly wrapped spiral arms that emanate from a somewhat asymmetric bar. In Figure 12b, the high-contrast grey-scale image, one can clearly see the ring-like structure that connects with the companion to the north-west.

Fig. 13.— 3 cm radio image of IC 5063 overlayed with the centroid positions of the H α knots (taken from Heisler & Vader, 1995) which are represented by crosses. (a) assumes that radio knot B is the nucleus and the H α knot and knot B are made to align. (b) assumes that radio knot A is the nucleus and the H α knot and knot A are made to align. The radio contours are 0.16 mJy beam⁻¹ x (-2.5,2.5,3,4,8,10,16,20,25,30,32,40,50,64,70,90,128,256,512,800).

This figure "sey_fig1.jpg" is available in "jpg" format from:

This figure "sey_fig2a_1.jpg" is available in "jpg" format from:

This figure "sey_fig2b_1.jpg" is available in "jpg" format from:

This figure "sey_fig2a_2.jpg" is available in "jpg" format from:

This figure "sey_fig2b_2.jpg" is available in "jpg" format from:

This figure "sey_fig2b_3.jpg" is available in "jpg" format from:

This figure "sey_fig3.jpg" is available in "jpg" format from:

This figure "sey_fig2b_4.jpg" is available in "jpg" format from:

This figure "sey_fig4.jpg" is available in "jpg" format from:

This figure "sey_fig5.jpg" is available in "jpg" format from:

This figure "sey_fig6.jpg" is available in "jpg" format from:

This figure "sey_fig7.jpg" is available in "jpg" format from:

This figure "sey_fig8.jpg" is available in "jpg" format from:

This figure "sey_fig9.jpg" is available in "jpg" format from:

This figure "sey_fig10.jpg" is available in "jpg" format from:

This figure "sey_fig11.jpg" is available in "jpg" format from:

This figure "sey_fig12a.jpg" is available in "jpg" format from:

This figure "sey_fig12b.jpg" is available in "jpg" format from:

This figure "sey_fig13.jpg" is available in "jpg" format from: