A FRESH LOOK AT THE ADHESION APPROXIMATION

T. BUCHERT

Theoretische Physik, Ludwig-Maximilians-Universität Theresienstr. 37, D-80333 München, Germany

I report on a systematic derivation of the phenomenological "adhesion approximation" from gravitational instability together with a brief evaluation of the related status of analytical modeling of large—scale structure.

Hectored by steadily improving computer power, N-body simulations suggestively establish themselves as the final answer to the modeling of large-scale structure in the Universe, admitting their limitations only in covered dynamical range and lack of "input physics". Still, the analytical approach provides for a complementary way of understanding. Ironically, "understanding" large-scale structure formation is itself becoming a discipline of N-body computing. Notwithstanding, a brief reminder of the line of development of analytical modeling is in order.

Well before we finally saw the structures, which are generically encountered around us, also in the Universe, Zel'dovich put forward a kinematical picture of large—scale structure formation into that what we nowadays call network, web, filaments, sheets and voids. Here, understanding preceded observation and finalized the necessity of departure of N-body simulations from spherical symmetry. Thereafter, almost the same picture is repeatedly presented. We learnt that N-body results down to the scales where we may consider them reliable can be understood, if not mirrored, by refining Zel'dovich's step (e.g. Lagrangian perturbation theory, now conquering the general—relativistic regime; optimized, i.e., high—frequency truncated Lagrangian schemes). One may consider it good or bad news that N-body results in gross terms agree with such simple models (for further reading see the review by Sahni & Coles¹ and references therein).

Analytical insight has concentrated on the issue of "shell–crossing" or "multi–streaming", which is not a problem of N–body computing, but of analytical modeling using "dust matter". Known for a long time outside of cosmology, Shandarin has particularly pushed the idea that this shortcoming after caustic formation can be overcome by Burgers–type equations for an "adhesive" modeling of structures²: an additional force proportional to the Laplacian of the peculiar–velocity is capable of stabilizing collapsed structures.

In a recent work³ we have taken up this line by demonstrating that a systematic derivation of the adhesion model from gravitational instability is feasible. It is important to realize that also collisionless matter features pressure—

like forces arising from the action of multi-streaming, a well-known fact in stellar systems theory. Forming velocity moments of the collisionless Boltzmann equation, we arrive at the following hierarchy of evolution equations:

$$\partial_t \varrho + (\varrho \bar{v}_i)_{,i} = 0 \quad , \tag{1}$$

$$\partial_t \bar{v}_i + \bar{v}_j \bar{v}_{i,j} = g_i - \frac{1}{\varrho} \Pi_{ij,j} \quad , \tag{2}$$

$$\partial_t \Pi_{ij} + \bar{v}_k \Pi_{ij,k} + \bar{v}_{k,k} \Pi_{ij} = -\Pi_{ik} \bar{v}_{j,k} - \Pi_{jk} \bar{v}_{i,k} - L_{ijk,k} \quad , \tag{3}$$

and so on, where $\Pi_{ij} = \varrho \langle (v_i - \bar{v}_i)(v_j - \bar{v}_j) \rangle$ is the tensor of velocity dispersion taking the multi–stream force into account, \bar{v}_i and g_i are the components of the mean velocity and the gravitational field strength obeying Newton's field equations, respectively, and ϱ is the mass density. We have suggested to close this hierarchy by assuming that velocity dispersion is small, but non–vanishing. Hence, beyond Jeans' equation (2) we only keep the evolution equation for the dispersion tensor (3), but neglect the contribution from the third–rank tensor $L_{ijk,k}$. Two further crude⁴ approximations then lead us to the key equation of the adhesion approximation: we only consider the isotropic part of the dispersion tensor, $\Pi_{ij} \approx p\delta_{ij}$ and, after transforming to peculiar–fields, we solve the resulting equation by iteration and find the first iterate to obey³:

$$\frac{d\tilde{\mathbf{u}}}{db} = \mu \Delta_{\mathbf{q}} \tilde{\mathbf{u}} \quad ; \quad \frac{d}{db} := \frac{\partial}{\partial b} + \tilde{\mathbf{u}} \cdot \nabla_{\mathbf{q}} \quad , \quad \tilde{\mathbf{u}} := \frac{\mathbf{u}}{a\dot{b}} \quad , \quad \mathbf{q} := \frac{\mathbf{x}}{a} \quad , \tag{4}$$

where **u** is the peculiar-velocity, a(t) is the scale factor of the background cosmology, b(t) is the growing mode solution of the linear theory, and μ is a function of time and density. This density dependence disappears for $p \propto \varrho^2$, whereas from (3) $p \propto \varrho^{5/3}$. In view of the restrictions done there is plenty of room for a generalization of the adhesion approximation, which may lead us to a deeper understanding of structure formation in the nonlinear regime.

Acknowledgments

This work is supported by the Sonderforschungsbereich SFB 375 für Astro-Teilchenphysik der Deutschen Forschungsgemeinschaft.

References

- 1. V. Sahni, P. Coles (1995) Physics Reports 262, 1.
- 2. S.N. Gurbatov, A.I. Saichev, S.F. Shandarin, (1989) M.N.R.A.S. 236, 385.
- 3. T. Buchert, A. Domínguez, A.& A., submitted (1997).
- 4. T. Buchert, A. Domínguez, J. Pérez-Mercader, Ap.J., submitted (1997).