On the Consistency between the Outer Rotation Curve of the Galaxy and the Microlensing Rate towards the Large Magellanic Cloud

Zong-Hong Zhu

Institute of Theoretical Physics, Chinese Academy of Sciences, Beijing 100080, China

and

Xiang-Ping Wu

Beijing Astronomical Observatory, Chinese Academy of Sciences, Beijing 100080, China

ABSTRACT

Recent determination of the outer rotation curve of the Galaxy suggests that the total Galactic mass is likely to concentrate into a region of radius ~ $2R_0$. If this finding is correct, the previously estimated optical depth of microlensing events towards the Large Magellanic Cloud (LMC) should be correspondingly reduced. It appears that a truncated Galactic halo model instead of the standard isothermal one extending to LMC accounts naturally for the reported microlensing rate by the MACHO collaboration. Such an explanation is indeed obvious yet has been overlooked in the past.

Subject headings: dark matter – Galaxy: halo – gravitational lensing

More than dozen microlensing events towards the Large Magellanic Cloud (LMC) have been so far detected by the MACHO and EROS collaborations (Alcock et al. 1997; Renault et al. 1997). An analysis of the first two year result of the MACHO collaboration gives rise to the microlensing rate of $\tau = 2.1^{+1.1}_{-0.7} \times 10^{-7}$. This is only about half of the value $\tau_G = 5 \times 10^{-7}$ for a standard spherical Galactic halo model (Paczyński 1986; Griest 1991), but is of an order of magnitude larger than the optical depth by the machos and stars in the LMC halo/disk itself (Wu 1994; Sahu 1994; Gould 1995). Such a discrepancy, together with the puzzle of the too large mass of $0.1-1M_{\odot}$ required for the machos in Galactic halo, has led many authors to make attempt at modifying the conventional models for the Galactic halo and the macho distributions between the Milky Way and LMC (e.g. Ansari, et al. 1996; De Paolis et al. 1996; Zhao 1997; Gates et al. 1997). It seems that many efforts are still needed before a better explanation of the microlensing result and a clear understanding of structure of the Galactic halo are achieved.

Here, we propose an alternative yet natural way to reconcile the microlensing rates observed towards LMC with the measurement of outer rotation curve in the Galaxy. This is primarily motivated by the recent discovery of the Keplerian rotation curves of some galaxies including the Milky Way (Honma & Sofue 1996, 1997; Binney & Dehnen 1997; Ryder et al. 1997; and references therein). These studies claimed that the declining rotation curves are observed in a great number of galaxies, implying that the possible truncation of galactic halos may occur at a certain radius of several disk scale lengths. Detection of such Keplerian rotation curves challenges the traditional point of view about the mass distribution in galaxies, which may have a significant impact on the study of dark matter in galactic halos including the ongoing searches for machos in the Galactic halo. Recall that the flat rotation curves beyond the optical disks are the unique convincing evidence for the presence of a large amount of dark matter in galactic halos (see Ashman 1992 for a review). In particular, the flat rotation curves indicate that the galactic halos can in principle extend to rather a large distance (> 100 kpc) with a density profile of r^{-2} . This uncomfortable "standard" model of dark galactic halo has been, nevertheless, accepted by astronomical society. Yet, it is a logical and also natural result that the dark halos should be truncated at some radii R_{cut} , no matter how large R_{cut} would be for various galaxies. In a sense, the recent detection of the Keplerian rotation curves of galaxies are not beyond expectation.

What would happen to the theoretical prediction of the optical depth τ_G if the galactic halo shows a declining tendency at a radius R_{cut} ? The analysis by Honma & Sofue (1996) and Binney & Dehnen (1997) indicates that $R_{cut} \approx 2R_0$, where R_0 is the distance from the Sun to the Galactic center. Their best-fit values of the the galactic constants Θ_0 , the circular velocity at R_0 , and R_0 are $\Theta_0 \approx 200 \text{ km s}^{-1}$ and $R_0 \approx 7.6 \text{ kpc}$, respectively. These parameters are somewhat smaller than the IAU standard values. Applying these values for a truncated isothermal Galactic halo model up to $R_{cut} = 15$ kpc instead of the traditional halo model extending to the LMC, we find the optical depth of microlensing towards LMC to be only $\tau_G \approx 2 \times 10^{-7}$. This estimate should be regarded as a low bound on the optical depth since the rotation curve does not show an abrupt decline at R_{cut} . Rather, as argued by Honma & Sofue (1996) and Binney & Dehnen (1997), the Keplerian velocity decreases very gently in the outer region, which may account for the fact that the flat rotation curves often turn out to be a good approximation for most of galaxies. In the case of the Milky Way, the circular velocity only reduces to 190 km s⁻¹ even at $3R_0$. So, a robust estimate of the optical depth of microlensing towards LMC depends on how a realistic model of the Galactic halo is constructed, which should take the variation of the rotation velocity along radius into account. We will not explore the detailed model in the present letter. Overall, a truncated Galactic halo model can easily provide an optical depth that is consistent with the result by the MACHO collaboration.

Does a truncated Galactic halo model affect our estimate of the macho mass ? To answer the question, we have computed the probability P that a microlensing event of duration T and maximum magnification μ_{max} be produced by a macho with mass m using the formulism of Jetzer & Massó (1994). The maximum of P gives rise to the most probably value of mr_E^2/v^2T^2 , where $r_E^2 = 4GM_{\odot}D_s/c^2$, D_s is the distance to the LMC and v is the relative velocity of traverse motion of the macho. Numerical computation shows that the position of the maximum of P shifts from $mr_E^2/v^2T^2 = 13.0$ to 17.9 when the extent of the Galactic halo is truncated at $2R_0$ instead of D_s . It turns out that the estimate of the most probably value of macho mass is nearly unaffected by the truncation of the Galactic halo if a circular velocity of $\Theta_0 = 200$ km s⁻¹ (Honma & Sofue 1996) or 180 km s⁻¹ (Olling & Merrifield 1997) instead of 220 km s⁻¹ is adopted. In other words, a truncated Galactic halo model does not help to release the difficulty of the unreasonably large macho mass (~ $0.5M_{\odot}$) inferred from the current MACHO experiment.

We emphasize in this brief note the consistency between the recent determination of the outer rotation curve of the Galaxy (Honma & Sofue 1996; Binney & Dehnen 1997) and the observed microlensing rate towards the LMC (Alcock et al. 1997). These measurements may significantly change our previous view of the dark Galactic halo. In any case, the straightforward inference that machos contribute a fraction of ~ 50% to the halo dark matter based on the standard isothermal model and the reported microlensing rate could be misleading.

This work was supported by the National Science Foundation of China.

REFERENCES

Alcock, C. et al. 1997, ApJ, 486, 697
Ansari, R. et al. 1996, A&A, 314, 94
Ashman, K. M. 1992, PASP, 104, 1109
Binney, J., & Dehnen, W. 1997, MNRAS, 287, L5
De Paolis, F., Ingrosso G., & Jetzer P. 1996, ApJ, 470, 493
Gates, E. I. Gyuk, G. Hold, G. P., & Turner, M. S. 1997, ApJ, submitted (astro-ph/971110)
Griest, K. 1991, ApJ, 366, 412
Gould, A. 1995, ApJ, 441, 77
Honma, M., & Sofue Y. 1996, PASJ, 48, L103
Honma, M., & Sofue Y. 1997, PASJ, 49, 453
Jetzer, Ph., & Massó E. 1994, Phys. Lett. B323, 347
Olling, R. P., & Merrifield, M. R. 1997, preprint astro-ph/9711157
Paczyński, B. 1986, ApJ, 304, 1
Renault, C. et al. 1997, A&A, 324, L69

Ryder, S. D. et al. 1997, MNRAS, in press
Sahu, K. C. 1994, Nature, 370, 275
Wu, X. P. 1994, ApJ, 435, 66
Zhao, H. 1997, MNRAS, submitted (astro-ph/9703097)

This 2-column preprint was prepared with the AAS ${\rm LATEX}$ macros v4.0.