SiO Maser Forest at the Galactic Center

Hideyuki Izumiura

Okayama Astrophysical Observatory, National Astronomical Observatory, Kamogata, Asakuchi, Okayama 719-02, Japan e-mail: izumiura@oao.nao.ac.jp

Shuji Deguchi

Nobeyama Radio Observatory, National Astronomical Observatory, Minamimaki, Minamisaku, Nagano 384-13, Japan e-mail: deguchi@nro.nao.ac.jp

and

Takahiro Fujii

Department of Astronomy, Faculty of Science, the University of Tokyo Bunkyo-ku, Tokyo 110, Japan e-mail: fujii@astron.s.u-tokyo.ac.jp

Received July 9, 1997; accepted December 5, 1997

(ApJL in press ;version 1.6, 11/17/97)

ABSTRACT

A moderately deep survey of stellar maser sources toward the Galactic center has been made in the SiO J=1-0 v=1 and v=2 transitions (~ 43 GHz) with the Nobeyama 45-m telescope of a beam size of about 40". At the Galactic center (Sgr A*), 7 (4 new) SiO maser sources were detected in one beam. In order to investigate the spatial distribution of SiO maser sources, we have also observed the offset positions by \pm 40" in Galactic longitude from the Galactic center. In total, 7 (4 new) SiO maser sources were detected at the offset positions. Taking into account the shorter integration time at the offset positions, we find that the surface number density of SiO maser sources is nearly constant at the Galactic center. The number density of SiO maser sources is found to be an order of magnitude higher than the density of OH 1612 MHz objects. A radial-velocity gradient in Galactic longitude was not detected. These facts indicate that the SiO maser sources seen toward the Galactic center are mostly associated with the stellar population of a Galactic stellar nuclear disk of more than a few arc minute radius.

Subject headings: galaxy:center — masers — radio lines:molecular :circumstellar — stars:late-type

1. Introduction

It is known that a cluster of evolved stars surrounds the Galactic center (Blum et al. 1996; Morris & Serabyn 1996)¹. These evolved stars very often exhibit molecular line masers (Winnberg 1996) and have been used for studying the structure of the Galactic center disk. OH 1612 MHz and SiO 43 GHz maser lines have been used to obtain radial velocities for sources in the Galactic center region (Lindqvist et al. 1991; Lindqvist et al. 1992a), for the Galactic bulge IRAS sources (Izumiura et al. 1995), and for sources toward the Sgr B2 molecular cloud (Shiki et al. 1997). Maser sources very close to the Galactic center are useful to establish a concordance between radio and infrared position-reference frames (Menten et al. 1997), identifying the Sgr A* in the infrared (Eckart et al. 1993), and studying proper motions (Eckart & Genzel 1997). With the Very Large Array (VLA), sensitive surveys of maser sources near Sgr A* have been made in the OH 1612 MHz line (e.g., Sjouwerman et al. 1997). A VLA survey in SiO/H₂O lines (Menten et al. 1997) was made with velocity resolution of 2.7 km s⁻¹ with 32 channels, resulting in two new SiO maser sources associated with known infrared objects very near Sgr A*. The detection limit of this VLA survey was at the level of ~0.05 Jy at 43 GHz.

In this paper, we report an attempt of surveying SiO maser sources near the Galactic center with the Nobeyama 45-m telescope. We have detected 7 SiO sources at the Galactic center. In order to investigate the source density near the Galactic center we have also observed at the two offset positions by $\pm 40''$ in Galactic longitude, resulting in detections of similar number of sources.

¹We refer to the dynamical center of our Galaxy as the radio continuum source, Sgr A^* , in this paper.

2. Observations

Simultaneous observations in the SiO J=1-0 v=1 and 2 transitions at 43.122 and 42.821 GHz, respectively, were made with the 45-m radio telescope at Nobeyama on April 25, May 10, 11, and 12, 1997. A cooled SIS receiver with a band width of about 0.4 GHz was used and the system temperature (including atmospheric noise) was 190-200 K. The aperture efficiency of the telescope was about 0.60 at 43 GHz. The half-power beam width (HPBW) was about 40" at 43 GHz. A factor, 3.6 Jy/K, was used to convert antenna temperature to flux density. An acousto-optical spectrometer array of a low-resolution (AOS-W) was used. Each spectrometer has a 250 MHz band width and 2048 frequency channels, giving the velocity coverage of about 1700 km s⁻¹ and the spectral resolution of 1.7 km s⁻¹ (per two binned channels). Observations were made in a position switching mode and the off position was chosen 10' away from the Galactic center in azimuth.

We observed three positions near the Galactic center: one at the position of Sgr A^{*}, (RA, Dec, Epoch)=(17h42m29.314s, -28° 59′ 18.3″, 1950) [(l,b)=(359.944°, -0.046°); Rogers et al. 1994], and the other two at the positions offset by one beam on either side of Sgr A^{*} in the Galactic longitude, $(\Delta l, \Delta b)=(\pm 40'', 0'')$. We performed the observations on clear nights without wind. Telescope pointing was carefully checked before and after each observation using a nearby strong SiO maser source, OH2.6-0.4. The average pointing accuracy was confirmed to be better than 5″. The total integration time was 6.8 h for the center and 3.0 h each for the offset positions. For the center, $(\Delta l, \Delta b)=(0'', 0'')$, we combined the spectra taken on April 25 and May 10, 1997. For the offset positions, data were taken on May 11 and May 12. Weather condition on May 12 was not perfect when the position (+40'', 0'') was observed; T_{sys} was about 220 K and the wind speed about 5 m s⁻¹ (which might have caused the pointing error of about 5″). Thus, the noise level on the spectrum at (+40'', 0'') was slightly higher than that on spectra taken on the other days.

Obtained spectra toward Sgr A^{*} exhibited a baseline offset of about 3.2 K from zero due to the continuum emission. This corresponds to a flux density of about 12 Jy at 43 GHz, which is consistent with the previous measurement of the Sgr $A^*(11.6 \text{ Jy})$ at 43 GHz with the 45-m system (Sofue et al. 1986). The spectra exhibited a baseline distortion (of about 0.3 K at the maximum) and weak ripples probably due to standing waves in the 45-m telescope system. The ripples in the velocity range of ± 350 km s⁻¹ of the SiO lines were relatively weak (of about 0.02 K on average). In order to remove complex ripple features from the spectra, we have taken running means of the spectra (average of 100 channels, or about the 80 km/s width) and averaged spectra were subtracted from originals. With this procedure, the baseline of the resulting spectra became quite flat. Because the SiO maser lines are quite narrow (the widths less than 10 km/s) and weak (T $_a < 0.2$ K), this method seems to work well. There are other molecular and atomic lines possibly contaminating the SiO J=1-0 v=2 line at 42.821 GHz: ²⁹SiO J=1-0, v=0 (42.879 GHz), H53 α (42.891 GHz), SiC_4 J=14-13 (42.945 GHz), and C_6H J=31/2-29/2 (42.970 and 42.970 GHz). These lines might appear as broad features at the Galactic center, and may overlap somewhat with the SiO J=1-0 v=2 line, causing a bad baseline. With this baseline removal, broadline features of the above mentioned molecules, even if present, were eliminated. The spectra finally obtained are shown in figure 1. The rms noise temperatures at the center, $(\Delta l, \Delta b) = (0'', 0'')$, are 0.004 and 0.009 K at 42.821 GHz (SiO J=1-0, v=2) and 43.122 GHz (SiO J=1-0, v=1), respectively. The spectrometer used for the v=1 line gave systematically higher rms value of noise. This is probably due to a calibration error of the photo-diode levels in the spectrometer.

EDITOR: PLACE FIGURE 1 HERE.

Detections of the SiO maser components were judged by the following criteria (both must be satisfied).

- Line intensities in either transition must be greater than 5 times the RMS noise level with line widths greater than three channels (2.5 km s^{-1}) .
- Emission peaks are seen at the same velocity in both the J=1-0, v=1 and 2 transitions (within 2 km s⁻¹), or at least, a peak is seen in one transition and a positive deflection (not trough) is seen in the other transition.

These criteria worked nicely in our experiences, guarding against identification as signal of anomalous noises known in AOS data. For the case of a short integration, the second criterion has been quite useful and follow-up integrations revealed the SiO lines very often.

SiO maser components detected at the three positions are shown in table 1. It is well known that the SiO maser velocity of a single source coincides with the stellar velocity, i.e., with the middle of the OH 1612 MHz double-peak velocities within a few km s⁻¹ (Jewell et al. 1991; Jiang et al. 1995). Therefore, if the velocity of the detected SiO feature is close to the velocity of an already known OH 1612 MHz source at the Galactic center, it is highly probable that the SiO source is associated with the OH source. With this velocity coincidence, we have assigned probable associations with OH/IR sources in table 1.

EDITOR: PLACE TABLE 1 HERE.

3. Discussion

The radial velocity of SiO maser emission indicates the stellar velocity, i.e., the velocity of a central star. Velocity widths are known to be less than 10 km s⁻¹ in most of SiO maser sources. Especially for the case of a source at the Galactic center, we can detect only a narrow emission peak and not a weak broad pedestal. From these facts, we can

safely assume that the detected SiO peaks, if separated more than 10 $\rm km\,s^{-1}$, come from individual sources in the telescope beam. In the following discussion, we regard each SiO component listed in table 1 as an individual source.

3.1. Associations with previously known objects at the Galactic center

Extensive surveys of the Galactic-center region have been made before with the VLA in OH 1612 MHz (Lindqvist et al. 1992b; Sjouwerman & van Langevelde 1996) and in H₂O and SiO lines (Levine et al. 1995; Menten et al. 1997). Because sources detected in this paper are within a 1' radius from the Galactic center, 4 sources in the list of Lindqvist et al. (1992b) are considered to be candidates for associations; OH359.939-0.052 ($V_{av} = 52.2$ km s⁻¹), OH359.946-0.047 ($V_{av} = -27.2$ km s⁻¹), OH359.952-0.036 ($V_{av} = 82.2$ km s⁻¹), and OH359.954-0.041 ($V_{av} = 70.2$ km s⁻¹). In addition, we take into account 5 SiO/H₂O sources listed by Menten et al. (1997), one of which coincides with OH359.946-0.047.

The judgment for association is based on the velocity coincidence. The position (\pm 20") and velocity (\pm 3 km s⁻¹) coincidences are shown as footnotes in table 1 for each SiO components. The radial velocities of the source No. 3 (-25.2 km s⁻¹) at (0", 0") and the source No. 3 ($V_{lsr} = -24.5$ and -28.3 km s⁻¹) at (+40",0"), are quite close. It is possible that they are the same source that is located at the middle of the two beams. However, the source No. 3 at (0",0") is probably associated with SiO(+7.7,+4.2) = IRS 10 EE in table 1 of Menten et al. (1997) and OH359.946-0.047 in Lindqvist et al. (1992b). In this case, the intensity of this source at (+40", 0") should be 7 % of the signal intensity at (0", 0"), or at most 14 % even if a pointing error of 5 " is taken into account, indicating that it is undetectable at (+40", 0"). Therefore, we conclude these two emissions at similar velocities come from different sources.

3.2. Velocity distribution

It is interesting to examine how the average velocity of sources varies with Galactic longitude, because fast rotating gas components of about 200-300 km/s have been found in the Galactic center spiral of an angular size of about 30" (Lo and Claussen 1983; Lacy et al. 1991). We have made a linear regression analysis of V_{lsr} with l. Source positions were assumed to be exactly at beam centers, i.e., -40", 0, and +40" in longitude. The least-square fit to the radial velocities of the detected sources gives $V_{lsr} = -25.5[\pm 32.2] - 0.098[\pm 1.14](\Delta l/")$ km s⁻¹, indicating a quite small velocity gradient.

It can be considered that the high-velocity source No. 7 at (0'', 0'') may influence the results. Because the SiO intensity of this source is quite strong, this high velocity source may be located closer to the Sun than the other sources on the same line of sight. We have also made the linear fit excluding this source and obtained $V_{lsr} = -0.9[\pm 21.6] - 0.010[\pm 0.734](\Delta l/'') \text{ km s}^{-1}$. The result does not differ significantly from the previous one.

The velocity dispersion (deviation from the regression line) is 111.4 km s⁻¹ for the set of all sources and 71.2 km s⁻¹ for the set excluding the high velocity source No. 7. This value is comparable with the velocity dispersion of ~125 km s⁻¹ of infrared stars within 20" from the center (McGinn et al. 1989). However, it seems significantly smaller than 154 (\pm 19) km s⁻¹ for the Helium-rich blue supergiants/Wolf-Rayet stars with distances of 1-12" from the Galactic center (Krabbe et al. 1995).

Lindqvist et al. (1992a) found that the linear regression of the (l-v) diagram for their 134 OH/IR sources is about $V_{lsr} = -7 + 180 (\Delta l/deg) \text{ km s}^{-1}$, or if expressed in the above unit, $V_{lsr} = -7 + 0.05 (\Delta l/'') \text{ km s}^{-1}$. The gradient is as small as the value obtained in this paper. Lindqvist et al. (1992a) mentioned that the smaller number of samples within 10 pc (about 4') from the Galactic center gave the larger velocity gradient. From their figure 9, we can estimate this gradient for the smaller sample as 0.83 km s⁻¹ /". McGinn et al. (1989) also gave the velocity gradient of ~1.0 km s⁻¹ /" from infrared stars within 100" from the center (estimated from figure 4 in their paper). These values are consistent with the present non-detection of rotational motion within the errors. Genzel et al. (1996) found that late-type stars within 5" from the center exhibit a gradient of about 5 km s⁻¹ /", which is somewhat larger than the present result.

If SiO sources near the Galactic center have the same kinematic properties as the stars detected by McGinn et al. (1989) and Lindqvist et al. (1992b), i.e., they belong to the nuclear disk of a size of more than a few arc minutes, the nondetection of the rotational motion in the present paper seems reasonable. The velocity dispersion obtained is also consistent with the previous measurements for the stellar disk.

3.3. Source number density

The source numbers we detected are 4, 7, and 3 at the three positions, (-40'',0''), (0'', 0''), and (+40'', 0''), respectively. The apparent decrease in the source number at the off-center positions is probably due to the larger noise level (less integration time) at these positions. In fact, if we correct the source counts for the noise levels, assuming that we try to detect the (0'', 0'') No. 1–7 sources with higher noise levels, the expected detections above 5 sigma are 3 and 1 with the noise levels at (-40'', 0'') and (+40'', 0''), respectively. Taking into account statistical uncertainties, we conclude that the source surface number density is not peaked at the Galactic center on a scale of about 40''.

It is interesting to compare the SiO source surface density with the density of OH/IR sources. Lindqvist et al. (1992a) gave the OH 1612 MHz source density of 1700 per square degree, which corresponds to about 0.47 sources per square arcminute. Apparently the

number of SiO sources at (0'', 0'') (=7 per 40 " beam) is about 40 times that expected from the OH/IR survey. The number density of SiO maser sources found in the present survey is an order of magnitude larger than the density of OH 1612 MHz maser sources which were detectable with the VLA. Hence we would like to call the cluster of SiO maser lines at the Galactic center a "SiO maser forest".

OH/IR stars with OH 1612 MHz emission are highly obscured red supergiants, while SiO maser stars are very often less massive semiregular or Mira variables which have weak or no OH 1612 MHz emission. For example, W Hya, a semiregular variable at the distance of 95 pc (Reid & Menten 1997), exhibits strong SiO maser emission of about 700 Jy in the J=1-0 v=1 and 2 transitions, but shows no OH 1612 MHz line. If it is placed at the Galactic center (at the distance of 8 kpc), the flux density of the same line would be 0.1 Jy, which is comparable with the flux densities observed in the present survey. Therefore, many of the SiO sources we have detected at the Galactic center might be Mira and semiregular variables. While the statistics of a sample of 172 bulge OH/SiO sources (observed by both OH and SiO lines; Jiang et al. 1995) shows that the number of SiO sources without OH is comparable with the number of OH sources with and without SiO. If the same statistics applies to the Galactic center sources, we should detect more OH 1612 MHz sources. Therefore, some violent mechanism might strip the outer (OH emitting) envelope of AGB stars near the Galactic center, causing the underdensity of OH 1612 MHz sources at the Galactic center.

4. Conclusions

We have detected 14 SiO maser sources at the Galactic center and at the 40" off-center positions; 6 of these sources are associated with already known OH 1612 MHz or SiO

sources, and 8 are new detections. ² We find that the radial velocity gradient with respect to Galactic longitude is small. The SiO source number density is an order of magnitude higher than the density of previously found OH 1612 MHz objects toward the Galactic center. The flat distribution of the source number density around the center and the velocity dispersion indicate that these SiO sources are associated with the Galactic nuclear stellar disk. The small radial velocity gradient of the sources also does not conflict with this interpretation.

We thank David Morris for reading the manuscript. We also thank Drs. N. Ukita, Y. Nakada, O. Kameya, S. Matsumoto, and H. Imai for discussions and encouragements. This research was partly supported by Scientific Research Grant (C) 08640337 from the Ministry of Education, Science, Sports, and Culture, Japan.

²Menten and Reid (1997) recently reported the detections of several more SiO sources [two of them are associated with No. 5 and 7 at (0'', 0'') in Table 1].

REFERENCES

- Blum, R. D., Sellgren, K., & DePoy, D. L. 1996, ApJ, 470, 864
- Eckart, A., & Genzel, R. 1997, MNRAS, 284, 576
- Eckart, A., Genzel, R., Hofman, R., Sams, B. J., & Tacconi-Garman, L. E. 1993, ApJ, 407, L77
- Genzel, R., Thatte, N., Krabbe, A., Kroker, H., & Tacconi-Garman, L. E. 1996, ApJ, 473, 153
- Izumiura, H., Deguchi, S., Hashimoto, O., Nakada, Y., Onaka, T., Ono, T., Ukita, N., & Yamamura, I. 1995, ApJ, 453, 837
- Jewell, P. R., Snyder, L. E., Walmsley, C. M., Wilson, T. L., & Gensheimer, P. D. 1991, A&A 242, 211
- Jiang, B. W., Deguchi, S., Izumiura, Y., Nakada, Y., & Yamamura, I. 1995, PASJ 47, 815
- Krabbe, A., Genzel, R., Eckart, A., Najarro, F., Lutz, D., Cameron, M., Kroker, H., Tacconi-Garman, L. E., Thattle, N., Weitzel, L., Draper, S., Geball, T., Stergnberg, A., & Kudritzki, R. 1995, ApJ, 447, L95
- Lacy, J.H., Achtermann, J.M., & Serabyn, E. 1991, ApJ, 380, L71
- Levine, D., Figer, D. F., Morris, M., & Mclean, I. S., 1995, ApJ, 447, L101
- Lindqvist, M., Winnberg, A., Johansson, L.E.B., & Ukita, N. 1991, A&A, 250, 431
- Lindqvist, M., Habing, H. J., & Winnberg, A. 1992a, A&A, 259, 118
- Lindqvist, M., Winnberg, A., Habing, H. J., & Matthews, H. E.1992b, A&AS, 92, 43
- Lo, K. Y., & Claussen, M. J. 1983, Nature 306, 647

- McGinn, M. T., Sellgren, K., Becklin, E. E., & Hall, D. N. B. 1989, ApJ, 338, 824
- Menten, K., & Reid, M. 1997, a poster paper presented in IAU Sympo. 184. held Aug 18-21, 1997, in Kyoto, Japan.
- Menten, K., Reid, M., Eckart, A., & Genzel, R. 1997, ApJ, 475, L111
- Morris, M., & Serabyn, E. 1996, ARAA, 34, 645
- Reid, M. J., & Menten, K. M. 1997, ApJ, 476, 327
- Rogers, A. E. E., et al. 1994, ApJ, 434, L59
- Shiki, S., Ohishi, M., & Deguchi, S. 1997, ApJ, 478, 206
- Sjouwerman, L. O., & van Langevelde, H. J. 1996, ApJ, 461, L41
- Sjouwerman, L. O., van Langevelde, H. J., Winnberg, A., & Habing, H. J. 1997, A&A, in press
- Sofue, Y., Inoue, M., Handa, T., Tsuboi, M., Hirabayashi, H., Morimoto, M., & Akabane, K. 1986, PASJ, 38, 475
- Winnberg, A. 1996, "The Galactic Center", ASP conf. ser. Vol.102, p294, ed. by Gredel et al. (Astron. Soc. Pacific; San Francisco)

This manuscript was prepared with the AAS LAT_EX macros v4.0.

Fig. 1.— Spectra of the SiO J=1-0 v=1 and v=2 transitions toward the Galactic center; top at $(\Delta l, \Delta b) = (+40'', 0'')$, middle at (0'', 0''), and bottom at (-40'', 0''). Detections except strongest ones are marked by ticks.

		v=2	I=1-0		v=1	J=1-0	
SiO l,b (°)	No	$\overline{V_{lsr}}$	I _{peak}	F	V=1 V _{lsr}	I _{peak}	F
$(\Delta l, \Delta b)$		$({\rm kms^{-1}})$	(Jy)	$\left(\mathrm{Jykms^{-1}}~\right)$	$\left(\mathrm{kms^{-1}}\right)$	(Jy)	$\left(Jykms^{-1} \right)$
SiO359.933-0.046	1	131.2	0.167	0.647	131.2	0.143	0.380
(-40", 0")	2				63.7	0.152	0.227
	3^{a}	15.9	0.179	0.497	15.9	0.289	0.942
	4	-105.1	0.273	0.921	-104.2	0.127	0.511
SiO359.944-0.046	1	53.1	0.169	0.712	54.4	0.177	0.738
(0", 0")	2^{b}	-14.6	0.121	0.435	-13.8	0.153	0.440
	$3^{\rm c}$	-25.2	0.164	0.797	-25.2	0.099	0.560
	4	-51.2	0.072	0.143			—
	5	-69.6	0.087	0.575			—
	$6^{\rm d}$	-114.9	0.094	0.497			—
	7	-341.9	0.333	0.195	-341.9	0.216	1.395
SiO359.955-0.046	$1^{\rm e}$	70.0	0.234	0.494			_
(+40", 0")	2^{f}	50.6	0.403	2.377	52.6	0.450	2.173
	3	-24.7	0.158	0.474			

Table 1. List of detected SiO peaks (sources)

^a=OH359.931-0.050, V_{av} =17 km s⁻¹ (Sjouwerman et al. 1997)

^b=IRS15NE =SiO(+1.2,+11.3), V_{av} =-15 km s⁻¹ (Menten et al. 1997)

^=OH359.946-0.047=IRS10 EE, $\mathrm{V}_{av}{=}{-}27~\,\mathrm{km\,s^{-1}}$ (Lindqvist et al. 1992b)

 $^{\rm d}{=}{\rm IRS7{=}SiO(0.0,{+}5.6)},\,{\rm V}_{av}{=}{-}120~\,{\rm km\,s^{-1}}$ (Menten et al. 1997)

 $^{\rm e}{=}{\rm OH359.954\text{-}0.041},\,{\rm V}_{av}{=}70~\,{\rm km\,s^{-1}}$ (Lindqvist et al. 1992b)

f=OH359.956-0.050, $\mathrm{V}_{av}{=}50~\mathrm{km\,s^{-1}}$ (Sjouwerman & van Langevelde 1996; Levine et al. 1995)

