

Instanton Cosmology

Roland E. Allen
Center for Theoretical Physics, Texas A&M University
College Station, Texas 77843

Abstract

A new cosmological model implies that dR/dt is constant for both $t > t_c$ and $t \rightarrow 0$, and that $dR^2/dt \propto n_s^{-1}$ for $t_c > t > 0$. Here $R(t)$ is the cosmic scale factor, t_c is a crossover time defined in the text, and n_s is approximately constant near t_c . The behavior at intermediate times is consistent with big-bang nucleosynthesis. The behavior at short and long times offers a solution to the smoothness, monopole, and flatness problems. In addition, the long-time behavior yields a comfortably large age for the universe and predicts that the deceleration parameter q_0 is exactly zero.

e-mail: allen@tamu.edu
fax : (409) 845-2590
tel. : (409) 845-4341

Recently a new fundamental theory was proposed [1]. Among its consequences is a specific cosmological model, which the present paper will explore in more detail. We will find that the cosmic scale factor has the following behavior:

$$R(t) = \eta t \quad , \quad t \rightarrow 0 \quad (1)$$

$$R(t) \propto \left[\int dt/n_s(t) \right]^{1/2} \quad , \quad t < t_c \quad (2)$$

$$dR(t)/dt = 2\sqrt{3} \quad , \quad t > t_c \quad (3)$$

where $\eta \gg 1$ and $n_s(t)$ is a decreasing function of t , with

$$n_s(t) \approx \bar{n}_s = \text{constant} \quad , \quad \bar{t} < t < t_c \quad (4)$$

The first prediction implies that the particle horizon in the early universe is nearly infinite. The second is consistent with the conclusion that [2]

$$0.55 < \alpha < 0.56 \quad (5)$$

if $R(t) \propto t^\alpha$ is assumed to hold during the period of big-bang nucleosynthesis [3]. The third implies that

$$q = 0 \quad (6)$$

where q is the deceleration parameter in the current epoch.

The theory of Ref. 1 begins with an Ising-like action S , which becomes a Ginzburg-Landau-like action in the continuum approximation. After a Wick rotation $x^0 \rightarrow ix^0$, one obtains the Lorentzian action

$$S_L = iS \quad (7)$$

$$= - \int d^D x \left[\frac{1}{2m} \partial^M \Psi^\dagger \partial_M \Psi - \mu \Psi^\dagger \Psi + \frac{1}{2} b (\Psi^\dagger \Psi)^2 \right]. \quad (8)$$

Here Ψ is a superfield, with both bosonic and fermionic parts. In the present context, however, it is not necessary to consider fermions or bosonic excitations. It is also convenient to initially ignore the $d = D - 4$ internal dimensions, so that we are left with the action for a GUT Higgs field in four-dimensional spacetime:

$$S_L = - \int d^4 x \Psi_s^\dagger \left(-\frac{1}{2m} \partial^\mu \partial_\mu + \frac{1}{2} b n_s - \mu \right) \Psi_s \quad (9)$$

where $n_s = \Psi_s^\dagger \Psi_s$. One can define a ‘‘superfluid velocity’’

$$v^\mu = -im^{-1}U^{-1}\partial^\mu U \quad (10)$$

where

$$\Psi_s = n_s^{1/2}U\eta_0 \quad (11)$$

and η_0 is a constant 2-component vector. The field v_α^μ , defined by

$$v^\mu = v_\alpha^\mu \sigma^\alpha, \quad (12)$$

is interpreted as the vierbein which determines the geometry of spacetime:

$$e_\alpha^\mu = v_\alpha^\mu. \quad (13)$$

(Here σ^0 and $\sigma^1, \sigma^2, \sigma^3$ are respectively the identity and Pauli matrices.) The curvature of spacetime is attributed to Planck-scale instantons. This picture leads to both Einstein gravity and an SO(10) grand unified theory of fermions and the other interactions [1].

The condensate Ψ_s can support specific kinds of topological defects, two of which are relevant in the present context: (a) SU(2) instantons, which are the four-dimensional generalization of vortices in an ordinary superfluid. With x^0 chosen to be the radial coordinate, v_α^μ has the form [4]

$$v_\alpha^k = \delta_\alpha^k a/mx^0 \quad (14)$$

$$v_\alpha^0 = 0 \quad (15)$$

where a is a constant and $k = 1, 2, 3$. From a four-dimensional point of view, it is as if there is a circulation of condensate around the origin. (b) U(1) monopole-like defects, which act as sources of the U(1) current

$$j_0^\mu = n_s v_0^\mu. \quad (16)$$

These defects can relieve the constraint

$$\partial_\mu j_0^\mu = 0 \quad (17)$$

in the same way that vortices in a superfluid can relieve the constraint $\vec{\nabla} \times \vec{v}_s = 0$. Suppose that such a defect has spherical symmetry, with j_0^μ pointing out from the origin. The equation of continuity (17) still holds for $r \neq 0$, where r is the radial coordinate, so

$$\partial(r^3 j_0)/\partial r = 0 \quad (18)$$

or

$$v_0 = a_1^3 / (r^3 n_s) \quad (19)$$

with a_1 a constant. From a four-dimensional point of view, it is as if there is an emanation of condensate from the origin.

These are the two natural point defects for a 2-component field Ψ_s . In the present cosmological model they are superposed, and the singularity at the origin is interpreted as the big bang [1]. Combining (13), (14), and (19) (with $r \rightarrow x^0$) we obtain

$$e_\alpha^k = \delta_\alpha^k a / m x^0 \quad (20)$$

$$e_\alpha^0 = \delta_\alpha^0 a_1^3 / (x^0)^3 n_s . \quad (21)$$

The usual relation

$$e_\mu^\alpha e_\beta^\mu = \delta_\beta^\alpha \quad (22)$$

gives the covariant vierbein

$$e_k^\alpha = \delta_k^\alpha m x^0 / a \quad (23)$$

$$e_0^\alpha = \delta_0^\alpha (x^0)^3 n_s / a_1^3 . \quad (24)$$

In addition, larger values of x^0 correspond to larger 3-spheres [1], so $dx^k = x^0 d\vec{x}^k$, where $d\vec{x}^k$ represents a distance within the unit 3-sphere. There are thus two contributions to the expansion of the universe in the present model, with both the metric tensor g_{kl} and the separation of comoving points increasing as functions of x^0 .

The vierbein of (23) and (24) leads to a metric with the Robertson-Walker form

$$ds^2 = \eta_{\alpha\beta} e_\mu^\alpha e_\nu^\beta dx^\mu dx^\nu \quad (25)$$

$$= - \left[\frac{(x^0)^3 n_s}{a_1^3} \right]^2 (dx^0)^2 + \left[\frac{m (x^0)^2}{a} \right]^2 d\vec{x}^k d\vec{x}^k \quad (26)$$

$$= -dt^2 + R(t)^2 \left(\frac{dr^2}{1-r^2} + r^2 d\Omega \right) \quad (27)$$

where

$$dt = (n_s / 4a_1^3) d(x^0)^4 \quad (28)$$

$$R(t) = \frac{2m}{a} \left(a_1^3 \int \frac{dt}{n_s} \right)^{1/2} . \quad (29)$$

Also, $\eta_{\alpha\beta} = \text{diag}(-1, 1, 1, 1)$ is the Minkowski metric tensor, and $d\Omega = d\theta^2 + \sin^2\theta d\phi^2$, with (r, θ, ϕ) representing the usual coordinates of the unit 3-sphere.

Notice that this metric is determined by the topology of the cosmological instanton, rather than the Einstein field equations. As usual, these equations follow from the variational principle $\delta S_L / \delta g^{\mu\nu} = 0$ if there are no constraints, where S_L is the full Lorentzian action [1]. In the present model, however, they are subordinate to the constraint imposed by topology on a cosmological scale. I.e., the Einstein field equations are only a very good approximation for *local* deformations of the geometry of spacetime.

The time dependence of the condensate density n_s is determined by the equation of motion for Ψ_s , which leads to the generalized Bernoulli equation [1]

$$-\frac{1}{2m}n_s^{-1/2}\partial^\mu\partial_\mu n_s^{1/2} + \frac{1}{2}mv_\alpha^\mu v_\mu^\alpha + bn_s = \mu \quad (30)$$

or [4]

$$\frac{1}{2m}n_s^{-1/2}(x^0)^{-3}\frac{d}{dx^0}\left[(x^0)^3\frac{d}{dx^0}n_s^{1/2}\right] + \frac{1}{2}mv_\alpha^k v_\alpha^k - \frac{1}{2}mv_\alpha^0 v_\alpha^0 + bn_s = \mu. \quad (31)$$

Substitution of (20) and (21) gives

$$\xi^2 n_s^{-1/2} (x^0)^{-3} \frac{d}{dx^0} \left[(x^0)^3 \frac{d}{dx^0} n_s^{1/2} \right] + \frac{3\xi^2 a^2}{(x^0)^2} - \frac{\xi^2 m^2 a_1^6}{(x^0)^6 n_s^2} + \frac{n_s}{\bar{n}_s} = 1 \quad (32)$$

where $\xi = (2m\mu)^{-1/2}$ is the coherence length and $\bar{n}_s = \mu/b$ is the condensate density in a uniform system. With the scalings

$$\rho = x^0/\xi \quad , \quad f = (n_s/\bar{n}_s)^{1/2} \quad (33)$$

we obtain an equation for f which is analogous to the standard equation for the condensate density near a superfluid vortex:

$$\frac{1}{\rho^3} \frac{d}{d\rho} \left(\rho^3 \frac{df}{d\rho} \right) + \frac{3a^2}{\rho^2} f - \frac{\bar{a}^6}{\rho^6 f^4} f + f^3 - f = 0 \quad (34)$$

where

$$\bar{a}^3 = a_1^3 m / \bar{n}_s \xi^2. \quad (35)$$

The asymptotic solutions are

$$f = 1 - 3a^2/2\rho^2 \quad , \quad \rho \rightarrow \infty \quad (36)$$

$$f = c/\rho \quad , \quad \rho \rightarrow 0 \quad (37)$$

with

$$c^6 + c^4 (3a^2 - 1) = \bar{a}^6 . \quad (38)$$

The forms (36) and (37) imply that there is a ρ_c such that

$$f(\rho_c) = 1 . \quad (39)$$

Suppose that

$$\rho_c^6 \gg \bar{a}^6 \quad (40)$$

and

$$\bar{a}^2 \gg 3a^2 . \quad (41)$$

One then expects that the first term of (34) is very small at ρ_c , and this expectation is confirmed by numerical calculations [5]. If this term is neglected, (34) and (39) give

$$\rho_c \approx \bar{a} \left(\bar{a}^2 / 3a^2 \right)^{1/4} . \quad (42)$$

One expects f to be a slowly decreasing function of ρ in the region $\bar{a} < \rho < \rho_c$, and this is again confirmed by the numerical calculations [5]. Then the condensate density n_s is a slowly decreasing function of the time t in a region $\bar{t} < t < t_c$, where \bar{t} and t_c are the times corresponding to $\rho = \bar{a}$ and ρ_c :

$$\bar{t} \sim \left(\bar{n}_s / 4a_1^3 \right) \xi^4 \bar{a}^4 \sim m \xi^2 \bar{a} \quad (43)$$

$$t_c \sim \left(\bar{n}_s / 4a_1^3 \right) \xi^4 \rho_c^4 \sim m \xi^2 (\bar{a}/a)^2 \bar{a} . \quad (44)$$

(See (28), (33), and (35). Also notice that (43) is consistent with (54).) Since $m \xi^2 \sim t_P$, where $t_P \sim 10^{-42}$ sec is a Planck time [1], we can also write

$$\bar{t} \sim \bar{a} t_P \quad (45)$$

$$t_c \sim \bar{\eta}^2 \bar{t} \quad (46)$$

where

$$\bar{\eta} = \bar{a}/a \gg 1 . \quad (47)$$

For example, $\bar{t} \sim 100$ sec and $\bar{\eta} \sim 100$ corresponds to $t_c \sim 10^6$ sec.

Starting with the action $S_L = \int dx^0 L$ of (9), one can follow the usual rules for obtaining the Hamiltonian $H = \int d^3x \mathcal{H}$. The result is

$$\mathcal{H} = \bar{n}_s \mu f \left[-\frac{1}{\rho^3} \frac{d}{d\rho} \left(\rho^3 \frac{df}{d\rho} \right) + \frac{3a^2}{\rho^2} f + \frac{\bar{a}^6}{\rho^6 f^4} f + \frac{1}{2} f^3 - f \right] . \quad (48)$$

Since the volume of a 3-sphere is $2\pi^2 (x^0)^3 = 2\pi^2 \xi^3 \rho^3$, (37) and (38) imply that

$$H = \pi^2 \bar{n}_s \mu \xi^3 (3c^4 + 12c^2 a^2) / \rho \approx 3\pi^2 \bar{n}_s \mu \bar{a}^4 \xi^3 / \rho \quad (49)$$

as $\rho \rightarrow 0$. (According to (2.5)-(2.7) of Ref. 1, there is a cutoff at $x^0 \sim \xi$, or $\rho \sim 1$, so that the initial value of H is comparable to $\bar{a}^4 m_P$.) In this same regime, (28), (29), (33), and (35) give

$$t = (c^2 m \xi^2 / 2\bar{a}^3) \rho^2 \quad (50)$$

$$n_s = (c^4 m \xi^2 / 2\bar{a}^3) \bar{n}_s / t \quad (51)$$

$$R(t) = \eta t \quad (52)$$

where

$$\eta = 2\bar{a}^3 / c^2 a \approx 2\bar{\eta} \gg 1 \quad (53)$$

with each equation holding in the limit $\rho \rightarrow 0$, or $t \rightarrow 0$. It follows that

$$t \approx \frac{m \xi^2}{2\bar{a}} \rho^2 \sim \frac{(\rho t_P)^2}{\bar{t}} \quad (54)$$

$$n_s \approx \frac{\bar{a} m \xi^2 \bar{n}_s}{2} \frac{\bar{n}_s}{t} \sim \frac{\bar{t}}{t} \bar{n}_s \quad (55)$$

$$R(t) \approx \eta t \quad (56)$$

for $t < \bar{t}$.

The particle horizon is determined as usual by

$$d_H(t) = R(t) \int_\epsilon^t \frac{dt'}{R(t')} \quad , \quad \epsilon \rightarrow 0. \quad (57)$$

In Friedmann models with $R(t) \propto t^\alpha$, and $\alpha < 1$, the horizon is finite:

$$d_H(t) = t / (1 - \alpha) \sim t. \quad (58)$$

In the present model, however, (52) implies that $\alpha \approx 1$ in the early universe, making the horizon nearly infinite: for $t < \bar{t}$,

$$d_H(t) \approx t \log(t/\epsilon) \rightarrow \infty. \quad (59)$$

Now let us turn to the behavior of the condensate at large times $t > t_c$. This is the regime already considered in Ref. 1, where the following

assumptions are made: (1) There are monopole-like defects which relieve the constraint (17) by acting as sources of the current j_0^μ , in the same way that vortices permit a superfluid to rotate by relieving the constraint $\vec{\nabla} \times \vec{v}_s = 0$. (See the discussion following (7.24) of Ref. 1.) These are Planck-scale versions of the cosmological U(1) monopole-like defect considered above. (2) At large times ($t > t_c$), the condensate density is constant in external spacetime. This requires that

$$\frac{1}{2}mv_\alpha^k v_\alpha^k - \frac{1}{2}mv_\alpha^0 v_\alpha^0 = 0 \quad , \quad t > t_c . \quad (60)$$

(See the discussion above (5.14) and below (8.40) in Ref. 1.) It follows that

$$v_\alpha^0 v_\alpha^0 = v_\alpha^k v_\alpha^k = 3 \left(a/mx^0 \right)^2 \quad (61)$$

or

$$e_0^\alpha = \delta_0^\alpha m x^0 / \sqrt{3} a . \quad (62)$$

Let us now consider the origin of the condition (60) in more detail, by including the internal space x_B of Ref. 1, with d coordinates x^m , as well as our external space with coordinates x^μ . The condensate wavefunction can always be written in the form

$$\Psi_s(x_A, x_B) = \Psi_A(x_A) \Psi_B(x_A, x_B) \quad (63)$$

with

$$n_s(x_A, x_B) = n_A(x_A) n_B(x_A, x_B) \quad (64)$$

$$n_A = \Psi_A^\dagger \Psi_A \quad , \quad n_B = \Psi_B^\dagger \Psi_B . \quad (65)$$

For $t > t_c$ it is assumed that the condensate has settled into a local energy minimum, with an internal instanton satisfying

$$\left[-\frac{1}{2m} \partial^m \partial_m + V(x_B) - \mu \right] \Psi_B(x_B) = 0 \quad (66)$$

$$V(x_B) = b n_A n_B(x_B) . \quad (67)$$

(See (5.14) of Ref. 1, and the comment below (8.40).) At all times, on the other hand, one has the equation of motion

$$\left(-\frac{1}{2m} \partial^\mu \partial_\mu - \frac{1}{2m} \partial^m \partial_m + b n_A n_B - \mu \right) \Psi_A \Psi_B = 0 . \quad (68)$$

(See (3.7) of Ref. 1.) In external spacetime this becomes

$$\left(-\frac{1}{2m}\partial^\mu\partial_\mu + \tilde{b}n_A - \tilde{\mu}\right)\Psi_A = 0 \quad (69)$$

where

$$\tilde{\mu} = \mu - T_B \quad (70)$$

$$T_B = N_B^{-1} \int d^d x \Psi_B^\dagger \left(-\frac{1}{2m}\partial^m\partial_m\right) \Psi_B \quad (71)$$

$$\tilde{b} = bN_B^{-1} \int d^d x n_B^2 \quad (72)$$

$$N_B = \int d^d x n_B. \quad (73)$$

It is reasonable to make the approximation that $\tilde{\mu}$ and \tilde{b} are slowly varying functions of t . We then regain the action of (9) and the previous results for $t < t_c$, with the replacements $\mu \rightarrow \tilde{\mu}$, $b \rightarrow \tilde{b}$, and $n_s \rightarrow n_A$.

In particular, the scaled equation (34) still holds as $t \rightarrow t_c$ from below. Since t_c is defined to be the point where $f = 1$, and the condensate is assumed to remain fully formed beyond this point, (33) implies that

$$n_A = \bar{n}_A = \tilde{\mu}/\tilde{b} \quad \text{for } t \geq t_c \quad (74)$$

with $\tilde{\mu}$ and \tilde{b} constant. Then (69) gives

$$\Psi_A^\dagger \left(-\frac{1}{2m}\partial^\mu\partial_\mu\right) \Psi_A = 0. \quad (75)$$

The arguments in Section 3 of Ref. 1 show that this condition is equivalent to

$$\frac{1}{2}mv_\alpha^\mu v_\mu^\alpha = 0 \quad (76)$$

which is the same as (60).

With e_0^α given by (62), and e_k^α still given by (23), (25) becomes

$$ds^2 = -\left(\frac{mx^0}{\sqrt{3a}}\right)^2 (dx^0)^2 + \left[\frac{m(x^0)^2}{a}\right]^2 d\vec{x}^k d\vec{x}^k \quad (77)$$

$$= -dt^2 + R(t)^2 \left(\frac{dr^2}{1-r^2} + r^2 d\Omega\right) \quad (78)$$

where

$$dt = (m/2\sqrt{3}a) d(x^0)^2 \quad (79)$$

$$R(t) = m(x^0)^2/a. \quad (80)$$

It follows that

$$\dot{R} = dR/dt = 2\sqrt{3}. \quad (81)$$

With the conventional definitions

$$H = \dot{R}/R \quad (82)$$

or

$$q = -\ddot{R}/RH^2 \quad (83)$$

we have

$$q = 0 \quad , \quad t > t_c. \quad (84)$$

The present model can be viewed from either of two perspectives:

In the Lorentzian spacetime of human observers, the universe is born containing a highly confined and coherent bosonic field Ψ_s , which is analogous to the coherent photon field in a laser cavity. The density n_s of this field is initially very large (at $x^0 \sim t_P$), but decreases to its minimum-energy value \bar{n}_s during a period $t < t_c$. At the same time, it releases energy to other fields, producing a hot big bang (although the coupling to other fields is omitted in the present treatment). For $t > t_c$, Ψ_s remains fully condensed, and can be properly interpreted as the condensate associated with a GUT Higgs field.

In Euclidean spacetime, which is regarded as more fundamental in Ref. 1, the big bang results from a topological defect centered on the point $t = 0$. The nature of this cosmological instanton is specified above (20). When one transforms the solution (37) to Euclidean spacetime, the action S is found to be positive, as is the Hamiltonian H in Lorentzian spacetime. S and H are also positive for the Planck-scale monopole-like defects postulated above (60).

Let us now consider the physical implications of the above results:

The fact that the particle horizon is nearly infinite at early times means that the observable universe was once causally connected, so its isotropy and homogeneity are not unnatural, as they are in standard cosmology without inflation [6-11].

In addition, since the Kibble mechanism requires a horizon [12], there is no longer a compelling mechanism for the production of unacceptably large quantities of cosmic relics like monopoles.

The curvature associated with (78) is proportional to $R(t)^{-2}$ [13]. The present model thus requires a universe that is asymptotically flat, because the topology of the cosmological instanton requires that $R(t) \propto t$ for $t \rightarrow \infty$. This is true even if

$$\Omega_m = \rho_m / \rho_m^c \neq 1 \tag{85}$$

where ρ_m^c is the critical mass density corresponding to a flat universe in conventional cosmology. Observations of the cosmic microwave background and of large scale structure indicate that spacetime is very nearly flat but that $\Omega_m \approx 0.3$ [14].

If the period $t < t_c$ is neglected, (81) implies that $R(t) \propto t$, so $H^{-1} = t$ rather than $\frac{3}{2}t$. The age of the universe is then given by $t_0 = H_0^{-1}$ rather than $\frac{2}{3}H_0^{-1}$. With the conventional definition $H_0 = 100h \text{ km s}^{-1}\text{Mpc}^{-1}$, a value of $h = 0.65$ [15] implies that $t_0 = 15 \text{ Gyr}$ rather than 10 Gyr . This larger value is more consistent with recent estimates of $10 - 14 \text{ Gyr}$ for the ages of the oldest globular clusters [16].

Perhaps the most directly testable prediction of the present theory is $q = 0$, which means that the universe is neither decelerating nor accelerating [17]. This feature clearly demarcates the present theory from standard cosmology with a flat or closed universe [18,19], in which $q_0 \geq 1/2$, and from Λ CDM models [20], in which $q_0 \sim -1/2$. Quantitative measurements of q_0 are now becoming attainable [21,22], so it should be possible to subject this prediction to convincing tests in the near future.

Acknowledgement

This work was supported by the Robert A. Welch Foundation. I have greatly benefitted from conversations with R.L. Arnowitt on new observational developments, and would also like to thank G. Steigman for informing me of the results of Ref. 2 before it was submitted for publication.

References

- [1] R.E. Allen, *Int. J. Mod. Phys. A* **12** (1997) 2385, hep-th/9612041.
- [2] M. Kaplinghat, G. Steigman, I. Tkachev, and T.P. Walker, astro-ph/9805114.
- [3] D.N. Schramm, *The Big Bang and Other Explosions in Nuclear and Particle Astrophysics* (World Scientific, Singapore, 1996).
- [4] The v_k^α are “ordinary” components, after the initial coordinates of Ref. 1 are transformed to four-dimensional spherical coordinates. See S. Weinberg, *Gravitation and Cosmology* (Wiley, New York, 1972), p. 109.
- [5] R.E. Allen, J. Dumoit, and A. Mondragon, to be published in the proceedings of PASCOS-98 (Sixth International Symposium on Particles, Strings and Cosmology; Northeastern University, Boston; March 22-27, 1998).
- [6] P.J.E. Peebles, *Principles of Physical Cosmology* (Princeton University Press, Princeton, 1993), and references therein.
- [7] E.W. Kolb and M.S. Turner, *The Early Universe* (Addison-Wesley, Redwood City, California, 1990), and references therein.
- [8] *The Early Universe: Reprints*, edited by E.W. Kolb and M.S. Turner (Addison-Wesley, Redwood City, California, 1988).
- [9] P.D.B. Collins, A.D. Martin, and E.J. Squires, *Particle Physics and Cosmology* (Wiley, New York, 1989).
- [10] A. Linde, *Particle Physics and Inflationary Cosmology* (Harwood, Chur, Switzerland, 1990).
- [11] *Unsolved Problems in Astrophysics*, edited by J.N. Bahcall and J.P. Ostriker (Princeton University Press, Princeton, 1997).
- [12] A. Vilenkin and E.P.S. Shellard, *Cosmic Strings and Other Topological Defects* (Cambridge University Press, Cambridge, 1994).
- [13] C.W. Misner, K.S. Thorne, and J.A. Wheeler, *Gravitation* (W.H. Freeman, San Francisco, 1973).
- [14] M. Webster, M.P. Hobson, A.N. Lasenby, O. Lahav, and G. Rocha, astro-ph/9802109, and references therein.

- [15] W. L. Freedman, J. R. Mould, R. C. Kennicutt, and B. F. Madore, astro-ph/9801080, and references therein.
- [16] See, for example, B. Chaboyer, P. Demarque, P. J. Kernan, and L. M. Krauss, astro-ph/9706128.
- [17] Other models predict $q_0 = 0$ for very different physical reasons. The first extensive study of such a model was by E.W. Kolb, *Astrophys. J.* **344** (1989) 543. See also M. Kamionkowski and N. Toumbas, *Phys. Rev. Lett.* **77** (1996) 587, and references therein.
- [18] Ya.B. Zel'dovich and I.D. Novikov, *Relativistic Astrophysics* (University of Chicago Press, Chicago, 1983).
- [19] A.R. Sandage, R.G. Kron, and M.S. Longair, *The Deep Universe* (Springer, Berlin, 1995).
- [20] M.S. Turner, *Nucl. Phys.* **A621** (1997) 522c, astro-ph/9704024, and references therein.
- [21] S. Perlmutter, G. Aldering, M. D. Valle, S. Deustua, R.S. Ellis, S. Fabbro, A. Fruchter, G. Goldhaber, A. Goobar, D.E. Groom, I. M. Hook, A.G. Kim, M.Y. Kim, R.A. Knop, C. Lidman, R. G. McMahon, P. Nugent, R. Pain, N. Panagia, C.R. Pennypacker, P. Ruiz-Lapuente, B. Schaefer, and N. Walton, *Nature* **391** (1998) 51, astro-ph/9712212.
- [22] P. M. Garnavich, R. P. Kirshner, P. Challis, J. Tonry, R. L. Gilliland, R. C. Smith, A. Clocchiatti, A. Diercks, A. V. Filippenko, M. Hamuy, C. J. Hogan, B. Leibundgut, M.M. Phillips, D. Reiss, A. G. Riess, B. P. Schmidt, J. Spyromilio, C. Stubbs, N. B. Suntzeff, and L. Wells, *Astrophys. J.* **493** (1998) L53, astro-ph/9710123.