Will a Black Hole Soon Emerge from SN 1997D?

Luca Zampieri¹, Stuart L. Shapiro^{1,2} and Monica Colpi³

ABSTRACT

Observations combined with theoretical modeling of the light curve of the recently discovered supernova 1997D in NGC 1536 suggest that it might host a black hole formed in the aftermath of the explosion. We consider some observable consequences of a black hole in SN 1997D and estimate the late-time accretion luminosity of the material which falls back onto the hole. We find that this luminosity, decaying with a characteristic power-law dependence on time, may emerge above the emission of the envelope in just a few years. Its detection would thus provide unmistakable evidence for the presence of a black hole.

Subject headings: Accretion — black holes — hydrodynamics — radiative transfer — relativity — supernovae: individual (SN 1987A, SN 1997D)

1. Introduction

The nature of the compact remnant formed in the aftermath of a supernova explosion depends on the mass of the iron core of the progenitor star, on the amount of material that falls back onto the core and on the maximum allowed neutron star mass ($M_{\rm max}$). The core is usually in the range 1.4–2.0 M_{\odot} , while the actual value of $M_{\rm max}$ is not conclusively established yet. According to recent results on the properties of dense nucleon matter, the maximum (nonrotating) neutron star mass falls in the interval 1.8–2.2 M_{\odot} (Akmal, Pandharipande & Ravenhall 1998), although smaller values are not yet completely ruled out. Consequently, the amount of matter fallback plays a crucial role in determining the fate of the collapsed remnant. An investigation of the fallback process for stars with different masses and different explosion energies has been performed by Woosley & Weaver (1995). They find that stars with main sequence mass larger than $\sim 25 M_{\odot}$ have rather massive cores ($\simeq 2M_{\odot}$) and undergo accretion of a significant amount of matter, so that the remnant mass is larger than $3M_{\odot}$, probably leading to the formation of a black hole (BH), after the launch of a successful shock. Fallback of a smaller amount of matter, $\sim 0.1 - 0.3 M_{\odot}$, may occur in explosions involving a progenitor of $19-25 M_{\odot}$, although the actual value depends sensitively on the energy of the explosion. Even in this case, a low mass (~ $1.5M_{\odot}$) BH may form if the equation for nuclear matter is not too stiff (Brown & Bethe 1994). The fact that a bright $(L \sim 10^{38} \text{ erg s}^{-1})$ accreting neutron star was not detected in SN 1987A in the early years following the explosion led to the hypothesis

¹Department of Physics, Loomis Laboratory of Physics, University of Illinois at Urbana–Champaign, 1110 West Green Street, Urbana, IL 61801–3080

 $^{^2 {\}rm Department}$ of Astronomy and National Center for Supercomputing Applications, University of Illinois at Urbana-Champaign, Urbana, IL 61801

³Dipartimento di Fisica, Università degli Studi di Milano, Via Celoria 16, I–20133 Milano, Italy

that a light underluminous BH may have formed (Chevalier 1989; Brown & Bethe 1994; Woosley & Timmes 1996). Woosely & Timmes (1996) first suggested that a signature of BH formation in the light curve would be a bright plateau that falls steeply to very low or zero luminosity because of the low radioactive emission from heavy nuclei present in the ejecta. Recently, Zampieri, Colpi, Shapiro & Wasserman (1998a; hereafter ZCSW) pointed out that a BH may leave a distinguishable imprint in the light curve, which could be revealed when the main sources of radioactive decay fade away (see also Zampieri, Colpi, Shapiro & Wasserman 1998b). Using a spherically symmetric, general–relativistic, radiation–hydrodynamic Lagrangian code, ZCSW showed that fallback may lead to an accretion luminosity decaying secularly with time as $L \propto t^{-25/18}$. For SN 1987A, the accretion luminosity turns out to be too dim to be detectable, at present, above the radioactive decay emission of ⁴⁴Ti.

Recently, a new supernova, SN 1997D, was discovered in NGC 1536 (De Mello & Benetti 1997). The very low absolute luminosity and the low inferred heavy element abundance in the ejecta make this supernova a promising candidate for the detection of a compact remnant. Following the suggestion by Turatto *et al.* (1998), we consider the consequences of a BH forming in SN 1997D. We give an estimate of the accretion luminosity and the time of its emergence above the radioactive decay emission of the envelope.

2. The peculiar Type II supernova 1997D

SN 1997D was serendipitously discovered on January 14, 1997 by De Mello & Benetti (1997) during an observation of the parent galaxy NGC 1536. Although the discovery of this supernova was made after maximum phase, a number of closely spaced observations of NGC 1536 set an upper limit to the optical luminosity ~ 2 magnitudes lower than that of a typical Type II supernova. This result makes SN 1997D possibly the least luminous Type II supernova ever observed. A comparison of theoretical models with observations seems to indicate that the epoch of the explosion probably occurred 50 days before the supernova was discovered (Turatto *et al.* 1998). During all the evolutionary stages, the bolometric luminosity of SN 1997D is orders of magnitude lower than that of the "prototype" SN 1987A (see Figures 1 and 2). At around 70–80 days after the explosion, the emission starts to be powered by γ -ray energy deposited into the ejecta and, as shown also by SN 1987A, the decline of the tail of the light curve follows closely the decay rate of ⁵⁶Co, the decay product of ⁵⁶Ni. If all the γ rays are thermalized, the light curve after few weeks is expected to follow the relation

$$L = L_0 \left(\frac{M_{\rm Co}}{M_\odot}\right) e^{-t/\tau_0} \,, \tag{1}$$

where $L_0 = 1.4 \times 10^{43}$ erg s⁻¹, $\tau_0 = 111$ days is the decay time of ⁵⁶Co and $M_{\rm Co}$ is the total mass of ⁵⁶Co present in the ejecta. Assuming for SN 1997D the same γ -ray deposition as in SN 1987A and using the ratio of the luminosities in the tails of the light curves, Turatto *et al.* (1998) estimate the mass of ⁵⁶Ni ejected by SN 1997D to be about 40 times less than that in SN 1987A, i.e. 0.001–0.004 M_{\odot} . Another peculiar property of SN 1997D is the low expansion velocity of the ejecta deduced from measurements of the shift of the absorption lines in the spectrum. It turns out to be ~ 1200 km s⁻¹.

Using a spherical, Newtonian, Lagrangian hydrodynamical code to simulate the explosion, Turatto *et al.* (1998) were able to produce a satisfactory fit to the light curve of SN 1997D. To fit the data, a progenitor model with a radius $r = 2 \times 10^{13}$ cm and a total mass $M = 26M_{\odot}$ was adopted. The explosion energy was taken to be $E = 4 \times 10^{50}$ ergs and the low luminosity tail from radioactive decay is reproduced by the ejection of $M_{\rm Ni} = 0.002M_{\odot}$ of ⁵⁶Ni. This figure is in agreement with that estimated by rescaling the

value of SN 1987A from the ratio of the luminosities of the tails. The small values of r and E/M account for both the observed low peak luminosity and expansion velocity.

The isotope ⁵⁶Ni is produced in the deep layers of the star, at the outer edge of the "iron core". In order to eject only a small amount of ⁵⁶Ni, a considerable fraction of the innermost stellar material has to fall back onto the central compact core where it is photodisintegrated before emitting γ -rays. The usual way to account for this effect in hydrodynamic simulations of the explosion is to impose a so-called "mass cut" in the post-explosion structure of the star. In the $26M_{\odot}$ model of SN 1997D of Turatto *et al.* (1998) the mass cut has been placed at 1.8 M_{\odot} . This guarantees the ejection of a mass of ⁵⁶ Ni as low as 0.002 M_{\odot} . The location of the mass cut implies that the mass of the central compact remnant is $\geq 1.8M_{\odot}$. Then, as suggested by Turatto *et al.* (1998), a BH could have formed and the low luminosity tail of SN 1997D might be a signature of the formation of such an object.

In the following we will need an estimate of the abundance of ⁵⁷Co and ⁴⁴Ti in the ejecta of SN 1997D. As an upper limit to the abundance of these two isotopes, we take the value inferred for SN 1987A ($\sim 5 \times 10^{-5} M_{\odot}$) and reduce it by a factor of 40, similarly to what estimated for the abundance of ⁵⁶Ni.

3. Late-time accretion luminosity

In a recent paper we have presented the first self-consistent, fully relativistic investigation of supernova fallback in presence of a BH (ZCSW). We computed the light curve from a hot expanding hydrogen "cloud" in which the emitted energy comes from the release of the heat residing in the original gas or generated by compression in the course of accretion onto a central BH. Radioactive energy sources were not included in these calculations. At the onset of evolution the cloud is optically thick and has homogeneous density ρ_0 ; it is set into homologous expansion with a velocity profile $u = r/t_0$, with t_0 the expansion timescale. The maximum velocity of the ejecta is $V_0 = u(r_{out}) = r_{out}/t_0$, where r_{out} is the outer radial boundary. During evolution, the unbound shells propagate outwards while those with an outward velocity smaller than the escape speed fall back after turnaround, causing accretion on a secular timescale. Initially, the cloud is radiation-dominated and photons diffuse outwards from the expanding stellar envelope. Hydrogen then recombines and the "cloud" becomes optically thin. Depending on the initial parameters, a significant amount of internal energy stored up into the gas is liberated, giving rise to a bump in the light curve. The density and velocity profiles display a self-similar character and are unaffected by the propagation of the recombination front. After the emission maximum, the gas internal energy is nearly exhausted and the luminosity falls off abruptly.

Soon after the recombination wave has propagated inward to the inner accreting region, the front stalls because the compressional heating due to accretion surpasses radiative cooling. From this moment on, the light curve is entirely powered by accretion: the central BH "becomes visible". The numerical calculations show that, at this stage, the accretion rate \dot{M} in the inner regions is nearly independent of radius. The evolution in the luminosity–accretion rate plane closely follows the analytic curve $L = L(\dot{M})$ derived by Blondin (1986) for stationary optically thick hypercritical spherical accretion onto BHs:

$$\frac{L}{L_{Edd}} \simeq 4 \times 10^{-7} \left(\frac{\mu}{0.5}\right)^{-4/3} \left(\frac{k_{es}}{0.4 \,\mathrm{cm}^2 \,\mathrm{g}^{-1}}\right)^{-1/3} \times \left(\frac{M_{bh}}{M_{\odot}}\right)^{-1/3} \left(\frac{\dot{M}}{\dot{M}_{Edd}}\right)^{5/6}, \qquad (2)$$



Fig. 1.— Bolometric light curves for SN 1987A (from ZCSW). The luminosity L is plotted as a function of time t. The *triangles* show the observed bolometric luminosity of SN 1987A. The *dashed lines* represent the expected contribution from the decay of radioactive elements (0.07 M_{\odot} of 56 Co and $5 \times 10^{-5} M_{\odot}$ of 57 Co and 44 Ti). The *dotted line* denotes the calculated bolometric luminosity emitted via accretion onto a putative BH in SN 1987A. The *arrow* marks the time of the BH emergence.



Fig. 2.— Same as in Figure 1 for SN 1997D. The data for SN 1997D *circles* are taken from Turatto *et al.* (1998). The *dashed lines* represent the expected contribution from the decay of radioactive elements (0.002 M_{\odot} of ⁵⁶Co and $1.25 \times 10^{-6} M_{\odot}$ of ⁵⁷Co and ⁴⁴Ti). Curves a, b, c and d denote the calculated accretion luminosity at late times for the following values of the initial density and expansion timescale: a) $t_0 = 2$ hr, $\rho_0 = 0.1$ g cm⁻³, b) $t_0 = 2$ hr, $\rho_0 = 2 \times 10^{-2}$ g cm⁻³, c) $t_0 = 1$ hr, $\rho_0 = 0.1$ g cm⁻³, d) $t_0 = 2$ hr, $\rho_0 = 2 \times 10^{-3}$ g cm⁻³. The *continuous line* is the total calculated bolometric light curve for case c and the arrow marks the time of the BH emergence for this case.

where M_{bh} is the BH mass, $L_{Edd} = 4\pi G M_{bh} c/k_{es} = 1.3 \times 10^{38} (k_{es}/0.4 \,\mathrm{cm}^2 \mathrm{g}^{-1})^{-1} (M_{bh}/M_{\odot})$ and $\dot{M}_{Edd} = L_{Edd}/c^2 = 2.3 \times 10^{-9} (k_{es}/0.4 \,\mathrm{cm}^2 \mathrm{g}^{-1})^{-1} (M_{bh}/M_{\odot}) M_{\odot} \,\mathrm{yr}^{-1}$ are the Eddington luminosity and accretion rate and k_{es} is the electron scattering opacity.

The late-time evolution of the accretion rate turns out to be in fair agreement with that of the hydrodynamic models computed by Colpi, Shapiro & Wasserman (1996; hereafter CSW) for polytropic flows with no net radiation flux. In fact, prior to recombination, the flow is radiation dominated and very optically thick to emission-absorption so that, from the dynamical point of view, it behaves like an adiabatic fluid gas with $\Gamma = 4/3$. On the other hand, by the time that recombination takes place and the flow becomes transparent, radiation and pressure forces at around the accretion radius $r_a = GM_{bh}/c_s^2$ are very small and the hydrodynamic evolution turns out to be close to that of a pressureless fluid (dust). Introducing the initial accretion timescale $t_{a,0} = GM_{bh}/c_{s,0}^3$ (where $c_{s,0}$ is the initial gas sound velocity) and the expansion timescale $t_0 = r_{out}/V_0$, CSW have shown that, for envelopes with $t_{a,0}/t_0 < 1$, pressure forces are important and \dot{M} evolves along a sequence of quasi-static Bondi-like states, with a slowly decreasing density at large distances. However, if the envelope has $t_{a,0}/t_0 > 1$, then the gas behaves like dust, and it can be shown that the late-time evolution of \dot{M} is given by (CSW; equation [29])

$$\frac{\dot{M}}{\dot{M}_{Edd}} \simeq \frac{4\pi^{2/3}}{9} \rho_0 t_0 k_{es} c \left(\frac{t}{t_0}\right)^{-5/3} \,. \tag{3}$$

Substituting equation (3) into equation (2), we obtain an analytic estimate of the late-time accretion luminosity for a pressureless fluid (ZCSW)

$$L = 5.7 \times 10^{36} \left(\frac{\mu}{0.5}\right)^{-4/3} \left(\frac{k_{es}}{0.4 \,\mathrm{cm}^2 \,\mathrm{g}^{-1}}\right)^{-1/2} \left(\frac{M_{bh}}{M_{\odot}}\right)^{2/3} \\ \left(\frac{\rho_0}{0.1 \,\mathrm{g} \,\mathrm{cm}^{-3}}\right)^{5/6} \left(\frac{t_0}{1 \,\mathrm{hr}}\right)^{20/9} \left(\frac{t}{1 \,\mathrm{yr}}\right)^{-25/18} \,\mathrm{erg} \,\mathrm{s}^{-1} \,.$$
(4)

We will use equation (3) to describe the evolution of the helium mantle of SN 1997D after the shock has passed through the envelope and after the reverse shock (formed at the hydrogen-helium interface) has reached the compact remnant (secondary fallback).

4. Emergence of the black hole in SN 1997D

The physical conditions of the He mantle after the passage of the shock, the amount of primary fallback and the mass of the central BH in SN 1997D depend sensitively on the explosion energy. Successful light curve and spectral fitting of SN 1997D show that the energy of the explosion was small (a few ×10⁵⁰ erg) and a significant fraction of the He mantle may have fallen back. Here we assume that, because of significant primary fallback, the mass of the BH is $M_{bh} \simeq 3M_{\odot}$ (of which ~ $1.4M_{\odot}$ represents the mass of the iron core and ~ $1.6M_{\odot}$ is accreted from the helium mantle). A few hours after the explosion, the internal energy in the He mantle of SN 1997D turns out to be typically $\tilde{E} \simeq 10^{15}$ erg g⁻¹, yielding a sound velocity $c_{s,0} = (\Gamma \tilde{E}/3)^{1/2} \simeq 2 \times 10^7$ cm s^{-1} (where the polytropic index is $\Gamma = 5/3$ since the flow is gas-pressure dominated) and an accretion timescale $t_{a,0} \simeq 3 \times 10^4$ s. The characteristic value of the expansion timescale of the helium mantle inferred from low energy explosion numerical models turns out to be $t_0 \leq 1-2$ hr (T.R. Young; private communication). Since $t_{a,0} > t_0$, the late time hydrodynamic evolution is described within the dust approximation. Accordingly, at late times the accretion luminosity of SN 1997D will be given by equation (4) with $\mu = 4/3$ and $k_{es} = 0.2$ cm² g⁻¹ to account for the helium composition. In Figures 1 and 2, we plot the estimated late-time accretion luminosity of the He mantle of SN 1987A and SN 1997D. For SN 1997D we plot four curves corresponding to different initial values for the average density and the expansion timescale, which span a physically plausible range of the parameter space. In order to bracket uncertainties in the value of the average density of the helium mantle a few hours after the explosion, we select three values of ρ_0 corresponding to outer radii in the interval $r_{He} \simeq 3 \times 10^{11}$ - 10^{12} cm. As can be seen from Figure 2, the putative BH in SN 1997D should emerge above the radioactive decay emission of the envelope approximately 3 years after the explosion. Since the luminosity by radioactive decay falls off exponentially with time, while the accretion luminosity drops only as a power law, this estimate is not very sensitive to the actual value of the initial parameters of the He mantle. An estimate of the time t_e of the BH emergence above the radioactive decay emission of the envelope can be obtained by equating equations (1) and (4). We get

$$\frac{t_e}{\tau_0} - \frac{25}{18} \ln\left(\frac{t_e}{\tau_0}\right) = 6.2 - \ln\left[\left(\frac{\mu}{0.5}\right)^{-4/3} \times \left(\frac{k_{es}}{0.4 \,\mathrm{cm}^2 \mathrm{g}^{-1}}\right)^{-1/2} \left(\frac{M_{bh}}{M_{\odot}}\right)^{2/3} \left(\frac{M_{\mathrm{Co}}}{10^{-3} M_{\odot}}\right)^{-1} \times \left(\frac{\rho_0}{0.1 \,\mathrm{gcm}^{-3}}\right)^{5/6} \left(\frac{t_0}{1 \,\mathrm{hr}}\right)^{20/9}\right].$$
(5)

Thus, $t_e \sim 6\tau_0 \sim 2$ yrs, and its value depends only logarithmically on the initial parameters. On the other hand, as shown in Figure 2, in the four cases considered, the accretion luminosity at emergence (equation [4]) varies significantly within the interval $10^{35}-5 \times 10^{36}$ erg s⁻¹. Thus, uncertainties in the inferred parameters of the mantle do not affect significantly the time of the emergence of the accretion luminosity at above the emission of 56 Co, but they do cause large uncertainties in the magnitude of the luminosity at emergence, since $L \propto \rho_0^{5/6} t_0^{20/9}$. Note that, for cases a, b, and c considered in Figure 2, the mass accreted during secondary fallback is \gtrsim a few M_{\odot} , leading eventually to the formation of a rather massive black hole.

The possible presence of other radioactive isotopes in the ejecta adds an element of uncertainty to the above discussion. As an upper limit in plotting the radioactive decay luminosity of ⁵⁷Co and ⁴⁴Ti in Figure 2 we have adopted the same rate of γ -ray deposition, at an abundance 40 times lower, as in SN 1987A (~ $1.25 \times 10^{-6} M_{\odot}$; see Section 2 for a discussion). As can be seen from the Figure, even in the least favorable case considered (case d), the accretion luminosity at emergence is as large as the decay luminosity of ⁵⁷Co.

By comparing Figures 1 and 2, it appears that SN 1997D is more promising than SN 1987A for detecting an accreting BH because of the lower amount of radioactive elements present in the envelope (which shifts the dashed curves downward) and the higher density of the He mantle (which moves the dotted curves upward).

5. Discussion and conclusions

We have shown that, if a BH has formed in SN 1997D, the late-time accretion luminosity may dominate the bolometric light curve ~ 3 years after the explosion. Depending on the actual value of the initial density and expansion timescale of the He mantle, the bolometric luminosity at emergence falls in the interval $L_e \simeq 2 \times 10^{35}$ - 10^{37} erg s⁻¹. After breakout, the light curve is expected to fall off as $t^{-25/18}$. To estimate the apparent magnitude at emergence, we assume a black body emission spectrum and extrapolate from the present temperature to get an effective temperature $T_{eff} = 4,000-6,000$ K. This is reasonable since we expect that the inner accreting region is ionized at $T \sim 10^4$ K and that dust may form in the outer ejecta reprocessing the radiation produced by accretion and re-emitting it at a lower temperature. Adopting a distance modulus $\mu = 30.64$ mag for NGC 1536, the estimated apparent visual, red and I-band magnitudes at emergence are: $m_v \simeq 27$, $m_r \simeq m_I \simeq 26$ if $L_e \simeq 10^{37}$ erg s⁻¹ and $m_v \simeq 31$, $m_r \simeq m_I \simeq 30$ if $L_e \simeq 2 \times 10^{35}$ erg s⁻¹. If $L_e \gtrsim 2 \times 10^{36}$ erg s⁻¹, the BH could be detectable for a few years in the red and I band with the Hubble Space Telescope, until the luminosity drops below threshold. Clearly, the precise value of the initial parameters is crucial in establishing the luminosity at emergence and hence whether or not the accreting BH can be detected. Even if it is only marginally visible ($L_e \sim 2 \times 10^{36}$ erg s⁻¹), subtracting off the ⁵⁶Co decay luminosity would reveal a power–law decay accretion luminosity which would confirm the presence of a BH. We note that a supernova with properties similar to SN 1997D exploding in a nearer galaxy would appear much brighter and could reveal the presence of a newly formed BH from the distinct behavior of the late–time light curve.

The $t^{-25/18}$ power law decay of the light curve is a signature of the presence of a black hole accreting from the stellar envelope and follows directly from equations (2) and (3). We note that this result is not valid for an accreting neutron star after the accretion shock has reached the trapping radius (Chevalier 1989). The reasons are that radiation pressure cannot be neglected at so early an epoch (\leq few years) and the accretion luminosity efficiency is quite different.

A number of caveats apply to our result. The formation and the actual mass of the BH depend on the amount of mass which falls back during the first hours after the explosion. This parameter is quite sensitive to the explosion energy, as shown by Woosley & Weaver (1995). In addition, the state of the flow in the He mantle is determined by the global hydrodynamics of the explosion. The estimate of the late-time accretion luminosity given in Section 3 assumes that the opacity in the accreting region is contributed mainly by electron scattering and that the thermal and dynamical effects of radioactive heating can be neglected. Early in the evolution, in the inner accreting region, the temperature is relatively high $(T \gtrsim 10^7)$ K) so that all elements are almost completely ionized. In these conditions, the dominant contribution to the Rosseland mean opacity comes from electron scattering and hence our assumption is reasonable. Later on, the temperature decreases and bound-free and bound-bound transitions (especially those of heavy elements) may be important. A lot of photons could remain trapped in the flow. Analogous considerations apply to the expanding envelope at late times. For example, if the mean opacity of the envelope is $\sim 10^3 k_{es}$. the time taken by photons to diffuse out of the He mantle would be ~ 20 years for the typical parameters of SN 1997D. Finally, in deriving our estimate we have assumed that, after 1-2 years, the amount of heat deposited by γ -rays from radioactive decay is not sufficiently large to alter the character of the flow. Given the small amount of heavy elements in the ejecta of SN 1997D, this assumption seems reasonable.

Improved radiation-hydrodynamic calculations are necessary to provide a theoretical guide for identifying the nature of the compact remnant.

It is a pleasure to thank Massimo Turatto, Tim Young and their collaborators for providing us with data for the helium mantle of SN 1997D in advance of publication and for several useful discussions. We thank the referee for useful comments. This work was supported in part by NSF grant AST 96–18524 and NASA grants NAG–5–2925 and NAG–5–3420 at the University of Illinois at Urbana–Champaign.

REFERENCES

- Akmal, A., Pandharipande, V.R., & Ravenhall, D.G. 1998, Phys. Rev. C, submitted; nucl-th/9804027
- Blondin, J.M. 1986, ApJ, 308, 755
- Brown, G.E., & Bethe, H.A. 1994, ApJ, 423, 659
- Chevalier, R.A. 1989, ApJ, 346, 847
- Colpi, M., Shapiro, S.L., & Wasserman, I. 1996, ApJ, 470, 1075 (CSW)
- De Mello, D., & Benetti, S. 1997, IAUC 6537
- Turatto et al. 1998, ApJ, 498, L129
- Woosley, S.E., & Timmes, F.X. 1996, Nucl. Phys. A, 606, 137
- Woosley, S.E., & Weaver T.A. 1995, ApJS, 101, 181
- Zampieri, L., Colpi, M., Shapiro, S.L., Wasserman, I. 1998a, ApJ, in press (ZCSW); astro-ph/9804122
- Zampieri, L., Colpi, M., Shapiro, S.L., Wasserman, I. 1998b, in the Proceedings of the 8th Annual Astrophysics Conference in Maryland, "Accretion Processes in Astrophysical Systems", eds. S.S. Holt & T. Kallman (New York: AIP Press), 417

This preprint was prepared with the AAS LATEX macros v4.0.