Propagation of a Huygens front through turbulent medium

M. Chertkov^a and V. Yakhot^b

Physics Department^a and Programm in Applied and Computational Mathematics^b,

Princeton University, Princeton, NJ 08544, USA

(September 25, 1997)

The dynamics of a thin Huygens front propagating through turbulent medium is considered. A rigorous asymptotic expression for the effective velocity v_F proportional to the front area is derived. The small-scale fluctuations of the front position are shown to be strongly intermittent. This intermittency plays a crucial role in establishing a steady state magnitude of the front velocity. The results are compared with experimental data.

The problem of propagation of a thin passive front (of flame, phase-transition, etc.) through turbulent flow has attracted a lot of attention since early forties [1,2]when it was realized that velocity fluctuations tend to generate strongly convoluted front, thus dramatically increasing their area. In premixed combustion processes the flame front area is directly related to the speed of the front propagation. The first expression for the flame front velocity $v_F \propto u_{rms}$, valid in the limit $u_{rms} \gg u_0$, where u_{rms} is the root-mean-square velocity of the integral eddy and u_0 is the laminar flame speed depending on the details of chemical kinetics, was proposed by Schelkin [2] (see [3,4] for modern reviews on the theory of turbulent combustion). The problem is also important for description of light propagation in the media with fluctuating dielectric constant, shock wave fronts etc. [6]. The renormalization group approach [7], which included some theoretically unjustified steps, yielded the expression, differing from the Schelkin result by a logarithmic factor, which agreed with experimental data [8–10] in a wide range of parameter variation. Still, despite substantial activity, no rigorous derivation of the front speed appeared and the question of the dependence of v_F on both u_0 and the Reynolds number Re remained open. It is shown in this work that recent advances in the theory of a passive scalar, advected by turbulence, enable one to accurately account for the small-scale intermittency of a scalar field, crucial for description of the front fluctuations and derivation of the effective velocity v_F .

We consider a problem of propagation of a passive front though turbulent flow. The front can be described by the equation for a passive scalar [3,11] (so called "G"equation)

$$\partial_t G + (\mathbf{v}\partial_\mathbf{r})G = u_0|\partial_\mathbf{r}G|,\tag{1}$$

where $G(t, \mathbf{r})$ is a scalar field whose level surface, say G = 0, represents the thin front position. Statistics of turbulent velocity \mathbf{v} is supposed to be known. The equation (1) with $\mathbf{v} = 0$ describes a front propagating with the constant speed u_0 (laminar front speed) normally to the local orientation of the front. For example, if the front at time t = 0, is defined at the x - y plane, it will propagate with a constant speed u_0 and constant area

 $S_0 = const$ in the z- direction. The role of the random field **v** is in generation of a strongly convoluted (" wrinkled") front with a substantially increased area $S_T > S_0$. When **v** = 0 the mass of the reagents (fuel) consumed per unit time is: $dm/dt = u_0S_0 = const$ does not change in time. In general, $dm/dt = u_0S_T \equiv v_FS_0$, where S_T is the area of the wrinkled front. This gives a definition of a turbulent or effective front velocity, $v_F = u_0S_T/S_0$. Assuming that a steady (both v_F and S_T do not depend on time after a long evolution) regime is realized, we introduce a new variable, $G(t, \mathbf{r}) \equiv v_F t - z + h(t, \mathbf{r})$. Then, (1) reads:

$$\partial_t h + (\mathbf{v}\partial_\mathbf{r})h = v_z + u_0\sqrt{\left(\partial_\mathbf{r}h\right)^2 + 1 - 2\partial_z h} - v_F.$$
 (2)

In the moving frame the fluctuations of the front position h are assumed to be in a statistically steady state, that fixes the value of the front speed v_F , which in the turbulent regime discussed ($v_F \gg u_0$, when $|\partial_r h|^2 \gg |\partial_z h| \gg 1$) is given by

$$v_F = u_0 \langle |\partial_{\mathbf{r}} h| \rangle \sim u_0 \langle |\delta h(r_0)| \rangle / r_0.$$
(3)

Here, $|\delta h(r_0)|$ is a magnitude of the velocity difference at the scale r_0 where the "chemical" (u_0 -dependent) and advective contributions to (2) balance each other. Averaging over the turbulent velocity is assumed in (3). Since $u_{rms} \gg u_0, r_0 \ll L$, where L is the scale of the turbulence source. This means that the scalar, injected at the scale L, is dissipated at the propagating front as a result of generation of very sharp cusps of the radius $r_0 << L$. Formation of such cusps was observed in numerical simulations [12,13]. Derivation of characteristic width r_0 , magnitude $|\delta h(r_0)|$ of these cusps and, as a result, v_F , is the goal of the theory.

The presence of the two characteristic scales η and r_0 defines three possible flame regimes:

A:
$$L \gtrsim \eta \gg r_0$$
, B: $L \gg r_0 \gg \eta$, C: $L \gg \eta \gg r_0$. (4)

Generation of the small-scale scalar fluctuations (direct cascade) [14,15] takes place in both inertial-convective, $L > r > \eta$ and the dissipative-convective, $\eta > r > r_0$ (which is valid in the A, C cases but not B) intervals. Thus, the problem is naturally divided into two: First,

we need to describe scalar (height of the flame brush) correlations in both convective ranges $L > r \gg r_0$. Once the solutions in the convective intervals are found, we will be ready to resolve the second and principal part of the problem: to calculate the value of the dissipative scale r_0 and, matching dissipative and convective intervals, find turbulent speed of the front, v_F . Therefore, naturally, we are starting from the first task, considering all the regimes (A - C) one after another.

<u>A.</u> Batchelor (a pure viscous - convective) regime. The case A corresponds to a well- studied situation, first discussed by Batchelor [16] and developed further in [17–19]. Without loss of generality and following [19] we will consider the velocity difference to be Gaussian in the case

$$\left\langle \delta v_r^{\alpha}(t) \delta v_r^{\beta}(t) \right\rangle = \frac{D}{\tau} \left[2\delta^{\alpha\beta} r^2 - r^{\alpha} r^{\beta} \right] \exp\left[-\frac{t}{\tau} \right], \quad (5)$$

where τ is turn-over time of the integral (*L*-size) eddy. The pair correlation function of the scalar obeys the famous logarithmic law in the convective interval of scales, $L \gg r \gg r_0$, [17–19]

$$\langle h_1 h_2 \rangle = DL^2 \ln \left[L/r_{12} \right] /\lambda, \tag{6}$$

where DL^2 in the nominator of the logarithmic prefactor describes the fluctuation of the "source" function v_z while λ stands for the Lyapunov exponent corresponding to the rate of Lagrangian stretching. Correlation between the source and convective terms in (2) does not contribute to (6). Accounting for these correlations slightly modifies the higher-order moments generating subleading contributions and a mere renormalization of bare coefficients. The Lyapunov exponent λ as a function of τ , D was found for the two-dimensional version of the model (5) in [19]. Asymptotic of the large and small τ were described explicitly in [19]. Generally, the problem of finding λ was reduced in [19] to a well-defined auxiliary quantum mechanics which was easy to solve numerically. An interpolation formula for λ , fitting well all the known asymptotic is:

$$\lambda = \sqrt{D/\tau} \tanh\left[\sqrt{D\tau}\right].$$
 (7)

This formula, valid in the space of arbitrary dimensionality d > 2, holds up to $\mathcal{O}(d)$ corrections. Statistics of the scalar fluctuations in the convective interval is shown to be Gaussian [19]. Therefore, the typical fluctuation of the height h at r_0 , estimated by the second moment (6) is

$$|\delta h_{r_0}| \sim L\sqrt{D/\lambda}.$$
 (8)

The expression (8) is derived without any spatial averaging over the large- scale ($\sim L$) structures, always present in a real flow where all macroscopic characteristics, including the Lyapunov exponents λ , are slightly modulated on the integral scale. Accounting for this spatial variation gives an estimate following directly from (6):

$$|\delta h_{r_0}| \sim L \sqrt{D/\lambda} \sqrt{\ln [L/r_0]}.$$
 (9)

The modified regime A, accounting for the large- scale averaging, will be denoted hereafter by A'.

<u>B</u>. Pure inertial-convective range. Inertial-convective range is realized at the scales $r \gg \eta \gg r_0$ where one cannot neglect small-scale advection contribution. In the case of a general non-smooth velocity one finds a strongly intermittent behavior, manifested in the anomalous scaling of the scalar structure functions:

$$S_{2n}(r) = \left\langle \left[h(\mathbf{r}_1) - h(\mathbf{r}_2) \right]^{2n} \right\rangle \sim L^{2n - \zeta_{2n}} r_{12}^{\zeta_{2n}}, \quad (10)$$

valid at the separations $r_{12} \ll L$. The fundamental origin of the anomalous scaling, $\zeta_{2n} < n\zeta_2$, was discovered recently [21–23]. It was understood that the anomalous exponents originate from zero modes of the eddy-diffusivity operator: ζ_{2n} are universal numbers, solely defined by the velocity statistics and independent on the properties of the pumping term. The exponents were analytically calculated for the case of the velocity field, rapidly varying in time, introduced by Kraichnan [20] in $1/d_{-}$, $2-\zeta_{2}$, and ζ_{2} -expansions [21], [22], [23], respectively. An instanton approach [24] yields yet another large n ($n \gg d$) asymptotic for $\zeta_{2n} \to \zeta_{\infty}(\zeta_2, d)$ when $n \to \infty$. The constant $\zeta_{\infty}(\zeta_2, d)$ was explicitly calculated. The saturation of exponents ζ_n was predicted also in [25]). The instanton consideration, applied to a general passive scalar problem, always results in the collapse of exponents $\zeta_{2n} \to \zeta_{\infty} < \infty$ with the asymptotic value ζ_{∞} to be a complicated functional of the velocity field statistics. It can be easily understood: the 2n-th moment of the scalar can be represented as a path integral over 2nfluid particles. The dominant contribution into the high (2n-th) order structure function originates from the most probable 2n- particle trajectory. In the incompressible world it is impossible to avoid a divergence of particles at least in one of the directions and it has been shown in [24] that contribution to the path integral from a single diverging trajectory is sufficient to cause saturation of the exponents ζ_n . This effect has a very strong and important influence on the parametric dependence of an effective front speed u_T calculated below.

<u>C</u>. Consecutive inertial-convective and dissipativeconvective ranges. The description of the scalar correlations in the inertial-convective regime (ICR) does not deviate from the one considered above for the case *B*. However, in the dissipative-convective interval (DCR), $L \gg \eta \gg r \gg r_0$, the behavior of the scalar is very different from that observed in the low-Reynolds- number Batchelor regime when $L \approx \eta$. The crucial difference stems from the essential non-Gaussianity and intermittency of the scalar field on the velocity dissipation scale η where the solutions in both intervals have to match. In this case the scale η is an integral scale for the DCR $r_0 \ll r \ll \eta$ where powerful injection of all higher- order integrals of motion $(\int d\mathbf{r} h^n)$ takes place. The point is that a 2n-2 moment defines a pumping for the next 2n-th order one. Thus, considering a generalization of the Kraichnan model for the case C(and neglecting the correlations between the source and convective terms, which can slightly renormalize some constants) one obtains the exact equations for the multi-point correlation functions $F_{2n}(\mathbf{r}_1, \cdots, \mathbf{r}_{2n}) \equiv \langle h(\mathbf{r}_1)h(\mathbf{r}_2)...h(\mathbf{r}_{2n}) \rangle$,

$$\hat{L}F_{2n} = V^{11}(r_{12})F_{2n-2}(\mathbf{r}_3,\cdots,\mathbf{r}_{2n}) + perm.,$$
 (11)

where $\hat{L} \equiv -\mathcal{K}^{\alpha\beta}(\mathbf{r}_{ij})\nabla_i^{\alpha}\nabla_j^{\beta}$, $V^{\alpha\beta}(r) = K^{\alpha\beta}(L) - K^{\alpha\beta}(r)$ and $K(r) \sim r^{2-\gamma}D$ at, $L > r > \eta$, while $K(r) \sim r^2\eta^{-\gamma}D$ at $\eta > r$. Operator \hat{L} has the $-\gamma$ dimensionality (it scales as $r^{-\gamma}$) in the inertial-convective range while it is $O(r^0)$ in the dissipative-convective range. As a result, the 2*n*-th moment F_{2n} , explicitly depending on all 2*n*-th points, is dominated by the forced "logarithmic" solution of (11) in the DCR. In the ICR, however, it is a zero mode of the operator \hat{L} which dominates the solution. Therefore, the 2*n*-th moment of the scalar difference, $S_{2n}(r) = \langle \delta h_r^{2n} \rangle$ is estimated by

$$S_{2n}(r) = a_n L^2 \ln\left[\frac{r}{r_0}\right] S_{2n-2},$$
 (12)

where a_n are dimensionless *n*-dependent constants. On the another hand, at the scale η (14) this expression should match the anomalous structure functions (10) from the upper (inertial-convective) interval giving:

$$S_{2n}(r) \sim \eta^{\zeta_{2n}} L^{2n-\zeta_{2n}} \frac{\ln^n [r/r_0]]}{\ln^n [\eta/r_0]}.$$
 (13)

Let us proceed now with the second task and estimate the value of the dissipative scale, r_0 , required for calculation of the front velocity u_T . The point of a crucial importance is a necessity to distinguish between the different moments of the front fluctuations at the dissipative scale, δh_{r_0} . Usually one estimates the typical fluctuations as a root-means-square value of the corresponding variable. This is fine in case of "normal scaling" or if the small-scale intermittency is not too strong (cases A and A'). However in the cases B and C intermittency in the convective intervals is extremely strong: estimates for a typical front fluctuation based on various moments, $\delta h_r^{(n)} \approx \langle \delta h_r^n \rangle^{1/n}$ are very different. As a result we get from (8,9,10,13) the following *n*-dependent estimate at the dissipative scale r_0

$$|\delta h_{r_0}^{(n)}| \sim L \begin{cases} L\sqrt{D/\lambda}, & A, \\ \sqrt{D/\lambda}\sqrt{\ln[L/r_0]}, & A', \\ [r_0/L]^{\zeta_n/n}, & B, \\ [\eta/L]^{\zeta_n/n}/\sqrt{\ln[\eta/r_0]}, & C, \end{cases}$$
(14)

for the cases A, A', B and C respectively. On the other hand, the value of $|\delta h_{r_0}|$ is defined by equilibration of the convective and "dissipative" terms in (2) giving:

$$|\delta h_{r_d}^{(n)}| \sim v_F T, \quad T \sim \begin{cases} \lambda^{-1} \ln [L/r_0], & A, A', \\ L/v_{rms}, & B, C. \end{cases}$$
 (15)

It shows that the linear dimension of the flame brush, calculated on the dissipative scale, is defined by the turbulent flame speed and the overall time of Lagrangian evolution which is a typical time for Lagrangian separation, initially equal to r_0 , to reach the integral scale. In the B, C regimes the overall time is r_0 - independent (note that the time of evolution from r_0 to η is neglected in comparison with one describing the inertial-convective stage of evolution, from η to L). Which number n is essential to define v_F via (15) is yet to be discussed. One should add (3) to (14,15). Collecting all the relations in one table we get

where, $\beta_n \equiv [1 - \zeta_1 + \zeta_n/n]^{-1}$. As was cited before, $\zeta_{2n}/[2n] \to 0$ and respectively, $\beta_n \to 1/[1 - \zeta_1] > 0$, at $n \to \infty$. Since, intermittency of the scalar is growing downscales from the scale of the pumping, the higher ratio of the integral scale to the dissipative one (of the turbulent velocity to the chemical one) the higher the moment-number *n* entering the actual stationary velocity of the flame (which is defining the actual width of the front). The highest value of $\delta h_{r_0}^{(n)}$ is reached at $n \to \infty$. That is to say that the stationary speed of the front is defined by such optimal and rare (instanton) configurations

which give the dominant contribution into the highest moments of the scalar field h. The optimal configurations define the maximal spreading of the flame brush and control (mainly) the dissipation and the stationary velocity of the brush finally. The necessity to account for the highest moments of the δh_{rd} has a simple Lagrangian explanation. The point of a crucial importance is a failure of 2n- particle description of the 2n-order correlation function of the scalar at the scales smaller than the dissipative one. This fact can be easily illustrated by a numerical procedure attempting to describe the initially plane front in terms of collection of n Lagrangian particles [26] each moving with velocity $u_0 \mathbf{n} + \mathbf{v}(\mathbf{x}, \mathbf{t})$ where **n** is a unit vector in the z-direction. The initial (t = 0) distance between these particles is $l_0 \approx 1/n$. Very soon after beginning of the simulation the sharp cusps started to form and one had to add more and more particles to preserve continuity of the front. The development of these very sharp gradients, where dissipation takes place, drives the necessary number of particle $n \to \infty$. We would like to reiterate a very important and profound peculiarity of the situation: the expression (16), taken at the largest n, corresponds to a very rare optimal (instanton) configuration, responsible for the highest moments of the scalar difference. This means that to describe the small-scale $(r \ll r_0)$ dynamics of the front height $h(t; \mathbf{r})$, one has to evaluate contributions from these configurations. We postpone the more accurate investigation (which requires some further development of the instanton technique of [24] accounting for the entire dynamics of the front surface) for a future publication.

One of the principle results of this work, given by (16)at $n \to \infty$, is that turbulent front speed strongly depends on the ratio $\Gamma = r_0/\eta$, which is a novel dimensionless parameter of the problem. Therefore in order to compare experimental data with theoretical predictions we must first estimate Γ and determine the front propagation regime. Recent experimental studies of a passive front propagating in turbulent media were performed in the flows generated by vibrating grids, capillary waves, Taylor-Couette flow and Hell-Show cells. All experiments (for the turbulent regime, $v_{rms} \gg v_0$) corresponded to the case C. For example, in the vibrating grid experiments [10] the values of $u_{rms} \approx 1 \ cm/sec$, $L \approx 1-10 \ cm$, $Re \approx 10^2 - 10^3$ and $u_0/u_{rms} \approx 10^{-2} - 10^{-3}$. (In the case of the Kolmogorov turbulence one has an estimation $\eta \approx 10 L R e^{-3/4}$, where Re is the Reynolds number and the factor $c \approx 10$ agrees with available experimental data). For the experimental conditions of [9] one gets $\Gamma \approx 10^{-3} - 10^{-4}$. The expression for the front velocity, following from the Table (16), evaluated at r_0 corresponding to $n \to \infty$ is:

$$v_F \sim \frac{u_{rms}}{\sqrt{\frac{2}{3}ln\left[\eta/L\right] + ln\left[U\right]}}_{U \to \infty} \to \frac{u_{rms}}{\sqrt{ln\left[U\right]}}, \qquad (17)$$

where $U \equiv u_{rms}/u_0$. The rhs of (17) (similar formula was derived in [7], see also [27], where the dynamic renormalization group has been used for evaluation of the turbulent flame speed v_F) agrees very well with experimental data on v_F in a variety of turbulent flows in a wide range of variation of the dimensionless turbulent intensity $U \leq 20 - 500$. This universality can be readily understood: in all the experimental situations the $O(\ln [\eta/L])$ - contribution is not large and can be neglected in comparison with the O(ln[U])-term. In a typical case $U \approx 20 - 100$ the transitional u_{rms} -based Reynolds number is: $Re_c \approx 10^3 - 10^4$, which is very high. This explains the relatively broad applicability of the large U asymptotic of (17) and therefore challenge a higher Re experiment to test the general structure of (17).

We acknowledge support of a R.H. Dicke fellowship (MC) and the ONR/URI grant (VY).

- G. Z. Damkohler, Z. Electrochem. Angew. Phys. Chem. 46, 601 (1940).
- [2] K.I.Schelkin, NACA Tech. Memo. 1110 (1947).
- [3] F. A. Williams, "Combustion theory", The Benjamin Publishing, 1985.
- [4] S. B. Pope, Ann. Rev. Fluid Mech 19, 237 (1987).
- [5] P.D. Ronney, in J. Buckmaster and T. Takeno Eds, Lecture notes in Physics, vol. 449 (Springer-Verlag, Berlin, 1995).
- [6] M.C. Cross, P.C. Hohenberg, Rev. of Mod. Phys. 65, 851 (1993).
- [7] V. Yakhot, Comb. Sci. and Tech. 60, 191 (1988).
- [8] S.S. Shy, P.D. Ronney, S.G. Buckley, and V. Yakhot, Proceedings of the 24th Symposium on Combustion, Combustion Institute, Pitsburgh 1992.
- [9] P.D. Ronney, B.D. Haslam, and N.O. Rhys, Phys.Rev. Lett. 74, 3804 (1995).
- [10] S.S. Shy, R. H. Jang, and P.D. Ronney, Combust. Sci. and Tech. 113-114, 329 (1996).
- [11] A. R. Kerstein, W.T. Ashurst, and F. A. Williams, Phys. Rev. A 37, 2728 (1988).
- [12] W.T. Ashurst, G. I. Sivashinsky, and V. Yakhot, Combust. Sci. and Tech. 62,273 (1988).
- [13] S. Osher, and J.A. Sethian, J. Comp. Phys. 79, 12 (1988).
- [14] A. M. Obukhov, Izv. Akad. Nauk SSSR, Geogr. Geophiz.13, 58 (1949).
- [15] S. Corrsin, J. Appl. Phys. 22, 469 (1951).
- [16] G.K. Batchelor, J. Fluid Mech. 5, 113 (1959).
- [17] R. Kraichnan, Phys. Fluids 10, 1417 (1967).
- [18] B. Shraiman and E. Siggia, Phys. Rev. E 49, 2912 (1994).
- [19] M. Chertkov, G. Falkovich, I. Kolokolov, and V. Lebedev, Phys. Rev. E 51, 5609 (1995).
- [20] R. H. Kraichnan, J. Fluid Mech. 64, 737 (1974).
- [21] M. Chertkov, G. Falkovich, I. Kolokolov, and V. Lebedev, Phys. Rev. E 52, 4924 (1995).
- [22] K. Gawedzki, and A. Kupianen, Phys. Rev. Lett. 75, 3834 (1995).
- [23] B. Shraiman, and E. Siggia, C.R. Acad. Sci. **321**, Serie II, 279 (1995).
- [24] M. Chertkov, Phys. Rev. E 55, 2722 (1997).
- [25] V. Yakhot, Phys.Rev. E 55, 329 (1997).
- [26] Agistein, A. Migdal, and V.Yakhot, unpublished (1991).
- [27] A. R. Kerstein, Combust. Sci. and Tech. 60,163 (1988).