Electrodynamics of helium with retardation and diffusion in the neighborhood of Wannier orbits...

Jayme De Luca Instituto de Química de Araraquara-UNESP Caixa Postal 355, Araraquara, S˜ao Paulo, 14801-970

We show that an extra constant of motion with an analytic form can exist in the neighborhood of some discrete Wannier orbits of helium, when one includes retardation and self-interaction effects. The energies of these discrete stable Wannier orbits are in the correct atomic magnitude. The highest frequency in the stable manifold of one such orbit agrees with the highest frequency sharp line of parahelium to within 2%. The generic term of the frequency in the stable manifold to higher orbits is also in agreement with the asymptotic form of quantum mechanics for helium.

PACS numbers: 31.15.Ct, 03.20.+i,05.45.+b

In this letter we explore some surprising consequences of the retardation effects of Maxwell's electrodynamics to atomic physics. We show that electrodynamics with retardation prescribes a discrete set of stable Wannier orbits for the helium atom. A Wannier orbit is one where the two electrons are in the same circular orbit and in phase opposition, that is, along a diameter [\[1\]](#page-4-0).

The linearized dynamics about a Wannier orbit has a one-dimensional stable direction, a one-dimensional unstable direction and ten neutrally stable oscillatory modes [\[1](#page-4-0)]. For an infinitely massive α -particle, helium is a six-degree of freedom Hamiltonian system with only four independent constants of motion, (namely, the energy and the three components of the total angular momentum). In the neighborhood of Wannier orbits, even if one restricts to the center manifold[[2\]](#page-4-0), one still has a five-degree of freedom system. Because there are only four constants, the dynamics in the neighborhood of a generic Wannier orbit can diffuse away to large distances of the orbit. Here we show that an extra complex constant can exist in the neighborhood of a discrete set of Wannier orbits, when one includes retardation effects.

To include the retardation, we use the work of Page (1918) [[3\]](#page-4-0) on the expansion of the Liénard-Wiechert formula [\[4](#page-4-0)]. This formula is complicated because of the retardation constraint, and one way to convert it into a useful differential equation is to develop the constraint in a Taylor series. This was done by Page up to the fifth orderin 1918 [[3\]](#page-4-0). Truncated to second order in (v/c) , this retarded interaction is described by the Darwin Lagrangian [\[4](#page-4-0)], which introduces the first retardation correction to the Coulomb interaction.

The Darwin Lagrangian is rotationally invariant, which generates an angular momentum-like constant of the motion according to Noether's theorem [\[6](#page-4-0)]. This constant is the angular momentum of the electrons plus the small correction of the angular momentum in the electromagnetic field. There is also an energy-like constant (because the Darwin Lagrangian is time independent). According to a recent result, the existence of a constant of motion analytic in the neighborhood of an equilibrium point requires the eigenvalues of the linearized dynamics to satisfy some resonance condition [\[7](#page-4-0)]. Since the eigenvalues in the presence of retardation depend on the particular orbit, an extra complex constant is possible for some discrete orbits, which is a genuine signature of the

nonlinear dynamics with retardation. We stress that it is absolutely essential to include the retardation to unfold a degeneracy of the Coulomb dynamics, and not to attain some better precision. Last, we consider only the dynamics of spinless pointlike charges, and issues like quantum mechanical spin are not discussed.

In this work we assume that the self-interaction effects are described by the Lorentz-Dirac equation (LDE) with a renormalized mass[[4,8](#page-4-0)]. The LDE equation for an electron of charge $-e$ can be written in the convenient, non-covariant form as

$$
\frac{d}{dt}(\gamma M_e \dot{\mathbf{x}}_e) = \mathbf{F}_{rad} + \mathbf{F}_{ext},
$$
\n(1)

where $\dot{\mathbf{x}}_e$ is the electron's velocity, M_e the renormalized electronicmass [[8\]](#page-4-0) and $\gamma = 1/\sqrt{1 - (|\dot{\mathbf{x}}_e|/c)^2}$. In (1), \mathbf{F}_{ext} is the external electromagnetic force and in this work we use $\mathbf{F}_{rad} = \frac{2e^2}{3c^3}$ $\frac{2e^2}{3c^3}$ $\dddot{\mathbf{x}}_e$, which is the lowest order approximation in powers of (v/c) to the exact relativistic radiaton force [\[8\]](#page-4-0). For a circular orbit, this is a term of order $(v/c)^3$.

We now introduce the expansion of the retardation constraint of the Liénard-Wiechert interaction, hence-forthcalled the Page series [[3\]](#page-4-0). Let x_1 be the position of electron 1, of charge $-e$, and β_1 its velocity vector divided by c. The formula for the electric field made by electron 1 at a position x_2 is

$$
\mathbf{E}_{12} = -e\frac{\hat{\mathbf{n}}}{R_{12}^2} + \mathbf{E}_{12}^{(2)}(\mathbf{x}_2, \mathbf{x}_1, \dot{\mathbf{x}}_1, \ddot{\mathbf{x}}_1) + \mathbf{E}_{12}^{(3)}(\dddot{\mathbf{x}}_1) + \dots (2)
$$

where

$$
\mathbf{E}_{12}^{(2)} \equiv -e\{\frac{(|\beta_1|^2 - 3(\hat{\mathbf{n}} \cdot \beta_1)^2)\hat{\mathbf{n}}}{2R_{12}^2} - \frac{\dot{\beta}_1}{2cR_{12}} - \frac{(\hat{\mathbf{n}} \cdot \dot{\beta}_1)\hat{\mathbf{n}}}{2cR_{12}}\},\tag{3}
$$

and

$$
\mathbf{E}_{12}^{(3)}(\ddot{\mathbf{x}}_1) \equiv -\frac{2e}{3c^3} \dddot{\mathbf{x}}_1,\tag{4}
$$

where $\hat{\mathbf{n}}$ is the unit vector in the direction of $\mathbf{x}_2 - \mathbf{x}_1$ and $R_{12} = |\mathbf{x}_2 - \mathbf{x}_1|$. Notice that all functions are evaluated at present time. For a circular orbit, in units of the zeroth order Coulomb term, (3) is of order $(v/c)^2$ and (4) is of order $(v/c)^3$. We assume that the Taylor series of (2) converges, since $(v/c) \ll 1$ in atomic physics. The first

term of the series for the magnetic field of electron 1 at \mathbf{x}_2 is [[3\]](#page-4-0)

$$
\mathbf{B}_{12}^{(1)}(\mathbf{x}_2, \mathbf{x}_1, \dot{\mathbf{x}}_1) = -\frac{e}{R_{12}^2} [\beta_1 \times \hat{\mathbf{n}}].
$$
 (5)

In this work we consider only the above terms of the electromagnetic interaction, as well as the relativistic correction to the electronic dynamics up to order $(v/c)^2$.

Along a Wannier orbit, the total force acting on electron 2 can be calculated, using (3) , (4) , (5) and the selfinteraction of electron 2, to be

$$
\mathbf{F}_2 = \frac{2e^2}{3c^3} (\dddot{\mathbf{x}}_1 + \dddot{\mathbf{x}}_2) - 7 \frac{e^2 \hat{\mathbf{n}}}{4R^2} (1 - \frac{3}{7} |\beta|^2),
$$
(6)

where R is the radius of the Wannier orbit. In (6) we have also added the Coulomb attraction of the α -particle (of charge 2e), that we suppose infinitely massive and resting at the origin. Along the Wannier orbit, because the electrons are in phase opposition, the force along the velocity cancels out, demonstrating that the Wannier orbit is a possible periodic solution of the electromagnetic equations up to third order in (v/c) .

Notice the appearance of the dipole term in (6),

$$
\ddot{\mathbf{D}} = -e(\ddot{\mathbf{x}}_1 + \ddot{\mathbf{x}}_2). \tag{7}
$$

The total far field caused by the electrons depends linearly on the quantity $\ddot{\mathbf{D}}$ of (7), up to quadrupole terms [[4\]](#page-4-0). If D is zero, the orbit does not radiate in dipole. The fifth-order quadrupole terms would be important only in a much longer time scale, of order $T/(v/c)^5$ [[9\]](#page-4-0). In this letter we consider only the dynamical system defined by truncating the Schott-Page interaction and selfinteraction to third order in (v/c) .

Next we consider the dynamics of helium in the neighborhood of a Wannier orbit, according to the Darwin Lagrangian. For the Coulomb dynamics, this stability problem has already been considered by many authors [[1,10\]](#page-4-0). Here we consider only the second order of the Page series, which is essential to unfold a degeneracy of the Coulomb dynamics. The detailed consideration of the dissipative effect of the third-order term of the Page series and the (also third-order) self-interaction term will be published elsewhere [\[9](#page-4-0)]. Here it suffices to say that these higher-order terms are important only in a much longer time-scale. To second order in (v/c) , the equations of motion for helium with retardation and relativistic correction are

$$
M_e \frac{d}{dt} [(1 + \frac{1}{2}|\dot{\mathbf{x}}_1|^2) \dot{\mathbf{x}}_1] = -\frac{2e^2}{R_1^3} (\mathbf{x}_1) - \frac{e^2}{R_{12}^3} (\mathbf{x}_2 - \mathbf{x}_1)
$$

+ $\mathbf{E}_{21}^{(2)} (\mathbf{x}_1, \mathbf{x}_2, \dot{\mathbf{x}}_2, \ddot{\mathbf{x}}_2) - \frac{e}{c} \dot{\mathbf{x}}_1 \times \mathbf{B}_{21}^{(1)} (\mathbf{x}_1, \mathbf{x}_2, \dot{\mathbf{x}}_2),$

$$
M_e \frac{d}{dt} [(1 + \frac{1}{2}|\dot{\mathbf{x}}_2|^2) \dot{\mathbf{x}}_2] = -\frac{2e^2}{R_2^3} (\mathbf{x}_2) - \frac{e^2}{R_{12}^3} (\mathbf{x}_1 - \mathbf{x}_2)
$$

+ $\mathbf{E}_{12}^{(2)} (\mathbf{x}_2, \mathbf{x}_1, \dot{\mathbf{x}}_1, \ddot{\mathbf{x}}_1) - \frac{e}{c} \dot{\mathbf{x}}_2 \times \mathbf{B}_{12}^{(1)} (\mathbf{x}_2, \mathbf{x}_1, \dot{\mathbf{x}}_1),$ (8)

where M_e is the electronic mass, $R_1 = |\mathbf{x}_1|, R_2 = |\mathbf{x}_2|,$ $R_{12} = |\mathbf{x}_1 - \mathbf{x}_2|, \, \mathbf{E}_{12}^{(2)}(\mathbf{x}_2, \mathbf{x}_1, \dot{\mathbf{x}}_1, \ddot{\mathbf{x}}_1)$ is defined in [\(3](#page-1-0)) and $\mathbf{B}_{12}^{(1)}(\mathbf{x}_2,\mathbf{x}_1,\dot{\mathbf{x}}_1)$ is defined in (5). The Wannier orbit is the following periodic solution of (8)

$$
x_1 = R\cos(\omega t), y_1 = R\sin(\omega t), z_1 = 0,
$$

\n $x_2 = -R\cos(\omega t), y_2 = -R\sin(\omega t), z_2 = 0.$ (9)

According to (8), the frequency of the orbit is related to R by

$$
M_e \omega^2 = \frac{7e^2}{4R^3} (1 - \frac{3}{7}|\beta|^2), \tag{10}
$$

which is correct to second order in (v/c) .

Linearizing (8) about the Wannier orbit (9), we obtain a parametric linear differential equation with coefficients periodic in time and period $T = \pi/\omega$ [[9\]](#page-4-0). If the Floquet exponents are all nondegenerate, one can find a complete set of solutions of the form [\[11](#page-4-0)]

$$
\delta \mathbf{x}_1 = \exp(2i\omega\mu t) \sum_n \mathbf{x}_n^1 \exp(2in\omega t),
$$

$$
\delta \mathbf{x}_2 = \exp(2i\omega\mu t) \sum_n \mathbf{x}_n^2 \exp(2in\omega t),
$$
 (11)

where the Floquet exponent μ is a complex number defined in the first Brillouin zone, $-1/2 < Re(\mu) < 1/2$. Notice that for the Floquet components we use an upper index, but to label coordinates as functions of time, as in (8), we use a lower index (to distinguish it from the Floquet components). To bring the variational equations to normal form, we define the coordinates

$$
\xi_n^{\kappa} \equiv \frac{1}{\sqrt{2}} (x_n^{\kappa} - iy_n^{\kappa}), \ \chi_n^{\kappa} \equiv \frac{1}{i\sqrt{2}} (x_n^{\kappa} + iy_n^{\kappa}). \tag{12}
$$

Next we calculate Hill's secular determinant[[11\]](#page-4-0) for the linearization of (8) about (9). As a simplification, let us define $\bar{n} \equiv n + \mu$ to be the running variable in the summations of (11). Last, to introduce physical intuition, it is convenient to define the radiation coordinate x_r and relative coordinate x_d as

$$
\mathbf{x}_r \equiv \mathbf{x}_1 + \mathbf{x}_2, \tag{13}
$$

$$
\mathbf{x}_d \equiv \mathbf{x}_1 - \mathbf{x}_2. \tag{14}
$$

It turns out, as one can easily verify, that the linearized dynamics for the δx_d variable is decoupled from the δx_r variation. Here, for lack of space, we only develop the dynamics of the δx_d variable along the $z = 0$ plane, correct tosecond order in (v/c) . We refer to [[1\]](#page-4-0) for the Coulomb eigenvalues of the δx_r dynamics. A complete study of the eigenvalues will be published elsewhere [\[9](#page-4-0)]. Before we write the planar variational equations, we need still another definition

$$
U_n^{\kappa} \equiv \frac{1}{\sqrt{2}} \xi_{n+1}^{\kappa} + \frac{i}{\sqrt{2}} \chi_{n-1}^{\kappa}
$$

$$
V_n^{\kappa} \equiv \frac{-i}{\sqrt{2}} \xi_{n+1}^{\kappa} - \frac{1}{\sqrt{2}} \chi_{n-1}^{\kappa}.
$$
 (15)

We make the quantities U_n^{κ} and V_n^{κ} into the x and y components of a bidimensional vector $\mathbf{K}_n^{\kappa} \equiv (U_n^{\kappa}, V_n^{\kappa}),$ and also define $\tilde{\mathbf{K}}_n^{\kappa} \equiv (-V_n^{\kappa}, U_n^{\kappa}), \mathbf{x}_n \equiv (x_n, y_n)$ and $\tilde{\mathbf{x}}_n \equiv (-y_n^{\kappa}, x_n^{\kappa}).$ The variational equations for planar motion along the variable δx_d can be written most simply in Floquet components as [\[9](#page-4-0)]

$$
-4\bar{n}^2 \mathbf{x}_n^d - \frac{1}{2} (\mathbf{x}_n^d + 3\mathbf{K}_n^d)
$$

$$
+ |\beta|^2 (P(\bar{n}) \mathbf{x}_n^d + Q(\bar{n}) \mathbf{K}_n^d - \frac{6i\bar{n}}{7} \tilde{\mathbf{x}}_n^d + \frac{18i\bar{n}}{7} \tilde{\mathbf{K}}_n^d) = 0. \tag{16}
$$

We recall that $\bar{n} = n + \mu$, and there is one equation for every value of n . In principle, this would lead to an infinite Hill's secular determinant[[11\]](#page-4-0). For the case of circular orbits, when we use the ξ and χ variables, we find that the ξ_n variables couple only to χ_{n+1} and vice versa. The variational problem for the coordinate $\delta \mathbf{x}_d$, as defined by (16), turns into the following 2×2 matrix equation

$$
\begin{bmatrix}\n-(\frac{1}{2} + 4\bar{n}^2) & \frac{-3i}{2} \\
\frac{3i}{2} & -(\frac{1}{2} + 4\bar{n}^2_+) \n\end{bmatrix}\n\begin{bmatrix}\n\xi_n^d \\
\chi_{n+1}^d\n\end{bmatrix} + |\beta|^2 \begin{bmatrix}\nP_2(\bar{n}) & iQ_2(\bar{n}) \\
-iQ_2(-\bar{n}_+) & P_2(-\bar{n}_+)\n\end{bmatrix}\n\begin{bmatrix}\n\xi_n^d \\
\chi_{n+1}^d\n\end{bmatrix} = 0.
$$
\n(17)

where $P_2(\bar{n}) \equiv \left(-\frac{4\bar{n}^2}{7} + \frac{6\bar{n}}{7} + \frac{3}{28}\right)$, $Q_2(\bar{n}) \equiv \left(\frac{18\bar{n}^2}{7} - \frac{18\bar{n}}{7} + \frac{1}{28}\right)$ and $\bar{n}_+ \equiv (1 + \bar{n})$. We now consider the secular equation associated with (17): At $|\beta|=0$, we obtain the eigenvalues of the Coulomb stability problem

$$
\bar{n} = 0, -1, -1/2, -1/2.
$$
 (18)

The next step is to consider the secular equation of (17) for finite values of $|\beta|$. The essential ingredient for us here is the degenerate root $\bar{n} = -1/2$ of (18). Because of this degeneracy, the double root $\bar{n} = -1/2$ acquires a correction proportional to $|\beta|$, instead of the generic second-order correction of the other nondegenerate roots. Solving the secular equation of (17), in the neighborhood of $\bar{n} = -1/2$ and in the lowest order approximation in $|\beta|$, we find

$$
\bar{n} \simeq -1/2 \pm \sqrt{\frac{51}{28}} |\beta|.
$$
 (19)

Accordingto ([11\)](#page-2-0), \bar{n} is a frequency of oscillation in units of 2ω . For example, along the linear mode of (17) defined by $\bar{n} = -1/2$, the frequencies in [\(11](#page-2-0)) are $-\omega = 2\omega\bar{n}$ and $\omega = 2\omega(1+\bar{n}).$

Furta's theorem gives the condition for the existence of an analytic constant in the neighborhood of a fixed point of an autonomous vector field[[7\]](#page-4-0). The Wannier orbit is a fixed point of the autonomous vector field describing the dynamics in the system rotating with the frequency of the Wannier orbit. To apply the theorem, we must move to this rotating system. In this rotating system, the new frequency for oscillation along the mode (19) is found by adding $1/2$ to (19). In units of 2ω , the frequency of Coulombic oscillations along the z direction in the mode δz_r δz_r δz_r is found in [[1\]](#page-4-0) to be

$$
\omega_1 \equiv \sqrt{\frac{2}{7}} \simeq 0.5345\ldots \tag{20}
$$

and for planar Coulombic oscillations along the mode $\delta \mathbf{x}_r$ $\delta \mathbf{x}_r$ $\delta \mathbf{x}_r$, in the rotating frame, is [[1\]](#page-4-0)

$$
\omega_2 \equiv \sqrt{\frac{3 + \sqrt{32}}{28}} \simeq 0.5560\ldots \tag{21}
$$

also in units of 2ω . A new resonance condition involving the root (19) is the easiest to be satisfied for small values of $|\beta|$. Therefore, we are looking for a resonance among the frequencies of (21) , (20) , and (19) , which in the rotating frame evaluates to

$$
\omega_3 \simeq \sqrt{\frac{51}{28}} |\beta| \simeq 1.3496 |\beta|,\tag{22}
$$

in units of 2ω . According to Furta's theorem [\[7](#page-4-0)], for an additional analytic constant to exist one needs a new resonance among the linear frequencies. By inspection, we find that the resonance involving the above three frequencies and with the minimal integer multipliers is of type

$$
\omega_1 - \omega_2 + 2n\omega_3 = 0,\tag{23}
$$

where n is an integer number and the factor of two multiplying ω_3 is because mode 3 comes from a degenerate eigenvalue. Condition (23) is satisfied for $|\beta|$ given by

$$
|\beta| \simeq \frac{0.007969423}{n}.\tag{24}
$$

The binding energies, in atomic units, for the above values of $|\beta|$ are

$$
E = -\frac{2.3853}{n^2} \tag{25}
$$

and the frequencies of the special "stable" Wannier orbits described by (23), in atomic units, are given by

$$
\omega = \frac{0.7442}{n^3}.\tag{26}
$$

Because we are involving a resonance among the planar oscillations $\delta \mathbf{x}_r$, $\delta \mathbf{x}_d$, and the δz_r oscillation, the leading term in the McLaurin series[[7\]](#page-4-0) of the constant is a product of δz_r times the planar coordinate variations. In reference [9] we show how to construct this extra constant and discuss the shape of its level surfaces.

The frequency ω_z for oscillations along the δz_r linear mode can be obtained, as prescribed by([20\)](#page-3-0), by multiplying the above frequency by $2\sqrt{2/7}$,

$$
\omega_z = \frac{0.7956}{n^3}.
$$
\n(27)

The inclusion of the third-order radiative terms makes this linear mode of [\(20](#page-3-0)) a stable mode [9], whereby the perturbation radiates energy and decays to the perfectly circular orbit. The frequency of this stable oscillation should then appear in the spectrum of sharp lines, given that the dynamics selects the neighborhood of the stable orbits. For example, the transition $2^1P \rightarrow 1^1S$ of parahelium corresponds to a frequency of $2.9037 - 2.1237 =$ 0.7799 atomic units [5,12], and (27) with $n = 1$ evaluates to 0.7956, which is a 2% difference. The transition $3^1P \rightarrow 2^1S$ in parahelium [12] has a frequency of $2.1459 - 2.0551 = 0.0908$ atomic units, and (27) with $n = 2$ evaluates to 0.0995, which is an 8% difference. The asymptotic form of the quantum mechanical energies of helium are [5]

$$
E = -\frac{Z^2}{2} - \frac{(Z-1)^2}{2(n_r + L + 1)^2},\tag{28}
$$

with $Z = 2$. For transitions with $\Delta L = 1$, the frequencies predicted from the above formula are given by

$$
\omega_q = \frac{1}{(n_r + L + 1)^3},\tag{29}
$$

which agrees with (27) to within 20%. We stress that we relied only on dynamics to calculate the frequency of (27). Equation (29), on the other hand, was obtained in the usual way of taking differences between quantum energies, and was included only for comparison.

A more detailed comparison with the spectrum of helium requires the consideration of elliptical orbits and zero-angular-momentum orbits [9]. In reference [9] we include the third-order terms and find that the resulting damping of the oscillation mode of (27) agrees well with the quantum calculations of the linewidth.

The extra complex constant is actually two real functions of phase space [7,9]. Together with the four global constants of helium, there are six functions that can produce a barrier in phase space, at least in the small region inside the resonance islands [14]. Projected on the plane of the stable and unstable Wannier directions, the oneperiod return map of the full dynamics has a hyperbolic fixed point. The resonance islands are possibly found inside the homoclinic tangle of the invariant manifolds, but this remains to be investigated in detail.

Historically, the Wannier instability, first found by Nicholson [10], led Bohr to postulate that, along a quantized set of orbits, some special mechanism would supersede the Wannier instability, thus stabilizing the atom, which proved to be a very fruitful intuition [13]. Some of the details of how the six constants stabilize the dynamics, in a time scale of order $T/(v/c)^2$, are discussed elsewhere [9].

The existence of this extra constant is a genuine effect of electrodynamics with retardation. It is surprising that classical electrodynamics with retardation can offer such a sensible approximation to the spectrum of sharp lines. We do not know of any prior similar application of dynamics to atomic physics.

I acknowledge discussions with R. Napolitano and J. C. Egues. This work was supported by Fapesp.

- [1] M. Poirier, *Phys. Rev. A* **40**, 3498 (1989).
- [2] J. Guckenheimer and P. Holmes, Nonlinear Oscillations, Dynamical Systems and Bifurcations of Vector Fields, Springer, NY (1983).
- [3] L. Page, Phys. Rev. 11, 376 (1918), see also G. A. Schott, Electromagnetic Radiation, Cambridge (1912).
- [4] J. D. Jackson, *Classical Electrodynamics*, Wiley, New York (1975).
- [5] H. A. Bethe and E. E. Salpeter, Quantum Mechanics of One- and Two-electron atoms, Plenum, NY (1977).
- [6] V. I. Arnold, Mathematical Methods of Classical Mechanics, Springer Verlag, New York (1978).
- [7] S. D. Furta, Z. angew. Math. Phys., 47, 112 (1996).
- [8] S. Coleman in Electromagnetism, Paths to Research, edited by D. Teplitz, Plenum, NY (1982).
- [9] J. De Luca (to be published).
- [10] J. W. Nicholson, *Monthly Notices of the Royal Astronom*ical Society LXXII, p.49, 139, 677, 693 and 729 (1912).
- [11] D. W. Jordan and P. Smith, Nonlinear Ordinary Differential equations, Claredon, Oxford (1977).
- [12] Ming-Keh Chen, J. Phys. B **27**, 4847 (1994).
- [13] N. Bohr, *Philosophical Magazine*, **26**, 1 (1913) and *Philo*sophical Magazine, 26, 476 (1913).
- [14] A. J. Lichtenberg and M. A. Lieberman, Regular and Chaotic Dynamics, second edition, Springer (1992).