# arXiv:chao-dyn/9811024v1 25 Nov 1998

# Continued Fraction Representation of Temporal Multi Scaling in Turbulence

David Daems<sup>\*,\*\*</sup>, Siegfried Grossmann<sup>†</sup>, Victor S. L'vov<sup>\*</sup> and Itamar Procaccia<sup>\*</sup>

\* Department of Chemical Physics, The Weizmann Institute of Science, Rehovot 76100, Israel

\*\* Center for Nonlinear Phenomena and Complex Systems, Université Libre de Bruxelles, 1050 Brussels, Belgium

<sup>†</sup>Fachbereich Physik, Philipps Universitaet Marburg, Renthof 5, Marburg D-35032, Germany

It was shown recently that the anomalous scaling of simultaneous correlation functions in turbulence is intimately related to the breaking of temporal scale invariance, which is equivalent to the appearance of infinitely many times scales in the time dependence of time-correlation functions. In this paper we derive a continued fraction representation of turbulent time correlation functions which is exact and in which the multiplicity of time scales is explicit. We demonstrate that this form yields precisely the same scaling laws for time derivatives and time integrals as the "multi-fractal" representation that was used before. Truncating the continued fraction representation yields the "best" estimates of time correlation functions if the given information is limited to the scaling exponents of the simultaneous correlation functions up to a certain, finite order. It is not that the derivation of a continued fraction representation obtained here for an operator which is not Hermitian or anti-Hermitian may be of independent interest.

### I. INTRODUCTION

It is commonly argued [1,2] that fully developed hydrodynamic turbulence exhibits simultaneous statistical objects whose scaling properties are anomalous. For example the so called structure functions satisfy scaling laws of the form

$$S_n(R) = \langle |\boldsymbol{u}(\boldsymbol{r} + \boldsymbol{R}, t) - \boldsymbol{u}(\boldsymbol{r}, t)|^n \rangle \sim R^{\zeta_n} , \qquad (1)$$

where  $\boldsymbol{u}(\boldsymbol{r},t)$  is the Eulerian velocity field, and  $\zeta_n$  are scaling exponents which are nonlinear functions of n. The nonlinear dependence is referred to as "anomalous scaling" or "multi-scaling", and the issue of evaluating these exponents from either phenomenological models or from first principles has attracted significant amount of effort in the last decade.

It has only recently been discovered [3] that also the time dependence of the *n*-th order correlation functions is multi-scaling, and that "dynamical scaling" is broken. This phenomenon seems to distinguish turbulence from other problems in which scaling is anomalous, like critical phenomena. In the latter case dynamical scaling is invoked by stating that a space time correlation function F(R, t) is a homogeneous function of its arguments in the sense that  $F(\lambda R, \lambda^z t) = \lambda^{\zeta} F(R, t)$ , where  $\zeta$  and z are the "static" and "dynamic" scaling exponents respectively. In turbulence such relations do not exist even when the same-time correlation functions are homogeneous functions of the spatial coordinates. The importance of this fact in determining the structure of the theory has been stressed in [4], and see also [5].

In this paper we address temporal multi-scaling on the basis of the continued fraction representation of turbulent correlation function [6,7]. This approach will lead us to a different point of view of temporal multi-scaling, in agreement with the conclusions as Ref. [3]. The advantage of the present formulation is three-fold: first, it is derived on the basis of an exact formulation of the time correlation functions and their dynamics. The phenomenon of temporal multi-scaling is related in this approach to the scaling properties of higher order temporal derivatives of correlation functions, computed at zero time. Second, this approach furnishes information not only about leading scaling exponents, but also about sub-leading ones. Third, a finite truncation of the continued fraction representation is in a sense the "best" possible representation when the information about the scaling of one-time correlation functions is limited to the low order scaling exponents. We will show that the scaling laws exhibited by the continued fraction representation are identical to those predicted by the "multi-fractal" representation of time-correlation function [4], adding weight and justification to the latter. Since the multi-fractal representation was used recently to estimate the scaling exponents from first principles [5], we ascribe some weight to being able to justify it further.

To keep the formalism minimal and the result clearest, we treat in this paper only the second order space-time correlation function of turbulent fields. The formalism can be used to generate representations of any higher order correlation function, but we do not elaborate on this in the present text. In Section II we review briefly the Zwanzig-Mori formalism [8,9] which was applied to time correlation functions in turbulence [6,7], and display the continued fraction representation of the second order time correlation function of the velocity field. We show that the coefficients in this representation can be written in terms of time derivatives at zero time of the same second order correlation function. In Section III we set up a theory for the evaluation of the scaling exponents of these coefficients. In Section IV we derive the scaling laws implied by the continued fraction representation for the time derivatives of the correlation function evaluated at time zero. In Section V we relate our results to the multi-fractal representation of the correlations functions, and explain the scaling-equivalence of the two representations. In Section VI we offer a summary and a short discussion.

### II. CONTINUED FRACTION REPRESENTATION

In thinking about dynamics one cannot deal with timecorrelation functions of the Eulerian field since these are dominated by the kinematic sweeping time scale. We need to consider Lagrangian or Belinicher-L'vov velocity differences. We prefer the latter since they obey Navier-Stokes like equations of motion which are local in time. In terms of the Eulerian velocity  $\boldsymbol{u}(\boldsymbol{r},t)$  Belinicher and L'vov defined [10] the field  $\boldsymbol{v}(\boldsymbol{r}_0,t_0|\boldsymbol{r},t)$  as

$$\boldsymbol{v}(\boldsymbol{r}_0, t_0 | \boldsymbol{r}, t) \equiv \boldsymbol{u}[\boldsymbol{r} + \boldsymbol{\rho}(\boldsymbol{r}_0, t), t]$$
(2)

where

$$\boldsymbol{\rho}(\boldsymbol{r}_0, t) = \int_{t_0}^t ds \boldsymbol{u}[\boldsymbol{r}_0 + \boldsymbol{\rho}(\boldsymbol{r}_0, s), s] .$$
 (3)

The observation of Belinicher and L'vov [10] was that the variables  $\mathcal{W}(\mathbf{r}_0, t_0 | \mathbf{r}, \mathbf{r}, t)$  defined as

$$\mathcal{W}(\boldsymbol{r}_0, t_0 | \boldsymbol{r}, \boldsymbol{r}', t) \equiv \boldsymbol{v}(\boldsymbol{r}_0, t_0 | \boldsymbol{r}, t) - \boldsymbol{v}(\boldsymbol{r}_0, t_0 | \boldsymbol{r}', t) , \quad (4)$$

exactly satisfy a Navier-Stokes like equation in the incompressible limit:

$$\left[\frac{\partial}{\partial t} + \overleftrightarrow{\mathcal{P}}\mathcal{W}(\boldsymbol{r}_0, t_0 | \boldsymbol{r}, \boldsymbol{r}_0, t) \cdot \boldsymbol{\nabla}_r + \overleftrightarrow{\mathcal{P}}'\mathcal{W}(\boldsymbol{r}_0, t_0 | \boldsymbol{r}', \boldsymbol{r}_0, t) \cdot \boldsymbol{\nabla}_r' - \nu(\boldsymbol{\nabla}_r^2 + \boldsymbol{\nabla}_r'^2)\right] \mathcal{W}(\boldsymbol{r}_0, t_0 | \boldsymbol{r}, \boldsymbol{r}', t) = 0.$$
(5)

We remind the reader that the application of the transversal projector  $\stackrel{\leftrightarrow}{\mathcal{P}}$  to any given vector field a(r) is non local, and has the form:

$$[\overset{\leftrightarrow}{\mathcal{P}}\boldsymbol{a}(\boldsymbol{r})]^{\alpha} = \int d\tilde{\boldsymbol{r}} \mathcal{P}^{\alpha\beta}(\boldsymbol{r} - \tilde{\boldsymbol{r}}) a^{\beta}(\tilde{\boldsymbol{r}}).$$
(6)

The explicit form of the kernel  $\mathcal{P}^{\alpha\beta}(\mathbf{r})$  can be found, for example, in [11]. In (5)  $\overleftrightarrow{\mathcal{P}}$  and  $\overleftrightarrow{\mathcal{P}}'$  are projection operators which act on fields that depend on the corresponding coordinates  $\mathbf{r}$  and  $\mathbf{r}'$ . For our purpose below we introduce the Liouville operator  $\hat{\mathcal{L}}(\mathbf{r}_0, t_0 | \mathbf{r}, \mathbf{r}')$  which is defined via the total time derivative of  $\mathcal{W}(\mathbf{r}_0, t_0 | \mathbf{r}, \mathbf{r}', t)$  at time  $t = t_0$ :

$$\frac{d\mathcal{W}(\boldsymbol{r}_{0},t_{0}|\boldsymbol{r},\boldsymbol{r}',t)}{dt}\bigg|_{t=t_{0}} \equiv \hat{\mathcal{L}}(\boldsymbol{r}_{0},t_{0}|\boldsymbol{r},\boldsymbol{r}')\mathcal{W}(\boldsymbol{r}_{0},t_{0}|\boldsymbol{r},\boldsymbol{r}',t_{0}).$$
(7)

The identity of the Liouville operator follows by definition,

$$\frac{d\boldsymbol{v}(\boldsymbol{r}_{0}, t_{0} | \boldsymbol{r} - \boldsymbol{\rho}(\boldsymbol{r}_{0}, t), t], t)}{dt} \equiv \frac{d\boldsymbol{u}(\boldsymbol{r}, t)}{dt}$$
$$= \frac{\partial \boldsymbol{u}(\boldsymbol{r}, t)}{\partial t} + [\boldsymbol{u}(\boldsymbol{r}, t) \cdot \boldsymbol{\nabla}] \boldsymbol{u}(\boldsymbol{r}, t)$$
(8)

Translating all coordinates by  $\rho(\mathbf{r}_0, t)$  we find after elementary algebra,

$$\hat{\mathcal{L}}(\boldsymbol{r}_{0}, t_{0} | \boldsymbol{r}, \boldsymbol{r}') = \frac{\partial}{\partial t} \bigg|_{t=t_{0}} + \boldsymbol{v}(\boldsymbol{r}_{0}, t_{0} | \boldsymbol{r}, t_{0}) \cdot \boldsymbol{\nabla}_{r} + \boldsymbol{v}(\boldsymbol{r}_{0}, t_{0} | \boldsymbol{r}', t_{0}) \cdot \boldsymbol{\nabla}_{r'} .$$
(9)

Consider now the time dependence of "fully fused" 2nd order correlation function

$$\mathcal{F}_{2}^{\alpha\beta}(\boldsymbol{r}_{0}|\boldsymbol{r},\boldsymbol{r}',\tau) = \left\langle \mathcal{W}^{\alpha}(\boldsymbol{r}_{0}|\boldsymbol{r},\boldsymbol{r}',t_{0})\mathcal{W}^{\beta}(\boldsymbol{r}_{0}|\boldsymbol{r},\boldsymbol{r}',t_{0}+\tau) \right\rangle.$$
(10)

By "fully fused" we mean here that the coordinates of the two velocity differences are the same, and they differ only in their time argument. The same-time counterpart of this correlation function, i.e.  $\mathcal{F}_2(\mathbf{r}_0|\mathbf{r},\mathbf{r}',\tau=0)$  is independent of  $\mathbf{r}_0$  and in an isotropic and homogeneous ensemble it is a function of  $|\mathbf{r}-\mathbf{r}'|$  only. Accordingly it differs from the standard structure function  $S_2(|\mathbf{r}-\mathbf{r}'|)$ only in having the full 2nd rank tensorial character. For this analysis we choose all the three vector distances to have the modulus in the inertial range, of the order of R. In the notation we will keep only this R but remember that the angular dependence is not shown explicitly.

To proceed, we consider the tensorial correlation function (10) as an inner product in the space of vectors  $\mathcal{W}$ , denoted as

$$\mathcal{F}_2(R,\tau) = \left( \mathcal{W}, e^{\hat{\mathcal{L}}\tau} \mathcal{W} \right)$$
 (11)

In particular we are interested in the Laplace transform of Eq. (11)

$$\tilde{\mathcal{F}}_2(R,z) = (\mathcal{W}, \frac{1}{z - \hat{\mathcal{L}}}\mathcal{W})$$
 (12)

It has been shown by Grossmann and Thomae [6] that the Zwanzig-Mori projection operator formalism [8,9] applies to turbulent systems described by Navier-Stokes like equations. In [7] it has been demonstrated that the contribution of the memory kernel and higher order continued fraction is considerable. The central idea is to decompose the resolvent

$$\hat{\mathcal{R}}(z) = \frac{1}{z - \hat{\mathcal{L}}} \tag{13}$$

by means of a projection operator  $\hat{P}$ . With  $\hat{Q} = \hat{1} - \hat{P}$  as projector orthogonal to  $\hat{P}$  one has the resolvent identity

$$\hat{P}\hat{\mathcal{R}}(z)\hat{P} = \hat{P}\frac{1}{z - \hat{P}\hat{\mathcal{L}}\hat{P} - \hat{P}\hat{\mathcal{L}}\hat{Q}\frac{1}{z - \hat{Q}\hat{\mathcal{L}}\hat{Q}}\hat{Q}\hat{\mathcal{L}}\hat{P}} .$$
 (14)

Since we note that the time correlation function of interest is the  $\mathcal{W} - \mathcal{W}$  matrix element of the resolvent, it is useful to choose  $\hat{P}$  to be the projector on  $\mathcal{W}$ ,

$$\hat{P} \equiv (\boldsymbol{\mathcal{W}}, .)(\boldsymbol{\mathcal{W}}, \boldsymbol{\mathcal{W}})^{-1}\boldsymbol{\mathcal{W}} .$$
(15)

Its basic properties are  $\hat{P}\hat{P} = \hat{P}$  (idempotent) and  $\hat{P}^{\dagger} = \hat{P}$  (self adjoint), characterizing an orthogonal projection. From Eq.(14) we have an expression for  $\hat{P}$ :

$$\hat{P} = \hat{P}\hat{\mathcal{R}}(z)\hat{P}\left[z - \hat{P}\hat{\mathcal{L}}\hat{P} - \hat{P}\hat{\mathcal{L}}\hat{Q}\frac{1}{z - \hat{Q}\hat{\mathcal{L}}\hat{Q}}\hat{Q}\hat{\mathcal{L}}\hat{P}\right]$$
(16)

Compute now the  $\mathcal{W} - \mathcal{W}$  matrix element, use  $\hat{P}\mathcal{W} = \mathcal{W}$ , as well as the definition of  $\hat{P}$  to compute

$$(\boldsymbol{\mathcal{W}}, \hat{\mathcal{R}}(z)\boldsymbol{\mathcal{W}})(\boldsymbol{\mathcal{W}}, \boldsymbol{\mathcal{W}})^{-1}[z(\boldsymbol{\mathcal{W}}, \boldsymbol{\mathcal{W}})$$
(17)  
$$-(\boldsymbol{\mathcal{W}}, \hat{\mathcal{L}}\boldsymbol{\mathcal{W}}) - (\boldsymbol{\mathcal{W}}, \hat{\mathcal{L}}\hat{Q}\frac{1}{z - \hat{Q}\hat{\mathcal{L}}\hat{Q}}\hat{Q}\hat{\mathcal{L}}\boldsymbol{\mathcal{W}})] = (\boldsymbol{\mathcal{W}}, \boldsymbol{\mathcal{W}}) .$$

We therefore conclude with

$$\tilde{\boldsymbol{\mathcal{F}}}_2(\boldsymbol{R}, \boldsymbol{z}) = \frac{\boldsymbol{k}_0(\boldsymbol{R})}{\boldsymbol{z} - \boldsymbol{\gamma}_0(\boldsymbol{R}) - \tilde{\boldsymbol{K}}_0(\boldsymbol{R}, \boldsymbol{z})} , \qquad (18)$$

where

$$\begin{aligned} \boldsymbol{k}_0(R) &= (\boldsymbol{\mathcal{W}}, \boldsymbol{\mathcal{W}}), \\ \boldsymbol{\gamma}_0(R) &= (\boldsymbol{\mathcal{W}}, \hat{\mathcal{L}} \boldsymbol{\mathcal{W}}) (\boldsymbol{\mathcal{W}}, \boldsymbol{\mathcal{W}})^{-1}, \\ \tilde{\boldsymbol{K}}_0(R, z) &= \left( \hat{Q}^{\dagger} \hat{\mathcal{L}}^{\dagger} \boldsymbol{\mathcal{W}}, \frac{1}{z - \hat{Q} \hat{\mathcal{L}} \hat{Q}} \hat{Q} \hat{\mathcal{L}} \boldsymbol{\mathcal{W}} \right) (\boldsymbol{\mathcal{W}}, \boldsymbol{\mathcal{W}})^{-1}. \end{aligned}$$
(19)

Here, of course,  $\hat{Q}^{\dagger} = \hat{Q}$ , but in the next steps of generating the continued fraction hermiticity will not hold. Realizing that the kernel  $\tilde{K}_0(R, z)$  has the same resolvent structure as  $\tilde{\mathcal{F}}_2(R, z)$  except that it now features  $\hat{Q}\hat{\mathcal{L}}\hat{Q}$ instead of  $\hat{\mathcal{L}}$  and the states are  $\hat{Q}^{\dagger}\hat{\mathcal{L}}^{\dagger}\mathcal{W}$ ,  $\hat{Q}\hat{\mathcal{L}}\mathcal{W}$  instead of  $\mathcal{W}$ ,  $\mathcal{W}$ , with  $\hat{\mathcal{L}}^{\dagger}$  the adjoint of  $\hat{\mathcal{L}}$  - one can continue the fraction by the same procedure. This is more transparent if we denote

$$\mathcal{W}_1 = \hat{Q}\hat{\mathcal{L}}\mathcal{W}, \quad \tilde{\mathcal{W}}_1 = \hat{Q}^{\dagger}\hat{\mathcal{L}}^{\dagger}\mathcal{W}, \quad \hat{\mathcal{L}}_1 = \hat{Q}\hat{\mathcal{L}}\hat{Q} \quad (20)$$

so that  $\tilde{\boldsymbol{K}}_0(\boldsymbol{R},z)$  takes on the form

$$\tilde{\boldsymbol{K}}_0(\boldsymbol{R}, \boldsymbol{z}) = \left(\tilde{\boldsymbol{\mathcal{W}}}_1, \frac{1}{\boldsymbol{z} - \hat{\mathcal{L}}_1} \boldsymbol{\mathcal{W}}_1\right) (\boldsymbol{\mathcal{W}}, \boldsymbol{\mathcal{W}})^{-1} .$$
(21)

Now we define a new projection operator

$$\hat{P}_1 \cdot \equiv (\tilde{\boldsymbol{\mathcal{W}}}_1, \cdot) (\tilde{\boldsymbol{\mathcal{W}}}_1, \boldsymbol{\mathcal{W}}_1)^{-1} \boldsymbol{\mathcal{W}}_1 .$$
(22)

As a result of  $\tilde{\mathcal{W}}_1$  being different from  $\mathcal{W}_1$  when  $\hat{\mathcal{L}}$  is not Hermitian or anti-Hermitian, this operator  $\hat{P}_1$  is not

Hermitian and performs accordingly non-orthogonal projections. But  $\hat{P}_1$  still is idempotent,  $\hat{P}_1\hat{P}_1 = \hat{P}_1$ , which is the essential property for deriving the analogous resolvent identity with  $\hat{\mathcal{L}}_1$  in (21) as for the original resolvent (12) with  $\hat{\mathcal{L}}$ . Defining  $\hat{Q}_1 \equiv 1 - \hat{P}_1$  we can repeat the argument leading to (18) and (19), and find

$$\tilde{\boldsymbol{K}}_0(\boldsymbol{R}, \boldsymbol{z}) = \frac{\boldsymbol{k}_1(\boldsymbol{R})}{\boldsymbol{z} - \boldsymbol{\gamma}_1(\boldsymbol{R}) - \tilde{\boldsymbol{K}}_1(\boldsymbol{R}, \boldsymbol{z})} , \qquad (23)$$

where

$$\boldsymbol{k}_{1}(R) = (\tilde{\boldsymbol{\mathcal{W}}}_{1}, \boldsymbol{\mathcal{W}}_{1})(\boldsymbol{\mathcal{W}}, \boldsymbol{\mathcal{W}})^{-1}, \qquad (24)$$
  
$$\boldsymbol{\gamma}_{1}(R) = (\tilde{\boldsymbol{\mathcal{W}}}_{1}, \hat{\mathcal{L}}_{1} \boldsymbol{\mathcal{W}}_{1})(\tilde{\boldsymbol{\mathcal{W}}}_{1}, \boldsymbol{\mathcal{W}}_{1})^{-1}, \qquad (24)$$
  
$$\tilde{\boldsymbol{K}}_{1}(R, z) = \left(\hat{Q}_{1}^{\dagger} \hat{\mathcal{L}}_{1}^{\dagger} \tilde{\boldsymbol{\mathcal{W}}}_{1}, \frac{1}{z - \hat{Q}_{1} \hat{\mathcal{L}}_{1} \hat{Q}_{1}} \hat{Q}_{1} \hat{\mathcal{L}}_{1} \boldsymbol{\mathcal{W}}_{1}\right) \times (\tilde{\boldsymbol{\mathcal{W}}}_{1}, \boldsymbol{\mathcal{W}}_{1})^{-1}.$$

Hence starting from

$$\tilde{\boldsymbol{K}}_{n}(\boldsymbol{R},\boldsymbol{z}) = \left(\tilde{\boldsymbol{\mathcal{W}}}_{n+1}, \frac{1}{\boldsymbol{z} - \hat{\mathcal{L}}_{n+1}} \boldsymbol{\mathcal{W}}_{n+1}\right) (\tilde{\boldsymbol{\mathcal{W}}}_{n}, \boldsymbol{\mathcal{W}}_{n})^{-1} ,$$
(25)

with  $n \ge 0$  one arrives at

$$\tilde{K}_{n}(R,z) = \frac{k_{n+1}(R)}{z - \gamma_{n+1}(R) - \tilde{K}_{n+1}(R,z)}, \quad (26)$$

where

$$\boldsymbol{k}_{n+1}(R) = (\tilde{\boldsymbol{\mathcal{W}}}_{n+1}, \boldsymbol{\mathcal{W}}_{n+1})(\tilde{\boldsymbol{\mathcal{W}}}_n, \boldsymbol{\mathcal{W}}_n)^{-1} ,$$
  
$$\boldsymbol{\gamma}_{n+1}(R) = (\tilde{\boldsymbol{\mathcal{W}}}_{n+1}, \hat{\mathcal{L}}_{n+1} \boldsymbol{\mathcal{W}}_{n+1})(\tilde{\boldsymbol{\mathcal{W}}}_{n+1}, \boldsymbol{\mathcal{W}}_{n+1})^{-1} .$$
(27)

Here we used the notation

$$\mathcal{W}_{n+1} = \hat{Q}_n \hat{\mathcal{L}}_n \mathcal{W}_n, \quad \tilde{\mathcal{W}}_{n+1} = \hat{Q}_n^{\dagger} \hat{\mathcal{L}}_n^{\dagger} \tilde{\mathcal{W}}_n, \\ \hat{\mathcal{L}}_{n+1} = \hat{Q}_n \hat{\mathcal{L}}_n \hat{Q}_n, \quad \mathcal{W}_0 = \tilde{\mathcal{W}}_0 = \mathcal{W}, \quad \hat{\mathcal{L}}_0 = \hat{\mathcal{L}}, \quad (28)$$

and defined new projection operators as

$$\hat{P}_{n} \cdot = (\tilde{\mathcal{W}}_{n}, \cdot)(\tilde{\mathcal{W}}_{n}, \mathcal{W}_{n})^{-1}\mathcal{W}_{n}$$
$$\hat{Q}_{n} = \hat{1} - \hat{P}_{n} .$$
(29)

From (18) and (26) it thus follows that the Laplace transform  $\tilde{\mathcal{F}}_2(R, z)$  of the correlation function (10) can be written in continued fraction representation:

$$\tilde{\mathcal{F}}_{2}(R,z) = \frac{\mathbf{k}_{0}(R)}{z - \gamma_{0}(R) - \frac{\mathbf{k}_{1}(R)}{z - \gamma_{1}(R) - \frac{\mathbf{k}_{2}(R)}{z - \gamma_{2}(R) - \ddots}}$$
(30)

The novelty when the operator  $\hat{\mathcal{L}}$  is not Hermitian or anti-Hermitian as is the case here is that the new projection operators introduced to continue the fraction perform non-orthogonal projections. Only the lowest order iterates of  $\hat{P}_n$  and  $\hat{Q}_n$ , i.e. the operators  $\hat{P}$  and  $\hat{Q}$ , perform orthogonal projections. In the following section we analyze the scaling properties of this representation.

## III. SCALING PROPERTIES OF THE CONTINUED FRACTION REPRESENTATION

In this section we determine the *leading* scaling exponents of the coefficients which appear in the continued fraction representation (30). This leading scaling behavior is given in terms of correlation functions of time derivatives of the velocity differences  $\mathcal{W}$  computed at zero time:

$$\boldsymbol{k}_{n}(R) \approx \langle \hat{\mathcal{L}}^{n} \boldsymbol{\mathcal{W}} \hat{\mathcal{L}}^{n} \boldsymbol{\mathcal{W}} \rangle / \langle \hat{\mathcal{L}}^{n-1} \boldsymbol{\mathcal{W}} \hat{\mathcal{L}}^{n-1} \boldsymbol{\mathcal{W}} \rangle, \qquad (31)$$
$$\boldsymbol{\gamma}_{n}(R) \approx \langle \hat{\mathcal{L}}^{n+1} \boldsymbol{\mathcal{W}} \hat{\mathcal{L}}^{n} \boldsymbol{\mathcal{W}} \rangle / \langle \hat{\mathcal{L}}^{n} \boldsymbol{\mathcal{W}} \hat{\mathcal{L}}^{n} \boldsymbol{\mathcal{W}} \rangle .$$

Here the symbol  $\approx$  means "leading scaling order". Eq.(31) yields the following explicit scaling

The rest of this section is a demonstration of this result. The reader who prefers to see the connection to the multi-fractal representation can go directly to the next section.

Let us define

$$\hat{J}_n \equiv \hat{Q}_0 \dots \hat{Q}_{n-1} \hat{Q}_n \hat{Q}_{n-1} \dots \hat{Q}_0 = \hat{Q}_n .$$
 (33)

Now

$$\hat{P}_i \hat{P}_j = \delta_{ij} \hat{P}_i , \qquad i, j \ge 0 \tag{34}$$

implies that

$$\hat{J}_n = \hat{Q}_0 \hat{Q}_1 \dots \hat{Q}_n = \hat{Q}_n \hat{Q}_{n-1} \dots \hat{Q}_0 = \hat{Q}_n$$
  
= 1 -  $\hat{P}_0 - \dots - \hat{P}_n$ . (35)

Hence Eq. (28) leads to

$$\boldsymbol{\mathcal{W}}_{n+1} = \hat{J}_n \hat{\mathcal{L}} \hat{J}_{n-1} \hat{\mathcal{L}} \hat{J}_{n-2} \dots \hat{\mathcal{L}} \hat{J}_0 \hat{\mathcal{L}} \boldsymbol{\mathcal{W}}, 
\tilde{\boldsymbol{\mathcal{W}}}_{n+1} = \hat{J}_n^{\dagger} \hat{\mathcal{L}}^{\dagger} \hat{J}_{n-1}^{\dagger} \hat{\mathcal{L}}^{\dagger} \hat{J}_{n-2}^{\dagger} \dots \hat{\mathcal{L}}^{\dagger} \hat{J}_0^{\dagger} \hat{\mathcal{L}}^{\dagger} \tilde{\boldsymbol{\mathcal{W}}}, 
\hat{\mathcal{L}}_{n+1} \boldsymbol{\mathcal{W}}_{n+1} = \hat{J}_n \hat{\mathcal{L}} \hat{J}_n \hat{\mathcal{L}} \hat{J}_{n-1} \dots \hat{\mathcal{L}} \hat{J}_0 \hat{\mathcal{L}} \boldsymbol{\mathcal{W}}.$$
(36)

It follows that

$$(\tilde{\boldsymbol{\mathcal{W}}}_{n+1}, \boldsymbol{\mathcal{W}}_{n+1}) = (\boldsymbol{\mathcal{W}}, \hat{\mathcal{L}}\hat{J}_{0}\hat{\mathcal{L}}\hat{J}_{1}\dots\hat{\mathcal{L}}\hat{J}_{n-1}\hat{\mathcal{L}}\hat{J}_{n}\hat{\mathcal{L}}\hat{J}_{n-1}\dots\hat{\mathcal{L}}\hat{J}_{0}\hat{\mathcal{L}}\boldsymbol{\mathcal{W}}), (\tilde{\boldsymbol{\mathcal{W}}}_{n+1}, \hat{\mathcal{L}}_{n+1}\boldsymbol{\mathcal{W}}_{n+1}) = (\boldsymbol{\mathcal{W}}, \hat{\mathcal{L}}\hat{J}_{0}\hat{\mathcal{L}}\hat{J}_{1}\dots\hat{\mathcal{L}}\hat{J}_{n}\hat{\mathcal{L}}\hat{J}_{n}\hat{\mathcal{L}}\hat{J}_{n-1}\dots\hat{\mathcal{L}}\hat{J}_{0}\hat{\mathcal{L}}\boldsymbol{\mathcal{W}}).$$
 (37)

In order to arrive at Eq. (31) we have to show that

$$(\tilde{\mathcal{W}}_{n+1}, \mathcal{W}_{n+1}) \approx (\mathcal{W}, \hat{\mathcal{L}}^{2n+2} \mathcal{W})$$
  
$$(\tilde{\mathcal{W}}_{n+1}, \hat{\mathcal{L}}_{n+1} \mathcal{W}_{n+1}) \approx (\mathcal{W}, \hat{\mathcal{L}}^{2n+3} \mathcal{W}) .$$
(38)

It suffices to use the following assertion which we prove below

$$(\mathcal{W}, \hat{\mathcal{L}}^k \hat{J}_q \hat{I}_q \mathcal{W}) \approx (\mathcal{W}, \hat{\mathcal{L}}^k \hat{I}_q \mathcal{W}), \qquad k > q \ge 0, \quad (39)$$

where  $\hat{I}_q$  is any composition of  $\hat{\mathcal{L}}$  and  $\hat{J}_i$  which has two properties. First, it contains at least q + 1 times the operator  $\hat{\mathcal{L}}$  which we indicate by the subscript q in  $\hat{I}_q$  and second, on the right hand of each  $\hat{J}_i$  there are at least i + 1 operators  $\hat{\mathcal{L}}$  in the composition. Since the compositions of  $\hat{\mathcal{L}}$  and  $\hat{J}_i$  appearing in (37) satisfy these two properties, (38) results straightforwardly.

We now proceed to the proof of (39) by induction. Assuming for p < q < k that

$$(\boldsymbol{\mathcal{W}}, \hat{\mathcal{L}}^k \hat{J}_p \hat{I}_p \boldsymbol{\mathcal{W}}) \approx (\boldsymbol{\mathcal{W}}, \hat{\mathcal{L}}^k \hat{I}_p \boldsymbol{\mathcal{W}}) ,$$
 (40)

we show that

$$(\boldsymbol{\mathcal{W}}, \hat{\mathcal{L}}^k \hat{J}_q \hat{I}_q \boldsymbol{\mathcal{W}}) \approx (\boldsymbol{\mathcal{W}}, \hat{\mathcal{L}}^k \hat{I}_q \boldsymbol{\mathcal{W}}) .$$
 (41)

Using Eq. (35) the LHS of Eq. (41) becomes

$$(\boldsymbol{\mathcal{W}}, \hat{\mathcal{L}}^k \hat{J}_q \hat{I}_q \boldsymbol{\mathcal{W}}) = (\boldsymbol{\mathcal{W}}, \hat{\mathcal{L}}^k \hat{J}_{q-1} \hat{I}_q \boldsymbol{\mathcal{W}}) - (\boldsymbol{\mathcal{W}}, \hat{\mathcal{L}}^k \hat{P}_q \hat{I}_q \boldsymbol{\mathcal{W}}) .$$

$$(42)$$

By Eq. (40) the first term on the RHS scales as

$$(\mathcal{W}, \hat{\mathcal{L}}^k \hat{J}_{q-1} \hat{I}_q \mathcal{W}) \approx (\mathcal{W}, \hat{\mathcal{L}}^k \hat{I}_q \mathcal{W}) .$$
 (43)

As for the second term, without specifying  $I_q$  we notice using Eq. (29) for  $\hat{P}_q$  that it is the product of the following three factors that feature only operators  $\hat{J}_p$  with p < q and with more than p times the operator  $\hat{\mathcal{L}}$  on their right hand so that by (40) one has

$$(\boldsymbol{\mathcal{W}}, \hat{\mathcal{L}}^{k} \boldsymbol{\mathcal{W}}_{q}) = (\boldsymbol{\mathcal{W}}, \hat{\mathcal{L}}^{k} \hat{J}_{q-1} \hat{\mathcal{L}} \hat{J}_{q-2} \dots \hat{\mathcal{L}} \hat{J}_{0} \hat{\mathcal{L}} \boldsymbol{\mathcal{W}})$$

$$\approx (\boldsymbol{\mathcal{W}}, \hat{\mathcal{L}}^{k+q} \boldsymbol{\mathcal{W}}),$$

$$(\tilde{\boldsymbol{\mathcal{W}}}_{q}, \boldsymbol{\mathcal{W}}_{q}) = (\boldsymbol{\mathcal{W}}, \hat{\mathcal{L}} \hat{J}_{0} \hat{\mathcal{L}} \hat{J}_{1} \dots \hat{\mathcal{L}} \hat{J}_{q-1} \dots \hat{\mathcal{L}} \hat{J}_{0} \hat{\mathcal{L}} \boldsymbol{\mathcal{W}})$$

$$\approx (\boldsymbol{\mathcal{W}}, \hat{\mathcal{L}}^{2q} \boldsymbol{\mathcal{W}}),$$

$$(\tilde{\boldsymbol{\mathcal{W}}}_{q}, \hat{I}_{q} \boldsymbol{\mathcal{W}}) = (\boldsymbol{\mathcal{W}}, \hat{\mathcal{L}} \hat{J}_{0} \hat{\mathcal{L}} \hat{J}_{1} \dots \hat{\mathcal{L}} \hat{J}_{q-1} \hat{I}_{q} \boldsymbol{\mathcal{W}})$$

$$\approx (\boldsymbol{\mathcal{W}}, \hat{\mathcal{L}}^{q} \hat{I}_{q} \boldsymbol{\mathcal{W}}). \qquad (44)$$

In order to compare the scaling behavior of the two terms on the RHS of Eq. (42) in the limit of infinite Reynolds number we recall that within the inner product (or equivalently the average operation) an operator  $\hat{\mathcal{L}}$ amounts to simply  $\mathcal{W}/R$  [3]. This follows from the convergence in the UV and in the IR (in the limit  $\text{Re} \to \infty$ ) of the integral implied by the terms  $\mathcal{P}\mathcal{W}\cdot\nabla_r$  in Eq. (5), so that the leading contribution comes from distances of the order of R. In other words,  $\hat{\mathcal{L}}$  in a correlation function, when it operates on  $\mathcal{W}$ , introduces a term of the order of  $\mathcal{W}\cdot\nabla\mathcal{W}$ , which, due to the demonstrated locality in scale space, can be estimated as adding to the correlation a factor of the order of  $\mathcal{W}/R$ . Hence one has

$$(\boldsymbol{\mathcal{W}}, \hat{\mathcal{L}}^k \boldsymbol{\mathcal{W}}) \approx \left\langle \frac{\boldsymbol{\mathcal{W}}^{k+2}}{R^k} \right\rangle \propto R^{\zeta_{k+2}-k} , \qquad (45)$$

where the last step follows from Eq. (1).

The composition  $I_q$  containing say  $j^*$  times the operator  $\hat{\mathcal{L}}$  with  $j^* > q$  and having at least *i* times the operator  $\hat{\mathcal{L}}$  on the right hand of each operator  $\hat{J}_i$ , one can write

$$\hat{I}_{q}\boldsymbol{\mathcal{W}} = \sum_{j=0}^{j^{*}} c_{j}(R) \hat{\mathcal{L}}^{j} \boldsymbol{\mathcal{W}} , \qquad (46)$$

where  $c_j(R)$  is a function of the separation distance Rresulting from all the contributions to  $\hat{\mathcal{L}}^j \mathcal{W}$ . Note in particular that  $c_{j^*}(R) = 1$ . Considering now the following quantity for arbitrary s one has

$$(\mathcal{W}, \hat{\mathcal{L}}_s \hat{I}_q \mathcal{W}) \propto \sum_j c_j(R) R^{\zeta_{s+j+2}-j-s}$$
. (47)

Because of the Hölder inequalities

$$\zeta_m - \zeta_{m-r} \le \zeta_n - \zeta_{n-r} \qquad m > n \ , \ r > 0 \ , \tag{48}$$

to which any two contributions of Eq. (47) can be reduced, the leading contribution is that for which the index of  $\zeta$  is maximum, i.e.  $s + j^* + 2$ . As a result

$$(\mathcal{W}, \hat{\mathcal{L}}_s \hat{I}_q \mathcal{W}) \propto R^{\zeta_{s+j^*+2} - j^* - s}$$
 (49)

It follows from Eqs. (43) and (49) that the scaling behavior of the first term on the RHS of Eq. (42) is

$$(\boldsymbol{\mathcal{W}}, \hat{\mathcal{L}}_k \hat{I}_q \boldsymbol{\mathcal{W}}) \propto R^{\zeta_{k+j^*+2} - k - j^*}$$
, (50)

whereas using Eqs. (44),(45) and (49) that of the second one is

$$(\boldsymbol{\mathcal{W}}, \hat{\mathcal{L}}^k \hat{P}_q \hat{I}_q \boldsymbol{\mathcal{W}}) \propto R^{\zeta_{k+q+2} - \zeta_{2q+2} + \zeta_{q+j^*+2} - k - j^*} .$$
(51)

Recalling that k > q and  $j^* > q$  one has by Eq. (48)

$$\zeta_{k+j^*+2} - \zeta_{k+q+2} \le \zeta_{q+j^*+2} - \zeta_{2q+2} \,. \tag{52}$$

Hence expression (50) which coincides with the RHS of Eq. (41) is the leading one. We have therefore proven that relation (40) implies relation (41). Now it remains to show that Eq. (40) is true for p = 0. But this follows straightforwardly by setting q = 0 in Eq. (42) and replacing  $\hat{J}_{-1}$  by 1. Hence assertion (39) is proven. This in turn implies as was shown at the beginning of this section that (31) is satisfied.

# IV. SCALING LAWS IMPLIED BY THE CONTINUED FRACTION REPRESENTATION: DERIVATIVES AT TIME ZERO

To prepare for the comparison between the continued fraction and the multi-fractal representations we identify in this Section the leading scaling exponents that characterize the *n*th order time derivative of the correlation function Eq. (11) at  $\tau = 0$ . We show that

$$\boldsymbol{\mathcal{F}}_{2}^{(n)}(R,0) \approx R^{\zeta_{2+n}-n} \quad \forall n \;.$$
 (53)

Here we use the shorthand notation

$$\boldsymbol{\mathcal{F}}_{2}^{(n)}(R,0) \equiv \left. \frac{\partial^{n} \boldsymbol{\mathcal{F}}_{2}(R,\tau)}{\partial \tau^{n}} \right|_{\tau=0} \,. \tag{54}$$

In the next Section we will show that the same exponents are predicted by the multi-fractal representation, making the prediction of the Taylor expansion of the correlation function the same from the point of view of scaling behavior.

From Eq. (18) one deduces by inverse Laplace transform the following equation for  $\mathcal{F}_2^{(1)}(R,\tau)$ 

$$\mathcal{F}_{2}^{(1)}(R,\tau) = \gamma_{0}(R)\mathcal{F}_{2}(R,\tau) \qquad (55)$$
$$+ \int_{0}^{\tau} \boldsymbol{K}_{0}(R,\tau')\mathcal{F}_{2}(R,\tau-\tau')d\tau' ,$$

where  $\mathbf{K}_0(R,\tau)$  is the inverse Laplace transform of  $\tilde{\mathbf{K}}_0(R,z)$  [Eq. (21)],

$$\boldsymbol{K}_{0}(\boldsymbol{R},\tau) = \frac{1}{\boldsymbol{k}_{0}(\boldsymbol{R})} (\boldsymbol{\mathcal{W}}_{1}, e^{\hat{\mathcal{L}}_{1}\tau} \boldsymbol{\mathcal{W}}_{1}) .$$
 (56)

Equation (55) is the so-called memory-function equation,  $\mathbf{K}_0(R, \tau)$  being the memory kernel. At  $\tau = 0$  Eq. (55) becomes

$$\mathcal{F}_{2}^{(1)}(R,0) = \gamma_{0}(R) \mathbf{k}_{0}(R) \propto R^{\zeta_{3}-1} ,$$
 (57)

where we used the scaling laws (32).

The higher order partial time derivatives are obtained by differentiating Eq. (55),

$$\boldsymbol{\mathcal{F}}_{2}^{(n)}(R,\tau) = \boldsymbol{\gamma}_{0}(R)\boldsymbol{\mathcal{F}}_{2}^{(n-1)}(R,\tau) + \sum_{k=0}^{n-2} \boldsymbol{K}_{0}^{(n-k-2)}(R,\tau)\boldsymbol{\mathcal{F}}_{2}^{(k)}(R,0) + \int_{0}^{\tau} d\tau' \boldsymbol{K}_{0}(R,\tau')\boldsymbol{\mathcal{F}}_{2}^{(n-1)}(R,\tau-\tau') .$$
(58)

At  $\tau = 0$  one obtains

$$\mathcal{F}_{2}^{(n)}(R,0) = \gamma_{0}(R)\mathcal{F}_{2}^{(n-1)}(R,0)$$

$$+ \sum_{k=0}^{n-2} \mathbf{K}_{0}^{(n-k-2)}(R,0)\mathcal{F}_{2}^{(k)}(R,0) .$$
(59)

Now one realizes that this leads us to consider a hierarchy of equations for  $\mathbf{K}_{i}^{(q)}(R,0), q \geq 1, i \geq 0$ . Noticing that Eq. (26) is formally the same as Eq. (18) one obtains in analogy with Eq. (55)

$$\boldsymbol{K}_{i}^{(1)}(R,\tau) = \gamma_{i+1}(R)\boldsymbol{K}_{i}(R,\tau)$$

$$+ \int_{0}^{\tau} \boldsymbol{K}_{i+1}(R,\tau')\boldsymbol{K}_{i}(R,\tau-\tau')d\tau' ,$$
(60)

where  $\mathbf{K}_i(R, \tau)$  is the inverse Laplace transform of Eq. (25)

$$\boldsymbol{K}_{i}(\boldsymbol{R},\tau) = (\tilde{\boldsymbol{\mathcal{W}}}_{i+1}, e^{\hat{\mathcal{L}}_{i+1}\tau} \boldsymbol{\mathcal{W}}_{i+1}) (\tilde{\boldsymbol{\mathcal{W}}}_{i}, \boldsymbol{\mathcal{W}}_{i})^{-1} .$$
(61)

Notice that at  $\tau = 0$  using also Eq. (27) yields

$$\boldsymbol{K}_i(R,0) = \boldsymbol{k}_{i+1}(R) \ . \tag{62}$$

In analogy with Eq. (59) one has for  $q \ge 1$ 

$$\boldsymbol{K}_{i}^{(q)}(R,0) = \boldsymbol{\gamma}_{i+1}(R) \boldsymbol{K}_{i}^{(q-1)}(R,0) \qquad (63)$$
$$+ \sum_{k=0}^{q-2 \ge 0} \boldsymbol{K}_{i+1}^{(q-k-2)}(R,0) \boldsymbol{K}_{i}^{(k)}(R,0) .$$

Note that denoting from now on  $\mathbf{K}_{-1} \equiv \mathbf{\mathcal{F}}_2$  and allowing  $i \geq -1$  in Eq. (63) one recovers Eq. (59).

It follows that to determine the scaling behavior of  $\mathbf{K}_{-1}^{(n)}(R,0)$  for  $n \geq 1$  in order to prove Eq. (53) one has to know  $\mathbf{K}_{i}^{(q)}(R,0) \forall q + 2i \leq n-2$  with  $q \geq 0$  and  $i \geq -1$ . It is sufficient to prove the following

$$\boldsymbol{K}_{i}^{(q)}(R,0) \propto R^{\zeta_{q+2i+4}-\zeta_{2i+2}-q-2+2\delta_{i,-1}}$$
, (64)

where  $q \geq 0$ ,  $i \geq -1$  and  $\delta_{i,-1}$  is a Kronecker delta, which we do by induction in the sequel. Setting i = -1and recalling that  $\zeta_0 = 0$  Eq. (53) results directly.

Let us assume that Eq. (64) is true  $\forall q + 2i \leq n-3$ with  $n \geq 1$ . Then we show that this expression is still valid for n + 1. By the scaling relations (32) we already know that Eq. (64) is true  $\forall q + 2i \leq n-3$  with n = 1. For q + 2i = n - 2 Eq. (63) becomes

$$\boldsymbol{K}_{i}^{(n-2-2i)}(R,0) = \boldsymbol{\gamma}_{i+1}(R) \boldsymbol{K}_{i}^{(n-3-2i)}(R,0) \qquad (65)$$
$$+ \sum_{k=0}^{n-4-2i\geq 0} \boldsymbol{K}_{i+1}^{(n-4-2i-k)}(R,0) \boldsymbol{K}_{i}^{(k)}(R,0) .$$

Note that for 2i = n - 2 (*n* even) one has to resort to Eq. (62) and that for 2i = n - 3 (*n* odd) the sum in Eq. (65) is empty yielding directly

$$\begin{aligned} \boldsymbol{K}_{\frac{n-2}{2}}(R,0) &= \boldsymbol{k}_{\frac{n}{2}}(R) \propto R^{\zeta_{n+2}-\zeta_n-2} , \qquad (66) \\ \boldsymbol{K}_{\frac{n-3}{2}}^{(1)}(R,0) &= \boldsymbol{\gamma}_{\frac{n-1}{2}}(R) \boldsymbol{k}_{\frac{n-1}{2}}(R) \propto R^{\zeta_{n+2}-\zeta_{n-1}-3+2\delta_{n,1}} , \end{aligned}$$

respectively, which are indeed of the form of Eq. (64). Now for  $n-4-2i \ge 0$  the only object in Eq. (65) whose scaling is not known by Eq. (64) for  $q + 2i \le n - 3$  is that corresonding to k = 0,  $\mathbf{K}_{i+1}^{(n-4-2i)}(R, 0)$ . Hence we prove also by induction, but now on i, that Eq.(64) holds for q + 2i = n - 2. Assuming that  $\mathbf{K}_{i+1}^{(n-4-2i)}(R, 0)$  scales according to Eq. (64) we show that it is also true for i - 1. By Eq. (66) we already know that this is true for 2i = n - 4 (n even) and 2i = n - 3 (n odd). We first show that for  $n - 4 - 2i \ge 0$  the term in k = 0 in Eq. (65) is leading with respect to the terms in higher k appearing in this equation. By hypothesis eq. (65) yields

$$\mathbf{K}_{i}^{(n-2-2i)}(R,0) \propto R^{-\zeta_{2i+4}-\zeta_{2i+2}-n+2i+2\delta_{i,-1}} \qquad (67) \\
\left(R^{\zeta_{2i+5}+\zeta_{n+1}} + \sum_{k=0}^{n-4-2i\geq 0} R^{\zeta_{n+2-k}+\zeta_{2i+4+k}}\right).$$

Hence one has to show that for k > 0

$$\zeta_{n+2} - \zeta_{n+2-k} \le \zeta_{2i+4+k} - \zeta_{2i+4} , \qquad (68)$$

which follows from the Hölder inequalities since  $k \leq n-4-2i$ . Secondly, we notice from Eq. (67) that the term in front of the sum scales exactly as a term in k = 1 so that by the preceding argument it is subleading with respect to the term in k = 0 for  $n - 4 - 2i \geq 0$ . It follows that the term in k = 0 is the leading one in Eq. (67) so that coming back to Eq. (65) one has

$$\boldsymbol{K}_{i}^{(n-2-2i)}(R,0) \approx \boldsymbol{K}_{i+1}^{(n-4-2i)}(R,0)\boldsymbol{K}_{i}^{(0)}(R,0) \qquad (69)$$
$$\propto R^{\zeta_{n+2}-\zeta_{2i+2}-n+2i+2\delta_{i,-1}}.$$

Hence Eq. (64) is true for q + 2i = n - 2 with  $q \ge 0$  and by induction on *i* for any  $i \ge -1$ . Now by induction on *n* it is also true for any *n*. We have therefore proven Eq. (64) and henceforth Eq. (53). As for Eq. (53) it results directly from Eqs. (66) and (69)

### V. THE MULTI-SCALING REPRESENTATION

The multi-scaling representation of  $\mathcal{F}_2(R,\tau)$  can be written as [3]:

$$\boldsymbol{\mathcal{F}}_{2}(R,\tau) = U^{2} \int d\mu(h) \left(\frac{R}{L}\right)^{2h+\mathcal{Z}(h)} \boldsymbol{f}_{2}\left(\frac{\tau}{\tau_{R,h}}\right), \quad (70)$$

where U is the characteristic magnitude of the velocity difference across the outer scale of turbulence,  $f_2$  is a function of the scaled time variable only, and

$$\tau_{R,h} \sim \frac{R}{U} \left(\frac{L}{R}\right)^h.$$
(71)

The function  $\mathcal{Z}(h)$  is related to the scaling exponents  $\zeta_n$  of the *n*th order structure functions through the saddle point requirement

$$\zeta_n = \min_h [nh + \mathcal{Z}(h)] . \tag{72}$$

To find the scaling exponents associated with the time derivatives of  $\mathcal{F}_2(R,\tau)$  at  $\tau = 0$  one computes the *n*-th order partial time derivative of Eq. (70) to obtain

$$\boldsymbol{\mathcal{F}}_{2}^{(n)}(\boldsymbol{R},\tau) \tag{73}$$

$$= \frac{U^{2+n}}{R^{n}} \int d\mu(h) \left(\frac{\boldsymbol{R}}{\boldsymbol{L}}\right)^{(2+n)h+\mathcal{Z}(h)} \boldsymbol{f}_{2}^{(n)}\left(\frac{\tau}{\tau_{\boldsymbol{R},h}}\right),$$

where

$$\boldsymbol{f}_{2}^{(n)}(s) = \frac{d^{n} \boldsymbol{f}_{2}(s)}{ds^{n}} .$$
(74)

At  $\tau = 0$  this gives

$$\mathcal{F}_{2}^{(n)}(R,0) = \mathbf{f}_{2}^{(n)}(0) \frac{U^{2+n}}{R^{n}} \int d\mu(h) \left(\frac{R}{L}\right)^{(2+n)h+\mathcal{Z}(h)} .$$
(75)

Computing the integral at the saddle point in the limit  $R/L \rightarrow 0$  and using (72) we find

$$\boldsymbol{\mathcal{F}}_{2}^{(n)}(R,0) \propto R^{\zeta_{2+n}-n} \,, \tag{76}$$

in correspondence with Eq. (53). We thus see that the continued fraction representation generates the same predictions regarding the multiplicity of time scales characterizing the time correlation functions as the multiscaling representation. We take this as an independent evidence for the correctness of the latter.

# VI. CONCLUSION

In conclusion, we showed that the formally exact continued fraction representation of the time correlation functions of Belinchier-L'vov velocity differences has the same Taylor expansion as the multi-scaling representation, at least in terms of the leading scaling exponents order by order.

It should be noted that the continued fraction representation can be used as an approximant for the correlation function when analytic forms of the time-correlation functions are needed. In the lowest approximation one takes in Eq. (30)  $\mathbf{k}_1(R) = 0$ , producing an exponential decay of the correlation function, with a typical decay rate  $\gamma_0(R)$ . In every successive approximation ( $\mathbf{k}_2 = 0$ ,  $\mathbf{k}_3 = 0$ , etc.) one introduces more and more characteristic scales, each one characterized by a different "dynamical exponent", taking progressively more information about the statistics of higher order correlation functions into account.

### ACKNOWLEDGMENTS

We thank Daniela Pierotti and Reuven Zeitak for discussions concerning the non-Hermiticity of the Liouvillian. This work was supported in part by the German Israeli Foundation, the European Commission under the Training and Mobility of Researchers program, The Basic Research Fund administered by the Israel Academy of Science and Humanities, the Minerva Center for Nonlinear Physics, and the Naftali and Anna Backenroth-Bronicki Fund for Research in Chaos and Complexity.

- K.R. Sreenivasan and R.A. Antonia, Ann. Rev. of Fluid Mech. 29, 435 (1997).
- [2] U. Frisch, *Turbulence: The Legacy of A.N. Kolmogorov* (Cambridge University Press, Cambridge 1995).
- [3] V.S. L'vov, E.V. Podivilov and I. Procaccia, Phys. Rev. E 55, 7030 (1997),
- [4] V.S. L'vov and I. Procaccia Physica A, 257 165 (1998).
- [5] V.I. Belinicher, V.S. L'vov, A. Pomyalov and I. Procaccia, "Computing the Scaling Exponents in Fluid Turbulence from First Principles: Demonstration of Multiscaling", J. Stat. Phys., in press.
- [6] S. Grossmann and S. Thomae, Z. Phys. B 49, 253 (1982).
- [7] S. Grossmann and C.Wiele, Z. Phys. B 103, 469 (1997).
- [8] R. Zwanzig, Phys. Rev. **124**, 983 (1961).
- [9] H. Mori, Prog. Theor. Phys. 34, 399 (1965).
- [10] A.S. Monin, Sov. Phys. Doklady 4,271 (1959).
   V.I. Belinicher and V.S. L'vov Zh. Eksp. Teor. Fiz. 93, 1269 (1987) [Sov. Phys. JETP, 66, 303 (1987)].
- [11] A.S. Monin and A.M. Yaglom, *Statistical Fluid Mechan*ics (MIT press, 1971).
- [12] V.S L'vov, E.V. Podivilov and I. Procaccia, Phys. Rev. Lett. 76, 3963 (1996).