Nonlinear Thermomagnetic Waves in Type II Superconductors

N.A. Taylanov

Theoretical Physics Department and Institute of Applied Physics, National University of Uzbekistan,

E-mail: taylanov@iaph.silk.org

bstract

This work is devoted to analysis of the nonlinear dynamics of interrelated thermal and electromagnetic perturbations in a superconductor caused by dissipative effects involved in the viscous motion of the magnetic flux. The structure and evolution of the stationary thermomagnetic waves propagating in the superconductor is studied based on a system of nonlinear equations describing dynamics of the instability development in the system studied. The apperance of these structures is qualititively described and the wave propagation velocity is estimated. The affect a thermoactivated flux creep to the structures of the nonlinear thermomagnetic wave propagation is studied. It is shown that taking into account a nonlinear dependency of the current density j on the electric field strength E would not qualitatively change the main results, since the character of the equilibrium state on the phase plane remains esentially the same.

At present superconducting systems with high critical field strengs and current densites are widely implemented in various advenced technologies. However, successful operation of the superconducting materials is only possible provided that special measures are taken to prevent a system from the thermal or magnetic breakage of superconductivity and transition to a normal (resistive)state [1]. For this reason, one of the main problems in the investigation of properties of superconductors is that of predicting the superconducting state breakage caused by dissipative and nonlinear effects related to viscous motions of the magnetic flux. This explains a considerable interest in the study of dissipative and nonlinear effects in superconductors that has apose in recent years.

This work is devoted to analysis of the nonlinear dynamics of interrelated thermal and electromagnetic perturbations in a superconductor caused by dissipative effects involved in the viscous motion of the magnetic flux. The structure and evolution of the stationary thermomagnetic waves propagating in the superconductor is studied based on a system of nonlinear equations describing dynamics of the instability development in the system is studied.

The evolution of thermal and electromagnetic perturbations in a superconductor is described by a set involving the thermal conductivity equation [2-5]

$$\nu \frac{dT}{dt} = \nabla(\kappa \nabla T) + \vec{j}\vec{E}, \qquad (1)$$

the Maxwell equation

$$\frac{4\pi}{c^2}\frac{d\vec{j}}{dt} = \Delta\vec{E}\,,\tag{2}$$

and the related equation of the critical state

$$\vec{j} = \vec{j}(T, \vec{H}) + \vec{j}_r(\vec{E})$$
. (3)

Here $\nu = \nu(T)$ is the heat capacity, $\kappa = \kappa(T)$ is the thermal conductivity coefficient, \vec{j}_c is the critical current density, and \vec{j}_r is the current density in the resistive state.

The above system is essentially nonlinear because the right-hand part of Eq.(1) contains a term describing the Joule heat evolution in the region of a resistive phase. Such a set (1)-(3) of nonlinear parabolic differential equations in partial derivatives has no exact analytical solution.

Let us consider a planar semi-infinite sample (x > 0) placed in an external magnetic field $\vec{H} = (0, 0, H_e)$ growing at a constant rate $\frac{d\vec{H}}{dt} = const$. According to Maxwel's equation (2), there is a vortex electric field $\vec{E} = (0, E_e, 0)$ in the sample, directed parallel to the current density \vec{j} : $\vec{E} \parallel \vec{j}$; where H_e is the amplitude of the external magnetic field and E_e is the amplitude of the external electric field.

For the automodel propagating waves of the type $\xi(x,t) = x - vt$ the initial differential Eqs. (1)-(3) acquire the following form

$$-\nu v \frac{dT}{d\xi} = \frac{d}{d\xi} \left[\kappa \frac{dT}{d\xi} \right] + jE \,, \tag{4}$$

$$\frac{dE}{d\xi} = -\frac{4\pi v}{c^2} j \,, \tag{5}$$

$$E = \frac{v}{c}H.$$
 (6)

A set of boundary conditions corresponding to Eqs. (4)-(6) is as follows:

$$T(\xi \to +\infty) = T_0, \qquad \frac{dT}{d\xi}(\xi \to -\infty) = 0,$$

$$E(\xi \to +\infty) = 0, \qquad E(\xi \to -\infty) = E_e,$$
(7)

Let us use the Bean-London model of the critical state for the dependence $j_c(T, H)$; $\frac{dj_c}{dH} = 0$ [6]. According to this model the dependence $j_c(T)$ can be represented as $j_c(T) = j_0[1 - a(T - T_0)]$: where j_0 is the equilibrium current density, a is the constant parameter and T_0 is the temperature of the cooling medium. A characteristic field dependence of $j_r(E)$ in the region of sufficiently strong electric fields $(E > E_f)$ can be approximated by a piecewise linear function $j_r \approx \sigma_f E$, where σ_f is the efficitive conductivity. In the field of small field strengths $(E < E_f)$ we assume a relationship $j_r(E) \approx j_1 \ln \frac{E}{E_0}$, to be valid with j_1 being a characteristic local current density scatter (related to a pinning force inhomogeneity) on the order of $j_1 \approx 0,01j_c$ and E_0 being a constant. This relationship between j_r and E is due to a thermoactivated early experiments (see, e.g. [1]). The thermoactivated flux creep is

principal mechanism of the energy dissipation during the magnetic flux penetration deep into the sample. This factor must affect the character of the nonlinear thermomagnetic wave propagation in the system studied.

Excluding the variables $T(\xi)$ and $H(\xi)$, using Eqs.(4) and (6), and taking into account the boundary conditions (7), we obtain an equation describing the electric field $E(\xi)$ distribution:

$$\frac{d^2 E}{dz^2} + \beta \left[1 + \frac{j_1}{\sigma_d E} \tau \right] \frac{dE}{dz} + \beta^2 \tau \left[\frac{j_0}{\sigma_d} + \frac{j_1}{\sigma_d} \ln \frac{E}{E_0} \right] = \frac{E^2}{2E_\kappa} \,. \tag{8}$$

The dimensionless variables used in Eq. (8) are defined as follows:

$$z = \frac{\xi}{L}, \quad \beta = \frac{vt_{\kappa}}{L}, \quad t_{\kappa} = \frac{\nu L^2}{\kappa}, \quad \tau = \frac{4\pi\sigma_d\kappa}{c^2\nu}, \quad E_{\kappa} = \frac{\kappa}{aL^2}, \quad L = \frac{cH_e}{4\pi j_0},$$

where L is the depth of the magnetic flux penetration into the sample, σ_d is the differential conductivity. The corresponding equation of state is obtained using a relationship

$$\Omega E^2 = X(E) = 1 + \frac{j_1}{j_0} \ln \frac{E}{E_0},$$
(9)

where $\Omega = \frac{\sigma_d}{2\beta^2 \tau j_0 E_{\kappa}}$. As seen from the plots of X(E) versus E (see figure), there exists a single point of intersection of the curves $y = \Omega E^2$ and $y = 1 + (j_1/j_0) \ln(E/E_0)$, which corresponds to a single stable equilibrium state. The stability of this state is determined by the sign of derivative d^2E/dz^2 in the vicinity of the equilibrium point. The wave velocity v_E can be determined from stationary $\left(\frac{dE}{dz}=0\right)$ Eq.(8) with an allowance for the boundary conditions [7]:

$$v_E = \frac{L}{t_\kappa} E_e \left[2\tau \frac{j_0 E_\kappa}{\sigma_d} \left(1 + \frac{j_1}{j_0} \ln \frac{E_e}{E_0} \right) \right]^{-1/2} \,. \tag{10}$$

Now we can use Eq. (6) and readily derive an equation describing the field distribution for a nonlinear H - wave. The wave velocity v_H of this H- wave is given by the formula

$$v_H = \frac{cE}{H_e} \exp\left[\frac{\sigma_d}{2\tau j_1} E_\kappa \left(\frac{LH_e}{ct_\kappa}\right)^2 - \frac{j_0}{j_1}\right].$$
(11)

As seen, the wave velocity v_H exponentially increases with the field amplitude H_e . For a sufficiently small amplitude $H_e < H_a$, where

$$H_a = \frac{ct_\kappa}{L} \left[\frac{2\tau j_0 E_\kappa}{\sigma_d} \right]^{1/2} \tag{12}$$

the wave velocity is negligible small, which corresponds to the case of the thermoactivated flux creep. For $H_e = H_a$, the wave propagates at a finite constant velocity. In this case, the maximum heating of a superconductor in the region immiadiately in front of the wave is described by the relationship

$$\frac{T - T_0}{T_0} = \frac{H_e^2}{8\pi\nu T_0} \,. \tag{13}$$

In conluding, we must note that taking into account a nonlinear dependency of the current density j on the electric field strength E would not qualitatively change the main results, since the character of the equilibrium state on the phase plane remains esentially the same (see [2]).

- [1] Mints R.G., Rakhmanov A.L. Instability in superconductors. .: Nauka, 1984.
- [2] Taylanov N.A., Maksimov I.L., Mastakov U.N.. Phys.Solid State (FTT, Russia) Vol. 28. N 6 (1986), pp. 2323-2328.
- [3] Adkhamov A.A., Taylanov N.A., Maksimov I.L., Phys.Solid State (FTT, Russia) Vol. 30. N 3 (1988). pp. 887-888.
- [4] N.A.Taylanov N.A., Kuchkarov S.P. Phys.Solid State (FTT, Russia)Vol. 33. N 6 (1991), pp. 1873-1875.
- [5] N.A.Taylanov N.A., Kuchkarov S.P. Technical Phys. (JTF, Russia) Vol. 61, N 7 (1991) pp.197-198.
- [6] Bean C.P. Phys. Rev. Lett. 8, 6, 250(1962).
- [7] .. Nonlinear waves in the dispersive media. .: Nauka, 1973.