## X-ray Dichroism and Orbital Anapoles

Paolo Carra<sup>1</sup>, Andrés Jerez<sup>2\*</sup>, and Ivan Marri<sup>1\*\*</sup>

<sup>1</sup>European Synchrotron Radiation Facility, B.P. 220, F-38043 Grenoble Cédex, France

<sup>2</sup>Institut Laue-Langevin, B.P. 156X, F-38042 Grenoble Cédex, France

(October 27, 2018)

We present a theoretical analysis of photo-absorption spectroscopies in noncentrosymmetric systems, covering both x-ray and optical regions. Integrated dichroic spectra are interpreted using microscopic effective operators, which are obtained by coupling the orbital angular momentum to the orbital anapole. Symmetry arguments afford a classification of valence-electron states in the presence of parity nonconserving hybridisation. Enantiomorphism is identified by a two-particle chiral operator.

PACS numbers:78.70.Dm, 33.55.Ad

The current paper aims at providing a theoretical interpretation of two recent experiments:

> X-ray natural circular dichroism (XNCD), probed in  $\alpha$ -LiIO<sub>3</sub> [1] and in Na<sub>3</sub>Nd(digly)<sub>3</sub>·2NaBF<sub>4</sub>·6H<sub>2</sub>O [2]. (The effect was observed at the iodine *L* edges and at the Nd *L*<sub>3</sub> edge.)

> X-ray nonreciprocal linear dichroism (XNRLD), detected at vanadium K edge in the low-temperature insulating phase of a Cr-doped V<sub>2</sub>O<sub>3</sub> crystal [3].

XNCD measures the difference in absorption between right and left circularly polarised radiations. XNRLD implies a difference in absorption between radiations with linear polarisation parallel or perpendicular to a local symmetry axis. Both phenomena stem from the interference between electric-dipole (E1) and electric-quadrupole (E2) transitions that raise an inner-shell electron to empty valence orbitals. Detecting a nonvanishing signal thus requires an ordered structure (crystal) and the breaking of space inversion.

As will be shown, Lie groups (or, alternatively, Lie algebras) furnish a powerful tool for interpreting x-ray dichroism in non-centrosymmetric crystals. In fact, effective microscopic operators, which express electronic properties revealed by the spectra, are readily deduced from the pertinent group generators: *The orbital angular momentum and orbital anapole*. Our approach hinges therefore on spectrum-generating algebras (SGAs), a concept originally introduced in nuclear [4] and particle physics [5,6].

We can switch from such an abstract formulation to a physical picture that is easy to grasp. This is achieved by relating the chosen SGA to the electron orbital current. To this end, Trammell's expansion [7] is exploited.

As no pseudoscalar is accessible to the E1-E2 interference, we must include E1-magnetic-dipole (M1) excitations in order to discuss orbital chirality. Our approach readily extends to the optical range. Results for the E1-M1 circular dichroism (observable also in powder samples) will thus be reported.

Our analysis is restricted to dichroic spectra integrated

over a finite energy range, corresponding to the two partners of a spin-orbit split inner-shell:  $j_{\pm} = l_c \pm \frac{1}{2}$ .

We follow Carra and Benoist [8] and consider the general framework of a de Sitter algebra: so(3, 2) [4,9]. A realisation of such an algebra is provided by the operators:  $\mathbf{A}^- = i(\mathbf{A} - \mathbf{A}^{\dagger})/2$ ,  $\mathbf{A}^+ = (\mathbf{A} + \mathbf{A}^{\dagger})/2$ ,  $\mathbf{L}$  and  $N_0$ , where  $\mathbf{A} = \mathbf{n} f_1(N_0) + \nabla_{\Omega} f_2(N_0)$ , with  $\mathbf{n} = \mathbf{r}/r$  and  $\nabla_{\Omega} = -i\mathbf{n} \times \mathbf{L}$ ;  $\mathbf{L}$  denotes the orbital angular momentum (in units of  $\hbar$ ). Furthermore,  $f_1(N_0) = (N_0 - 1/2)f_2(N_0)$  and  $f_2(N_0) = \sqrt{(N_0 - 1)/N_0}$ , where  $N_0|lm\rangle = (l + \frac{1}{2})|lm\rangle$ , with  $|lm\rangle$  a spherical harmonic.  $\mathbf{A}$  and  $\mathbf{A}^{\dagger}$  are known as shift operators as their action on  $|lm\rangle$  changes l into  $l \pm 1$ .

As will be seen, discussing x-ray dichroism in noncentrosymmetric systems only requires L (rotations) and  $A^-$  (boosts), which themselves generate a homogeneous Lorentz group: SO(3, 1). We observe that

$$\boldsymbol{L} \cdot \boldsymbol{A}^{-} = \boldsymbol{A}^{-} \cdot \boldsymbol{L} = 0, \qquad (1)$$

and that a physical interpretation of  $A^-$  is provided by the relation

$$\boldsymbol{\Omega}^{-} = (\boldsymbol{n} \times \boldsymbol{L} - \boldsymbol{L} \times \boldsymbol{n})/2 = i \left[\boldsymbol{n}, \boldsymbol{L}^{2}\right]/2 \qquad (2)$$
$$= \frac{i}{2} \left(\boldsymbol{\nabla}_{\Omega}^{\dagger} - \boldsymbol{\nabla}_{\Omega}\right) = \frac{1}{2} \frac{1}{\sqrt{N_{0}}} \left[N_{0}, \boldsymbol{A}^{-}\right]_{+} \frac{1}{\sqrt{N_{0}}},$$

where  $[...]_+$  denotes an anticommutator. Eq. (2) defines the (purely angular) orbital anapole [10–12]:  $\Omega^-$ . As pointed out by Dothan *et al.* [6],  $\boldsymbol{L}$  and  $\Omega^-$  also generate SO(3, 1). [ $\boldsymbol{L}$  and  $\boldsymbol{A}^+$  generate another SO(3, 1) subgroup of SO(3, 2). Being space-odd and time-even,  $\boldsymbol{A}^+$  does not enter our theory of dichroism in noncentrosymmetric systems.]

Our formulation is based on a localised (atomic) model and exploits the Racah-Wigner calculus to relate integrated dichroic spectra to the ground-state expectation value of effective orbital operators. These operators are constructed by coupling the hermitean vector operators  $\boldsymbol{L}$  (space-even) and  $\boldsymbol{\Omega}^-$  (space-odd) to obtain space-odd irreducible tensors. In such a framework, one-electron tensors are given by ( $\iota$  is an electron label)

$$\sum_{\iota} [\boldsymbol{L}_{\iota} \otimes \boldsymbol{\Omega}_{\iota}^{-}]_{q}^{(k)l \leftrightarrow l'} \equiv \sum_{\iota} [\boldsymbol{L}_{\iota} \otimes \boldsymbol{\Omega}_{\iota}^{-}]_{q}^{(k)l,l'} + \text{H.c.}$$
$$= \sum_{mm'} \langle l'm' | [\boldsymbol{L} \otimes \boldsymbol{\Omega}^{-}]_{q}^{(k)} | lm \rangle a_{l'm'}^{\dagger} a_{lm} + \text{H.c.}$$
(3)

Here,  $l' = l \pm 1$ ;  $a_{lm}^{\dagger}$  and  $a_{l'm'}$  create and annihilate valence electrons. (As usual, tensor couplings are defined via Clebsch-Gordan coefficients:  $[U^{(p)} \otimes V^{(\kappa)}]_q^{(k)} \equiv \sum_{\mu,\nu} C_{p\mu;\kappa\nu}^{kq} U_{\mu}^{(p)} V_{\nu}^{\kappa}$ .) Extending Eq. (3) to define twoelectron space-odd irreducible tensors is straightforward. (Point-group symmetry can be added to our analysis by introducing *point-group coordinates* [13] and retaing the pertinent totally-symmetric representations. Full details will be given elsewhere.)

E1-E2 interference: XNCD [1,2] The effect is characterised by an orbital pseudodeviator, i.e. a space-odd and time-even rank-two irreducible tensor [14]. In our formalism, its microscopic expression is given by a coupling of  $\boldsymbol{L}$  to the orbital anapole; it reads:  $[\boldsymbol{L} \otimes \boldsymbol{\Omega}^{-}]_{q}^{(2)l \leftrightarrow l'}$ . For the integrated XNCD spectrum, we thus find

$$\int_{j_{-}+j_{+}} \frac{\sigma_{\text{XNCD}}(\omega)}{(\hbar\omega)^{2}} d(\hbar\omega) = \frac{8\pi^{2}\alpha}{3\hbar c} (2l_{c}+1)$$

$$\sum_{\substack{l=l_{c}\pm 1\\l'=l\pm 1}} R_{l_{c}l}^{(1)} R_{l_{c}l'}^{(2)} a_{l'}(l_{c},l) \sum_{q} \sqrt{\frac{3}{2}} T_{q}^{(2)*}(\boldsymbol{\epsilon}^{-},\boldsymbol{\epsilon}^{+},\hat{\mathbf{k}})_{2} \quad (4)$$

$$\langle \psi_{0} | \sum_{\iota} [\boldsymbol{L}_{\iota} \otimes (\boldsymbol{L} \times \boldsymbol{n} - \boldsymbol{n} \times \boldsymbol{L})_{\iota}]_{q}^{(2)l \leftrightarrow l'} | \psi_{0} \rangle.$$

In the foregoing equation,  $\sigma_{\text{XNCD}}(\omega) = \sigma_{\text{X}}^{\epsilon^+}(\omega) - \sigma_{\text{X}}^{\epsilon^-}(\omega)$ denotes the cross section for E1-E2 circular dichroism in the x-ray region (X), with

$$\sigma_{\mathbf{X}}^{\boldsymbol{\epsilon}}(\omega) = 2\pi^{2}\alpha\hbar\omega[i\sum_{f}\sum_{\iota\iota'}\langle g \mid \boldsymbol{\epsilon}^{*}\cdot\mathbf{r}_{\iota}\mid f\rangle \qquad (5)$$
$$\langle f \mid \boldsymbol{\epsilon}\cdot\mathbf{r}_{\iota'}\mathbf{k}\cdot\mathbf{r}_{\iota'}\mid g\rangle + \text{c.c.}]\delta(E_{f}-E_{g}-\hbar\omega);$$

 $\hbar\omega$ , **k** and  $\epsilon^{\pm} = \mp (i/\sqrt{2})(\epsilon_1 \pm i\epsilon_2)$  represent energy, wave vector, and circular polarisations of the photon, respectively;  $\alpha = e^2/\hbar c$ . The radial integrals are given by  $R_{l_cl}^{(L)} = \int_0^\infty dr \varphi_{l_c}(r) r^{L+2} \varphi_l(r)$ , where  $\varphi_{l_c}(r)$  and  $\varphi_l(r)$ denote inner-shell and valence radial wave functions, respectively [15,16].

A general expression can be given for the wave-vector and polarisation responses used throughout this work. It reads:  $i^z T_q^{(k)}(\boldsymbol{\epsilon}, \boldsymbol{\epsilon}^*, \hat{\mathbf{k}})_z$ , where

$$T_q^{(k)}(\boldsymbol{\epsilon}, \boldsymbol{\epsilon}^*, \hat{\mathbf{k}})_z = \sum_y \sqrt{(2z+1)(2y+1)}(-1)^{1+k} \qquad (6)$$

$$\left\{ \begin{array}{cc} 1 & k & y \\ 1 & 1 & z \end{array} \right\} \sum_{\eta\beta} C^{kq}_{y\eta;1\beta} \sum_{\alpha\delta} C^{y\eta}_{1\delta;1\alpha} Y_{1\delta}(\boldsymbol{\epsilon}^*) Y_{1\alpha}(\boldsymbol{\epsilon}) Y_{1\beta}(\hat{\mathbf{k}}) \,,$$

with z = 1, 2 denoting E1-M1 (imaginary) and E1-E2 (real) responses, respectively;  $\hat{\mathbf{k}} = \mathbf{k}/k$ . Explicit forms are given in Table I.

E1-E2 Interference: XNRLD [3]. One effective operators is found in this case: a space and time odd tensor of rank three, i.e. an orbital septor. This unusual magnetic moment stems from a coupling of the orbital anapole with the quadrupolar moment of  $\boldsymbol{L}$  and takes the form  $[[\boldsymbol{L} \otimes \boldsymbol{L}]^{(2)} \otimes \boldsymbol{\Omega}^{-}]_{q}^{(3)l \leftrightarrow l'}$ .

The integrated XNRLD spectrum can thus be written as

$$\int_{j_{-}+j_{+}} \frac{\sigma_{\text{XNRLD}}(\omega)}{(\hbar\omega)^{2}} d(\hbar\omega) = \frac{8\pi^{2}\alpha}{\hbar c} (2l_{c}+1) \sum_{\substack{l=l_{c}\pm 1\\l'=l\pm 1}} R_{l_{c}l'}^{(1)}$$
$$R_{l_{c}l'}^{(2)} b_{l'}(l_{c},l) \sum_{q} \left( T_{q}^{(3)*}(\boldsymbol{\epsilon}^{\parallel},\hat{\mathbf{k}})_{2} - T_{q}^{(3)*}(\boldsymbol{\epsilon}^{\perp},\hat{\mathbf{k}})_{2} \right)$$
(7)
$$\langle \psi_{0} | \sum_{\iota} \left[ [\boldsymbol{L}_{\iota} \otimes \boldsymbol{L}_{\iota}]^{2} \otimes (\boldsymbol{L} \times \boldsymbol{n} - \boldsymbol{n} \times \boldsymbol{L})_{\iota} \right]_{q}^{(3)\, l \leftrightarrow l'} | \psi_{0} \rangle ,$$

where  $\sigma_{\text{XNRLD}}(\omega) = \sigma_{\text{X}}^{\parallel}(\omega) - \sigma_{\text{X}}^{\perp}(\omega)$  stands for the cross section for E1-E2 linear dichroism; here,  $\parallel$  and  $\perp$  denote two orthogonal linear-polarisation states [16].

Notice that the E1-E2 interference contains a further magnetic term, which vanishes in the geometry of Eq. (7). It is an *orbital vector*, with polarisation response  $T_q^{(1)}(\boldsymbol{\epsilon}, \hat{\mathbf{k}})_2$ . In our formulation, it is given by  $\sum_{\iota} [(\boldsymbol{L}_{\iota} \times \boldsymbol{n}_{\iota} - \boldsymbol{n}_{\iota} \times \boldsymbol{L}_{\iota})_q]^{l \leftrightarrow l'}$ , i.e. by the orbital anapole. This toroidal contribution [10] to the orbital current (see below) could be detected by the following experiment. Absorb linearly polarised x rays at the 'magic angle'  $\widehat{\mathbf{kB}} \simeq 39.23^{\circ}$  for parallel and antiparallel magnetoelectric annealing [3] of the sample;  $\boldsymbol{B} \equiv$  external magnetic field. Subtract the two spectra.

Stemming from parity nonconserving hybridisation of the valence electrons, the effect should be easier to detect than its nuclear counterpart [17], as the work of Ref. [3] appears to indicate.

E1-M1 interference. As previously stated, our approach readily extends to cover the visible range, where the E1-M1 interference is predominant. In this case, the cross section for optical natural circular dichroism (ONCD) takes the form:  $\sigma_{ONCD}(\omega) = \sigma_{O}^{\epsilon^{+}}(\omega) - \sigma_{O}^{\epsilon^{-}}(\omega)$ , with [18]

$$\sigma_{\rm O}^{\boldsymbol{\epsilon}}(\omega) = \frac{2\pi^2 \alpha \hbar}{m} \left[ \sum_{f} \sum_{\iota\iota'} \langle g \mid \boldsymbol{\epsilon}^* \cdot \mathbf{r}_{\iota} \mid f \rangle \right]$$

$$\langle f \mid \boldsymbol{\epsilon} \times \mathbf{k} \cdot \boldsymbol{l}_{\iota'} \mid g \rangle + \text{c.c.} \delta(E_f - E_g - \hbar \omega),$$
(8)

with  $\boldsymbol{l} = \hbar \boldsymbol{L}$ .

For the integrated ONCD spectrum we obtain

$$\int_{j_-+j_+} \frac{\sigma_{\text{ONCD}}(\omega)}{\hbar\omega} d(\hbar\omega) = \frac{4\pi^2 \alpha}{mc} \sum_{l,l'=l\pm 1} R_{ll'}^{(1)} R_{l'l'}^{(0)}$$
$$d(l,l') \langle \psi_0 | \left\{ \sqrt{\frac{2}{3}} T_0^{(0)}(\boldsymbol{\epsilon}^+, \boldsymbol{\epsilon}^-, \hat{\mathbf{k}})_1 \right.$$
(9)

$$\begin{split} &\sum_{\iota\neq\iota'} \left[ \boldsymbol{L}_{\iota} \cdot (\boldsymbol{L} \times \boldsymbol{n} - \boldsymbol{n} \times \boldsymbol{L})_{\iota'} \right]^{l\leftrightarrow l'} - \sum_{q} \sqrt{2} T_{q}^{(2)} (\boldsymbol{\epsilon}^{+}, \boldsymbol{\epsilon}^{-}, \hat{\mathbf{k}})_{1} \\ &\sum_{\iota,\iota'} \left[ \boldsymbol{L}_{\iota} \otimes (\boldsymbol{L} \times \boldsymbol{n} - \boldsymbol{n} \times \boldsymbol{L})_{\iota'} \right]_{q}^{(2)l\leftrightarrow l'} \\ \end{bmatrix} |\psi_{0}\rangle \,, \end{split}$$

with d(l, l + 1) = 1/(l + 1) and d(l, l - 1) = -1/l. Two irreducible tensors are thus associated to ONCD: a two-particle orbital pseudoscalar, which identifies chirality (see below), and a orbital pseudodeviator; the latter generalises the results of Eq. (4) to the two-particle case and will not be dissussed any further. (There is an important difference between E1-E2 and E1-M1 integrated dichroic spectra. E1-E2 implies excitations from inner shells, which are filled in the ground state and can therefore be 'integrated out'. As a result, one-electron properties of the valence states are probed in the x-ray region. This is not the case in the optical range; E1-M1 involves intra-shell valence excitations and two-electron quantities are measured.)

Interpretation of the results. The choice of L and  $\Omega^$ as the building blocks of our analysis is physically motivated as follows. As both generators are magnetic, they are expected to contribute to the electron orbital current, a conjecture which is readily verified. To this purpose, consider Trammell's expansion of the Fourier transform of the atomic orbital magnetisation density [7,19]

$$\mathcal{L}(\mathbf{k}) = \frac{1}{2} \langle \psi_0 | \sum_{\iota} \left[ \mathbf{L}_{\iota} f(\mathbf{k} \cdot \mathbf{r}_{\iota}) + f(\mathbf{k} \cdot \mathbf{r}_{\iota}) \mathbf{L}_{\iota} \right] | \psi_0 \rangle, \quad (10)$$

and perform a recoupling with use of [20]:

$$f(\mathbf{k} \cdot \mathbf{r}) = 4\pi \sum_{lm} i^l g_l(kr) Y_{lm}(\boldsymbol{n}) Y_{lm}^*(\hat{\mathbf{k}}), \qquad (11)$$

with  $g_l(x) = (2/x^2) \int_0^x d\xi \xi j_l(\xi)$ , where  $j_l(\xi)$  denotes a spherical Bessel function.

We have

$$\mathcal{L}(\mathbf{k}) = \sum_{\iota} \langle \psi_0 | \{ g_0(kr_{\iota}) \, \boldsymbol{L}_{\iota} + i \, 3g_1(kr_{\iota})/2 \\ (\hat{\mathbf{k}} \times (\boldsymbol{L}_{\iota} \times \boldsymbol{n}_{\iota} - \boldsymbol{n}_{\iota} \times \boldsymbol{L}_{\iota}) / 2 + \cdots \} | \psi_0 \rangle , \qquad (12)$$

displaying the orbital angular momentum and anapole contributions, and thus providing a physical basis to our chosen so(3,1) SGA, which is realised by L and  $\Omega^-$ .

The representations of the homogeneous Lorentz group, which are identified by a pair of indexes,  $\nu$  and  $\rho$ , afford a classification of electronic states in noncentrosymmetric systems. Unitary (unirreps) and nonunitary (nonunirreps) irreducible representations will be considered.

It is readily seen [21] that  $\Omega^-$  corresponds to having  $\nu = \rho = 0$ , yielding the unirrep  $D(\nu = 0, \rho = 0) = \sum_{l=0,1,\dots}^{\infty} \oplus D^l$  ("supplementary" series), with basis  $\sum_{l=0,1,\dots}^{\infty} \oplus |lm\rangle$ , thus confirming that we are dealing with a spinless case. (Here,  $D^l$  identifies the representations of the rotation group.) Notice that the spherical harmonics are eigenstates not only of the so(3) invariant (Casimir)  $L^2$ , but also of the corresponding so(3, 1)invariants  $L^2 - (\Omega^-)^2$  and  $L \cdot \Omega^-$ . The latter, with eigenvalues  $i\nu\rho$ , identifies the chiral operator and evinces the deep interweaving of the Lorentz group with chirality. As  $L \cdot \Omega^- = 0$  [Eq. (1)], orbital chirality cannot manifest itself at the one-particle level.

We can build a two-particle basis using the same so(3,1) SGA. The generators are given by  $\mathbf{L}_T = \mathbf{L}_1 + \mathbf{L}_2$ and  $\mathbf{\Omega}_T^- = \mathbf{\Omega}_1^- + \mathbf{\Omega}_2^-$ , in this case. In such a basis, the twoparticle states are eigenstates of the so(3,1) invariants; in particular of  $\mathbf{L}_T \cdot \mathbf{\Omega}_T^-$ , the chiral operator that emerges from the E1-M1 processes [Eq. (9)]. The Clebsch-Gordan series for the direct product of two D(0,0) representation takes the form [22]:  $D(0,0) \otimes D(0,0) =$  $\sum_{\nu=0}^{\infty} \int_{-\infty}^{\infty} D(\nu, \rho) d\mu(\nu, \rho)$ . The direct product thus decomposes into a direct sum of representations some of which characterised by  $\nu \rho \neq 0$ , i.e. orbital chirality appears naturally when considering two-particle properties.

From the relations between the so(3, 1) Clebsch-Gordan coefficients [23], it is readily seen that states with  $\nu \rho \neq 0$  have mixed parity. For a given value of  $\nu |\rho| \neq 0$ , there are two inequivalent representations with basis states of the form  $|\text{even}\rangle + i|\text{odd}\rangle$  and  $|\text{even}\rangle - i|\text{odd}\rangle$ , i.e. with opposite chiralities. Furthermore, for a given pair  $(l_1, l_2)$ , only chiral stes are allowed to have components with angular momenta  $(l_1 \pm 1, l_2)$  and  $(l_1, l_2 \pm 1)$ (two-particle hybridisation).

We can switch to nonunirreps by introducing the transformation:  $\mathbf{N} = (\mathbf{L} + i\mathbf{\Omega}^{-})/2$ , and  $\mathbf{N}^{\dagger} = (\mathbf{L} - \mathbf{U})/2$  $i\Omega^{-})/2$  [24,25]. **N** is a normal operator;  $N_q$  and  $N_q^{\dagger}$ independently obey the Lie algebra of SU(2). Therefore  $SO(3,1) \cong SU(2) \times SU(2)$ . The invariants are given by  $N^2$  and  $(N^{\dagger})^2$ , with eigenvalues j(j+1) and j'(j'+1), respectively; j, j' take integer or half-integers values. These nonunirreps are denoted by  $D^{(j,j')}$ . The representation space is spanned by the (2j + 1)(2j' + 1) basis vectors  $|jm\rangle|j'm'\rangle$  (canonical basis). [All nonunirreps for which j + j' is an integer are true representations of SO(3, 1), while those for which j + j' is half-integer are doublevalued.] Observe that N and  $N^{\dagger}$  are not independent as they can be interchanged by space inversion. This is equivalent to swapping  $j \leftrightarrow j'$ . Notice that  $L \cdot \Omega^- = 0$ implies j = j'. Again, orbital chirality does not appear at the one-particle level.

To illustrate the use of nonunirreps in constructing two-electron states, we resort to the simple case  $D^{(\frac{1}{2},\frac{1}{2})} \otimes D^{(\frac{1}{2},\frac{1}{2})}$ . We have

$$\Psi_{MM'}^{LL'}(1,2) = \sum_{\text{all}\sigma} C_{\frac{1}{2}\sigma_1;\frac{1}{2}\sigma_2}^{LM} C_{\frac{1}{2}\sigma'_1;\frac{1}{2}\sigma'_2}^{LM} \psi_{\sigma_1,\sigma'_1}^{\frac{1}{2}}(1) \psi_{\sigma_2,\sigma'_2}^{\frac{1}{2}}(2) \,,$$

with  $\psi_{\sigma,\sigma'}^{\frac{1}{2}} \equiv |\frac{1}{2}\sigma\rangle|\frac{1}{2}\sigma'\rangle$ . In this basis, the twoparticle chiral operator takes the form:  $L_T \cdot \Omega_T^- =$   $2i \left[ \mathbf{N}(1) \cdot \mathbf{N}(2) - \mathbf{N}^{\dagger}(1) \cdot \mathbf{N}^{\dagger}(2) \right]$ . Given  $\Psi_{MM'}^{LL'}(1,2)$ , we can set:  $(sp)P = \Psi^{01}$ ,  $(ps)P = \Psi^{10}$  and similarly for  $(s^2)S$ ,  $(p^2)S$ ,  $(p^2)P$ ,  $(p^2)D$ . It is straightforward to verify that (sp)P and (ps)P are eigenstates of  $\mathbf{L}_T \cdot \mathbf{\Omega}_T^$ with opposite eigenvalues, and that the remaining states are eigenstates of the two-particle chiral operator with zero eigenvalue. A link between orbital chirality and hybridisation of the valence electrons is thus established.

The linear combination  $\Psi_R = \Psi^{01} \oplus \Psi^{10}$  has definite parity and  $\langle \Psi_R | \mathbf{L}_T \cdot \mathbf{\Omega}_T^- | \Psi_R \rangle = 0$ , furnishing a simple example of a *racemic mixture*. (Notice the similarity between our two-particle racemic mixture and a Dirac electron, which is made out of two Weyl spinors:  $D^{(\frac{1}{2},0)} \oplus D^{(0,\frac{1}{2})}$  [26].)

To summarise: The work has discussed a theoretical approach to photo-absorption spectroscopies in noncentrosymmetric systems.

A symmetry analysis based on homogeneous Lorentz group has indicated that, in the presence of inversionsymmetry-breaking hybridisation, valence electrons develop an orbital anapole moment. The coupling of such a moment to the orbital angular momentum yields a set of microscopic effective operators, which identify the electronic properties probed by x-ray and optical dichroism (E1-E2 and E1-M1 interferences).

It has also been shown that orbital chirality (enantiomophism) is described by a two-electron so(3, 1) invariant (pseudoscalar), which is measured by optical natural circular dichroism.

Stimulating discussions with E. Katz are gratefully acknowledged.

- \* Address from September 1, 2001: European Synchrotron Radiation Facility, B.P. 220, F-38043 Grenoble Cédex, France
- \*\* Permanent address: INFM and Dipartimento di Fisica dell'Università degli Studi di Modena e Reggio Emilia, 41100 Modena, Italy.
- [1] Goulon J. et al., J. Chem. Phys. 108, 6394 (1998).
- [2] Alagna L. et al., Phys. Rev. Lett. 80, 4799 (1998).
- [3] J. Goulon et al. Phys. Rev. Lett., 85, 4385 (2000).
- [4] S. Goshen and H. J. Lipkin, Ann. Phys. (N. Y.) 6, 301 (1959); and in Spectroscopic and Group Theoretical Methods in Physics - Racah Memorial Volume, F. Bloch et al. Eds., North-Holland, Amsterdam, 1968.
- [5] Y. Dothan, M. Gell-Mann, and Y. Ne'eman, Phys. Lett. 17, 148 (1965); N. Mukunda, L. O'Raifeartaigh and E.C.G. Sudarshan, Phys. Rev. Lett. 15, 1041 (1965).
- [6] Y. Dothan and Y. Ne'eman, Band Spectra Generated By Non-Compact Algebra, reprinted in F.J. Dyson, Symmetry Groups in Nuclear and Particle Physics - A Lecture-Note and Reprint Volume, Bejamin, New York, 1966.

- [7] G. T. Trammell, Phys. Rev. 92, 1387 (1953).
- [8] P. Carra and R. Benoist, Phys. Rev. B 62, R7703 (2000).
- [9] M. J. Englefield, Group Theory and the Coulomb Problem, Wiley-Interscience, New York, 1972.
- [10] Ya. B. Zel'dovich, Zh. Éksp. Teor. Fiz. 33, 1531 (1957)
   [Sov. Phys. JETP 6, 1184 (1958)].
- [11] V. V. Flambaum and I. B. Khriplovich, Zh. Éksp. Teor. Fiz. **79**, 1656 (1980) [Sov. Phys. JETP **52**, 835 (1980)].
- [12] M.-A. Bouchiat and C. Bouchiat, Rep. Prog. Phys. 60, 1351 (1997).
- [13] P. Carra and B. T. Thole, Rev. Mod. Phys. 66, 1509 (1994).
- [14] C. R. Natoli et al., Eur. Phys. J. B 4, 1 (1998).
- [15] Relativistic corrections to the radial part of the atomic wave functions are neglected.
- [16] In Eq. (4) and (7), the coefficients are given by

 $a_{l'}(l_c, l) = b_{l'}(l_c, l)[6 + 3l_c(l_c + 1) - 2l(l + 1) - l'(l' + 1)],$ with

$$b_{l'}(l_c, l) = \frac{2(2l+1)(2l'+1)}{h(l, l')(l_c - 3l' + 2l)(l_c + 3l' - 2l + 1)\bar{h}(l_c, l)}$$

h(l, l') = l + l' + 1 and  $\bar{h}(l_c, l) = (l_c + l)^2 (l_c + l + 2)^2$ .

- [17] C. S. Wood *et al.*, Science **275**, 1759 (1997).
- [18] Equation (8) corrects a misprint in Eq. (16) of Ref. [8].
- [19] M. Blume and D. Gibbs, Phys. Rev. B 37, 1779 (1988).
- [20] S. W. Lovesey, J. Phys. C 2, 470 (1969).
- [21] A. Bohm, Quantum Mechanics: Foundations and Applications, Springer-Verlag, Heidelberg, 1993; Ch. V.
- M.A. Naimark, Tr. Mosk. Mat. Obshch. 8, 121 (1959);
   9 237 (1960); 10, 181 (1961) [Amer. Math. Soc. Transl., Ser. 2, 36, 101-229 (1964)].
- [23] R.L. Anderson, R. Raczka, M.A. Rashid, and P. Winternitz, J. Math. Phys. **11**, 1050 (1970); **11**, 1059 (1970), and references therein.
- [24] Y. Frishman and C. Itzykson. Phys. Rev. 180, 1556 (1969).
- [25] L. Fonda and G.C. Ghirardi, Symmetry Principles in Quantum Physics, Marcel Dekker, New York, 1970; Ch. III.
- [26] P. Ramond, Field Theory: A Moderm Primer, Addison-Wesley, Redwood City, California, 1989.

TABLE I. Polarisation responses of E1-E2 and E1-M1 interferences.

$$\begin{split} iT_{0}^{(0)}(\boldsymbol{\epsilon}^{*},\boldsymbol{\epsilon},\hat{\mathbf{k}})_{1} &= -iT_{0}^{(0)}(\boldsymbol{\epsilon},\boldsymbol{\epsilon}^{*},\hat{\mathbf{k}})_{1} = -i/\sqrt{6}, \\ iT_{q}^{(2)}(\boldsymbol{\epsilon}^{*},\boldsymbol{\epsilon},\hat{\mathbf{k}})_{1} &= -iT_{q}^{(2)}(\boldsymbol{\epsilon},\boldsymbol{\epsilon}^{*},\hat{\mathbf{k}})_{1} = -i/2[[\boldsymbol{\epsilon}^{*}\otimes\boldsymbol{\epsilon}]^{(1)}\otimes\hat{\mathbf{k}}]_{q}^{(2)}, \\ T_{q}^{(1)}(\boldsymbol{\epsilon}^{\parallel},\hat{\mathbf{k}})_{2} &= T_{q}^{(1)}(\boldsymbol{\epsilon}^{\perp},\hat{\mathbf{k}})_{2} = -1/2\sqrt{\frac{3}{5}}\hat{\mathbf{k}}_{q}, \\ T_{q}^{(2)}(\boldsymbol{\epsilon}^{*},\boldsymbol{\epsilon},\hat{\mathbf{k}})_{2} &= -T_{q}^{(2)}(\boldsymbol{\epsilon},\boldsymbol{\epsilon}^{*},\hat{\mathbf{k}})_{2} = \frac{\sqrt{3}}{2}[[\boldsymbol{\epsilon}^{*}\otimes\boldsymbol{\epsilon}]^{(1)}\otimes\hat{\mathbf{k}}]_{q}^{(2)}, \\ T_{q}^{(3)}(\boldsymbol{\epsilon}^{\parallel},\hat{\mathbf{k}})_{2} &= [[\boldsymbol{\epsilon}^{\parallel}\otimes\boldsymbol{\epsilon}^{\parallel}]^{(2)}\otimes\hat{\mathbf{k}}]_{q}^{(3)} \neq T_{q}^{(3)}(\boldsymbol{\epsilon}^{\perp},\hat{\mathbf{k}})_{2} \\ &= [[\boldsymbol{\epsilon}^{\perp}\otimes\boldsymbol{\epsilon}^{\perp}]^{(2)}\otimes\hat{\mathbf{k}}]_{q}^{(3)} \end{split}$$